


North Carolina Shrimp Fishery Management Plan – Draft Amendment 1

North Carolina Division of Marine Fisheries

SEDAR-PW6-RD26

7 May 2014



DRAFT

North Carolina
Shrimp
Fishery Management Plan

Draft Amendment 1

By

North Carolina Division of Marine Fisheries



North Carolina Department of Environment and Natural Resources
North Carolina Division of Marine Fisheries
3441 Arendell Street
P. O. Box 769
Morehead City, N.C. 28557

February 2014

Sept 2011	Timeline begins
July 2012	Internal review with DMF comments
Aug 2012	Draft Revision 1 approved by MFC for public comment
Nov 2012	Management changes proposed begin Draft Amendment 1
Nov 2013	Draft Amendment 1 approved by MFC for public comment
Feb 2014	MFC selects preferred management options
Mar 2014	Submit draft to DENR Secretary

DRAFT

1.0 ACKNOWLEDGEMENTS

The Shrimp FMP Amendment 1 was prepared by the North Carolina Department of Environment and Natural Resources Division of Marine Fisheries (DMF) under direction of the North Carolina Marine Fisheries Commission (MFC) to address bycatch issues in the commercial and recreational shrimp fisheries with the advice of the Shrimp Advisory Committee.

DMF Plan Development Team

Trish Murphey Co-Lead
Chris Stewart Co-Lead
Alan Bianchi
Kevin Brown
John Hadley
Kevin Hart
Jack Holland
Dean Nelson
Greg Judy
Blake Price (NMFS)
David Taylor
Holly White
Katy West

Marine Fisheries Commission

Paul C. Rose, Chairman
Anna Barrios Beckwith, Vice-Chairman
Mikey Daniels
Kelly L. Darden, Jr.
Mark L. Gorges
Charles Laughridge III
Allyn Powell
Joe Shute
Joseph J. Smith Jr.

Shrimp FMP Advisory Committee

Joe Albea
Julian Anderson
Frank Brown
Ray Brown
Nancy Edens
P.D. Mason
Steve Parrish
Kenny Rustick
Scott Whitley

DRAFT

2.0 TABLE OF CONTENTS

1.0	ACKNOWLEDGEMENTS	2
2.0	TABLE OF CONTENTS	3
2.1	LIST OF TABLES	8
2.2	LIST OF FIGURES	12
2.3	LIST OF ACRONYMS	16
3.0	EXECUTIVE SUMMARY	20
3.1	STATUS OF THE FISHERIES	21
3.1.1	Commercial Fishery.....	21
3.1.2	Recreational Fishery.....	22
3.1.3	Shrimp Trawl Bycatch.....	22
3.2	PROTECTED RESOURCES.....	23
3.3	ECONOMIC STATUS.....	23
3.4	SOCIOECONOMIC CHARACTERISTICS.....	23
3.5	ENVIRONMENTAL FACTORS	24
3.6	MANAGEMENT STRATEGIES	24
3.6.1	Shrimp Management by Size.....	25
3.6.2	Shrimp Management by Area	25
3.6.3	Shrimp Management in the Southern District.....	25
3.6.4	Shrimp Management in the Central District	25
3.6.5	Shrimp Management in Pamlico District	25
3.6.6	Shrimp Management in the Northern District	26
3.6.7	Atlantic Ocean	26
3.7	BYCATCH IN THE SHRIMP FISHERY AND MANAGEMENT OPTIONS	26
3.7.1	Marine Fisheries Commission Preferred Management Strategies and Required Actions	27
4.0	INTRODUCTION.....	29
4.1	LEGAL AUTHORITY FOR MANAGEMENT.....	29
4.2	RECOMMENDED MANAGEMENT PROGRAM	30
4.2.1	Goals and Objectives	30
4.2.2	Sustainable Harvest	31
4.2.3	Management Strategy	31
4.3	DEFINITION OF MANAGEMENT UNIT.....	31
4.4	GENERAL PROBLEM STATEMENT	31
4.4.1	New River Trawl Fishery.....	31
4.4.2	Use of Other Fishing Gears	31
4.4.3	Gear Modifications.....	32
4.4.4	Live Bait Shrimp Fishery.....	32
4.4.5	Effort Management.....	32
4.4.6	Area Restrictions	32
4.5	EXISTING PLANS STATUTES, AND RULES	32
4.5.1	Plans	32
4.5.2	Statutes.....	34
4.5.3	Marine Fisheries Commission Rules	34
4.5.3.1	General.....	34
4.5.3.2	Nets, Pots, Dredges, and Other Fishing Devices	35
4.5.3.3	Oysters, Clams, Scallops and Mussels	38
4.5.3.4	Shrimp, Crabs, and Lobster	38
4.5.3.5	Licenses, Leases, Franchises, and Permits	39
4.5.5	Federal Regulations	40

DRAFT

5.0	STATUS OF THE STOCK.....	41
5.1	GENERAL LIFE HISTORY	41
5.1.1	Brown Shrimp.....	41
5.1.2	Pink Shrimp.....	42
5.1.3	White Shrimp.....	42
5.1.4	Movement	43
5.1.5	Predation.....	44
5.1.6	Parasites and Disease.....	45
5.2	STOCK STATUS.....	46
6.0	STATUS OF FISHERIES	47
6.1	COMMERCIAL	47
6.1.1	History.....	47
6.1.2	State Landings and Effort	49
6.1.3	Landings by Waterbody.....	52
6.1.4	Landings by Gear	53
6.1.5	Landings by Species	54
6.1.2.5	Regional Summary	58
6.2	RECREATIONAL FISHERY	58
6.2.1	Recreational Commercial Gear License (RCGL) History	59
6.2.2	RCGL Survey	59
6.2.3	RCGL Survey Methodology	60
6.2.4	RCGL Survey Results	61
6.2.5	Regional RCGL Characterization for Shrimp Trawls.....	63
6.2.6	Contribution of RCGL Harvest compared to Other Fisheries	65
6.3	SHRIMP TRAWL BYCATCH.....	67
6.3.1	History of Bycatch Management in North Carolina	68
6.3.2	Incidental Catch.....	69
6.3.3	Discarded Catch.....	76
6.3.4	Biological Implications of Bycatch.....	81
6.3.5	Bycatch Impacts on Stock Assessment and Prediction.....	83
6.3.6	Bycatch Impacts on the Stock Assessment of Non-Target Species	85
6.3.6.1	Blue Crab Status	86
6.3.6.2	Weakfish Status.....	87
6.3.6.3	Atlantic Croaker Status	88
6.3.6.4	Spot Status.....	90
6.3.6.5	Southern Flounder Status.....	91
6.3.6.6	Summary	92
6.3.7	North Carolina Management Strategies to Reduce Bycatch	92
6.3.7.1	Tailbag Mesh Size	92
6.3.7.2	Bycatch Reduction Devices	93
6.3.7.3	Turtle Excluder Devices.....	98
6.3.7.4	Alternate Gears	99
6.3.7.5	Catch Restrictions.....	100
6.3.7.6	Harvest Seasons	101
6.3.7.7	Time Restrictions	101
6.3.7.8	Area Restrictions	101
6.3.7.9	Limited Entry.....	102
7.0	PROTECTED RESOURCES.....	103
7.1	PROTECTED RESOURCES LEGISLATION.....	103
7.1.1	Federal Endangered Species Act (ESA).....	103
7.1.2	Marine Mammal Protection Act (MMPA).....	104

DRAFT

7.1.3 North Carolina Endangered Species Act (Chapter 113 Article 25) 105

7.2 SPECIES THAT MAY INTERACT WITH THE SHRIMP FISHERY 106

7.2.3 Bottlenose Dolphin 106

7.2.4 Sea Turtles 107

7.2.4.1 Sea Turtles and the Shrimp Fishery 109

7.2.4.2 NCDMF Programs 110

8.0 ECONOMIC STATUS 113

8.1 COMMERCIAL FISHERY 113

8.1.1 Harvesting sector 113

8.1.1.1 Ex-vessel value and price 113

8.1.1.2 Gear 118

8.1.1.3 Water bodies 120

8.1.1.4 Participants and trips 127

8.1.1.5 Processing 132

8.1.1.6 Marketing and distribution 132

8.1.1.7 Economic impact of commercial fishery 133

8.1.2 Recreational fishery economics 134

8.1.2.1 Marine Recreational Information Program (MRIP) 134

8.1.2.2 Recreational use of commercial gear (RCGL) 134

8.1.2.3 Other Recreational Fisheries 135

9.0 SOCIOECONOMIC CHARACTERISTICS 136

9.1 SOCIAL IMPORTANCE OF THE FISHERY 136

9.1.1 Commercial fishermen 136

9.1.1.1 Historical importance 136

9.1.1.2 Community reliance on the commercial fishery 136

9.1.1.3 Perceived conflicts 138

9.1.1.4 Perception of important issues 139

9.2 RECREATIONAL FISHERY 140

9.2.1 Historical importance 140

9.2.2 Community reliance on the recreational fishery 140

9.2.3 Perceived conflicts 140

9.2.4 Perception of important issues 140

9.3 DEMOGRAPHIC CHARACTERISTICS 141

9.3.1 Commercial fishermen 141

9.3.2 Recreational fishermen 142

9.4 DEFINITIONS 143

10.0 ENVIRONMENTAL FACTORS 145

10.1 HABITAT 145

10.1.1 Water column 145

10.1.2 Wetlands 146

10.1.3 Soft bottom 147

10.1.4 Submerged Aquatic Vegetation (SAV) 153

10.1.5 Shell bottom 154

10.1.6 Hard Bottom 156

10.2 HABITAT CONCERNS 156

10.2.1 Wetland Loss 156

10.2.2 Soft Bottom Impacts 157

10.2.3 Submerged Aquatic Vegetation Impacts 159

10.3 WATER QUALITY 162

10.3.1 Hydrological modifications 162

10.3.2 Low oxygen 163

DRAFT

10.3.3 Toxins..... 165

10.3.4 Tiger Shrimp..... 166

10.4 HABITAT AND WATER QUALITY PROTECTION..... 167

10.4.1 MFC Authority 167

10.4.2 Authority of Other Agencies..... 167

10.4.3 Nursery Area Protection 168

10.4.4 Coastal Habitat Protection Plan..... 170

10.5 STATUS OF 2006 SHRIMP FMP ENVIRONMENTAL FACTORS RECOMMENDATION.....171

10.5.1 Coastal Habitat Protection Plan Actions..... 172

11.0 MANAGEMENT STRATEGIES..... 175

11.1 HISTORY OF SHRIMP MANAGEMENT 175

11.2 SHRIMP MANAGEMENT BY SIZE 176

11.3 SHRIMP MANAGEMENT BY AREA..... 177

11.3.1 Shrimp Management in the Southern District..... 178

11.3.1.1 Brunswick County..... 178

11.3.1.2 Cape Fear River Complex 180

11.3.1.3 Intracoastal Waterway and Sounds from Carolina Beach to Rich’s Inlet 182

11.3.1.4 Intracoastal Waterway and sounds from Rich’s Inlet to New River..... 184

11.3.1.5 New River 184

11.3.1.6 Chadwick Bay..... 186

11.3.2 Shrimp Management in the Central District 187

11.3.2.1 White Oak River 187

11.3.2.3 Newport River..... 189

11.3.2.4 North River 191

11.3.2.5 Jarrett Bay 191

11.3.2.6 Core Sound 191

11.3.3 Shrimp Management in Pamlico District..... 193

11.3.3.1 Neuse River..... 194

11.3.3.2 South Side of Neuse River..... 196

11.3.3.3 Bay River..... 196

11.3.3.4 Pamlico River 196

11.3.3.5 Pungo River..... 198

11.3.3.6 Pamlico Sound 198

11.3.3 Shrimp Management in the Northern District 198

11.3.3.1 Croatan Sound 200

11.3.3.2 Roanoke Sound..... 201

11.3.4 Atlantic Ocean..... 201

11.4 SUMMARY OF MANAGEMENT ACTIONS..... 201

11.4.1 Rules (new, modifications, or technical changes) 201

11.4.2 Legislative Action (new, modifications, or technical changes)..... 201

11.4.3 Actions by Other Agencies 201

11.4.4 Management Related Research Needs..... 203

11.4.5 Biological Research Needs..... 203

11.4.6 Social and Economic Research Needs..... 204

11.4.7 Data Needs 204

12.0 BYCATCH IN THE SHRIMP FISHERY AND MANAGEMENT OPTIONS..... 205

12.1 TRAWLING IN THE NEW RIVER ABOVE THE HIGHWAY 172 BRIDGE 206

12.2 EVALUATION OF THE SKIMMER TRAWL AND OTHER GEARS USED FOR SHRIMPING IN NORTH CAROLINA..... 221

12.3 RESEARCH RESULTS – THE USE OF TURTLE EXCLUDER DEVICES (TEDS) IN COMMERCIAL SKIMMER TRAWL OPERATIONS 247

DRAFT

12.4 CONSIDERATION FOR A COMMERCIAL LIVE BAIT SHRIMP FISHERY IN NORTH CAROLINA.....270

12.5 GEAR MODIFICATIONS IN NORTH CAROLINA SHRIMP TRAWLS TO REDUCE FINFISH BYCATCH276

12.6 EFFORT MANAGEMENT FOR BYCATCH REDUCTION IN THE NORTH CAROLINA SHRIMP TRAWL FISHERY.....296

12.7 CHARACTERIZATION OF THE NORTH CAROLINA COMMERCIAL SHRIMP TRAWL FLEET309

12.8 AREA RESTRICTIONS TO REDUCE SHRIMP TRAWL BYCATCH IN NORTH CAROLINA’S INTERNAL COASTAL WATERS.....334

12.9 REMOVAL OF THE SHRIMP TRAWL FROM THE RECREATIONAL COMMERCIAL GEAR LICENSE.....348

12.10 BYCATCH MANAGMENT RECOMMENDATIONS.....354

 12.10.1 Trawling in the New River above the Highway 172 Bridge354

 12.10.2 Evaluation of the skimmer trawl and other gears used for shrimping in North Carolina.....354

 12.10.3 The use of TEDs in commercial skimmer trawl operations.....354

 12.10.4 Consideration of a commercial live bait shrimp fishery in North Carolina355

 12.10.5 Gear Modifications in North Carolina shrimp trawls to reduce finfish bycatch355

 12.10.6 Effort Management for bycatch reduction in the North Carolina shrimp trawl fishery.....357

 12.10.7 Characterization of the North Carolina commercial shrimp trawl fleet357

 12.10.8 Area restrictions to reduce shrimp trawl bycatch in North Carolina’s internal coastal waters358

12.11 PROPOSED RULES358

 12.11.1 Trawling in the New River above the Highway 172 Bridge358

 12.11.2 Evaluation of the Skimmer Trawl and other gears used for shrimping in North Carolina.....358

 12.11.3 Gear Modifications in the North Carolina Shrimp Trawl to Reduce Finfish Bycatch359

 12.11.4 Characterization of the North Carolina Commercial Shrimp Trawl Fleet..... 364

 12.11.5 Area Restrictions to Reduce Shrimp Trawl Bycatch in North Carolina’s Internal Coastal Waters.....365

 12.11.6 Additional Rule Changes to Address Clarity and Consistency.....366

13.0 LITERATURE CITED369

14.0 APPENDICES.....389

14.1 APPENDIX1- AMENDMENT 1- PUBLIC COMMENT389

14.2 APPENDIX 2 – REVISION REVIEW PROCESS407

14.3 APPENDIX 3 – BYCATCH REDUCTION DEVICES AND TURTLE EXCLUDER DEVICES474

DRAFT

2.1 LIST OF TABLES

Table 5.1	Environmental Requirements of three shrimp species found in North Carolina. .43
Table 5.2	Migration studies in North Carolina of three shrimp species found in North Carolina.44
Table 6.1	Annual number of trips reported for shrimp in inside and ocean waters [†] , 1978-2010 (Detailed Shrimp Program and Trip Ticket Program).51
Table 6.2	Percent contribution of landings (1999 – 2010) by waterbody and species.53
Table 6.3	Shrimp landings in pounds from the South Atlantic, 1999-2010.56
Table 6.4	Number of fiscal license sales of Recreational Commercial Gear Licenses, 2002 through 2010 (fiscal year, July 1 through June 30).61
Table 6.5	Number of trips by shrimp trawl by region, 2002 through 2008.63
Table 6.6	Harvest (lb) and pounds per trip of shrimp by RCGL gear from 2002 through 2008.63
Table 6.7	Top four species harvested (lb) by RCGL shrimp trawls in the Southern Region, 2002-2008.63
Table 6.8	Top species harvested (lb) by RCGL shrimp trawls in the Central Region, 2002-2008.64
Table 6.9	Top species harvested (lb) by RCGL shrimp trawls in the Pamlico Region, 2002-2008.64
Table 6.10	Top species harvested (lb) by RCGL shrimp trawl in the Northern Region, 2002-2008.65
Table 6.11	Contribution in percent (pounds) of RCGL harvest to the overall harvest of finfish and shellfish based on the average yearly harvest from each sector during the period 2002 through 2008.66
Table 6.12	Percent shrimp trawl landings (lb) [†] of major market groups for North Carolina, 1994-2010.70
Table 6.13	Percent shrimp trawl landings (lb) [†] of major market groups for North Carolina by waterbody, 1994-2010.71
Table 6.14	Yearly finfish landings (lb) [†] from shrimp trawls all North Carolina Waters combined, 1994-2010.72
Table 6.15	Percent shrimp trawl landings (lb) [†] of top five finfish groups by waterbody, 1994-2010.73
Table 6.16	Percent shrimp trawl landings (lb) [†] of major market groups landed for North Carolina (all waters combined) by month, 1994-2010.74
Table 6.17	Percent shrimp trawl landings (lb) [†] of top five finfish groups by month, 1994-2010.75
Table 6.18	Author (year published), years sampled, area sampled, number of trips sampled, number of tows sampled, number of species observed , percent bycatch, percent finfish, percent shrimp, finfish to shrimp ratio (F:S), and bycatch to shrimp ratio (BC:S) of previous bycatch characterization work conducted in North Carolina and the South Atlantic.80
Table 8.1.	Detail values of pounds landed, total value, inflation adjusted value, price per pound, and percent change from year to year for shrimp landed in North Carolina, 1972 – 2010 (DMF Trip Ticket Program). 116
Table 8.2.	Gear type, pounds, price per pound, and total value of shrimp landings by gear in all North Carolina waters, 1994 – 2010 (DMF Trip Ticket Program). 119
Table 8.3.	Pounds and value of shrimp landed from North Carolina water bodies from 1994 – 2010 (DMF Trip Ticket Program). 121
Table 8.4.	Number of participants in the shrimp fishery by value of landings and year in North Carolina, 1999 – 2010 (DMF Trip Ticket Program). 127

DRAFT

Table 8.5.	Number of participants and the number of trips taken that landed shrimp in North Carolina, 1999 – 2010 (DMF Trip Ticket Program).....	128
Table 8.6.	Number of seafood dealer and pounds of shrimp purchased by North Carolina fish dealers from North Carolina fishermen, 1994 – 2010 (DMF Trip Ticket Program).....	130
Table 8.7.	Economic impact of the commercial shrimp fishery in North Carolina, 2010.....	134
Table 8.8.	Economic impact of RCGL fishing trips for shrimp in 2007 (DMF RCGL Survey).	135
Table 9.1.	Most frequently cited communities where shrimp fishermen live (DMF Socioeconomics Program).	137
Table 9.2.	Other prevalent species targeted by shrimp fishermen and average percent of fishing income made from non-target species (DMF Socioeconomics Program).	137
Table 9.3.	Fishing related issues considered most important to shrimp fishermen (DMF Socioeconomics Program).	139
Table 9.4.	Demographic characteristics of commercial shrimp fishermen (DMF Socioeconomics Program).	141
Table 9.5.	Demographic characteristics of RCGL holders who targeted shrimp in 2007. (DMF RCGL Survey Program).	142
Table 10.1	Spawning seasons for Penaeid shrimp species in North Carolina (Pattilo et al. 1997).	146
Table 10.2	Fishing gears used in North Carolina identified as potentially damaging to submerged aquatic vegetation habitat. (Source: MSC 1996).....	160
Table 10.3	Reported observations of tiger shrimp in NC since 2008.....	167
Table 12.1	Catch and effort data on shrimp and landed bycatch for channel nets in New River, 1994-2011 (courtesy of DMF trip ticket program).	212
Table 12.2	Catch and effort data on shrimp and landed bycatch for otter trawls in New River, 1994-2011 (courtesy of DMF trip ticket program).	212
Table 12.3	Catch and effort data on shrimp and landed bycatch for skimmer trawls in New River, 1994-2011 (courtesy of DMF trip ticket program).	213
Table 12.4	Catch and effort data on shrimp and landed bycatch for miscellaneous gear (cast nets, gill nets, etc.) in New River, 1994-2011 (courtesy of DMF trip ticket program). *Data confidential due to less than three participants reporting landings.	213
Table 12.5	Catch and effort data on crab and landed bycatch for crab trawls in New River, 1994-2010 (courtesy of DMF trip ticket program). *Data confidential due to less than three participants reporting landings.	214
Table 12.6	Statewide catch and effort data on shrimp and landed bycatch for channel nets, 1994-2011 (courtesy of DMF trip ticket program).	214
Table 12.7	Statewide catch and effort data on shrimp and landed bycatch for otter trawls, 1994-2011 (courtesy of DMF trip ticket program).	215
Table 12.8	Statewide catch and effort data on shrimp and landed bycatch for skimmer trawls, 1994-2011 (courtesy of DMF trip ticket program).	216
Table 12.9	Statewide catch and effort data on shrimp and landed bycatch for miscellaneous gear (cast nets, gill nets, etc.), 1994-2011 (courtesy of DMF trip ticket program).	216
Table 12.10	Statewide catch and effort data on crab and landed bycatch for crab trawls, 1994-2011 (courtesy of DMF trip ticket program).	217
Table 12.11	Number and average pounds, trips, and value in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in state estuarine waters, 1994-2011.	241
Table 12.12	Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Southern District*, 1994-2011.....	242
Table 12.13	Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Central District*, 1994-2011.....	243

DRAFT

Table 12.14 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Pamlico District*, 1994-2011.244

Table 12.15 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Northern District*, 1994-2011.245

Table 12.16 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in Pamlico Sound, 1994-2011.246

Table 12.17 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch (bycatch + shrimp catch), bycatch, and shrimp catch (kg) by vessel for 2008 skimmer trawl TED testing conducted in MS and AL.257

Table 12.18 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, teleost fish, and rays) (kg) by vessel for 2008 skimmer trawl TED testing conducted in MS and AL.257

Table 12.19 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch (bycatch + shrimp catch), bycatch, and shrimp catch (kg) by vessel for 2009 skimmer trawl TED testing conducted in MS and AL.258

Table 12.20 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, teleost fish, and rays) (kg) by vessel for 2009 skimmer trawl TED testing conducted in MS and AL.258

Table 12.21 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch (bycatch + shrimp catch), bycatch, and shrimp catch (kg) by vessel for 2010 skimmer trawl TED testing conducted in NC.259

Table 12.22 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, teleost fish, and rays) (kg) by vessel for 2010 skimmer trawl TED testing conducted in NC.260

Table 12.23 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch, bycatch, and shrimp catch (kg) for each experimental TED type in the 2011 NC skimmer trawl TED testing.261

Table 12.24 Summary statistics, results of paired t tests, power analyses, percent differences (kg) and corresponding 95% confidence intervals for groups (crustaceans, debris, invertebrates, teleost fish) for each experimental TED type in the 2011 NC skimmer trawl TED testing.261

Table 12.25 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch, bycatch, and shrimp catch (kg) for each experimental TED type in the 2012 NC skimmer trawl TED testing.262

Table 12.26 Summary statistics, results of paired t tests, power analyses, percent differences (kg) and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, fish, debris, rays and sharks) for each experimental TED type in the 2012 NC skimmer trawl TED testing.262

Table 12.27 Number of Pounds of Live Bait Shrimp (dozens, numbers) 1994 through 2011.271

Table 12.28 Number and average pounds, trips, and value in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in state estuarine waters, 1994-2011. 297

Table 12.29 Number and pounds, trips, and participants in shrimp trawls, skimmer trawls, channel nets, cast nets in state estuarine waters, 1994-2011.298

DRAFT

Table 12.30	Average pounds of shrimp landed in North Carolina's inside waters per month by species, 2001-2011.....	300
Table 12.31	Average landings and effort by month of all shrimp species by waterbody* from otter and skimmer trawls combined), 2002-2011.....	303
Table 12.32	Maximum commercial, bait, and recreational shrimp trawl configurations by state	322
Table 12.33	North Carolina vessel and shrimp trawl configuration by area and year, 2010-2011.....	323
Table 12.34	Number of pounds, trips, value and participation in the estuarine shrimp trawl fishery.....	336
Table 12.35	Weighted CPUE, standard error (SE), total number collected (n), mean size (mm), and size range for select species during 2011 in the Pamlico Sound Survey (DMF 2012).....	338
Table 12.36	Current and past designated special secondary nursery areas.....	342
Table 12.37	Number of license sales of Recreational Commercial Gear Licenses, 2002 though 2011. (fiscal year, July 1 through June 30).....	349
Table 12.38	Number of trips by shrimp trawl by region, 2002-2008.	350
Table 12.39	Harvest (lb) and pounds per trip of shrimp by RCGL gear from 2002-2008.....	350
Table 12.40	Commercial estuarine shrimp trawl harvest and Recreational Commercial shrimp trawl harvest, 2002-2008.....	350
Table 12.41	Recreational Commercial trawl discard numbers and trips, 2002-2008.	351

DRAFT

2.2 LIST OF FIGURES

Figure 6.1	Annual shrimp landings (lb) for North Carolina: 1962 – 2010.	50
Figure 6.2	Landings (lb) and trips for 1978-2010.....	50
Figure 6.4	North Carolina landings of shrimp by species 1978-1993.....	55
Figure 6.5	North Carolina landings of shrimp by species: 1994 – 2010.	56
Figure 6.6	Regions used to summarize findings from the RCGL surveys.	60
Figure 6.7	Distribution of RCGL residents	62
Figure 6.8	Diagram of Florida Fish Excluder (FFE) tested in North Carolina waters.....	94
Figure 6.9	Diagram of large mesh extended funnel BRD (LMEF) tested in North Carolina.	95
Figure 6.10	Diagram of modified large mesh funnel excluder (LMFE) tested in North Carolina.	96
Figure 6.11.	Diagram of “Sea Eagle” fish excluder tested in North Carolina.....	97
Figure 8.1.	Value of shrimp landings in North Carolina, 1972 – 2010 (DMF Trip Ticket Program).....	113
Figure 8.2.	Average price per pound of shrimp landings in North Carolina, 1972 – 2010 (DMF Trip Ticket Program).	114
Figure 8.3.	Percent of landings by gear used to harvest shrimp in all North Carolina waters, 1994 – 2010 (DMF Trip Ticket Program).....	118
Figure 8.4.	Number of seafood dealers who purchased shrimp from 1994 to 2010 (DMF Trip Ticket Program).	129
Figure 9.1.	Reported conflicts of North Carolina Shrimp Fishermen (DMF Socioeconomics Program).....	139
Figure 10.1a	Shrimp nursery areas, including MFC designated nursery areas and SAV beds, in Pamlico Sound NC.....	149
Figure 10.1b	Shrimp nursery areas, including MFC designated nursery areas and SAV beds, for the Tar/Pamlico and Neuse Rivers.....	150
Figure 10.1c	Shrimp nursery areas, including MFC designated nursery areas and SAV beds, for the Core Banks to Topsail Island.	151
Figure 10.1d	Shrimp nursery areas, including MFC designated nursery areas and SAV beds, for the Topsail Island to South Carolina.	152
Figure 11.1	Map of shrimp management areas in Brunswick County.....	179
Figure 11.2	Map of shrimp management areas in the Cape Fear River Complex.	181
Figure 11.4	Map of shrimp management areas in the Intracoastal Waterway and sounds from Rich’s Inlet to New River.....	185
Figure 11.5	Shrimp management areas in the White Oak River and Bogue Sound.	188
Figure 11.6	Map of the shrimp management areas in the Newport River, North River, Ward Creek, The Straits, and Back Sound.	190
Figure 11.7	Map of shrimp management areas in Core Sound and its surround estuaries..	192

DRAFT

Figure 11.8	Map of the shrimp management areas in The Neuse River and its tributaries.	195
Figure 11.9	Map of shrimp management areas in the Pamlico and Pungo Rivers.....	197
Figure 11.10	Map of shrimp management areas in the Pamlico Sound and its tributaries...	199
Figure 11.11	Map of shrimp management areas in the Croatan and Roanoke Sounds.	200
Figure 12.1	Map of the New River showing the areas for the PNAs and SSNA as well as the otter trawl closure line.....	218
Figure 12.2	Shrimp trips by gear in New River, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.	219
Figure 12.3	Mean catch of shrimp in New River, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.	219
Figure 12.4	Total catch of shrimp in pounds by gear in the New River, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.....	220
Figure 12.5	Number of trips and participants in the New River crab trawl fishery, 1994-2010.	220
Figure 12.6	Pounds of crab landed and number of pounds of crab landed per trip in the New River crab trawl fishery, 1994-2010.....	221
Figure 12.7	NCDMF Fishery Management Districts	233
Figure 12.8	Skimmer Trawl Locations in Pender County.....	234
Figure 12.9	Skimmer Trawl Locations in Onslow County.....	235
Figure 12.10	Skimmer Trawl Locations in New River.	236
Figure 12.11	Skimmer Trawl Locations in Bogue Sound, White Oak River and Queens Creek.	237
Figure 12.12	Skimmer Trawl Locations in Newport River, North River, Back Sound and Lower Core Sound.....	238
Figure 12.13	Skimmer Trawl Locations in Neuse River and Upper Core Sound.....	239
Figure 12.14	Skimmer Trawl Locations in Pamlico Sound.....	240
Figure 12.15	Diagram of typical skimmer trawl operation (Source: Hein and Meier 1995).	263
Figure 12.16	Skimmer trawl TED testing locations during 2008 in Mississippi and Alabama.	263
Figure 12.17	Skimmer trawl TED testing locations during 2009 in Mississippi and Alabama.	264
Figure 12.18	Skimmer trawl TED testing locations in 2010 in North Carolina.	264
Figure 12.19	Skimmer trawl track lines in the 2011 NC skimmer trawl TED testing.	265
Figure 12.20	Trawl locations for Vessel 1 (TODC TED configuration) in the 2012 NC Skimmer Trawl TED testing.	266
Figure 12.21	Trawl locations for Vessel 2 (BODC TED configuration) in the 2012 NC Skimmer Trawl TED testing.	267
Figure 12.22	Percent difference for total catch, bycatch, and shrimp (kg) for each experimental TED type (TODC and BODC) in the 2012 NC skimmer trawl TED testing.....	268

DRAFT

Figure 12.23 Percent difference for species groups by each experimental TED type (TODC and BODC) in the 2012 NC skimmer trawl TED testing.....268

Figure 12.24 Percent difference for species groups by each experimental TED type (TOSC, BOSC, and TODC) in the 2011 LA skimmer trawl TED testing.....269

Figure 12.25 Diagram of typical otter trawl (NOAA 2008).279

Figure 12.26 Topless trawl (Northeast Fisheries Science Center)280

Figure 12.27 Standard trawl (top) and T90 trawl (bottom) (Knuckey et al. 2008).286

Figure 12.28 T90 extension constructed by Mikey Daniels.....287

Figure 12.29 Schematic of an otter trawl and its components (from NMFS 2012).324

Figure 12.30 Schematic of a skimmer trawl and its components (from NMFS 2012).....324

Figure 12.31 Length frequency distribution of vessels using otter trawls in the Pamlico Sound, 2010-11.325

Figure 12.32 Length frequency distribution of total headrope length (ft) of otter trawls in the Pamlico Sound, 2010-11.....325

Figure 12.34 Length frequency distribution of total headrope length (ft) of skimmer trawls in the Pamlico Sound, 2010-11.....326

Figure 12.35 Length frequency distribution of vessels (ft) using otter trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.327

Figure 12.36 Length frequency distribution of total headrope length (ft) of otter trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.....327

Figure 12.37 Length frequency distribution of vessels (ft) using skimmer trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.328

Figure 12.38 Length frequency distribution of total headrope length (ft) of skimmer trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.328

Figure 12.39 Length frequency distribution of vessels (ft) using otter trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11.329

Figure 12.40 Length frequency distribution of total headrope length (ft) of otter trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11. ...329

Figure 12.41 Length frequency distribution of vessels (ft) using skimmer trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11.330

Figure 12.42 Length frequency distribution of total headrope length (ft) of skimmer trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11. ...330

Figure 12.43 Length frequency distribution of vessels (ft) using otter trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11. 331

Figure 12.44 Length frequency distribution of total headrope length (ft) of otter trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.331

Figure 12.45 Length frequency distribution of vessels (ft) using skimmer trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.332

DRAFT

Figure 12.46	Length frequency distribution of total headrope length (ft) of skimmer trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.	332
Figure 12.47	Length frequency distribution of vessels (ft) using otter trawls in the Atlantic Ocean, 2010-11.	333
Figure 12.48	Length frequency distribution of total headrope length (ft) of otter trawls in the Atlantic Ocean, 2010-11.....	333
Figure 12.49	Scatter plot of vessel length (ft) and total headrope length (ft) of the North Carolina shrimp trawler fleet (all water bodies included), 2010-11.....	334
Figure 12.50	Weighted CPUE of Atlantic croaker, spot, and weakfish from the NCDMF Pamlico Sound Survey and number of shrimp trawl trips (effort lagged 1 year) in Pamlico Sound and Neuse, Pamlico and Pungo rivers (DMF 2012).	339
Figure 12.52	Brunswick County Shrimp Management Areas.....	343

2.3 LIST OF ACRONYMS

AC – Advisory Committee

AEC – Areas of Environmental Concern

AFSA – Anadromous Fish Spawning Areas

ASMFC – Atlantic States Marine Fisheries Commission

BDTRT – Bottlenose Dolphin Take Reduction Team

BMP – Best Management Practices

BOD – Biological Oxygen Demand

BRD – Bycatch Reduction Device

CAHA – Cape Hatteras National Seashore

CAMA – Coastal Area Management Act

CEIP – Coastal Energy Impact Program

CHPP – Coastal Habitat Protection Plan

COE – United States Army Corp of Engineers

CPI – Consumer Price Index

CPUE – Catch Per Unit Effort

CRC – North Carolina Coastal Resources Commission

CRFL – Coastal Recreational Fishing License

DCM – North Carolina Division of Coastal Management

DENR – North Carolina Department of Environment and Natural Resources

DMF – Division of Marine Fisheries, The

DO – Dissolved Oxygen

DOT – North Carolina Department of Transportation

DPS – Distinct Population Segments

DWQ – North Carolina Division of Water Quality

E – Endangered

DRAFT

EDC – Endocrine Disrupting Chemicals

EEP – Ecosystem Enhancement Program

EEZ – Exclusive Economic Zone

EFH – Essential Fish Habitat

EMC – North Carolina Environmental Management Commission

EPA – United States Environmental Protection Administration

ESA – Endangered Species Act

FDA – United States Food and Drug Administration

FED – Fish Escape Device

FFE – Florida Fish Excluder

FMP – Fishery Management Plan

FRA – Fishery Reform Act

FRG – Fishery Resource Grant

FSC – Federal Species of Concern

GIS – Geographical Information System

GS – General Statute

GSAFDF – Gulf and South Atlantic Fisheries Development Foundation

HQW- High Quality Waters

IGNS – Independent Gill Net Study

ITP – Incidental Take Permit

IWW – Intracoastal Waterway

JHA - Juvenile Hormone Analogs

JLCSA – Joint Legislative Commission for Seafood and Aquaculture

LMEF – Large Mesh Extended Funnel

LMFE – Large Mesh Funnel Excluder

MAFMC – Mid Atlantic Fisheries Management Council

DRAFT

MFC – North Carolina Marine Fisheries Commission

MLMFE – Modified Large Mesh Funnel Excluder

MMAP – Marine Mammal Authorization Program

MMPA – Marine Mammal Protection Act

MRFSS – Marine Recreational Fisheries Statistical Survey

MRIP – Marine Recreational Information Program

MRT – Management Review Team

MSC – Moratorium Steering Committee

MSFCMA – Magnuson-Stevens Fishery Conservation and Management Act

NC – North Carolina

NCAC – North Carolina Administrative Code

NCDA – North Carolina Department of Agriculture

NCDACS – North Carolina Department of Agriculture and Consumer Services

NCDMF – North Carolina Division of Marine Fisheries

NCSU – North Carolina State University

NCTTP – North Carolina Trip Ticket Program

NEFSC – Northeast Fisheries Science Centers

NMFS – National Marine Fisheries Service

NOAA – National Oceanic and Atmospheric Administration

NWL – Normal Water Level

PDT – Plan Development Team

PNA – Primary Nursery Area

PPI – Producer Price Index

PPT – Parts Per Thousand

PSE – Proportional Standard Error

PSGNRA – Pamlico Sound Gill Net Restricted Area

DRAFT

RAT – Rules Advisory Team

RCGL – Recreational Commercial Gear License

RDD – Random Digit Dialing

RSCFL – Retired Standard Commercial Fishing License

SAFMC – South Atlantic Fishery Management Council

SAV – Submerged Aquatic Vegetation

SC – Species of Concern

SCFL – Standard Commercial Fishing License

SHA – Strategic Habitat Area

SNA – Secondary Nursery Area

SSB – Spawning Stock Biomass

SSA – Southern Shrimp Alliance

SSNA – Special Secondary Nursery Area

SSR – Stock Status Report

STAC – Sea Turtle Advisory Committee

STSSN – Sea Turtle Stranding and Salvage Network data

T – Threatened

TED – Turtle Excluder Device

TNC – The Nature Conservancy

TRP – Take Reduction Plan

USFWS – United States Fish and Wildlife Service

VPA – Virtual Population Analysis

WRC – North Carolina Wildlife Resources Commission

WS – Water Supply

YOY – Young of the Year

DRAFT

3.0 EXECUTIVE SUMMARY

The first North Carolina Shrimp Fishery Management Plan (FMP) was developed and approved by the Marine Fisheries Commission (MFC) in 2006. The plan set forth measures necessary to address bycatch, habitat, and conflict among shrimp fishermen as well as with other user groups. NC FMPs are reviewed at a minimum of every five years. A Shrimp FMP Plan Development Team (PDT) met beginning in the fall of 2011 to review the 2006 Shrimp FMP and determine whether it should be amended or revised. The PDT concluded that current management strategies in the plan continue to meet the goals and objectives of the Shrimp FMP and recommended to the Director of the Division of Marine Fisheries that the 2011 Shrimp FMP should proceed as a revision. After review by the MFC, the revision was taken to the MFC regional advisory committees, the MFC Habitat and Water Quality Advisory Committee and the MFC Crustacean/Shellfish Advisory Committee for review and public comment. Based on the concerns voiced at these meetings by the public on bycatch in the shrimp trawl fishery, the DMF recommended amending the 2006 Shrimp FMP. The MFC, at its November 2012 meeting, directed the DMF to amend the Shrimp Plan but to limit the scope of the amendment to bycatch issues in the commercial and recreational fisheries.

The goal of the North Carolina Shrimp FMP is to utilize a management strategy that provides adequate resource protection, optimizes the long-term commercial harvest, maximizes social and economic value, provides sufficient opportunity for recreational shrimpers, and considers the needs of all user groups. To achieve this goal, it is recommended that the following objectives be met:

1. Minimize waste and enhance economic value of the shrimp resource by promoting more effective harvesting practices.
2. Minimize harvest of non-target species of finfish and crustaceans and protected, threatened, and endangered species.
3. Promote the protection, restoration, and enhancement of habitats and environmental quality necessary for enhancing the shrimp resource.
4. Maintain a clear distinction between conservation goals and allocation issues.
5. Reduce conflicts among and within user groups, including non-shrimping user groups and activities.
6. Encourage research and education to improve the understanding and management of the shrimp resource.

There are three shrimp species that make up the shrimp fishery in North Carolina. These are the brown shrimp, *Farfantepenaeus aztecus*, the pink shrimp, *F. duorarum* and the white shrimp, *Litopenaeus setiferus*. Collectively, they are commonly referred to as penaeid shrimp. These species, considered annual crops, have similar life histories and are susceptible to similar predation, parasites and disease.

Population size is regulated by environmental conditions, and while fishing reduces the population size over the season, fishing is not believed to have a major impact on subsequent

DRAFT

year class strength unless the spawning stock has been reduced below a minimum threshold level by environmental conditions. Because of high fecundity and migratory behavior, shrimp are capable of rebounding from a very low population size in one year to a large population size in the next, provided environmental conditions are favorable. Fluctuations in abundance resulting from changes in environmental conditions will continue to occur. Shrimp stocks of all three species in North Carolina are considered viable.

3.1 STATUS OF THE FISHERIES

3.1.1 Commercial Fishery

Between the Civil War and the end of the first decade of the twentieth century, shrimp were caught with dip nets, cast nets and seines. The introduction of otter trawl technology in North Carolina seems to have first involved sampling nets used by the U.S. Bureau of Fisheries in Beaufort in 1912. The use of this technology prompted the development of trawl vessels. The type that was first used in the fishery involved open skiffs from 15 to 20 foot in length that were powered by small gasoline engines. As the fishery expanded during the 1930s, the construction of larger vessels specifically designed for shrimp trawling expanded. Technological advances in the shrimping industry have increased the catching efficiency of larger boats, particularly in Pamlico Sound. Modern safety and navigation equipment have allowed North Carolina shrimpers to steam longer distances, for longer periods of time to shrimp; and also to engage in a constantly changing variety of harvesting activities other than shrimping throughout the calendar year.

Landings in the North Carolina shrimp fishery vary from year to year and are dependent primarily on environmental conditions. The annual average was 6,460,849 lb for the period 1962-2010; 75% were harvested from inshore waters and 25% from the Atlantic Ocean. About 70-93% of shrimp trips occur in estuarine waters, with the remainder in ocean waters, primarily within state territorial seas (<3 mi offshore) off the central and southern coast of North Carolina. Total annual shrimping effort has decreased from a high of 40,000 trips in 1982 to a low of 6,500 trips in 2005. An examination of harvest by water body for the most recent twelve year period shows that 56% of the landings are from Pamlico Sound, 24% from the Atlantic Ocean and 6% from Core Sound. No other water bodies contribute more than 4% to the state's total landings.

The vast majority of the shrimp harvest (92%) is taken by otter trawls however, there has been a slight shift in the types of gear used to harvest shrimp in North Carolina in recent years. There has been an increase in number of vessels in Carteret, Onslow, and Pender counties that have switched from otter trawls to skimmers with skimmers accounting for 3% of the average annual state landings. Channel nets are stationary nets that fish the surface and middle depths on an outgoing tide. They resemble a staked-out trawl anchored to the bottom to keep it open. Channel nets account for 5% of the average annual shrimp landings.

North Carolina brown shrimp commercial landings have averaged 3.8 Mlb since 1999 with fluctuations from a high of 6.5 Mlb in 2000 to a low of 1.5 Mlb in 2005. Generally, 85% of all brown shrimp landed are caught in estuarine waters with Pamlico Sound, Core Sound, New River, and Neuse River accounting for most of the harvest.

Pink shrimp have historically (1978-1993) accounted for about 27% of the shrimp landings. North Carolina commercial pink shrimp landings averaged 1.8 Mlb from 1978 to 1993. However, since 1999, pink shrimp landings have averaged only 0.2 Mlb. despite mild winters in

DRAFT

the late 1990s and early 2000s. Pink shrimp have accounted for 4% of the state's harvest during the last 12 years. Core Sound accounts for 45% of the landings, followed by Pamlico Sound (29%), and the ocean (13%).

During the period 1978-1993, North Carolina commercial white shrimp landings averaged 0.5 Mlb with landings fluctuating from a high of 1.7 Mlb in 1993 to a low of 11,000 pounds in 1981. The landings increased significantly for the most recent 12 years to an average of 2.5 Mlb. The percentage of the white shrimp taken in the ocean is higher (40%) than the other two species, reflecting its greater abundance in the southern part of the state where the majority of the ocean fishery occurs. Since 1999 the majority of white shrimp have been harvested from the Ocean (40%), Pamlico Sound (35%) and New River (6%).

3.1.2 Recreational Fishery

Shrimp are harvested recreationally throughout the state by otter trawls, skimmer trawls, seines, cast nets, shrimp pots and shrimp pounds. As of July 1, 1999, anyone wishing to harvest shrimp recreationally with commercial gear is required to purchase a Recreational Commercial Gear License (RCGL). RCGL allow recreational fishermen to use limited amounts of commercial gear to harvest seafood for their personal consumption. Seafood harvested under this license cannot be sold. RCGL holders are limited to the same bag and size limits as Coastal Recreational Fishing License (CRFL) holders.

On average the highest number of RCGL trips using shrimp trawls from 2002 to 2008 occurred in the Pamlico region, followed by the southern region, the central region, and the northern region. In the Pamlico region, the number of trips ranged from 1,127 (2005) to 2,384 (2002), averaging 1,642 per year from 2002 to 2008. In the southern region, the number of trips ranged from 355 (2007) to 1,123 (2002), averaging 586 trips per year. An average of 413 trips a year were made in the central region, ranging from 132 (2008) to 1,070 (2002). In the Northern region, the number of trips ranged from 50 (2006) to 911 (2004). Overall, the highest number of trips made by RCGL using shrimp trawls was observed in 2002; the lowest was observed in 2007.

RCGL holders harvested an average of 52,352 pounds of shrimp a year from 2002 to 2008 with the highest landings occurring in 2002 (101,766 lb), followed by 2008 (54,359 lb) and 2003 (50,961 lb). RCGL holders harvested an average of 16.8 pounds of shrimp per trip from 2002 to 2008. The highest pounds of shrimp per trip was observed in 2009 (22.3 lb/trip), followed by 2006 (20.3 lb/trip) and 2002 (19.1 lb/trip).

3.1.3 Shrimp Trawl Bycatch

Bycatch can be divided into two components: incidental catch and discarded catch. Incidental catch refers to retained catch of non-targeted species. Discarded catch is that portion of the catch returned to the sea as a result of economic, legal, or personal considerations. While it is becoming increasingly apparent to scientists, natural resource managers, and much of the general public that bycatch is an important issue that must be addressed, characterizing the nature and extent of bycatch and its impact on fish stocks has proven extremely difficult. Although many species are caught as bycatch in the estuarine shrimp trawl fishery, four species, blue crab, weakfish, Atlantic croaker, and spot have, since the first studies were conducted in the 1950s and continuing to the present, accounted for the bulk of the bycatch.

DRAFT

Trawl minimum mesh size regulations are the principal method used to regulate fishing mortality on fish stocks and is the preferred management tool in lieu of other more stringent regulations. Bycatch reduction devices (BRDs) and turtle excluder devices (TEDs) are required in shrimp trawls for use in reducing bycatch of finfish and sea turtles. Other gears such as shrimp pots, pounds and cast nets also reduce finfish bycatch; minimize environmental concerns and conflicts with other fisheries. Catch restrictions have been used by fisheries managers to maintain fish stocks, extend fishing seasons, allocate resources, and reduce bycatch. In North Carolina this method is being used to reduce the targeting of marketable finfish with shrimp trawls. Area restrictions for trawling are also used to deal with allocation, resource, bycatch, habitat, and safety issues in North Carolina.

3.2 PROTECTED RESOURCES

Of the federal and state protected species listed, only bottlenose dolphins, and sea turtles interact with the shrimp fishery. Otter trawls and skimmer trawls are the predominant gear in the shrimp fishery. Both trawls are active gears that focus on the estuarine bottom, and are restricted to areas without submerged aquatic vegetation; interactions with protected species are plausible. Channel nets used less extensively in the shrimping fishery are a passive gear and use tide flow and current to fish. There is no information on interactions with protected species and channel nets.

3.3 ECONOMIC STATUS

The annual nominal (inflated) value of shrimp landings typically has been volatile with large changes between years. The lowest nominal value was \$3.5 million in 1972. The highest nominal value was \$25.4 million in 2000. Landings value in 1981 dropped 69% from 1980. The fishery rebounded in 1982 with a 210% increase in the nominal value of landings over 1981. The value of the fishery dropped by 53% in 2001 from the record high 2000 value. In 2002, the value increased 54% over the 2001 value, but it remained considerably lower than the 2000 value. The nominal value hit a 20 year low in 2005 (\$4.4 million), dropping 50% over the previous year's value; however, the fishery recovered to over \$19.2 million in 2008. Ex-vessel value of landings in 2010 was \$10.7 million.

3.4 SOCIOECONOMIC CHARACTERISTICS

DMF surveys (2005, 2007, 2008, and 2009) asked the fishermen for their opinion as to how historically important they think commercial fishing is to their community. On a scale of one to ten, with one being not at all important to ten being extremely important, the average rating across all 175 persons interviewed was 9.7, indicating almost universal agreement that fishing has been historically important to their community. When asked how much their community supports commercial fishing now (using the same 10-point scale), the rating was 8.1, indicating they largely feel supported.

North Carolina coastal communities rely significantly less on commercial fishing now than in the past. This is the result of the development of the communities as multiple use zones, with retirement, light industry, recreation, and tourism becoming the dominant domains of the local economies. Fewer and fewer native born residents make a full time living as fishermen like those in previous generations. DMF studies found that among commercial shrimp fishermen, the average fisherman earned about 76% of his income from commercial fishing. More specifically the studies found that just over half (51%) were totally reliant on fishing for their

DRAFT

incomes. This compares with data gathered in the late 1980s where nearly all full time fishermen captains were committed to fishing for nearly all (95%) of their incomes.

3.5 ENVIRONMENTAL FACTORS

Penaeid shrimp use a variety of estuarine and coastal ocean habitats with variations in habitat preference due to location, season, and ontogenetic stage. Penaeid shrimp are found in the water column, wetlands, submerged aquatic vegetation (SAV), soft bottom, and shell bottom. Each of these habitats is part of a larger habitat mosaic, which plays a vital role in the overall productivity and health of the coastal ecosystem. Although penaeid shrimp are found in all of these habitats, the usage varies by habitat. Additionally, these habitats provide the appropriate physicochemical and biological conditions necessary to maintain and enhance the penaeid shrimp population. Each habitat provides ecological services that aid in maintaining and enhancing shrimp stock sustainability, and also influences the functioning of the ecosystem overall. Protecting the integrity of the entire system is therefore necessary to manage this species.

Adequate water quality is also necessary to maintain the chemical properties of the water column that are needed by shrimp, as well as sustain SAV, shell bottom, and soft bottom habitats that support shrimp. Human activities that degrade water quality or alter water flow can negatively impact shrimp growth or survival. The common causes of water quality use support impairment in North Carolina's coastal river basins are excessive sediment loading and low dissolved oxygen (DO). Hydrological modifications, low DO and toxin contamination are probably the greatest water quality concerns for penaeid shrimp in North Carolina.

3.6 MANAGEMENT STRATEGIES

There were several major issues identified as being pertinent to the shrimp fishery in the 2006 FMP. These included trawling (bycatch, habitat), competition among shrimp fishermen as well as with other user groups and insufficient bycatch data. Management strategies were developed to address these issues and will remain in place in the Shrimp FMP Amendment 1. Shrimp trawling is a controversial topic and has been the subject of much debate. Strategies from the 2006 plan that address both bycatch and habitat concerns include area closures and restrictions in gear size in specific water bodies. Gear studies and shrimp trawl characterization studies have been carried out and will continue to be addressed in the future.

Protection of vulnerable habitats from the effects of trawling was also achieved through implementation of the 2006 FMP. Additional closures along with increased gear restrictions and gear more friendly to habitat continue to be implemented in this plan.

The use of trawls by RCGL holders and the significance of its impact on the shrimp fishery was examined. Management strategies include; a 48-quart limit on recreational shrimp catches, allowing skimmer trawls as a RCGL gear and defined dimensions of a shrimp pound for use as a RCGL gear.

Many of the management strategies are water body specific and address user conflicts through area and gear restrictions. A 90 foot headrope limit in internal waters, with the exception of Pamlico Sound and portions of the Neuse, Pamlico and Pungo rivers, also reduces conflict as well as decrease bycatch.

3.6.1 Shrimp Management by Size

Shrimp grow at different rates depending on water temperature and salinity. As growth increases, shrimp migrate to deeper, saltier waters of the sound and eventually to the ocean. As shrimp migrate to the ocean, they enter areas that are open or may be opened by DMF to the harvest of shrimp. Sampling is conducted by DMF staff to determine if an area should be opened or closed, based primarily on size and count. Over time, target sizes for opening different waterbodies have evolved and allow for better flexibility of management for both recreational and commercial shrimping.

3.6.2 Shrimp Management by Area

Historically, DMF has used a number of criteria to determine if trawling should be allowed in estuarine waters. These criteria include habitat issues such as aquatic vegetation, water depth and bottom types; shrimp size and abundance; economic and social factors; user conflicts; and bycatch issues. DMF uses rules and proclamations to manage trawling in internal coastal waters. The intention of these rules and proclamations has been to allow the harvest of shrimp and crabs in estuarine waters but prohibit directed finfish trawling. Openings and closings of specific areas are based primarily on the size of the shrimp.

3.6.3 Shrimp Management in the Southern District

The areas that can be opened to shrimping are typically located either in or landward of the Intracoastal Water Way (IWW) which runs the entire length of the Onslow, Pender, New Hanover and Brunswick counties coastline. In Brunswick and portions of New Hanover counties, where shrimp migrate at smaller sizes, DMF attempts to open on a 40-50 count shrimp. In Onslow and parts of Pender counties, sampling has shown that a 20-30 count can be attained before migration occurs. Channels that connect the IWW with the Atlantic Ocean are normally left open at all times to allow some harvest of shrimp as they migrate from closed areas to the ocean.

3.6.4 Shrimp Management in the Central District

Management of shrimping in the Central District takes place from the White Oak River on the Onslow/Carteret County line to Core Sound in Carteret County. The Central District also manages the south side of the Neuse River in Craven County. Areas that are open and closed to shrimping through proclamation include: West Bay/Long Bay, Thorofare Bay, several tributaries in Core Sound and Adams Creek, located on the south side of the Neuse River. Target counts vary dependent on the waterbody and range from 26 to 30 count to 31 to 35 count (heads-on).

3.6.5 Shrimp Management in Pamlico District

Management of shrimping in the Pamlico District occurs in the Neuse, Pamlico, Pungo and Bay rivers as well as Pamlico Sound. These areas with the exception of Bay River have permanent closure lines and requires little sampling. As sampling dictates, lines may be moved downstream by proclamation to protect small shrimp until they are large enough to harvest. The target count size is ranges from a 26-30 count or 31-35 count (heads-on). When sampling

DRAFT

indicates that the majority of the shrimp in a closed area have reached this target size, the area is opened by proclamation.

3.6.6 Shrimp Management in the Northern District

Species specific shrimp sampling occurs in the Northern District only when necessary (during banner shrimp years). An exception is data collected in Stumpy Point Bay for brown shrimp. The low relative abundance of white and pink shrimp in the Northern District requires minimal sampling effort except during times of extreme environmental conditions. In such cases, sampling efforts may be initiated on demand in order to provide the foundation for shrimp management decisions of DMF.

3.6.7 Atlantic Ocean

Since shrimp that migrate from the estuaries are usually large, DMF does not actively manage the ocean waters. However, in the past and exclusively off the Brunswick County coast, DMF has been requested by the fishermen to take a more active role in the management of the ocean shrimp fishery. These requests were precipitated as result of the heavy hurricane or tropical storm induced rains that have impacted southeastern North Carolina with regularity since the mid-1990s. Fresh water from these heavy rains dramatically reduces salinities in the estuaries causing the shrimp to prematurely migrate from the estuaries into the ocean. When this occurs, DMF generally closes the impacted ocean and estuarine waters to shrimp trawling.

3.7 BYCATCH IN THE SHRIMP FISHERY AND MANAGEMENT OPTIONS

The DMF, at the direction of the MFC, presented the 2012 Shrimp FMP revision to the MFC Southern Regional AC, the MFC Northern Regional AC, the MFC Habitat and Water Quality AC and the MFC Shellfish/Crustacean AC and also took public comment at each of these committees. With the exception of the Southern AC, all of the committees voted to revise the Shrimp FMP. However, due to the overwhelming public comment concerning the issue of bycatch in the shrimp trawl fishery and the acknowledgement that bycatch is an issue in the shrimp fishery; the DMF changed its recommendation to the MFC to move forward with amending the Shrimp FMP. The MFC then directed the division amend the plan but to limit the scope of the amendment to bycatch issues in the commercial and recreational fisheries.

Twenty-nine different management options were brought forward to a Shrimp FMP Advisory Committee (AC) to address eight different issues during monthly meetings from January through September 2013. Management strategies that were discussed included: Alternative fishing gears, Turtle Excluder Devices (TEDs) in skimmer trawls, gear modifications, effort management, head rope lengths, number of nets and vessel lengths, and area restrictions. In addition, at the request of the Southern AC, during the public comment review of the 2012 Shrimp FMP revision, the New River trawl fishery and the consideration of a live bait shrimp fishery was also addressed through Amendment 1.

The MFC, at its November 2013 meeting approved Amendment 1 for review by the public, regional and standing MFC committees and selected preferred management strategies during its February 2014 meeting.

DRAFT

3.7.1 Marine Fisheries Commission Preferred Management Strategies and Required Actions

The Commission’s preferred management strategies and required actions based on input are listed in table 3.1 and are identified under each bycatch issue addressed. An overview of the Shrimp Advisory Committee’s, the division’s, and the commission’s recommendations to reduce bycatch as provided to each regional and standing committee for input are provided in Appendix 1. Recommendations from each regional and standing committee as well as public input may also be found in Appendix 1.

Table 3.1 The Marine Fisheries Commission preferred management strategies, and required actions to reduce bycatch.

Management Strategy	Required Actions
Status quo (continue to prohibit otter trawls in the special secondary nursery area above the Highway 172 Bridge).	Rule change required in 15A NCAC 03J .0208
Allow hand cast netting of shrimp in all closed areas and increase the limit to four quarts per person, heads on.	Rule change required in 15A NCAC 03L .0105
Status quo on a license requirement to fish a cast net for shrimp.	No action required
Upon federal adoption of TEDs in skimmer trawls, the division will support the federal requirement.	No action required
Establish a permitted live shrimp bait fishery and for DMF to craft the guidelines and permit fees after reviewing permitted operations in other states, and allow live bait fishermen with a permit to fish until 12 p.m. (noon) on Saturday.	Based on review of other state operations, future rule changes will be required and include 15A NCAC 03J .0104, 03L .0102, 03O .0105, 03O .0503
Allow any federally certified BRD in all NC internal and offshore waters.	Existing proclamation authority
Update the scientific testing protocol for the state BRD certification program.	Existing authority
<p>Convene a stakeholder group to initiate industry testing of minimum tail bag mesh size, T-90 panels, skylight panels, and reduced bar spacing in TEDs to reduce bycatch to the extent practicable with a 40 percent target reduction.</p> <ul style="list-style-type: none"> • Upon securing funding, testing in the ocean and internal waters will consist of three years of data using test nets compared to a control net with a Florida fish eye, a federally approved TED and a 1.5-inch mesh tail bag. • Results should minimize shrimp loss and maximize reduction of bycatch of finfish. Promising configurations will be brought back to the commission for consideration for mandatory use. • The stakeholder group may be 	Existing authority

DRAFT

<p>partnered with the division and Sea Grant.</p> <ul style="list-style-type: none"> Members should consist of fishermen, net/gear manufacturers and scientific/gear specialists. 	
<p>Require either a T-90/square mesh tailbag or other applications of square mesh panels (e.g., skylight panel), reduced bar spacing in a TED, or another federal or state certified BRD in addition to existing TED and BRD requirements in all skimmer and otter trawls.</p>	<p>Existing proclamation authority Rule change required in 15A NCAC 03I .0101</p>
<p>Status quo on effort management (no change in season, weekend, or night time fishing).</p>	<p>No action required</p>
<p>Cap fleet capacity as a management tool, and establish a maximum combined headrope length of 220 feet in all internal coastal waters where there are no existing maximum combined headrope requirements with a two year phase out period.</p>	<p>Rule change required in 15A NCAC 03L .0103</p>
<p>Prohibit shrimp trawling in the IWW channel from Sunset Beach to the SC state line, including Eastern Channel, lower Calabash River and Shallotte River.</p>	<p>Rule change required in 15A NCAC 03R .0114</p>
<p>Recommend the Habitat and Water Quality Advisory Committee consider changing designation of special secondary nursery areas that have not been opened to trawling since 1991 to permanent secondary nursery areas.</p>	<p>Based on review of the advisory committee, rule changes will be required and include 15A NCAC 03R .0104, 03R .0105</p>

DRAFT

4.0 INTRODUCTION

4.1 LEGAL AUTHORITY FOR MANAGEMENT

Fisheries management includes all activities associated with maintenance, improvement, and utilization of the fisheries resources of the coastal area, including research, development, regulation, enhancement, and enforcement.

Many different state laws (General Statutes - G.S.) provide the necessary authority for fishery management in North Carolina. General authority for stewardship of the marine and estuarine resources by the North Carolina Department of Environment and Natural Resources (DENR) is provided in G.S. 113-131. The Division of Marine Fisheries (DMF) is the arm of the Department that carries out this responsibility. Enforcement authority for DMF enforcement officers is provided by G.S. 113-136. General Statute 113-163 authorizes research and statistical programs. The North Carolina Marine Fisheries Commission (MFC) is charged to “manage, restore, develop, cultivate, conserve, protect, and regulate the marine and estuarine resources of the State of North Carolina” (G.S. 143B-289.51). The MFC can regulate fishing times, areas, fishing gear, seasons, size limits, and quantities of fish harvested and possessed (G.S. 113-182 and 143B-289.52). General Statute 143B-289.52 allows the MFC to delegate authority to implement its regulations for fisheries “which may be affected by variable conditions” to the Director of DMF by issuing public notices called “proclamations”. Thus, North Carolina has a very powerful and flexible legal basis for coastal fisheries management. The General Assembly has retained for itself the authority to establish commercial fishing licenses and mandates that there will be no fees charged for permits unless specifically authorized. It has delegated to the MFC authority to establish permits for various commercial fishing activities.

The Fisheries Reform Act of 1997 (FRA) establishes a process for preparation of coastal fisheries management plans in North Carolina (G.S. 113-182). The Act was amended in 1998 and again in 2004. The FRA states that “the goal of the plans shall be to ensure the long-term viability of the State’s commercially and recreationally significant species or fisheries. Each plan shall be designed to reflect fishing practices so that one plan may apply to a specific fishery, while other plans may be based on gear or geographic areas. Each plan shall:

- a. Contain necessary information pertaining to the fishery or fisheries, including management goals and objectives, status of the relevant fish stocks, stock assessments for multi-year species, fishery habitat and water quality considerations consistent with Coastal Habitat Protection Plans adopted pursuant to G.S. 143B-279.8, social and economic impact of the fishery to the State, and user conflicts.
- b. Recommend management actions pertaining to the fishery or fisheries.
- c. Include conservation and management measures that will provide the greatest overall benefit to the State, particularly with respect to food production, recreational opportunities, and the protection of marine ecosystems, and will produce a sustainable harvest.
- d. Specify a time period, not to exceed two years from the date of the adoption of the plan, for ending overfishing. This subdivision shall only apply to a plan for a fishery that is not producing a sustainable harvest.

DRAFT

- e. Specify a time period, not to exceed 10 years from the date of adoption of the plan, for achieving a sustainable harvest. This subdivision shall not apply if the Fisheries Director determines the biology of the fish, environmental conditions, or lack of sufficient data make implementing the requirement of this subdivision incompatible with professional standards for fisheries management.
- f. Include a standard of at least fifty percent (50%) probability of achieving sustainable harvest for the fishery or fisheries. This subdivision shall not apply if the Fisheries Director determines the biology of the fish, environmental conditions, or lack of sufficient data make implementing the requirement of this subdivision incompatible with professional standards for fisheries management.

Sustainable harvest is defined in the FRA as “The amount of fish that can be taken from a fishery on a continuing basis without reducing the stock biomass of the fishery or causing the fishery to become overfished”.

Overfished is defined as “The condition of a fishery that occurs when the spawning stock biomass of the fishery is below the level that is adequate for the recruitment class of a fishery to replace the spawning class of the fishery”.

Overfishing is defined as “Fishing that causes a level of mortality that prevents a fishery from producing a sustainable harvest”.

4.2 RECOMMENDED MANAGEMENT PROGRAM

4.2.1 Goals and Objectives

The goal of the North Carolina Shrimp Fishery Management Plan is to utilize a management strategy that provides adequate resource protection, optimizes the long-term commercial harvest, maximizes social and economic value, provides sufficient opportunity for recreational shrimpers, and considers the needs of all user groups. To achieve this goal, it is recommended that the following objectives be met:

1. Minimize waste and enhance economic value of the shrimp resource by promoting more effective harvesting practices.
2. Minimize harvest of non-target species of finfish and crustaceans and protected, threatened, and endangered species.
3. Promote the protection, restoration, and enhancement of habitats and environmental quality necessary for enhancing the shrimp resource.
4. Maintain a clear distinction between conservation goals and allocation issues.
5. Reduce conflicts among and within user groups, including non-shrimping user groups and activities.
6. Encourage research and education to improve the understanding and management of the shrimp resource.

DRAFT

4.2.2 Sustainable Harvest

Sustainable harvest for the penaeid shrimp fishery in North Carolina is defined as the amount of harvest that can be taken by fishermen without reducing the spawning stock below the level necessary to ensure adequate reproduction. This is appropriate for an annual crop such as shrimp when recruitment is dependent largely on environmental conditions rather than female biomass. That is, a relatively small number of mature shrimp can provide sufficient recruits for the subsequent year's production. The sustainable harvest for the Shrimp FMP in North Carolina is the annual harvest of the three species of shrimp combined.

4.2.3 Management Strategy

The management strategy for the shrimp fisheries in North Carolina is to continue to 1) optimize resource use over the long-term, and 2) minimize waste. The first strategy is accomplished by protection of critical habitats, and gear and area restrictions to protect the stock. Minimization of waste is accomplished by gear modifications, bycatch reduction devices, area closures, and harvest restrictions.

4.3 DEFINITION OF MANAGEMENT UNIT

The management unit includes the three major shrimp species of shrimp: brown (*Farfantepenaeus aztecus*), pink (*Farfantepenaeus duorarum*), and white (*Litopenaeus setiferus*) and its fisheries in all coastal fishing waters of North Carolina, which includes the Atlantic Ocean offshore to three miles.

4.4 GENERAL PROBLEM STATEMENT

During preparation of Amendment 1, bycatch in the commercial and recreational shrimp fishery was addressed. Management options were separated into 1) gear modifications; 2) effort management; 3) area restrictions; and 4) other fishing gears. In addition, at the request of the Southern AC, during the public comment review of the 2012 Shrimp FMP revision, the New River trawl fishery and the consideration of a live bait shrimp fishery was also addressed through Amendment 1.

4.4.1 New River Trawl Fishery

At the request of the Southern AC, the prohibition of otter trawls in the New River Special Secondary Nursery Area (SSN) was discussed. The use of otter trawls upstream of the Highway 172 Bridge was phased out in 2010 following the adoption of the 2006 Shrimp FMP. Those who wished to continue to harvest shrimp in the waters above the Highway 172 Bridge were allowed a four year grace period to convert to skimmers trawls.

4.4.2 Use of Other Fishing Gears

The majority (89%) of the estuarine shrimp harvest in North Carolina comes from otter trawls. However, major concerns associated with otter trawls are the capture and discard of various amounts of other non-target species and discard mortality associated with otter trawls. Skimmer trawls, channel nets, pound nets and cast nets are other gears that are used to harvest shrimp; however factors that impact these gears' effectiveness have to be considered.

4.4.3 Gear Modifications

One available management measure to reduce bycatch is gear modification requirements. Potential gear modifications requirements such as requiring TEDs in skimmer trawls, mesh size changes, bycatch reduction devices (BRDs), and square mesh panels modifications are considered to further reduce bycatch in the shrimp fishery. Also considered by the AC were testing of gear by the industry and the creation of a stakeholder group as well as updating the division's BRD certification program.

4.4.4 Live Bait Shrimp Fishery

At the request of the Southern AC, the consideration of a regulatory process for live bait shrimping was addressed. The division's current management for larger shrimp causes areas where live bait shrimping occurs to close due to the presence of small shrimp. Bycatch occurs in the smaller bait trawls but at-net bycatch mortality is generally low due to short tow times, and culling times. However as temperatures increase, mortality usually increases as well.

4.4.5 Effort Management

In considering ways to reduce bycatch, reducing effort in the shrimp trawl fishery with seasonal closures, closing trawling an additional day of the week, nighttime closures, and restricting tow times were discussed by the AC. Also considered were ways to make the fishery less efficient by looking at restricting headrope lengths, vessel size, and net size.

4.4.6 Area Restrictions

Since 1978 almost one million acres of estuarine waters have been closed to trawling through fishery nursery area designations (primary and secondary nursery areas), military danger zones and restricted areas, and trawl net prohibited areas. This is approximately 45 percent of the estuarine waters. Another 65,000 acres of estuarine waters are closed some time during the year, either due to shrimp size management or areas classified as SSNAs. Several area restrictions ranging from closing all internal waters to closing a small area in Brunswick County was discussed as another way to reduce bycatch.

4.5 EXISTING PLANS STATUTES, AND RULES

4.5.1 Plans

The South Atlantic Fishery Management Council (SAFMC) Shrimp Fishery Management Plan allows for concurrent closures of Federal waters in conjunction with State closures through emergency action, following severe winter cold weather that results in an 80% or more reduction in the population of overwintering shrimp or if water temperatures are 8°C (46° F) for a minimum of one week and was implemented through Amendment 9. This cooperative plan allows maximum protection of the remaining adult population. The Council plan and its amendments have had minimal impact on the NC shrimp fishermen until the addition of Amendment 6. In December 2003, as part of Amendment 6, the Council voted to establish a control date of December 10, 2003 for the shrimp fishery in the Atlantic Exclusive Economic Zone (EEZ). This control date was set to place the industry on notice that a limited access

DRAFT

program may be developed. The amendment also includes options to monitor and measure bycatch within the fishery. The Council's preferred monitoring option is the implementation of the Atlantic Cooperative Statistics Program Release, Discard and Protected Species Module. This module establishes a minimum set of standard data to be collected to characterize and estimate levels of bycatch.

The first North Carolina Shrimp Fishery Management Plan was developed and approved by the MFC in 2006. The plan set forth measures necessary to address bycatch, habitat, and competition among shrimp fishermen as well as with other user groups. NC FMPs are reviewed at a minimum of every five years. A Shrimp FMP PDT met beginning in the fall of 2011 to review the 2006 Shrimp FMP and determine whether it should be amended or revised. The PDT discussed several management issues that had developed since the implementation of the 2006 FMP and included discussion of:

- restricted trawl areas offshore of Bogue Banks;
- a permanent shrimping line in Newport River;
- trawling in New River above the highway 172 bridge; and
- volumetric measurement of shrimp.

Each issue was addressed by the PDT in issue papers, providing background information as well as management options and their potential positive and/or negative impacts on the fishery (see Appendix 1). After thorough discussion, no changes in management strategies were recommended for three of the four issues listed above. The group did make recommendations to change from a count of 100 shrimp per person per day to a volumetric measure of shrimp of two quarts per person per day that may be taken while fishing in a closed area with a cast net. This recommendation is due to the nature of this fishery, where several fishermen will work together and combine their harvest in a single large container. This often leads to one Marine Patrol Officer having to count shrimp while surrounded by numerous fishermen, putting that Officer's safety at risk. This does result in a rule change for the benefit of Officer's Safety and it was concluded that this rule change will not have an impact on the shrimp stock and will remain consistent with the 2006 Shrimp FMP.

To insure the public was aware that the 2006 Shrimp FMP was under review, a press release was sent out on November 7, 2011 requesting public comment as part of the review process to determine whether to proceed with an amendment or a revision of the 2006 Shrimp FMP. The Division received six comments which were reviewed and addressed by the PDT (see Appendix 1). The PDT concluded that these comments and recommendations were appropriately addressed within the 2006 Shrimp FMP and that current management strategies in place are continuing to meet the goals and objectives of the 2006 Shrimp FMP. Therefore, it was recommended by the PDT to the Director that the 2012 Shrimp FMP should proceed as a revision.

The DMF, at the direction of the MFC, presented the 2012 Shrimp FMP revision to the MFC Southern Regional AC, the MFC Northern Regional AC, the MFC Habitat and Water Quality AC and the MFC Shellfish/Crustacean AC and also took public comment at each of these committees. With the exception of the Southern AC, all of the committees voted to revise the Shrimp FMP. However, due to the overwhelming public comment concerning the issue of

DRAFT

bycatch in the shrimp trawl fishery, the DMF changed its recommendation to the MFC to move forward with amending the Shrimp FMP. Meeting minutes from each committee and public comment may be found in Appendix 1 of this plan.

4.5.2 Statutes

All management authority for North Carolina's shrimp fishery is vested in the State of North Carolina. Statutes that have been applied to the shrimp fishery include:

- It is unlawful to engage in a commercial fishing operation in coastal fishing waters without holding a standard commercial fishing license (SCFL) [G.S. 133-1682].
- Individuals who are 65 years of age or older and are eligible for a SCFL may apply for a retired standard commercial fishing license (RSCFL) [G.S. 133-168.3].
- It is unlawful to fish in the ocean from vessels or with a net within 750 feet of a properly licensed and marked fishing pier [G.S. 113-185].
- It is unlawful to engage in trash or scrap fishing (the taking of young of edible fish before they are of sufficient size to be of value as individual food fish) for commercial disposition as bait, for sale to any dehydrating or nonfood processing plant, or for sale or commercial disposition in any manner. The MFC's rules may authorize the disposition of the young of edible fish taken in connection with the legitimate commercial fishing operations, provided it is a limited quantity and does not encourage "scrap fishing" [G.S. 113-185].
- It is unlawful to willfully take, disturb or destroy any sea turtles including green, hawksbill, loggerhead, Kemp's ridley, and leatherback turtles, or their nests or eggs. It shall be unlawful to willfully harm or destroy porpoises [G.S. 113-189].
- It is unlawful for any person without the authority of the owner of the equipment to take fish from nets, traps, pots, and other devices to catch fish which have been lawfully placed in the open waters of the State [G.S. 113-268 (a)].
- It is unlawful for any vessel in the navigable waters of the State to willfully, wantonly, and unnecessarily do injury to any seine, net or pot [G.S. 113-268 (b)].
- It is unlawful for any person to willfully destroy or injure any buoys, markers, stakes, nets, pots, or other devices or property lawfully set out in the open waters of the state in connection with any fishing or fishery [G.S. 113-268 (c)].

4.5.3 Marine Fisheries Commission Rules

4.5.3.1 General

- Channel net is defined as a net used to take shrimp which is anchored or attached to the bottom at both ends or with one end anchored or attached to the bottom and the other end attached to a boat [15A NCAC 31 .0101 (3)(b)].
- Headrope is defined as a support structure for the mesh or webbing of a trawl that is

DRAFT

nearest to the water surface when in use. [15A NCAC 3I .0101 (3)(i)].

- Nursery areas are defined as areas in which for reasons such as food, cover, bottom type, salinity, temperature and other factors, young finfish and crustaceans spend the major portion of their initial growing season [15A NCAC 3I .0101 (4)(f)].
- There is a cooperative agreement between the DENR, the MFC, and the Wildlife Resources Commission (WRC) that the WRC will have regulatory jurisdiction over any species of sea turtle and their eggs and nests consistent with the designation of endangered or threatened. Law enforcement officers of both the DMF and the WRC have jurisdiction to enforce any state laws and rules relating to endangered or threatened species of sea turtles [15A NCAC 3I .0107 (a)].
- The Fisheries Director may close or restrict by proclamation any coastal waters with respect to taking or attempting to take any marine resources when the method used is a serious threat to an endangered or threatened species [15A NCAC 3I .0107 (b)].
- Military danger zones and restricted areas are designated in 15A NCAC 3R .0102 and are enforced by the appropriate federal agency [15A NCAC 3I .0110 (a)].
- Maps or charts showing the boundaries of areas identified by rule or in proclamations are available for inspection [15A NCAC 3I .0121 (a)].
- The DMF shall mark boundaries with signs insofar as may be practical. No removal or relocation of signs shall have the effect of changing the classification or affect the applicability of any rule pertaining to that body of water [15A NCAC 3I .0121 (b)].

4.5.3.2 Nets, Pots, Dredges, and Other Fishing Devices

- It is unlawful to use or set a fixed or stationary net in the Intracoastal Waterway where it may be a hazard to navigation, block more than two-thirds of any natural or manmade waterway, in the middle third of any marked navigation channel [15A NCAC 3J .0101 (1)(2)(3)].
- It is unlawful to possess aboard a vessel while using a trawl in internal waters more than 500 pounds of finfish from December 1 through February 28 and 1,000 pounds of finfish from March 1 through November 30 [15A NCAC 3J .0104 (a)].
- It is unlawful to use trawls nets in internal coastal waters from 9:00 p.m. on Friday through 5:00 p.m. on Sunday, except for the areas described in the next bullet [15A NCAC 3J .0104 (b) (1)].
- It is unlawful to use trawl nets from December 1 through February 28 from one hour after sunset to one hour before sunrise in portions of the Pungo, Pamlico, Bay, Neuse, and New rivers [15A NCAC 3J .0104 (b) (5)(A)(B)(C)(D)(E)].
- Trawls cannot be used to take oysters [15A NCAC 3J.0104 (2)].
- It is unlawful to use trawl nets in Albemarle Sound and its tributaries [15A NCAC 3J .0104 (b) (3)].

DRAFT

- The Director may by proclamation, require bycatch reduction devices or codend modifications in trawl nets to reduce the catch of finfish that do not meet size limits or are unmarketable as individual foodfish by reason of size [15A NCAC 3J .0104 (d)].
- It is unlawful to use trawl nets in designated pot areas opened to the use of pots by 15A NCAC 3J .0301(a)(2) within an area bound by the shoreline to the depth of six feet [15A NCAC 3J .0104 (6)].
- It is unlawful to use shrimp trawls for the taking of blue crabs in internal waters, except that it shall be permissible to take or possess blue crabs incidental to commercial shrimp trawling provided that the weight of the crabs shall not exceed 50 percent of the total weight of the combined crab and shrimp catch; or 300 pounds, whichever is greater [15A NCAC 3J .0104 (f)(2)].
- For RCGL trawling, 50 crabs, not to exceed 100 blue crabs if two or more RCGL holders are on board [15A NCAC 3J .0104 (f)(1)].
- It is unlawful to use shrimp trawls for recreational purposes unless the trawl is marked with a pink buoy on the tailbag [15A NCAC 3J .0104 (e)].
- The Fisheries Director may, by proclamation, close any area to trawling for specific time periods in order to secure compliance with this rule [15A NCAC 3J .0104 (g)].
- It is unlawful to use a channel net until the Director specifies by proclamation when and where channel nets and other fixed nets for shrimping can be used [15A NCAC 3J .0106 (a)(1)].
- It is unlawful to set a channel net without yellow light reflective tape on the staffs, stakes and buoys [15A NCAC 3J .0106 (a)(2)].
- Channel nets cannot be set with any portion of the set within 50 feet of the center line of the Intracoastal Waterway (IWW) channel or in the middle third of any navigation channel marked by the Corps of Engineers or the Coast Guard. Fishermen must attend channel nets by being no more than 50 yards from the set at all times [15A NCAC 3J .0106 (a)(3)(4)(5)].
- The maximum corkline length of a channel net that can be used or possessed is 40 yards. No channel net, net buoys or stakes can be left in coastal waters from December 1 through March 1. From March 2 through November 30, cables and any attached buoy must be connected together with non-metal line when not attached to the net. Metallic floats or buoys to mark sets are unlawful [15A NCAC 3J .0106 (b)(c)(d)(e)].
- Channel nets must be properly marked with yellow light reflective tape and the owner's identification on each buoy. Identification includes one of the following: owner's NC motorboat registration number or the US vessel documentation number or owner's last name and initials. Channel nets, anchor lines or buoys are not to be used in any way that constitutes a hazard to navigation [15A NCAC 3J .0106 (f) and (g)].
- It is unlawful to use channel nets to take blue crabs in internal waters, except that it shall

DRAFT

be permissible to take or possess blue crabs incidental to channel net operations provided that the weight of the crabs do not exceed 50% of the total weight of crab and shrimp or 300 lbs, whichever is greater [15A NCAC 3J .0106 (h)(1)(A)(B)].

- The Director may, by proclamation, close any area to channel net use for specific time periods in order to secure compliance with the above bullet [15A NCAC 3J .0106 (h)(2)].
- It is unlawful to use nets from June 15 through August 15 in the waters of Masonboro Inlet or in the ocean within 300 yards of the beach between Masonboro Inlet and a line running 138° through the water tank on the northern end of Wrightsville Beach, a distance parallel with the beach of 4,400 yards. It is unlawful to use trawls within one-half mile of the beach between the Virginia line and Oregon Inlet [15A NCAC 3J .0202 (1)(2)].
- It is unlawful to use a trawl with a mesh length less than four inches in the body and three inches in the extension and on and three-fourths inches in the cod end or tail bag from the west side of Beaufort Inlet Channel to the shore off Salter Path within a half mile of shore [15A NCAC 3J .0202 (3)].
- From December 1 through March 31 it is unlawful to possess finfish caught incidental to shrimp and crab trawling in the Atlantic Ocean unless the weight of the combined catch of shrimp and crabs exceeds the weight of finfish; except that crab trawlers working south of Bogue Inlet may keep up to 300 pounds of kingfish, regardless of their shrimp or crab catch weight [15A NCAC 3J .0202 (5)].
- It is unlawful to use shrimp trawls in all waters west of a line beginning at the southeastern tip of Baldhead Island at a point 33° 50.4833'N – 77° 57.4667 W; running southerly in the Atlantic Ocean to a point 33° 46.2667'N – 77° 56.4000 W from 9:00 PM through 5:00 AM [15A NCAC 3J .0202 (8)].
- It is unlawful to use trawl nets upstream of the Highway 172 Bridge in New River from 9:00 p.m. through 5:00 a.m. when opened by proclamation from August 15 through November 30 (15A NCAC 3J .0208).
- It is unlawful to use any commercial fishing gear in the Southport Boat Harbor, Brunswick County and to use any commercial fishing gear in the Progress Energy Intake Canal between the fish diversion screen and the Brunswick nuclear power plant (15A NCAC 3J .0206, 15A NCAC 3J .0207).
- It is unlawful to use shrimp pots with mesh lengths smaller than one and one-fourth inches stretch or five-eighths inch bar [15A NCAC 3J .0301(e)].
- It is unlawful to use pots with leads or leaders to take shrimp. Leads are defined as any fixed or stationary net or device used to direct fish into any gear [15A NCAC 3J .0301(l)].
- It is unlawful for a RCGL holder to use pots, including shrimp pots unless each pot is marked by attaching one hot pink floating buoy; the buoy should be engraved with the gear owners boat registration number or US vessel documentation name [15A NCAC 3J .0302(a)(1)(2)].

DRAFT

- In Dare County commercial fishing gear may not be used within 750 feet of licensed fishing piers when opened to the public. Commercial fishing gear may not be used in the Atlantic Ocean off of portions of Onslow, Pender, and New Hanover counties during specified time frames [15A NCAC 3J .0402(a)(1)(A)(ii)(2)(A)(B)(i)(ii)(3)(A)(B)(i)(iii)(4)].
- Shrimp pound net set is defined as a pound net set constructed of stretch mesh equal to or greater than one and one-fourth inches and less than or equal to two inches [15A NCAC 3J .0501(a)(6)].
- A permit is required to deploy a pound net set and must be operational for a minimum of 30 consecutive days during the permit period. Each pound required the permittee's identification on a sign attached to a stake at the permitted ends of each set at all times. They must have yellow light reflective tape or yellow light reflective devices on each pound and have a marked navigational opening at least 25 feet wide at the end of every third pound and marked with yellow light reflective tape or yellow light reflective devices [15A NCAC 3J .0501 (b)(c)].
- It is unlawful to use a RCGL shrimp pound net unless it is marked by attaching to the offshore lead, one hot pink floating buoy. The owner shall be identified on the buoy by engraving the gear owner's current boat registration number or the owners US vessel documentation name. Each shrimp pound must be set a minimum of 100 yards from a RCGL pound net set or 300 yards from an operational permitted shrimp pound net set [15A NCAC 3J .0501(d)(1)(2)].
- It is unlawful within 30 days of abandonment of a permitted pound net set to fail to remove all stakes and associated gear from coastal fishing waters [15A NCAC 3J .0501(g)].
- Pound net permit applications, renewals and transfers are to comply with the permitting procedures and requirements for obtaining all DMF-issued permits. Application process, criteria for the granting of the permit, operational requirements and other elements of the shrimp pound net set permits are found in 15A NCAC 3J .0502, 15A NCAC 3J .0503, 15A NCAC 3J .0504 and 15A NCAC 3J .0505.

4.5.3.3 Oysters, Clams, Scallops and Mussels

- It is unlawful to use a trawl net in any designated Shellfish or Seed Management area [(15A NCAC 03K .0103 (b))].
- It is unlawful to use a trawl in any designated Seed Oyster Management Area [15A NCAC 03K .0208 (b)].
- It is unlawful to use a trawl in Oyster Sanctuaries [15A NCAC 03K .0209 (a)].

4.5.3.4 Shrimp, Crabs, and Lobster

- It is unlawful to take shrimp with nets until the Director opens the season in various waters by proclamation (15A NCAC 03L .0101).
- It is unlawful to take shrimp by any method from 9:00 PM on Friday through 5:00 p.m. on

DRAFT

Sunday except in the Atlantic ocean or with the use of fixed and channel nets, hand seines, shrimp pots and cast nets [15A NCAC 03L .0102 (1)(2)].

- It is unlawful to take shrimp with mesh lengths less than one and one-half inches in trawls, one and one-fourth inches in fixed nets, channel nets, float nets, butterfly nets and hand seines [15A NCAC 03L .0103)(a)(1)(2)].
- It is unlawful to take shrimp with a net constructed in a manner as to contain an inner or outer liner of any mesh size. Net material used as chafing gear shall be no less than four inches mesh length [15A NCAC 03L .0103) (b)].
- It is unlawful to take shrimp with trawls which have a combined headrope of greater than 90 feet in internal coastal waters except in Pamlico Sound, Pamlico River downstream of Pamlico Point/ Willow Point and Neuse River downstream of Winthrop Point/Windmill Point [15A NCAC 03L .0103)(c)(1)(2)(3)].
- It is unlawful to use a shrimp trawl in the Pungo River, upstream of Wades Point/Abels Bay, Pamlico River upstream of the entrance to Goose Creek/Wades Point and Neuse River upstream of Cherry Point/Wilkerson Point 15A [NCAC 03L .0103)(d)].
- It is unlawful to use a shrimp trawl that does not conform with the federal requirements for TEDs [15A NCAC 03L .0103)(g)].
- It is unlawful to possess more than 48 quarts, heads-on or 30 quarts heads-off of shrimp per person per day or per vessel per day for recreational purposes [15A NCAC 03L .0105)(1)].
- It is unlawful to take or possess shrimp taken from any area closed to the taking of shrimp except for 2 quarts per person per day may be taken with a cast net in a closed area [15A NCAC 03L .0105)(2)].
- It is unlawful to use trawls in the crab spawning sanctuaries from March 1 through August 31 [15A NCAC 03L .0205)(a)].
- It is unlawful to use a trawl net in any primary or permanent secondary nursery area [15A NCAC 3N .0104, 3N .0105 (a)].
- Special secondary nursery areas may be opened to shrimp and crab trawling from August 16 through May 14 [15A NCAC 3N .0105)(b)].

4.5.3.5 Licenses, Leases, Franchises, and Permits

- RCGL gear includes one shrimp trawl with a headrope not exceeding 26 feet in length per vessel, five shrimp pots, skimmer trawls, not exceeding 26 feet in total combined width and one shrimp pound net with each lead 10 feet or less in length and with a minimum lead net mesh of 1 ½ inches and enclosures constructed of net mesh of 1 ¼ inches or greater and with all dimensions being 36 inches or less. Attendance is required at all times for shrimp pounds [15A NCAC 3O .0302(a)(2)(3)(7)(8)].
- It is unlawful to possess more than 48 quarts, heads-on, or 30 quarts, heads-off, of

DRAFT

shrimp when only one person aboard a vessel possesses a valid RCGL and recreational commercial fishing equipment [15A NCAC 3N .0303(e)].

- It is unlawful to possess more than 96 quarts, heads on or 60 quarts, heads off of shrimp if more than one person aboard a vessel possesses a valid RCGL and recreational commercial fishing equipment [15A NCAC 3N .0303(f)].
- It is unlawful to trawl for shrimp in the Atlantic Ocean without TEDs within one nautical mile of shore from Browns Inlet to Rich's Inlet without a valid permit to waive the requirement to use TEDs in the Atlantic Ocean when allowed by proclamation from April 1 through November 30. It is unlawful to tow more than 55 minutes from April 1 through October 31 and 75 minutes from November 1 through November 30. It is unlawful to not fully empty the contents of each net after each tow. It is unlawful to refuse to take observers. It is unlawful to fail to report any sea turtle captured [15A NCAC 03O .0503 (d) (1)(2)(3)(4)(5)].

4.5.5 Federal Regulations

33 CFR 334.410 through 334.450

These rules designate prohibited and restricted military areas, including locations within North Carolina coastal fishing waters, and specify activities allowed in these areas.

50 CFR 223.206 - Exceptions to prohibitions relating to sea turtles.

The incidental taking of sea turtles in the shrimp trawl fishery is exempted from section 9 of the Endangered Species Act (ESA) if conservation regulations are followed and include the installation of NOAA Fisheries approved TEDs and alternative tow times for skimmer trawls, pusher-head trawls and butterfly trawls.

50 CFR 223.207 – Approved TEDs

This lists NOAA Fisheries approved TEDs such as the single-grid hard TEDs, hooped hard TEDs, special hard TEDs and soft TEDs, along with materials and gear specifications. Testing protocols for TEDs are also included in this rule.

50 CFR 229.7 – Monitoring of incidental mortalities

This requires that fishermen who participate in a Category I or II fishery are required to accommodate an observer onboard your vessel(s) up on request

50 CFR 622, Appendix D – Approved BRDs

This lists NOAA Fisheries approved BRDs and provides technical specifications for the construction and subsequent legal enforcement of these BRDs.

DRAFT

5.0 STATUS OF THE STOCK

5.1 GENERAL LIFE HISTORY

There are three shrimp species that make up the shrimp fishery in North Carolina. These are the brown shrimp, *Farfantepenaeus aztecus*, the pink shrimp, *F. duorarum* and the white shrimp, *Litopenaeus setiferus*. The lifecycle of these three species are similar in that the adults spawn offshore and eggs are hatched into free-swimming larvae. These larvae develop through several stages into post-larvae. Once post-larval shrimp enter the estuaries, growth is rapid and is dependent on salinities and temperatures. After reaching sub-adult sizes between 70 - 120 mm TL, they migrate seaward. It is hypothesized that as shrimp increase in size, they seek higher more stable salinities because of a decrease in the ability to osmoregulate (Bishop et al. 1980). In low salinity environments the growth rates of juvenile shrimp have been found to be significantly reduced because energy that would be allocated to somatic growth is used for osmoregulation (Rozas and Minello 2011). In general, shrimp are omnivorous, feeding primarily on sediment, detritus, algae, and benthic organisms. Feeding occurs mostly at night, although some daytime feeding will occur in turbid water. Shrimp are dioecious (separate sexes) with females growing larger than males. Shrimp copulate with the male depositing spermatophore onto the female's thelycum. Fertilization takes place when the female expels ova and spermatozoa simultaneously. Shrimp are very fecund with females expelling between 500,000 to 1,000,000 eggs. Spawning occurs before they reach 12 months old. Environmental requirements for the three species are listed in Table 5.1.

5.1.1 Brown Shrimp

Brown shrimp occur from Massachusetts to the Florida Keys and into the Gulf of Mexico to northwestern Yucatan. Highest abundances occur in the Gulf of Mexico, off Mississippi, Louisiana, and Texas. The species supports a major commercial fishery along the South Atlantic coast, primarily in North and South Carolina.

Brown shrimp reach sexual maturity at 140-145 mm and spawn in the ocean in deep water during February and March. Brown shrimp are thought to have the potential to spawn more than once within a year in the Gulf of Mexico (Calillouet Jr. et al. 2008); however, St. Amant et al. (1966) suggest that brown shrimp die after spawning once. After the eggs are hatched, larvae are then transported by wind and currents from the high salinity ocean waters to the estuaries. Ten to 17 days later, the larval shrimp have grown into postlarvae and are approximately between 8 and 14 mm. They generally enter the inlets on a flood tide. They are then carried by wind driven currents to the upper reaches of the estuaries beginning in February with peaks occurring in mid-March through mid-April (Williams 1955a, 1965). It takes approximately 4-6 weeks for postlarvae to grow to the juvenile stage. Rapid development into sub-adults begins to occur with reported growth rates ranging from 1 to 2.5 mm per day and is dependent on temperature and salinities (Williams 1955; Steele 2002). Significant growth occurs between 11°C and 18°C (Zein-Eldin and Aldrich 1965; Steele 2002). Growth is enhanced if salinities are greater than 10 ppt (Amant et al. 1966; Steele 2002) and reduced if salinities are less than 4 ppt (Saoud and Davis 2003). As the individuals increase in size, they move to the deeper, saltier waters of the sound and return to the sea in late fall. Brown shrimp are omnivorous, and feed on different plants and animals and organic debris (Steele 2002). Juveniles between 25 and 65 mm feed on detritus and microorganisms from the top layer of sediment while larger shrimp (65-104 mm) become active predators feeding on polychaetes, amphipods, nematodes as well as detritus and algae (Jones 1973; Steele 2002). Brown shrimp prefer peat and muddy bottoms but are also found on sand, silt, or clay mixed with shell and

DRAFT

rock fragments (Steele 2002). They also are found on bottoms covered with plant debris (Williams 1959). They are often more active in open waters at night than in daytime. Brown shrimp have a maximum life span of 18 months and may reach a size of 7 to 9 in.

5.1.2 Pink Shrimp

Pink shrimp are found from southern Chesapeake Bay to the Florida Keys, and around the coast through the Gulf of Mexico to Yucatan. The largest population of pink shrimp is off southwestern Florida in the Tortugas and Sanibel as well as in the southeastern portion of Golfo de Campeche. However, significant quantities of pink shrimp are also found off North Carolina, and along the northeast Florida coast (Steele 2002).

Spawning occurs in ocean waters from April to July with post larvae being carried into the estuary on wind-driven currents from May through November (Williams 1965). Histological examination of the ovaries of pink shrimp in Florida indicates year-round spawning; however, seasonal differences in water temperatures may inhibit spawning in the northern most regions (Kennedy and Barber 1981). The northernmost breeding population of pink shrimp is off North Carolina (Williams 1955a). Once in the nursery areas, the shrimp undergo rapid growth (1 to 1.8 mm/day). As they grow and develop, they move toward the deeper waters of the sound and eventually into the ocean. Browder et al. (2002) noted that pink shrimp growth is optimal at a salinity of 30 ppt and decreases as salinity increases or decreases around this mark; however growth was found to increase with temperature up to 35°C. Pink shrimp are active at night and burrow into the bottom during the day. A significant number of pink shrimp overwinter in the North Carolina estuaries before moving into the ocean the following spring. Pink shrimp are bottom feeders and feed primarily in shallow waters among marine plants. As with brown shrimp, the majority of feeding occurs at night, but feeding may also occur during the day when the water is turbid. Stomach content analysis of pink shrimp in Tampa Bay revealed sand, debris, algae, diatoms, seagrass particles, dinoflagellates, foraminiferans, nematodes, polychaetes, ostracods, copepods, mysids, isopods, caridean shrimp, caridean eggs, mollusks and fish scales. Female pink shrimp reach sexual maturity at 85 mm while males are sexually mature at 74 mm. They have a maximum life span of 24 months and can reach a size of 10 to 11 in.

5.1.3 White Shrimp

White shrimp occur along the Atlantic coast from Fire Island, New York to Saint Lucie Inlet Florida (Steele 2002). They also are found in the Gulf of Mexico from the mouth of the Ochlockonee River, Florida to the Golfo de Campeche to the vicinity of Ciudad Campech usually in depths less than 90 ft (Muncy 1984; Steele 2002).

Spawning occurs in the ocean at depths greater than 30 ft and within five miles of shore from March to November, peaking from April to October. White shrimp are capable of spawning more than once in a year (Nance et al. 2010); however, it is thought that they may only spawn once in North Carolina waters (Williams 1965). Spawning appears to be triggered by increasing bottom water temperatures in the spring and decreases with decreasing water temperature in the fall (Muncy 1984). In South Carolina, extremely cold spring water temperatures were found to delay sexual maturation while slightly warmer temperatures promoted maturation (DeLancey et al. 2005). Planktonic postlarvae move inshore with tidal currents, entering estuaries two to three weeks after hatching where they then become benthic. Shallow muddy bottoms in waters of low to moderate salinity serve as optimum nursery grounds for juvenile white shrimp. Juveniles reach lengths of about 20-31 mm by July, and move from shallow marshes into

DRAFT

deeper creeks, rivers, bays and sounds. White shrimp migrate out of the estuaries and southward during fall and early winter, and make up the valuable spring fishery for adult females in Georgia, South Carolina, and southern North Carolina. Some of the slower-growing individuals overwinter in the estuaries, but usually do not survive in North Carolina. White shrimp mortality has been reported at water temperatures of 46° F and lower, with total mortality occurring at 37° F or lower. Winter water temperatures in North Carolina sometimes are lethal for white shrimp. DeLancey et al. (2005) noted that the relative abundance of white shrimp was strongly influenced by winter water temperature, indicating that periods of milder winters yielded higher relative abundances of white shrimp in South Carolina. White shrimp are omnivorous, selective particulate feeders that search the sand grains and pass bits of food forward to the mouth. Gut content analysis findings include inorganic and organic debris, as well as fragments of different animals including nematodes, annelids, mollusks, crustaceans, particles of higher plants and a variety of diatoms and algae (Steele 2002). Soft muddy bottoms are the preferred habitat of white shrimp with highest abundances in areas of extensive brackish marshes. White shrimp have a maximum life span of 16 months and can reach a size of 7 to 8 in.

Table 5.1 Environmental Requirements of three shrimp species found in North Carolina.

Species	Salinity	Temperature	Oxygen	Juvenile recruitment	Season
Brown Shrimp	2-35 ppt	7° to 37° C (44.6° to 98.6° F)	< 2 ppm causes stress	February - March	Summer and fall
Pink Shrimp	0-45 ppt	6° to 38° C (42.8° to 100.4° F)	0.2 to 6.0 ppm	June - October	Spring Late
White Shrimp	2-35 ppt	7° to 38° C (44.6° to 100.4° F)	< 2 ppm causes stress	April - May	Summer and fall

5.1.4 Movement

DMF conducted several tagging studies on the three species of shrimp in the 1960s through the early 1970s (Table 5.2). Shrimp were marked with biological stains and fluorescent pigments and released throughout this time period within different areas of Pamlico Sound, Core Sound, Bogue Sound, New River and Cape Fear River. These shrimp were recovered in shrimp houses throughout the coastal counties. Rewards ranged from 0.50 cents to \$1.00 per returned shrimp.

McCoy and Brown (1967) marked brown and pink shrimp from Jarrett Bay and North River in Core Sound and white shrimp in Dutchman Creek-Elizabeth River and Cape Creek of the lower Cape Fear River. A combined average of 65% of all returned shrimp were recaptured before reaching the Atlantic ocean with resulting movement toward the higher salinity areas of Beaufort Inlet from Core Sound and Cape Fear Inlet from Cape Fear River.

White shrimp did move upriver in the Cape Fear River. However, this was caused by the strong tidal influences in the river. Of those shrimp that made it to the Atlantic Ocean, all three species had a pronounced southward coastal migration. It was concluded in this study that the brown and pink shrimp are more endemic to North Carolina while the white shrimp from the

DRAFT

southeastern coastal NC contribute to the shrimp fishery of South Carolina, Georgia, and Florida.

Table 5.2 Migration studies in North Carolina of three shrimp species found in North Carolina.

Study	Year	Waterbodies	Species	Release number	Percent return
McCoy and Brown (1967)	April-Oct, 1966	Core Sound, Lower Cape Fear	Brown, Pink, White	26,989	6.2
McCoy (1968)	June-Sept, 1967	Pamlico Sound	Brown, Pink	11,414	10.5
McCoy (1972)	May, July, 1968	Core and Bogue Sound, New River	Brown, Pink	9,231	42.4
Purvis and McCoy (1974)	1971-1972	Pamlico Sound	Brown	7,325	19.1

McCoy (1968) marked pink shrimp from West Bay that moved to the Atlantic Ocean through Core Sound and through Drum and Beaufort inlets. Pink shrimp from Adams Creek moved toward Beaufort Inlet and through southern Pamlico Sound to Drum Inlet and Bardens Inlet. This suggests that a significant portion of Pamlico Sound pink shrimp reach the ocean through Beaufort and Bardens inlets by migrating through Core Sound.

Brown shrimp marked by McCoy (1968) in Swan Quarter Bay and Jones Bay generally moved toward the central and southern Pamlico Sound area. Data were unclear as to the most probable route to the Atlantic Ocean but it did suggest that few shrimp from the northern and western sound reached the ocean. Brown shrimp randomly released in Pamlico Sound in 1972 generally moved toward the nearest inlet (Ocracoke). However, no mass migration from the sound to the ocean occurred to any appreciable degree resulting in the conclusion that the Pamlico Sound brown shrimp fishery is a self contained fishery with shrimp growing to large sizes (16-30 count heads-off) before migrating to the ocean (Purvis and McCoy 1972).

Pink shrimp marked in Core Sound moved to the ocean through Barden and Beaufort inlets with the majority of the movement through Beaufort Inlet. Bogue Sound pink shrimp moved toward the ocean via Beaufort and Bogue inlets with the largest number of recaptures occurring from the western half of the sound. There appeared to be no significant movement of pink shrimp between Core and Bogue Sound. Brown shrimp released in New River moved to the ocean in a southerly direction along the coast (McCoy 1972).

5.1.5 Predation

Shrimp are preyed upon by numerous species of finfish and invertebrates at various stages of their life cycle (Bielsa et al. 1983; Muncy 1984; Larson et al. 1989; Minello et al. 1989). Facendola and Scharf (2012) found that penaeid shrimp made up 30.7% of the diet (by weight) of age 0-1 juvenile red drum and 1.1% (by weight) of the diet of age 1-2 red drum in the New River; indicating as red drum grow, their diets shift from shrimp and crabs to primarily fish. Penaeid shrimp also have been reported to make up a large portion of the diets of other sciaenids as well as the diets of numerous finfish commonly found in marine and estuarine environments (Carr and Adams 1973; Minello and Zimmerman 1983). Additionally, a wide variety of coastal and wading birds are also known to prey upon shrimp. Given that penaeid shrimp are such an important food source for multiple species of organisms it is hard to quantify

DRAFT

exactly how much is consumed by each species and what affect it has on the year-to-year fluctuations in shrimp abundance.

5.1.6 Parasites and Disease

Diseases and parasites in penaeid shrimps come in the forms of viruses, bacteria, fungi, protozoa, flatworms and nematodes. Johnson (1978) noted that penaeid shrimp are vulnerable to numerous diseases which may be caused by microbes (bacteria, fungi, viruses), protozoa (microsporidians, gregarines, apostome ciliates, ectocommensal protozoa), as well as physical and chemical factors (lack of oxygen, poisons, low temperatures, salinity extremes). Disease ranks second only to predation and mass kills of natural populations in the Gulf and South Atlantic (Couch 1978) in shrimp mortality. The *Baculovirus* infects larval and adult shrimp and is associated with mortality, especially in larval shrimp. The effect of bacteria on mortality is unclear; however *Vibrio*, *Beneckeia*, and *Leucothrix* are associated with disease in penaeid shrimps. Several types of fungi can be very destructive to tissue of larval shrimp. There are several types of protozoa that are parasitic and commensal and include Microsporidia which cause the condition commonly known as “cotton shrimp” or “milk shrimp” and Ciliata which causes black gill disease. Flatworms and nematodes can also be found in muscles and viscera of penaeid shrimp (Couch 1978).

Cotton disease is widespread and is found in all three species of shrimp on the South Atlantic and Gulf coasts (Johnson 1978; Bielsa 1983; Muncy 1984; Larson 1989). There are several species of *Microsporidia* that infects the tail muscle of the shrimp, as well as the organs and tissues with masses of spores. These spores cause the white discoloration of muscle giving infected shrimp a cotton or paper-white color. These types of infections can also cause black banding throughout the abdomen of infected shrimp and can render shrimp incapable of reproduction (Johnson 1978). This parasite kills shrimp. A typical catch of wild shrimp contains a few infected individuals. Infected shrimp are not thought to be harmful to humans; however they are often discarded due to appearance and texture of the infected tissue.

Black gill disease results from infection by a single-celled protozoan called a Ciliata (SC DNR 2002). It attaches itself to a thin area around the gills of the shrimp's shell. This attachment either causes structural damage or erodes a hole through the shell, causing inflammation. The black pigmentation of the gill results from an immune response to the inflammation. Black gill is thought to inhibit respiration, slowing growth and potentially making shrimp more prone to predation. However, the infestation of black gill disease does not result in any noticeable mortality in the wild and appears to attach in mass when shrimp are stressed (SC DNR 2002). Black gill has been observed in pink, brown, and white shrimp (Johnson 1978). Black gill poses no threat to humans.

Several penaeid shrimp viruses may be carried by imports from Asia and South America as well as from expanding aquaculture. These viruses enter processing facilities and aquaculture facilities through infected brood stock, contaminated feed, infected transport containers or by migratory birds. These viruses may infect our three species of native shrimp but there is little information on the presence of exotic shrimp viruses in populations of our native shrimp in North Carolina. There is currently one permitted, *Penaeus vannamei* (Pacific White Shrimp) farm in Vass, NC and another under review in Morrisville (C. Hardy. NCDMF, personal communication).

5.2 STOCK STATUS

All three species of shrimp included in this FMP are essentially annual crops. Population size is regulated by environmental conditions, and while fishing reduces the population size over the season, fishing is not believed to have any impact on subsequent year class strength unless the spawning stock has been reduced below a minimum threshold level by environmental conditions. Estimates of population size are not available but since the fishery is considered to be fished at near maximum levels, annual landings are probably a good indication of relative abundance. Annual variations in catch are presumed to be due to a combination of prevailing environmental conditions and fishing effort. More recently, landings are showing the effects of changes in the economics of the fishery.

Because of high fecundity and migratory behavior, the three species are all capable of rebounding from a very low population size in one year to a large population size in the next, provided environmental conditions are favorable. Fluctuations in abundance resulting from changes in environmental conditions will continue to occur. Perhaps the most serious threat to the stocks is loss of habitat due to pollution or physical alteration. Especially vulnerable and critical to shrimp production is the salt marsh (for white and brown shrimp) and inshore seagrass habitat (especially for pink shrimp) which comprise the nursery areas for juvenile shrimp. Since the inception of the 2006 Shrimp FMP, shrimp stocks of all three species in North Carolina are still considered viable.

DRAFT

6.0 STATUS OF FISHERIES

6.1 COMMERCIAL

6.1.1 History

Between the Civil War and the end of the first decade of the twentieth century, shrimp were caught with dip nets, cast nets and seines. Most were consumed locally but some were used as bait and fertilizer. Distant markets were limited because of little interest in shrimp for food, production capability, few transportation options, and the lack of refrigeration (Maiolo 2004; Maiolo et al. 1980).

Just after the turn of the twentieth century, the South Atlantic and Gulf states became the center of the commercial shrimp fishery in the United States. Interest in the fishery developed rapidly in the Southport, N.C. area. The adoption of the otter trawl completely changed the means of harvesting, which fit nicely with the earlier innovations in power boating at the end of the previous century and market stimulation from the New York area. The creation of canning factories in Southport followed (Maiolo 2004; Maiolo et al. 1980).

The introduction of the otter trawl technology in North Carolina seems to have first involved sampling nets used by the U.S. Bureau of Fisheries in Beaufort in 1912. Even with this new and efficient capture technology, interest in the fishery was not uniform among coastal fishing villages. As late as the 1920s many fishermen still referred to shrimp as “pests” that fouled their nets and many residents, both coastal and inland, did not consider the animals suitable to eat (Maiolo 1981).

In March of 1916, a New Jersey fisherman brought a shrimp trawler to Southport and taught local fishermen how to use the otter trawl in the near shore ocean waters. Interest among fishermen expanded quickly in spite of a sluggish local market. By 1925, over 300 North Carolina fishermen were engaged in the shrimp fishery, mostly in Brunswick County (Maiolo 2004).

The use of otter trawl net technology prompted the development of trawl vessels. The type that was first used in the fishery involved open skiffs from 15 to 20 ft in length that were powered by small gasoline engines. “Decked” trawlers were introduced in the 1920s. Refrigeration (in the form of production of ice for shipment of fishery products), rail and truck transportation, and a close proximity to the eastern markets (as opposed to Florida and the Gulf states) began to make the North Carolina shrimp fishery lucrative. More than two hundred seasonal and part time workers found employment in the Southport packinghouses where many headed shrimp for a nickel per five gallon bucket. The majority of shrimp were shipped to markets in northeastern New York because local markets were still not developed (Maiolo 2004; Maiolo et al. 1980).

The first shrimp trawling in Carteret County occurred around 1930 after local fishermen learned how to harvest the resource from the Southport fishermen. At first, shrimping only occurred in between finfishing seasons. At the same time, a channel net fishery was developing near Harkers Island and in other communities in eastern Carteret County. A series of local customs developed among the fishermen by which the fishery was prosecuted. Many remain in place even today. A similar fishery has recently developed near Snead’s Ferry, but without the same kinds of local customs (Maiolo 2004).

DRAFT

Pamlico County fishermen began landing shrimp caught in pound nets about this time as well, and shrimp trawling caught on in northern Pamlico Sound in the late 1930s when a Louisiana fisherman demonstrated the use of the otter trawl. Shrimping in the northern counties was conducted both nearshore and in the Pamlico Sound. In 1934, the Pamlico Sound was closed to trawling to prevent finfish bycatch. But the following year, the regulation was modified to allow shrimp trawling from 15 August to 1 December (Maiolo et al. 1980; Maiolo 2004).

Like fishermen in other coastal communities in North Carolina who stitched shrimp harvesting into their patterns of annual rounds, fishermen in the northern part of the state pursued shrimping during the summer between oyster dredging and fall finfishing. Just as today, in the southern part of the state, some fishermen followed the shrimp south into South Carolina and Georgia in late summer and into fall (Maiolo 2004; Maiolo et al. 1980).

As the fishery expanded during the 1930s, the construction of larger vessels specifically designed for shrimp trawling expanded. Two of the most common vessels were the “Florida trawler” for ocean trawling in the southern part of the state, and the “Core Sounder” for estuarine trawling. Along with this, masts and booms, or masts and “A” frames, were developed. Additionally, power winches replaced retrieval of the nets by hand. The construction of trawls and doors locally which, up to then had occurred in Florida and Louisiana, began during this period (Maiolo 2004; Maiolo et al 1980).

Difficulties in organizing production and distribution capacity, along with the failure to expand markets into the interior of the state, resulted in inconsistencies in the demand for North Carolina shrimp prior to the outbreak of World War II. Additionally, poor ex-vessel prices hampered development of the state’s fishery. The War created a jolt in the popularity of the shrimp with consumers, because, unlike meat products, seafood was not rationed. There were still problems in the industry. The supply of seafood products, including shrimp decreased because of the war effort. There were fewer fishermen, boats, and equipment, until about 1944 when restrictions on strategic materials were eased. Also during this period trawling was restricted to inside waters because of the threat of German submarine attacks outside of the inlets. One result of this was increased effort in Pamlico Sound (Maiolo 2004).

Quick freezing technology was developed during the war years as well. Shrimp was no longer a perishable product, but a relatively stable commodity that the producer could control by freezing and holding for better prices when the market changed. However, this seems to have had a limited effect on North Carolina harvesting and distribution. Frozen shrimp from other regions had an impact on the markets, but most of North Carolina’s product was still shipped fresh to Northern markets (Maiolo 2004).

When the war concluded, and a recovering economy was redirected toward domestic matters, the fishing industry benefited along with the rest of the nation. There was a boom in construction of diesel-powered, large trawlers, and a considerable increase in shrimping effort. Prices increased dramatically, and North Carolina’s contribution to the Southeast shrimp landings became significant. Vessels were equipped with radar, fathometers, radios, steel cables and drum hoists (Maiolo 2004; Maiolo et al 1980).

Technological advances in the shrimping industry have increased the catching efficiency of larger boats, particularly in Pamlico Sound. In the 1940s and early 1950s, a 45 to 60 foot vessel pulled a single trawl with a headrope length of 60 to 65 feet. Now, with “four-barreled rigs” the same vessel can pull four nets with a combined headrope length of up to 200 feet. Four-barreled rigs allow fishermen to pull two nets from each outrigger. Conventional two-seam otter

DRAFT

trawls are used for the bottom-hugging pink and brown shrimp, while four-seam and tongue trawls with floats on the headrope are used for the white shrimp which have the ability to jump over two-seam trawls when disturbed. In Pamlico Sound, these large vessels stay out four or five days and tow from one to three hours, often working day and night. Smaller vessels make daily trips and employ shorter tow times. In the Core Sound area, the fishery occurs mainly at night, with trips lasting one night. In the southern area, fishing is conducted on a day-trip basis, mostly during daylight hours (Maiolo 2004).

Modern safety and navigation equipment have allowed North Carolina shrimpers to steam longer distances, for longer periods of time to shrimp; and also to engage in a constantly changing variety of harvesting activities other than shrimping throughout the calendar year. This widely recognized diversity of fishing activity occurs all along the Atlantic coastline and in the Gulf of Mexico. It is a continuation of adaptive strategies to changing resource opportunities and regulations as well as technology that dates back before the shrimp industry was born. In this respect, the history and development of harvesting activity in the shrimp industry may be seen as one more addition to the annual cycle of North Carolina's commercial (and to some extent, recreational) fishermen (Maiolo 2004; Orbach and Johnson 1988).

6.1.2 State Landings and Effort

Five different data sources are analyzed to describe the trends in the commercial shrimp industry. The first data source covers the years from 1962 to 1971 and was collected by NMFS. This older data set contains annual summarized landings of shrimp by county of landing, gear type used and water body harvested. The second data source covers the years from 1972 to 1977 and was also collected by NMFS. It also contains summarized landings of shrimp by the same categories as the older data set but it also contains the month of landing. The third data set analyzed covers the years from 1978 to 1993 and was collected under a cooperative statistics program between DMF and NMFS. This file is more detailed than the previous data sets as it contains summarized landings by county of landing, gear type used, water body harvested, month harvested and dealer landed. Another cooperative program between the DMF and NMFS was also started in 1978 that was designed to capture the number of trips and vessels in the shrimp fishery (commonly called the Detailed Shrimp Program). The Detailed Shrimp Program covers the years from 1978 to 1992. The last data source is the NC Trip Ticket Program and covers the years of 1994 to 2010. The data collected in the NC Trip Ticket Program is the most detailed and the most reliable of all the data collection programs as it contains the actual trip level commercial catch for all commercial landings in the state. However, from 1994 to 1998 the species composition of shrimp was not recorded in the NC Trip Ticket Program so analysis of species composition will be from 1978 to 1993 and 1999 to 2010.

DRAFT




Figure 6.1 Annual shrimp landings (lb) for North Carolina: 1962 – 2010.

Landings in the North Carolina shrimp fishery vary from year to year and are dependent primarily on environmental conditions (Figure 6.1 and Table 6.1). The annual average was 6,460,849 lb for the period 1962-2010; 75% were harvested from inshore waters and 25% from the Atlantic Ocean. This management plan will concentrate on landings from 1978 to 1993 and from the most recent 17 year period, 1994-2010, to evaluate trends in the fishery. The information from the earlier period will be used to illustrate historical trends while that from the later period will demonstrate changes in the fishery, especially species composition. Total landings from 1994 to 2010 have averaged 6,875,737 lb per year (range 2.4-10.3 Mlb) caught on an average of 14,256 annual trips (range 7,770-23,891 trips). The contribution to the landings continues to be 75% for inshore waters and 25% for the Atlantic Ocean (Figure 6.2).




Figure 6.2 Landings (lb) and trips for 1978-2010.

DRAFT

Annual effort for commercial shrimp harvest in North Carolina waterbodies is shown in Table 6.1 from 1978 to 2010 (Detailed Shrimp Program and TT Data). About 70-93% of the shrimp trips occur in estuarine waters, with the remainder in ocean waters, primarily within state territorial seas (<3 mi offshore) off the central and southern coast of North Carolina. Total annual shrimping effort has decreased from a high of 40,000 trips in 1982 to a low of 6,500 trips in 2005 while in 2010 7,800 trips were recorded. Annual shrimping effort has fluctuated with shrimp abundance, but it appears to have gradually declined since 1994 (Figure 6.2). This is due to a number of things including cheaper imported shrimp prices, increasing fuel prices, increased regulations, and fishermen retiring out of the industry.

Regionally, shrimping effort has generally been greatest in Core and Bogue sounds and associated estuaries [1,692-22,998 trips/year (Table 6.1)]. The Southern estuaries account for the second largest number of inside trips per year, ranging from 599 to 7,999 trips/year. In ocean waters, shrimping is highly concentrated in the southern portion of the state [Onslow through Brunswick counties (716-3,645 trips/year)], primarily in the summer (Table 6.1). In contrast, the annual effort in the central district (Carteret County) has ranged from 120 to 1,871 trips per year, and in the northern district (Virginia line through Hyde County) has ranged from 0 to 52 trips per year. Commercial shrimping effort has remained relatively stable over time in the southern ocean waters of the state.

Table 6.1 Annual number of trips reported for shrimp in inside and ocean waters[†], 1978-2010 (Detailed Shrimp Program and Trip Ticket Program).

Year	River and Sounds				Ocean Waters (<3 miles)				Percent Inside	Percent Ocean
	Albemarle	Core/Bogue	Pamlico	Southern	Northern	Central	Southern	Total		
1978	0	8,393	3,015	599	0	571	1,593	14,171	84.73	15.27
1979	0	9,031	2,391	3,665	9	777	2,120	17,993	83.85	16.15
1980	0	17,235	6,924	7,803	13	692	2,568	35,235	90.71	9.29
1981	0	15,854	3,654	4,195	14	383	1,450	25,550	92.77	7.23
1982	37	22,998	5,441	7,943	0	1,128	2,505	40,051	90.93	9.07
1983	0	22,274	4,912	7,999	5	1,236	2,890	39,317	89.49	10.51
1984	0	15,330	3,042	7,873	37	1,197	3,254	30,733	85.40	14.60
1985	0	12,207	8,075	3,911	22	1,246	1,056	26,517	91.24	8.76
1986	0	15,151	5,170	3,648	25	1,197	1,224	26,415	90.74	9.26
1987	0	13,348	2,574	2,797	0	1,322	1,406	21,447	87.28	12.72
1988	0	15,162	4,347	4,216	7	1,677	2,314	27,723	85.58	14.42
1989	0	18,403	4,997	4,901	0	1,871	2,402	32,574	86.88	13.12
1990	0	7,784	4,160	6,302	1	855	1,925	21,027	86.77	13.23
1991	0	12,497	5,277	6,859	0	591	2,266	27,490	89.61	10.39
1992	0	5,042	2,278	2,207	0	145	716	10,388	91.71	8.29
1993										
1994	0	9,494	4,603	3,893	3	332	3,439	21,764	82.66	17.34
1995	0	9,965	5,091	4,814	52	505	3,465	23,892	83.17	16.83
1996	1	7,615	2,817	3,412	19	420	2,802	17,086	81.03	18.97
1997	0	8,189	4,515	4,530	27	319	2,864	20,444	84.30	15.70
1998	0	6,006	1,750	3,630	7	550	3,026	14,969	76.06	23.94

DRAFT

Table 6.1 (continued). Annual number of trips reported for shrimp in inside and ocean waters[†], 1978-2010 (Detailed Shrimp Program and Trip Ticket Program).

Year	River and Sounds				Ocean Waters (<3 miles)				Percent Inside	Percent Ocean
	Albemarle	Core/Bogue	Pamlico	Southern	Northern	Central	Southern	Total		
1999	0	6,933	3,959	4,738	21	525	3,645	19,821	78.86	21.14
2000	4	5,490	5,385	4,409	16	342	2,795	18,441	82.90	17.10
2001	7	5,110	3,123	3,095	10	165	2,562	14,072	80.55	19.45
2002	1	6,579	4,837	4,276	7	231	2,411	18,342	85.56	14.44
2003	0	5,804	1,721	3,537	2	430	2,563	14,057	78.69	21.31
2004	0	3,835	2,746	2,377	7	367	2,550	11,882	75.39	24.61
2005	0	2,555	853	1,565	2	208	1,398	6,581	75.57	24.43
2006	0	2,386	1,887	1,330	1	334	2,083	8,021	69.85	30.15
2007	0	2,338	3,129	1,569	12	418	1,824	9,290	75.74	24.26
2008	0	1,993	2,841	1,471	33	231	1,513	8,082	78.01	21.99
2009	1	2,064	2,251	1,616	12	186	1,640	7,770	76.34	23.66
2010	0	1,692	2,105	2,440	13	120	1,491	7,861	79.34	20.66
Avg	2	9,336	3,746	3,988	12	643	2,243	19,969	83.49	16.51

† Albemarle Area: Albemarle Sound, Currituck sound, and all tributaries of Albemarle Sound.

Pamlico Area: Pamlico, Croatan, and Roanoke sounds; Pamlico, Bay, Neuse, and Pungo rivers.

Core/Bogue Area: Core and Bogue sounds; Newport, White Oak, and North rivers.

Southern Area: Masonboro, Stump, and Topsail sounds; Cape Fear, New, Shallotte, and Lockwood Folly rivers; IWW.

Northern district ocean waters: Virginia line through Hyde County.

Central district ocean waters: Carteret County.

Southern district ocean waters: Onslow County to the South Carolina line.

A trip may consist of multiple days in Pamlico Sound and the Atlantic Ocean.

6.1.3 Landings by Waterbody

An examination of harvest by waterbody for the most recent twelve year period shows that 56% of the landings are from Pamlico Sound, 24% from the Atlantic Ocean and 6% from Core Sound (Table 6.2). No other water bodies contribute more than 4% to the state's total landings. The totals for some water bodies have been combined for purposes of this discussion. For example, some of the water bodies in the southern part of the state where shrimp trawling is not allowed have been combined into the Inland Waterway; the shrimping activity took place in the Waterway that runs through the waterbody where the landings were recorded. It must also be taken into consideration that species composition was not noted on trip tickets for the years 1994 – 1998.

DRAFT

Table 6.2 Percent contribution of landings (1999 – 2010) by waterbody and species.

Waterbody	Percent brown	Percent pink	Percent white	Percent unclassified	Percent total
Pamlico Sound	70.84	29.22	35.18	60.51	55.95
Ocean	14.85	13.29	39.66	15.82	23.76
Core Sound	5.50	45.47	2.48	11.57	6.19
New River	1.12	4.46	5.55	1.91	2.87
Newport River	1.21	0.48	4.51	1.60	2.39
Neuse River	2.82	2.52	1.01	3.58	2.20
North River-Carteret	0.86	2.91	3.88	0.37	1.99
Other	0.79	0.52	1.28	1.25	0.98
Cape Fear River	0.66	0.46	1.39	0.91	0.93
Inland Waterway	0.04	0.02	2.26	1.12	0.90
White Oak River	0.09	0.03	1.75	0.10	0.68
Bogue Sound	0.40	0.61	0.98	0.14	0.60
Pamlico River	0.60	0.02	0.05	0.62	0.38
Bay River	0.16	0.01	0.01	0.50	0.12
Pungo River	0.06	0.00	0.00	0.00	0.04

6.1.4 Landings by Gear

The vast majority of the shrimp harvest (92%) is taken by otter trawls however, there has been a slight shift in the types of gear used to harvest shrimp in North Carolina in recent years (Figure 6.3). A type of trawl that has gained wide popularity in the central and southern areas since about 1991 is the skimmer trawl. This gear originated in the Gulf Coast states and is very effective at capturing white shrimp. Skimmers are modified wing nets sewn to an aluminum or steel pipe frame. The bottom of each outside pipe has a skid that rides over the bottom. The vessel can work in depths from two to fifteen feet and the tailbags can be hauled in more often without stopping to haul back. This increases the efficiency of the harvest and allows the bycatch to be released more frequently, thus reducing mortality. An increasing number of vessels in Carteret, Onslow, and Pender counties are switching from otter trawls to skimmers as their efficiency on brown shrimp harvest is improved. Skimmer nets account for 3% of the average annual state landings.

Channel nets are stationary nets that fish the surface and middle depths on an outgoing tide. They resemble a trawl anchored and staked to the bottom to keep it open. The nets are set at night on an ebb tide across a channel or slough in the path of seaward-migrating shrimp. The mouth of the net is oriented toward the direction of the oncoming current. The tailbag of the channel net is emptied into a skiff every 15 to 30 minutes. The net is retrieved from the water before the tide changes to prevent it from being turned inside out. The channel net must be set near inlets where the current is strong and where shrimp have concentrated to move out to sea. This activity is concentrated in estuarine waters from Beaufort Inlet to Rich's Inlet. Channel nets account for 5% of the average annual shrimp landings. Although not a significant contributor to shrimp landings, shrimp pound nets have recently been developed and employed in the taking of primarily brown shrimp. Shrimp pound nets are trap nets with a V-shaped lead that directs a shrimp to a funnel connected to a box-shaped pound. One of the leads extends to the shoreline

DRAFT

and the other extends out towards a channel or deeper water. Shrimp enter the nets at night as they migrate. The larger shrimp are trapped in the pound while the smaller ones are allowed to pass through. Interest in the use of shrimp pounds has increased since 2003 and issues raised were addressed in the 2006 FMP.

The cast net is another type of gear used to harvest shrimp. A few pink and brown shrimp are captured around the marshes and shallows during the summer with this circular net weighted around the perimeter that is thrown out over the shrimp. The weighted edges of the cast net sink to the bottom entrapping the shrimp, and they are pulled into the catcher by a line attached to the top of the net. The cast net is most successful on white shrimp in the fall as they school in large concentrations and leave the creeks and tributaries and head for the sounds and, eventually, the ocean. Throwing from boats or bridges over creeks is productive when they are migrating.




Figure 6.3 Inshore shrimp landings by gear for North Carolina: 1962 – 2010.

6.1.5 Landings by Species

The North Carolina shrimp fishery harvests three species: brown, pink, and white. Data on the species composition of the shrimp catch were collected from 1978 to 1982 through the Detailed Shrimp Program, from 1983 to 1993 through the DMF/NMFS Cooperative Statistics Program and from 1999 to 2010 through the DMF Trip Ticket program. Species composition was not collected from 1994 through 1998 so discussion of the contribution of each species to the total landings will concentrate on the time periods 1978-1993 and 1999-2010. Historically (1978-1993) brown shrimp accounted for 66% of the state total, averaged 4.5 Mlb and annual totals ranged from 1.1 Mlb in 1987 to 10.4 Mlb in 1985 (Figure 6.4). North Carolina brown shrimp commercial landings have averaged 3.8 Mlb since 1999 (Figure 6.5). During this time, landings have fluctuated from a high of 6.5 Mlb in 2000 to a low of 1.5 Mlb in 2005. Environmental factors, principally temperature and salinity, have a major influence on the yearly harvest. Generally, 85% of all brown shrimp landed are caught in estuarine waters with Pamlico Sound, Core Sound, New River, and Neuse River accounting for most of the harvest (Table 6.2). Since 1999, over 96% of all brown shrimp landed are caught by shrimp trawls. Channel nets and skimmer trawls account for the remaining landings.

Pink shrimp have historically (1978-1993) accounted for about 27% of the shrimp landings. North Carolina commercial pink shrimp landings averaged 1.8 Mlb from 1978 to 1993 (Figure 6.4). Environmental factors especially severity of winter temperatures, have a significant influence on the yearly harvest. However, since 1999, pink shrimp landings have averaged only 0.2 Mlb, despite a series of mild winters in the late 1990s and early 2000s. Pink shrimp have accounted for 4% of the state's harvest during the last 12 years (Figure 6.5). The cause of this decrease is not known. The majority of pink shrimp landed are caught in estuarine waters (87%). There are two seasonally distinct fisheries, one from late April through June, and the fall fishery that runs from September through November. Core Sound accounts for 45% of the landings, followed by Pamlico Sound (29%), and the ocean (13%) (Table 6.2). Since 1999, over 87% of all pink shrimp landed are caught by shrimp trawls. Channel nets (11%) and skimmer trawls (2%) account for the remainder.




Figure 6.4 North Carolina landings of shrimp by species 1978-1993.

During the period 1978-1993, North Carolina commercial white shrimp landings averaged 0.5 Mlb (Figure 6.4). Landings fluctuated from a high of 1.7 Mlb in 1993 to a low of 11,000 pounds in 1981 (Figure 6.4). The landings increased significantly for the most recent 12 years to an average of 2.5 Mlb that was 36% of the state landings (Figure 6.5). These fluctuations are not unusual for a species so vulnerable to environmental conditions, especially low winter water temperatures. The percentage of the white shrimp catch taken in the ocean is higher (40%) than the other two species, which reflects its greater abundance in the southern part of the state where the majority of the ocean fishery occurs. Since 1999, over 82% of white shrimp landed were caught in shrimp trawls. The other 18% were captured in channel nets (3%) or skimmer trawls (15%). On average, during 1978-1990, 60% of all white shrimp were landed in the southern coastal area (Onslow, Pender, New Hanover and Brunswick counties) and 26% in the central area (Pamlico and Carteret counties), and the remaining 12% were taken in the northern area. Since 1999 the majority of white shrimp have been harvested from the Ocean (40%), Pamlico Sound (35%) and New River (6%); which reflects the effects of a series of mild winters that has allowed white shrimp populations to be abundant in the northern portion of the state (Table 6.4).

DRAFT




Figure 6.5 North Carolina landings of shrimp by species: 1994 – 2010.

There are two seasonal fisheries for white shrimp in North Carolina. The spring fishery lasts from late April until June and the fall fishery that begins in late August and may last through December. In the spring fishery, trawlers primarily target "roe" (female) white shrimp. The majority of white shrimp landed come from the fall fishery, where it is the target species in the southern coastal area and other areas if they are abundant.

North Carolina's shrimp fishery is unusual in the southeast because all three species are taken here and the majority of the effort, about 83%, is expended in internal waters. While South Carolina, Georgia and Florida allow limited inside shrimping, the majority of their fisheries are conducted in the Atlantic Ocean and white shrimp comprise the most of their harvest (Table 6.3). North Carolina's landings for the period 1999-2010 were 34% of the total for the South Atlantic followed by Florida (25%), Georgia (23%) and South Carolina (19%).

Table 6.3 Shrimp landings in pounds from the South Atlantic, 1999-2010.

Area		Brown Shrimp	Pink Shrimp	White Shrimp	Grand Total
Florida East Coast	1999	1,014,167	1,035,004	3,947,723	5,996,894
	2000	639,573	905,375	2,455,165	4,000,113
	2001	1,221,035	482,736	2,386,053	4,089,824
	2002	1,008,681	615,462	3,316,938	4,941,082
	2003	884,123	209,988	2,098,503	3,192,614
	2004	1,037,684	456,313	3,813,020	5,307,017
	2005	393,985	408,183	3,801,199	4,603,367
	2006	606,855	642,531	3,964,873	5,214,258
	2007	1,284,146	210,949	3,633,139	5,128,234
	2008	641,537	379,926	3,952,565	4,974,028
	2009	701,369	256,923	3,264,738	4,223,029
	2010	1,093,991	777,657	4,383,569	6,255,217
Florida Total		10,527,146	6,381,046	41,017,485	57,925,677

DRAFT

Table 6.3 (continued).

Area		Brown Shrimp	Pink Shrimp	White Shrimp	Grand Total
Georgia	1999	1,352,545	0	5,340,885	6,693,430
	2000	772,932	0	4,599,183	5,372,115
	2001	1,471,975	0	2,789,070	4,261,045
	2002	683,818	0	4,246,202	4,930,020
	2003	1,407,018	0	4,015,844	5,422,862
	2004	568,241	0	4,410,584	4,978,825
	2005	1,421,386	0	3,013,279	4,434,665
	2006	207,816	0	3,467,257	3,675,073
	2007	510,169	0	2,211,691	2,721,860
	2008	378,332	0	2,642,896	3,021,228
	2009	326,382	0	2,594,351	2,920,733
2010	599,068	0	3,869,213	4,468,281	
Georgia Total		9,699,682	0	43,200,455	52,900,137
North Carolina	1999	1,672,959	10,060	3,659,302	5,342,321
	2000	6,489,495	161,422	3,214,862	9,865,779
	2001	3,923,540	211,858	863,153	4,998,551
	2002	6,029,219	879,894	2,514,342	9,423,455
	2003	4,828,513	219,010	1,100,128	6,147,651
	2004	2,749,009	143,954	1,923,460	4,816,423
	2005	1,523,028	43,489	780,169	2,346,686
	2006	1,944,380	65,232	3,682,529	5,692,141
	2007	3,110,266	84,168	6,339,883	9,534,317
	2008	5,502,793	830,488	3,076,444	9,409,725
	2009	3,804,694	250,213	1,347,561	5,402,468
2010	4,233,181	52,657	1,658,681	5,944,519	
North Carolina Total		45,811,078	2,952,446	30,160,513	78,924,037
South Carolina	1999	1,253,824	9,836	3,858,202	5,121,862
	2000	887,302	28,431	3,001,515	3,917,248
	2001	1,445,911	1,111	1,360,590	2,807,612
	2002	919,621	508	2,423,729	3,343,858
	2003	1,469,998	66	2,449,051	3,919,115
	2004	1,139,895	0	4,485,856	5,625,751
	2005	1,213,979	0	2,742,780	3,956,759
	2006	368,326	0	3,319,573	3,687,899
	2007	845,687	0	1,970,594	2,816,281
	2008	688,416	0	2,478,418	3,166,834
	2009	375,719	0	2,343,203	2,718,922
2010	968,916	0	2,988,253	3,957,169	
South Carolina Total		11,577,594	39,952	33,421,764	45,039,310
Grand Total		77,615,500	9,373,444	147,800,217	234,789,161

DRAFT

6.1.2.5 Regional Summary

The shrimp fishery in the northern portion of the state is conducted in Pamlico, Croatan, and Roanoke sounds and Pamlico, Pungo, Bay and Neuse rivers. The otter trawl is the predominant gear used in this portion of the state. Commercial activity occurs in all waters, while recreational activity usually occurs in the rivers and nearshore areas of the sounds.

The shrimp fishery in the central coastal area of the state occurs in Neuse River, Core Sound, North River, Newport River, Bogue Sound, and White Oak River. A variety of methods are used to catch shrimp including trawls, skimmers, channel nets, shrimp pounds, and cast nets. Trawls are used on all three species in both the estuary and the ocean with two seam trawls used for brown and pink shrimp and four seam and tongue trawls for white shrimp, which tend to swim higher in the water column and have the ability to jump to the surface when disturbed. Most trawling in the central portion of the state is conducted at night. Channel nets are popular around Harkers Island in the Straits and North River while skimmer trawling is very popular in Newport River.

In the southern portion of the state, the fishery is characterized by a large number of small boats fishing internal waters (primarily the Intracoastal Waterway, New and Cape Fear rivers) and larger craft fishing the Atlantic Ocean primarily off New River, Carolina Beach, and Brunswick County. Many of the small boats are fished by individuals who shrimp part-time or for personal consumption. Use of gears other than trawls has increased primarily in the area from New River to Rich's Inlet. Channel, float, and butterfly nets make use of tidal currents to push shrimp into the nets and offer the advantages of less fuel consumption and less bycatch than traditional shrimp trawls. Channel nets are fished extensively in the areas around New River and Topsail inlets. To shrimp with a "float net", fishermen attach large floats to the doors and top lines of trawls to make the net fish up in the water column and are pulled slowly forward to harvest shrimp that are migrating to the inlets at night. Butterfly nets use this same harvest strategy but are attached to a metal frame and are held stationary in the water column to capture shrimp as the current carries them into the net. Skimmer trawls have become more popular around New River and Topsail Sound. These alternative gears are employed very little in areas south of Rich's Inlet, however tidal conditions seem favorable for their use. Cast nets and seines are also used to harvest shrimp primarily for recreational uses, personal consumption, and to provide live shrimp for the commercial bait fishery.

6.2 RECREATIONAL FISHERY

Shrimp are harvested recreationally throughout the state by otter trawls, skimmer trawls, seines, cast nets, shrimp pots and shrimp pounds. As of July 1, 1999, anyone wishing to harvest shrimp recreationally with commercial gear is required to purchase a RCGL. RCGL holders are restricted to using otter and skimmer trawls with a headrope length of up to 26 feet, a 100 foot seine, five shrimp pots and one shrimp pound. Seines measuring less than 30 feet long and cast nets are exempt from this license. Cast nets are the only gear allowed in areas closed to other commercial methods of shrimping with a limit of 100 shrimp per person. This limit will change to two quarts beginning June 2013 due to concerns of law enforcement officer safety while enforcing this rule.

DRAFT

6.2.1 Recreational Commercial Gear License (RCGL) History

On August 14, 1997, the Fisheries Reform Act (FRA) was signed into law. One aspect of this law was the creation of the RCGL. According to the Fisheries Moratorium Steering Committee (MSC), a group that provided the recommendations for the FRA, the purpose of creating this license was to: (1) allow individuals and families who have traditionally accessed the State's public trust fishery with commercial gear to supply themselves with fresh seafood; (2) limit the effort that may be expended by this class of fishermen both individually and as a group; and (3) implement the principle that all persons who harvest state public trust resources pay for that privilege by investing in coastal fisheries conservation and management (Moratorium Steering Committee, 1996). DMF began selling this license July 1, 1999.

The MSC also recommended that the MFC be authorized to establish specific gear limits with "standing advisory committees" and those limits could vary by region. The MFC should be required to re-examine and revise the gear limitations on a recurring basis. The MSC further recommended that the RCGL be restricted to the use of the following gears and amounts during the period final gear limitation rules are being developed by the MFC: one – 100 yards of gill net; 2) five crab/fish pots and 3) a single trawl with a headrope less than or equal to 26 feet. These limits were meant to serve as the starting point for the MFC rule development on RCGL gear and were the result of extensive public input and deliberation by the MSC.

The FRA provided that the MFC: 1) shall adopt rules authorizing the use of a limited amount of commercial fishing equipment or gear for recreational fishing under a RCGL (G.S.113-173(c)); 2) may authorize the limited use of commercial gear on a uniform basis in all coastal fishing waters or may vary the limited use of commercial gear within specified areas of the coastal fishing waters; and 3) shall periodically evaluate and revise the authorized use of commercial gear for recreational fishing.

RCGL allow recreational fishermen to use limited amounts of commercial gear to harvest seafood for their personal consumption. Seafood harvested under this license cannot be sold. RCGL holders are limited to the same bag and size limits as CRFL holders. The 2006 Shrimp FMP added two new allowable RCGL gears, one shrimp pound and a 26 foot skimmer trawl. The FMP also limited all recreational harvesters, including RCGL holders to 48 quarts of head-on (32 quarts of head-off) shrimp per day, greatly reducing the harvest in some areas. If there are two valid license holders on board a vessel, then the shrimp possession limit may be doubled. The MFC also passed a rule allowing mechanical retrieval gear as long as a TED was properly installed in the trawl; prior to the FMP shrimp trawls could only be retrieved by hand.

6.2.2 RCGL Survey

Many of the species taken by recreational users of commercial gear are included in fisheries management plans. Until 2002, the influence that RCGL holders may have on these species was unknown. Two survey strategies were used to collect information from RCGL holders; a socioeconomic survey, conducted in 2001, 2004, and 2007, and catch and effort surveys conducted monthly from 2002 through 2008. Both of these surveys were terminated in 2008 due to budget constraints. While the harvest of RCGL holders has been shown to be minimal, the lack of current data could foster further debate over the impact of the use of commercial gear by recreational fishermen. Findings from these surveys are summarized by regions, using the DMF Fisheries Management District boundaries (Figure 6.6).




Figure 6.6 Regions used to summarize findings from the RCGL surveys.

6.2.3 RCGL Survey Methodology

Catch and effort survey questionnaires were designed to determine the number of trips taken and type and quantities of gear used during the month of the survey. Participants were also requested to provide estimates for the numbers and pounds of each species caught and retained as well as the number of each species discarded.

Participants for the survey were randomly selected using two different rates of sampling. A 30.0% coverage rate by county of residence for the period May through December was used. This is the period when the bulk of RCGL holders are actively fishing and is sufficient for the gears used and majority of the species targeted.

To estimate the total number of trips taken by all RCGL holders, the monthly survey data were extrapolated for each monthly sample period and gear combination by:

- Calculating the level of participation by dividing the total number of participants actively using a specific gear by the total number of returned questionnaires,
- Calculating the mean number of trips taken by the participants indicating actively using a specific gear, and
- The effort estimate was the product of the mean number of trips, level of participation, and the total number of RCGL holders for the given sample period.

Determinations of the estimated catch for each species were also calculated for each sample period and gear level by:

DRAFT

- Summing the total catch by species, sample period, and gear combination,
- Summing the total number of trips taken by sample period and gear combination,
- Dividing total catch by the total number of trips to determine the mean catch for each species for every sample period and gear combination, and
- The catch estimate was the product of the mean catch and the estimated effort.

Participants were also asked to specify the average amount of gear used. Quantities were categorized into ranges of values for head rope length of trawls, and length of seine. RCGL holder use of shrimp pots, shrimp pounds, and seines is negligible and only information gathered from RCGL holders that use otter trawls is presented.

6.2.4 RCGL Survey Results

With the exception of 2002, the number of RCGLs sold on a fiscal basis has declined each year from 2002 through 2010 (Table 6.4); with 24% overall decline from the first to last year in this period. The largest single year decline occurred in 2001 (8%) followed by 2006 (5%). In 2009 and 2010 there was an average of 3.1% increase in sales. This increase is probably due to the downward trend of the economy, thus, increasing the need of seafood for personal consumption. Twenty-five counties consistently comprise approximately 85% of the total number of RCGLs purchased each year. Southern counties such as Onslow, Pender, New Hanover, and Brunswick consistently rank in the top ten counties each year.

Table 6.4 Number of fiscal license sales of Recreational Commercial Gear Licenses, 2002 though 2010 (fiscal year, July 1 through June 30).

Fiscal Year	Number of RCGLs Sold	Percent Change from Previous Sales Year
2000	6,740	
2001	6,202	-8.0%
2002	6,300	1.6%
2003	6,157	-2.3%
2004	5,868	-4.7%
2005	5,653	-3.7%
2006	5,368	-5.0%
2007	5,134	-4.4%
2008	5,113	-0.4%
2009	5,268	2.9%
2010	5,451	3.3%

Typical RCGL holders were married Caucasian males with an average age of 56. Findings from license sales statistics and the three socioeconomic surveys conducted in 2001, 2004, and 2007 indicated that coastal counties, in particular, southern coastal counties, substantially contributed to the overall number of RCGL holders (Figure 6.7).




Figure 6.7 Distribution of RCGL residents.

The top three gears utilized by RCGL holders fishing in all regions were crab pot, small mesh gill net, and large mesh gill net. Shrimp trawls were the fourth most common gear utilized in the Pamlico, Southern, and Central Regions while fish pots were the fourth most common gear utilized in the Northern Region. On average the highest number of trips using shrimp trawls from 2002 to 2008 occurred in the Pamlico region, followed by the southern region, the central region, and the northern region (Table 6.5). In the Pamlico region, the number of trips ranged from 1,127 (2005) to 2,384 (2002), averaging 1,642 per year from 2002 to 2008. In the southern region, the number of trips ranged from 355 (2007) to 1,123 (2002), averaging 586 trips per year. An average of 413 trips a year were made in the central region, ranging from 132 (2008) to 1,070 (2002). In the Northern region, the number of trips ranged from 50 (2006) to 911 (2004). Overall, the highest number of trips made by RCGL using shrimp trawls was observed in 2002; the lowest was observed in 2007.

RCGL holders harvested an average of 52,352 pound of shrimp a year from 2002 to 2008 (Figure 6.6). The highest landings occurred in 2002 (101,766 lb), followed by 2008 (54,359 lb) and 2003 (50,961 lb). RCGL holders harvested an average of 16.8 pounds of shrimp per trip from 2002 to 2008 (Figure 6.6). The highest pounds of shrimp per trip was observed in 2009 (22.3 lb/trip), followed by 2006 (20.3 lb/trip) and 2002 (19.1 lb/trip).

DRAFT

Table 6.5 Number of trips by shrimp trawl by region, 2002 through 2008.

Year	Region				Total
	Southern	Central	Pamlico	Northern	
2002	1,123	1,070	2,384	742	5,319
2003	711	246	1,448	348	2,753
2004	392	318	2,122	911	3,743
2005	553	365	1,127	387	2,432
2006	471	464	1,441	50	2,426
2007	355	295	1,510	69	2,229
2008	500	132	1,464	337	2,433
Mean	586	413	1,642	406	3,048

Table 6.6 Harvest (lb) and pounds per trip of shrimp by RCGL gear from 2002 through 2008.

Year	Pounds	Pounds/trip
2002	101,766	19.1
2003	50,961	18.5
2004	43,698	9.3
2005	32,542	13.4
2006	49,362	20.3
2007	33,778	15.2
2008	54,359	22.3
Mean	52,352	16.8

6.2.5 Regional RCGL Characterization for Shrimp Trawls

Southern Region

The top species harvested by RCGL shrimp trawls in the Southern Region from 2002 to 2008 were shrimp, blue crab, flounder, and spot (Table 6.7). On average, shrimp made up 88.0% of the harvest, blue crab 5.2%, flounder 2.9% and spot 2.8%. Shrimp harvests ranged from 2,400 pounds (2007) to 25,642 pounds (2002), averaging 11,900 pounds annually. Overall, 22.7% of the total RCGL harvest was landed by shrimp trawls in the Southern Region.

Table 6.7 Top four species harvested (lb) by RCGL shrimp trawls in the Southern Region, 2002-2008.

Species	2002	2003	2004	2005	2006	2007	2008	Average
Shrimp	25,642	14,897	5,810	9,585	20,041	2,400	4,928	11,900
Blue crab	1,271	1,363	826	640	221	339	268	704
Flounder	603	383	365	1,151	121	15	143	397
Spot	0	29	667	65	789	256	839	378
Other*	58	9	80	15	704	0	36	129
Total	27,574	16,681	7,748	11,456	21,876	3,010	6,214	13,508

DRAFT

* Other includes: Atlantic menhaden, croaker, pigfish, pinfish, sharks and rays, sheepshead, shellfish (misc.), Spanish mackerel, weakfish

Central Region

The top five species harvested by shrimp trawls in the Central Region from 2002 to 2008 were shrimp, blue crab, flounder, croaker, and pigfish (Table 6.8). On average, shrimp made up 95.5% of the harvest, blue crab 3.5%, flounder 0.6%, croaker 0.3% and pigfish <0.1%. Shrimp harvests ranged from 2,175 pounds (2008) to 19,095 pounds (2002), averaging 7,501 pounds annually. Overall, 14.3% of the total RCGL harvest was landed by shrimp trawls in the Central Region.

Table 6.8 Top species harvested (lb) by RCGL shrimp trawls in the Central Region, 2002-2008.

Species	2002	2003	2004	2005	2006	2007	2008	Average
Shrimp	19,095	4,100	6,966	7,213	9,280	3,677	2,175	7,501
Blue crab	927	189	0	581	200	7	15	274
Flounder	246	41	0	0	51	14	0	50
Croaker	0	0	0	78	61	0	0	20
Pigfish	0	0	0	0	25	21	0	7
Total	20,268	4,330	6,966	7,888	9,617	3,719	2,190	7,852

Pamlico Region

The top five species harvested by shrimp trawl in the Pamlico Region were shrimp, blue crab, flounder, spot, and croaker (Table 6.9). On average, shrimp made up 82.1% of the harvest, blue crab 16.8%, flounder 0.2%, spot 0.2%, croaker <0.1%. Shrimp harvests ranged from 10,764 pounds (2005) to 48,982 pounds (2002), averaging 27,739 pounds annually. Overall, 53.0% of the total RCGL harvest was landed by shrimp trawls in the Pamlico Region; the highest among the four regions.

Table 6.9 Top species harvested (lb) by RCGL shrimp trawls in the Pamlico Region, 2002-2008.

Species	2002	2003	2004	2005	2006	2007	2008	Average
Shrimp	48,983	24,622	19,963	10,764	19,536	26,574	43,734	27,739
Blue crab	11,815	6,792	10,808	4,195	3,268	800	1,844	5,646
Flounder	283	17	18	0	110	68	0	71
Spot	0	0	48	0	137	170	0	51
Croaker	0	0	20	0	0	136	0	22
Other*	0	0	36	0	0	67	21	18
Total	61,081	31,431	30,893	14,959	23,051	27,815	45,599	33,547

* Other includes: Atlantic menhaden, pigfish, pinfish, sharks and rays, sheepshead, shellfish (misc.), Spanish mackerel, weakfish

DRAFT

Northern Region

The top species harvested by shrimp trawl in the Northern Region were shrimp, blue crab, croaker, flounder, and catfish (Table 6.10). On average, shrimp made up 72.8% of the harvest, blue crab 12.1%, croaker 7.9%, flounder 6.4% and catfish 0.7%. Shrimp harvests ranged from 57 pounds (2007) to 9,374 pounds (2004), averaging 3,914 pounds annually. Overall, 7.5% of the total RCGL harvest was landed by shrimp trawls in the Northern Region.

Table 6.10 Top species harvested (lb) by RCGL shrimp trawl in the Northern Region, 2002-2008.

Species	2002	2003	2004	2005	2006	2007	2008	Average
Shrimp	7,875	5,172	9,374	1,952	118	57	2,852	3,914
Blue crab	1,404	1,112	488	1,227	2	251	66	650
Croaker	0	78	2,815	65	0	0	0	423
Flounder	433	134	1,500	41	0	316	0	346
Catfish	0	0	276	0	0	0	0	39
Total	9,712	6,496	14,453	3,285	120	624	2,918	5,373

6.2.6 Contribution of RCGL Harvest compared to Other Fisheries

When compared to North Carolina's commercial harvest statistics from the NCTTP and recreational angling harvest estimates from the Marine Recreational Information Program (MRIP), the average yearly RCGL harvest has been shown to contribute only minimally to the overall harvest of those species encountered using RCGL gears (Table 6.11). From 2002 thru 2008 there was an average 52,352 pounds of shrimp that were landed by RCGL holders using shrimp trawls. In comparison, the total percent of RCGL shrimp landings account for 0.87% of the total commercial shrimp harvest.

The MRIP is a survey of marine and estuarine finfish species. The vast majority of interviews conducted each year are from angling trips; therefore species such as menhaden, striped mullet, and anadromous species are not encountered frequently enough to provide precise estimates.

Resource or conflict issues related to the RCGL since its implementation have been minimal. There have been instances, as with all gear, where the user was not acting responsibly. Reports to the DMF have ranged from shrimpers harvesting over the legal limit, improperly marked gear, and the illegal sale of RCGL harvested shrimp.

DRAFT

Table 6.11 Contribution in percent (pounds) of RCGL harvest to the overall harvest of finfish and shellfish based on the average yearly harvest from each sector during the period 2002 though 2008.

	Recreational Angling Harvest (lb) MRIP ¹	RCGL Harvest (lb) RCGL Surveys	Commercial Harvest (lb) NCTTP	Percent contribution from RCGL Harvest
Crustacean and Shellfish Species				
Shrimp		60,334	6,868,230	0.87
Blue Crab		116,797	31,392,856	0.37
All Shellfish		169,445	40,294,392	0.42
Finfish Species				
Bluefish	1,081,016	17,022	2,778,336	0.44
Catfish		6,864	405,198	1.67
Croaker, Atlantic	194,940	14,534	10,286,338	0.14
Drum, Black	313,684	6,101	189,932	1.2
Drum, Red	207,967	7,522	142,492	2.1
Flounder	535,996	65,059	6,086,025	0.97
Herring, River		10,873	132,193	7.6
Mackerel, Spanish	544,071	3,611	490,265	0.35
Menhaden, Atlantic		5,959	26,404,767	0.02
Mullet, Striped		41,197	1,788,300	2.25
Perch, White		15,531	272,052	5.4
Pigfish	51,777	1,263	36,327	1.41
Pinfish	121,754	268	43,224	0.16
Seatrout, Spotted	612,409	13,207	229,927	1.54
Shad, American		14,623	247,917	5.57
Shad, Hickory		12,053	91,260	11.67
Sheepshead	326,030	1,298	67,130	0.33
Spot	1,397,217	203,535	1,605,764	6.35
Striped bass	1,908,784	5,225	610,673	0.21
Weakfish	154,301	602	641,914	0.08
All finfish	21,656,437	453,065	62,021,830	0.54

6.3 SHRIMP TRAWL BYCATCH

Over the last two decades, bycatch has remained an important and controversial topic in fisheries management and marine conservation both in the United States and around the world (Alverson et al. 1994; Alverson and Hughes 1996; Crowder and Murawski 1998; Diamond 2003; Kelleher 2005; Davies et al. 2009). Interest in bycatch has shifted from its potential commercial use to concerns about impacts on finfish populations, biodiversity, and ecosystem trophic structure (Murray et al. 1992; Hall et al. 2000; Davies et al. 2009). In spite of increased public awareness, greater management scrutiny, and significant research efforts, many basic issues remain unresolved. Only recently has the term bycatch been defined in any standard manner, and important information on the magnitude of bycatch is severely lacking for many fisheries. Given this situation, it is not surprising that little is known of the impacts of bycatch on specific fisheries, fish populations, and marine communities. Although more information is needed to fully assess the effect of bycatch on fish populations and the ecosystem, continued concern and public policy dictates that bycatch be either eliminated or reduced to insignificant levels (Crowder and Murawski 1998). As perhaps the prime example of the new policy positions, the re-authorized Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA) contains a National Standard (#9) requiring bycatch minimization (USDOC 1996). National Standard 9 states: "Conservation and management measures shall, to the extent practicable, (A) minimize bycatch and (B) to the extent bycatch cannot be avoided, minimize the mortality of such bycatch." Additionally, in 1991 the MFC adopted a policy directing the DMF to establish the goal of reducing bycatch losses to the absolute minimum and to consciously incorporate that goal into all of its, management considerations (Murray et al. 1991).

Bycatch is defined by the Atlantic States Marine Fisheries Commission (ASMFC) as "the portion of a catch taken incidentally to the targeted catch because of non-selectivity of the fishing gear to either species or size differences" (ASMFC 1994). In the MSFCMA, bycatch is defined as "fish which are harvested in a fishery, but which are not sold or kept for personal use" (USDOC 1996). Bycatch can be divided into two components: incidental catch and discarded catch. Incidental catch refers to retained catch of non-targeted species. Discarded catch is that portion of the catch returned to the sea as a result of economic, legal, or personal considerations. Differences in market prices for a given size-class of species or limited storage space can also lead to "high grading", where less valuable species and size classes are discarded to make space for more valuable fish (Bellido et al. 2011). The biological significance of bycatch can be judged from a number of different perspectives, including those of the populations (e.g., of a particular species), of the fishery or fisheries that target or otherwise encounter the species, and of the general biological community (Murawski 1995).

During the late 1980s the DMF initiated gear testing to reduce bycatch in the shrimp trawl fishery (Pearce et al. 1988; Holland 1988). Due to growing concern over bycatch in shrimp trawl fisheries the MSFCMA was amended in 1990 to include bycatch research. Congress mandated that the US Secretary of Commerce conduct a three year research program to assess the impact of the incidental harvest by the shrimp trawl fishery on fishery resources in the South Atlantic and the Gulf of Mexico areas. The National Marine Fisheries Service (NMFS), along with the Gulf and South Atlantic Fisheries Development Foundation (GSAFDF), began a cooperative bycatch research program to: (1) update and expand bycatch estimates temporally and spatially; (2) identify, develop and evaluate gear options for reducing bycatch; (3) develop an information transfer and education program on bycatch; and (4) develop and operate a standardized data management system for centralized dissemination and access (NMFS 1995). Starting in 1992, observers were placed aboard cooperating vessels to characterize bycatch and to test BRDs during normal commercial shrimp trawling.

DRAFT

While it is becoming increasingly apparent to scientists, natural resource managers, and much of the general public that bycatch is an important issue that must be addressed, characterizing the nature and extent of bycatch has proven extremely difficult. These difficulties are generally attributed to inadequate monitoring of many pertinent characteristics, including actual bycatch levels, effort of the directed fishery, distribution of the bycatch species, and the mortality rate of the discarded species. The problem is exacerbated by the patchy distribution of effort and juvenile finfish in both time and space. The amount of bycatch in a particular trip is usually skewed, with many tows having some bycatch and fewer tows with high bycatch. Additionally, available effort data are often inadequate. Although research indicates that tow duration is often a significant factor when estimating bycatch losses, the DMF and most other agencies typically record effort data by trip without any accompanying information on tow duration or the number of tows made during a trip. Mortality of bycatch captured in trawls varies considerably, not only by species, but also in response to factors such as water temperature, tow time, fishing location, time of year, and gear configuration.

The lack of reliable discard estimates has not stopped researchers from investigating assessment impacts, but it has prevented increases in precision. Most assessments address the range of bycatch estimates through sensitivity analyses by comparing basic assessment results over the range of bycatch estimates and assumptions. If none of the results seem plausible, the assessment may proceed without the bycatch estimates included but with the caveat that results may be biased or contain additional uncertainties due to unknown levels of missing catch. However, the omission of discard data may result in an underestimation of fishing mortality and can lead to a biased assessment (Bellido et al. 2011).

6.3.1 History of Bycatch Management in North Carolina

Shrimp trawling in North Carolina began in the southern coastal area in the mid-1900s and by 1925 there were 300 fishermen participating in the shrimp trawl fishery. By the 1930s, trawling had spread into Core Sound and Pamlico Sound with the center of the industry in Carteret County. Concerns of bycatch began to be raised in the 1950s after experiencing serious declines in the catch of commercial fish in North Carolina waters with attention being focus on the shrimp fishery in Pamlico Sound.

In 1951 the ASMFC published a report on bycatch from trawling and its ultimate effect on abundance and weight of fish when they reached commercial size and what its impacts were on fishing mortality on top of natural mortality. Findings at that time were there was doubt of significant effects because finfish landings varied greatly and species such as spot, croaker and weakfish experience high natural mortality. ASMFC recommended additional studies on natural mortality and other causes of landing fluctuations.

During the 1960s and early 1970s, there was a primary concern of bycatch from directed ocean finfish trawling for bait and pet food. Rules were established by the MFC to prohibit directed scrap fishing. Nursery area designation also began during this time. The Albemarle Sound was closed to trawling in 1987 due to conflicts between crab potters and trawlers and in 1988 a subgroup of the Tar-Pamlico River Foundation called for the elimination of trawling because of bycatch and habitat concerns. By 1993, the MFC requested that the division prepare an issue paper on estuarine trawling. This paper was a comprehensive report on the knowledge of the issues at the time and addressed bycatch, overfishing, habitat and water quality concerns as well as conflict and competition. The division proposed a 50 ft headrope limit for a single trawl and 70 ft total headrope limit in all internal waters except Pamlico Sound. In Pamlico Sound, a

DRAFT

160 ft headrope limit with a two year phase in was proposed. Resulting rules in 1994 were the prohibition of trawling in the Outer Banks sea grass beds, the elimination of weekend trawling, and special secondary nursery areas could be opened by proclamation only from August 16 through May 14.

In 1997, a proposed net ban bill was introduced in the North Carolina General Assembly which proposed banning inside trawling and all net fishing with the exception of cast nets, dip nets and seines less than 12 ft long. It also proposed a buy-back program. However, a 1998 amendment to the 1997 FRA directed the Joint Legislative Commission on Seafood and Aquaculture (JLCSA) to study the biological, habitat, and socioeconomic impacts of the use of trawl nets in the sounds, river and estuaries.

In the meantime, the MFC Inland AC requested that the division examine estuarine trawling and determine the possibility of banning shrimp and crab trawling. Also, a 1999 petition was sent to the MFC and the General Assembly along with a letter requesting the management of estuarine trawling go through the FMP process. This all culminated into a 1999 division report on trawling and its effects on bycatch and habitat and again summarized the current knowledge of the time.

The JLCSA requested an analysis of research and information needs to address bycatch and habitat issues. Academia and division staff summarized current knowledge on trawling impacts and data needs necessary to make decisions. These included an accurate depiction of area and frequency of trawling by season and type of habitat involved, and the effect so bycatch on subsequent population yields of species. Other important data include rate of recovery of trawled bottom based on habitat type and trawl intensity and socioeconomic consequence of various alternative management options. An eight year study was presented to the JLCSA with budget needs of one to two million dollars a year. Although the JLCSA supported the concept, of the study, no funding was made available.

6.3.2 Incidental Catch

Total annual landings in the North Carolina shrimp trawl fishery have averaged 7.3 million pounds, ranging from 2.1 to 9.9 million pounds (DMF Trip Ticket data 1994-2010; Table 6.12). Shrimp (brown, pink, and white) account for 93% of the total landings followed by finfish (4%), crabs [3% (blue, stone, and horseshoe crabs)] and mollusks [0.19% (conchs/whelks, squid, and octopus)].

On average 255,776 pounds of finfish are landed and sold annually by shrimp trawls (Table 6.12). Eighty-nine percent of the total finfish landings were reported in the ocean (< 3 miles) and the Pamlico Sound (Table 6.13). Six groups; sea mullet [whiting, and kingfish 46.14%, 1,925,720 pounds), flounder [summer and southern (17%, 711,590 pounds), spot (17%, 697,715 pounds), Atlantic croaker (6%, 256,741 pounds), weakfish (4%, 166,669 pounds), and butterfish (4%, 156,131 pounds) account for 94% of the finfish landings (Table 6.14). Ninety-seven percent of sea mullet, flounder, spot, Atlantic croaker, and weakfish were reported from two areas, the Pamlico Sound and the ocean (Table 6.15). The ocean accounts for 85% of the croaker, 62% of the sea mullet, and 54% of the spot, 32% of the flounder, and 9% of the weakfish landed from shrimp trawls (Table 6.14). The Pamlico Sound accounts for 88% of the weakfish, 62% of the flounder, 41% of the spot, 37% of the sea mullet, and 14% of the croaker landings for this gear. The peak months for finfish landings from shrimp trawls are in October (22%) and November (22%), with the period of August through December accounting for 80% of all finfish landings (Table 6.16). The peak month for sea mullet landings from shrimp trawls is November (32%), while the period from July through December accounts for 84% of the

DRAFT

landings (Table 6.17). Eighty-one percent of the flounder are landed from July through November, with October accounting for 23% of the landings. The period of August through November accounts for 95% of the spot landings, with October accounting for 44% of the landings. Sixty-nine percent of the Atlantic croaker are landed in December, and 94% are landed from September through January. Weakfish landings peaked in August (26%), while 92% of the landings occur from July through November.

An average of 171,523 pounds of crabs were landed and sold annually by shrimp trawls from 1994 to 2010 (Table 6.12). Fifty-nine percent of the crab landings were reported in Core Sound and 27% in the Pamlico Sound (Table 6.13). The period of April through August accounts for 82% of the crab landings. Seventy-one percent of the mollusk landings were reported from the ocean (< 3 miles) and 14% from the Pamlico Sound. The peak month for mollusk in shrimp trawls is November (22%), while the period of May through December accounts for 94% of the landings.

Table 6.12 Percent shrimp trawl landings (lb)[†] of major market groups for North Carolina, 1994-2010.

Year	Shrimp		Fish		Crabs		Mollusk		Total pounds
	Pounds	% total	Pounds	% total	Pounds	% total	Pounds	% total	
1994	6,888,784	89.46	391,585	5.09	394,817	5.13	25,066	0.33	7,700,252
1995	7,903,144	90.77	562,058	6.46	203,379	2.34	38,285	0.44	8,706,866
1996	4,874,017	85.81	530,605	9.34	266,296	4.69	9,225	0.16	5,680,143
1997	6,451,315	91.51	317,716	4.51	264,656	3.75	16,008	0.23	7,049,696
1998	4,270,740	85.62	197,277	3.95	508,457	10.19	11,574	0.23	4,988,048
1999	8,108,209	92.34	411,973	4.69	247,198	2.82	13,063	0.15	8,780,443
2000	9,442,710	94.90	320,997	3.23	169,906	1.71	16,449	0.17	9,950,063
2001	4,749,564	93.86	141,304	2.79	161,169	3.18	8,256	0.16	5,060,293
2002	8,879,729	95.87	231,457	2.50	143,367	1.55	7,481	0.08	9,262,034
2003	5,432,418	92.85	142,410	2.43	266,528	4.56	9,687	0.17	5,851,042
2004	4,351,064	92.65	185,373	3.95	147,715	3.15	12,136	0.26	4,696,287
2005	2,046,274	95.46	34,746	1.62	58,178	2.71	4,445	0.21	2,143,643
2006	4,862,890	97.24	93,963	1.88	36,224	0.72	8,086	0.16	5,001,163
2007	8,781,019	97.18	216,117	2.39	27,984	0.31	10,345	0.11	9,035,464
2008	8,789,623	96.56	296,496	3.26	4,246	0.05	12,305	0.14	9,102,671
2009	5,039,827	96.50	168,523	3.23	4,827	0.09	9,605	0.18	5,222,782
2010	5,532,780	97.80	105,587	1.87	10,936	0.19	7,895	0.14	5,657,198
Average	6,259,065	93.32	255,776	3.72	171,523	2.77	12,936	0.19	6,699,299

[†]Single gear Trip Tickets

DRAFT

Table 6.13 Percent shrimp trawl landings (lb) †of major market groups for North Carolina by waterbody, 1994-2010.

Area	Shrimp		Fish		Crabs		Mollusk		Total	
	Pounds	% of total	Pounds	% of total	Pounds	% of total	Pounds	% of total	Pounds	%
Pamlico Sound	61,577,667	57.87	1,852,608	42.61	773,966	26.54	30,911	14.06	64,235,152	56.40
Ocean < 3 Miles	26,123,378	24.55	2,001,168	46.02	9,630	0.33	155,853	70.87	28,290,028	24.84
Core Sound	7,578,742	7.12	60,120	1.38	1,730,141	59.34	2,536	1.15	9,371,539	8.23
Ocean > 3 Miles	3,272,314	3.08	332,977	7.66	1,911	0.07	19,586	8.91	3,626,789	3.18
Neuse River	2,245,066	2.11	20,597	0.47	256,138	8.78	437	0.20	2,522,238	2.21
Inland Waterway	1,443,181	1.36	17,160	0.39	17,217	0.59	6,702	3.05	1,484,260	1.30
Cape Fear River	1,296,961	1.22	16,738	0.38	6,721	0.23	1,786	0.81	1,322,207	1.16
New River	928,421	0.87	30,189	0.69	35,553	1.22	1,222	0.56	995,385	0.87
Newport River	501,687	0.47	682	0.02	4,349	0.15	133	0.06	506,850	0.45
Pamlico River	472,582	0.44	7,535	0.17	9,665	0.33	*	*	489,868	0.43
North River/Back Sound	313,448	0.29	1,791	0.04	8,669	0.30	396	0.18	324,304	0.28
White Oak River	216,472	0.20	896	0.02	113	0.00	*	*	217,578	0.19
Bay River	152,702	0.14	1,369	0.03	13,271	0.46	*	*	167,360	0.15
Croatan Sound	142,549	0.13	2,993	0.07	30,102	1.03	0	0.00	175,644	0.15
Roanoke Sound	94,879	0.09	1,204	0.03	16,539	0.57	0	0.00	112,623	0.10
Pungo River	31,429	0.03	*	*	*	*	0	0.00	33,324	0.03
Shalotte River	11,847	0.01	149	0.00	*	*	*	*	12,157	0.01
Lockwood Folly	765	0.00	0	0.00	0	0.00	0	0.00	765	0.00
Albemarle Sound	*	*	0	0.00	0	0.00	0	0.00	18	0.00
Total	106,404,108	100.00	4,348,187	100.00	2,915,886	100.00	219,908	100.00	113,888,089	100.00

*Confidential

†Single gear Trip Tickets

DRAFT

Table 6.14 Yearly finfish landings (lb) †from shrimp trawls all North Carolina Waters combined, 1994-2010.

Year	Sea Mullet	Flounders	Spot	Croaker	Weakfish	Butterfish	Other Species	Sheepshead	Harvestfish	Spanish Mackerel	Pigfish
1994	93,244	131,247	57,835	14,305	47,385	8,710	29,957	4,206	1,722	850	2,123
1995	226,595	74,176	78,795	18,642	40,312	50,685	56,802	4,326	6,658	3,287	1,781
1996	132,953	70,688	72,924	190,251	18,492	18,905	14,681	3,155	4,077	2,273	2,208
1997	105,149	63,457	76,050	15,695	13,786	7,142	20,806	3,265	4,813	5,043	2,512
1998	78,843	39,143	43,493	1,857	5,014	6,657	12,874	2,749	3,199	1,911	1,538
1999	231,075	68,648	45,351	6,956	17,304	10,167	16,376	4,366	8,627	2,271	832
2000	154,700	38,810	80,608	1,129	7,190	7,347	18,440	4,911	5,327	1,439	1,097
2001	47,414	30,419	43,176	2,254	1,793	2,316	8,868	1,811	2,040	497	717
2002	113,705	48,581	36,013	1,661	2,983	6,925	12,926	4,315	2,534	1,183	632
2003	67,859	24,257	33,884	994	1,360	1,638	7,582	2,622	1,234	164	815
2004	107,529	24,223	27,090	705	2,917	9,230	8,479	3,684	869	173	473
2005	14,399	5,427	3,578	78	596	1,154	7,041	1,556	643	211	64
2006	45,688	16,080	15,740	449	1,959	3,800	8,125	1,418	554	45	107
2007	129,316	21,588	20,714	208	1,561	6,916	11,529	16,579	7,437	151	120
2008	211,946	24,873	21,609	519	2,787	8,325	16,931	5,445	3,228	583	252
2009	86,553	19,367	36,725	603	752	4,585	13,290	2,658	2,663	1,200	128
2010	78,750	10,605	4,130	436	480	1,631	6,313	1,329	1,457	307	150
Total lbs	1,925,720	711,590	697,715	256,741	166,669	156,131	96,436	68,390	57,079	21,585	15,547
% of total	46.14	17.05	16.72	6.15	3.99	3.74	2.31	1.64	1.37	0.52	0.37

†Single gear Trip Tickets

DRAFT

Table 6.15 Percent shrimp trawl landings (lb) †of top five finfish groups by waterbody, 1994-2010.

Waterbody	Sea Mullet		Flounders		Spot		Croaker		Weakfish		Total	
	Pounds	% sea mullet	Pounds	% flounder	Pounds	% spot	Pound	% croaker	Pounds	% weakfish	Pounds	%
Ocean	1,191,867	61.90	231,021	32.47	372,837	53.51	218,132	85.02	14,213	8.54	2,028,070	53.96
Pamlico Sound	709,326	36.84	437,684	61.51	287,497	41.26	35,329	13.77	146,633	88.15	1,616,469	43.01
Core Sound	9,269	0.48	16,457	2.31	6,363	0.91	366	0.14	2,019	1.21	34,473	0.92
New River	2,101	0.11	7,843	1.10	14,055	2.02	1,161	0.45	107	0.06	25,266	0.67
Neuse River	6,521	0.34	4,894	0.69	3,482	0.50	444	0.17	1,701	1.02	17,040	0.45
Inland Waterway	1,405	0.07	3,855	0.54	7,769	1.11	553	0.22	94	0.06	13,675	0.36
Cape Fear River	2,430	0.13	4,736	0.67	2,352	0.34	290	0.11	185	0.11	9,993	0.27
Pamlico River	2,107	0.11	2,837	0.40	860	0.12	100	0.04	384	0.23	6,287	0.17
Croatan Sound	204	0.01	574	0.08	1,087	0.16	113	0.04	452	0.27	2,430	0.06
North River/Back Sound	78	0.00	135	0.02	*	*	*	*	*	*	1,296	0.03
Bay River	224	0.01	366	0.05	131	0.02	54	0.02	455	0.27	1,229	0.03
Roanoke Sound	18	0.00	452	0.06	389	0.06	29	0.01	101	0.06	989	0.03
White Oak River	*	*	603	0.08	*	*	0	0.00	0	0.00	833	0.02
Newport River	*	*	108	0.02	*	*	*	*	*	*	233	0.01
Shalotte River	0	0.00	*	*	*	*	0	0.00	0	0.00	143	0.00
Pungo River	0	0.00	*	*	0	0.00	*	*	0	0.00	12	0.00
Total	1,925,720	100.00	711,590	100.00	697,715	100.00	256,741	100.00	166,669	100.00	3,758,435	100.00

*Confidential

†Single gear Trip Tickets

DRAFT

Table 6.16 Percent shrimp trawl landings (lb) [†]of major market groups landed for North Carolina (all waters combined) by month, 1994-2010.

Month	Shrimp		Fish		Crabs		Mollusk		Total	
	Pounds	% total	Pounds	% total	Pounds	% total	Pounds	% total	Pounds	%
January	663,541	0.62	71,145	1.64	11,235	0.39	2,305	1.05	748,226	0.66
February	458,803	0.43	86,844	2.00	6,317	0.22	3,708	1.69	555,672	0.49
March	308,600	0.29	90,926	2.09	49,284	1.69	1,272	0.58	450,082	0.40
April	598,610	0.56	54,255	1.25	271,902	9.32	6,733	3.06	931,499	0.82
May	2,306,314	2.17	105,169	2.42	531,624	18.23	16,205	7.37	2,959,311	2.60
June	7,243,268	6.81	130,633	3.00	616,100	21.13	21,785	9.91	8,011,786	7.03
July	28,302,130	26.60	351,103	8.07	663,163	22.74	24,838	11.29	29,341,234	25.76
August	22,540,167	21.18	505,297	11.62	298,053	10.22	29,606	13.46	23,373,123	20.52
September	15,816,897	14.86	566,800	13.04	131,437	4.51	16,298	7.41	16,531,432	14.52
October	17,060,394	16.03	965,699	22.21	134,967	4.63	31,018	14.10	18,192,078	15.97
November	8,996,863	8.46	975,606	22.44	152,002	5.21	48,802	22.19	10,173,273	8.93
December	2,108,521	1.98	444,712	10.23	49,802	1.71	17,338	7.88	2,620,374	2.30
Total	106,404,108	100.00	4,348,187	100.00	2,915,886	100.00	219,908	100.00	113,888,089	100.00

[†]Single gear Trip Tickets

DRAFT

Table 6.17 Percent shrimp trawl landings (lb) †of top five finfish groups by month, 1994-2010.

Month	Sea Mullet		Flounders		Spot		Croaker		Weakfish		Total	
	Pounds	% sea mullet	Pounds	% flounder	Pounds	% spot	Pounds	% croaker	Pounds	% weakfish	Pounds	%
January	46,808	2.43	5,331	0.75	33	0.00	9,691	3.77	289	0.17	62,151	1.65
February	79,560	4.13	2,679	0.38	20	0.00	10	0.00	401	0.24	82,670	2.20
March	65,996	3.43	7,231	1.02	557	0.08	1,000	0.39	636	0.38	75,420	2.01
April	30,151	1.57	16,118	2.27	992	0.14	391	0.15	393	0.24	48,044	1.28
May	30,595	1.59	35,175	4.94	2,263	0.32	339	0.13	2,288	1.37	70,658	1.88
June	49,894	2.59	44,655	6.28	3,072	0.44	661	0.26	4,198	2.52	102,481	2.73
July	171,832	8.92	71,455	10.04	25,665	3.68	4,206	1.64	27,593	16.56	300,750	8.00
August	217,017	11.27	89,113	12.52	74,100	10.62	9,300	3.62	42,543	25.53	432,073	11.50
September	107,601	5.59	163,400	22.96	163,107	23.38	16,316	6.36	27,360	16.42	477,785	12.71
October	323,726	16.81	166,200	23.36	306,548	43.94	18,031	7.02	31,539	18.92	846,043	22.51
November	635,332	32.99	84,050	11.81	117,991	16.91	18,770	7.31	24,630	14.78	880,772	23.43
December	167,211	8.68	26,182	3.68	3,367	0.48	178,027	69.34	4,802	2.88	379,588	10.10
Total	1,925,720	100.00	711,590	100.00	697,715	100.00	256,741	100.00	166,669	100.00	3,758,435	100.00

†Single gear Trip Tickets

DRAFT

6.3.3 Discarded Catch

In 1998 the NMFS completed a report summarizing the results from their Southeastern United States Shrimp Trawl Bycatch Program in response to a Congressional requirement imposed by the Sustainable Fisheries Act of 1996. The shrimp trawl bycatch program was initiated in 1992 as part of Section 405(e) of The Magnuson-Stevens Fishery Conservation and Management Act. In that report, more than 150 taxa were identified in shrimp trawl catches in the South Atlantic, with an average overall catch rate of 57.33 pounds per hour (Nance 1998). Finfish made-up 54% of the catch by weight, shrimp 18%, other invertebrates 18%, and the remaining 13% was composed of crustacean. Seasonal distribution of finfish bycatch in the south Atlantic indicates that the highest percentage by weight occurs in the summer, while numerically finfish bycatch is highest in the spring. The top ten species by weight were: cannonball jelly (14%), white shrimp, spot, and Atlantic menhaden each at 9%, brown shrimp and other jellyfish at 6% each, Atlantic croaker contributes 6%, southern kingfish, and blue crab each at 4%, and star drum at 3%.

In the Gulf of Mexico over 450 taxa were identified in shrimp trawls (Nance 1998). The average hourly catch was approximately 59 pounds per hour of towing. Finfish made-up 67% of the catch by weight, shrimp 16%, crustacean 13%, and the remaining 4% was composed of other invertebrates. Seasonally, finfish bycatch was highest, by weight, in the fall. The 10 most abundant species by weight were: longspined porgy (15%), brown shrimp (9%), Atlantic croaker (9%), inshore lizardfish (6%), pink shrimp (3%), gulf butterfish, and lesser blue crab, white shrimp, longspined swimming crab, and brown rock shrimp each comprising 2% of the catch.

In 1950 sampling was conducted aboard commercial shrimp trawlers working in Core and Pamlico sounds (Roelofs 1950). Although only total weights were reported for shrimp and finfish, Roelofs (1950) indicated that for Core Sound "85 to 90% of the fish taken were croakers and spot, with croaker predominating; while in late August, hogfish, pinfish and other trash species increased until they made up over 50 per cent of the catch". Seven tows were sampled in Pamlico Sound during September of 1950. Atlantic croaker comprised 73% of the finfish taken, with spot and trout each accounting for 10% (Roelofs 1950).

Prior to the work done by Diamond-Tissue (1999) and Johnson (2003; 2006), there was little information characterizing the bycatch in the North Carolina shrimp trawl fishery. Diamond-Tissue's (1999) 1995 characterization study examined 52 tows conducted over 15 trips. Samples from Pamlico Sound (n=16 tows) and the Cape Fear River (n=24 tows) were collected monthly from July through October 1995. Additionally, four tows were sampled in Core Sound in August 1995, and eight tows were examined off Carolina Beach during July and August. Sampled boats had one or two nets, and all nets contained the required TED and BRD. A total of 92 different species, including 66 species of finfish, 10 species of crabs, and 13 other invertebrates were identified. For all areas combined, market-size penaeid shrimp made up 44.3% of the organisms by number and 30.8% by weight. The top finfish species by number were star drum, Atlantic croaker, weakfish, and spot, while Atlantic croaker, weakfish, spot, and star drum were the top finfish species by weight. In Pamlico Sound, 38 species were identified in the catches, 37 were identified in Core Sound, and 50 species were identified in Cape Fear River. Market-size penaeid shrimp were the top species in terms of both numbers and weight for all areas combined, as well as for all individual areas by number, and all areas by weight except Core Sound. The composition of finfish in the bycatch varied by area, with Atlantic croaker, spot, and weakfish accounting for 53% of the total catch by number and 56% by weight in Pamlico Sound. In Core Sound, pigfish, spot, and Atlantic croaker were the most abundant

DRAFT

finfish species in terms of number and weight. Star drum, weakfish, and Atlantic croaker were the most abundant species in the Cape Fear River.

Johnson (2003) quantified the catch of shrimp trawlers working in Core Sound (n=46 tows) and the Neuse River (n=8 tows) during the summers of 1999 and 2000. Overall, blue crabs accounted for 26% by weight of the total combined catch. Spot accounted for 17% of the total catch and 40% of the total finfish bycatch. Core Sound catches were dominated by invertebrates, crabs, and shrimp, which accounted for 71% of the total catch. Three species of finfish; spot (48%), Atlantic croaker (13%), and pinfish (12%) accounted for 73% of the finfish bycatch from this area. In the Neuse River, invertebrates made up 24% of the sampled catches. Atlantic croaker (44%) and spot (33%) accounted for 77% of the finfish bycatch.

During the spring and summer of 1999 and 2000 Johnson (2006) also characterized the bycatch of inshore commercial shrimp trawlers working in Core Sound, Southern Pamlico Sound and Back Sound. All nets sampled were fitted with BRDs and TEDs, the Florida Fish Eye (FEE) excluder was cited as the most commonly used BRD. A total of 52 trawls were sampled, however only 50 trawls were analyzed for species composition. Overall, shrimp accounted for 21% of the catch by weight. Fish, blue crabs, and other organisms, such as jellyfish, horseshoe crabs, and other species of crabs made up 27%, 33%, and 20% of the catch by weight, respectively. The majority of the bycatch and discards were made up of juvenile estuarine fish and juvenile and adult blue crabs. Spot (21%), Atlantic croaker (8%) and pinfish (4%) were the most abundant finfish by weight. The average CPUE for shrimp was 15.3 kg/hr and 19.0 kg/hr for fish (1 kilogram [kg] = 2.20 pounds).

Logothetis and McCuiston (2004) characterized the bycatch of the inshore commercial shrimp fishery in southeastern North Carolina during the 2004 shrimp season. From April through November, 64 trips were observed, consisting of 132 tows in five regions (IWW of Brunswick, New Hanover, Onslow, and Pender Counties, and the Cape Fear River). Fishing took place on a 24 foot shrimp trawler using single-rig otter trawls; trawl type depended on the target species. All tailbags were 1 1/2" stretch mesh, all nets were fitted with diamond shaped FFEs and an aluminum TED. Shrimp (brown, pink, white) made up 55% of the total catch. Bycatch made up roughly 45% of the total catch and consisted of 84 different species. Blue crabs accounted for 9% by weight of the total combined catch. Atlantic croaker (8%), weakfish (4%), pinfish (4%), spot (4%), and flounder species (southern and summer flounder) accounted for (2%) by weight of the total catch. One Kemp's ridley sea turtle was also caught during the study. The length frequencies of the bycatch indicated that nearly all of the bycatch were juvenile to subadult species. Roughly 50% of the blue crab, 100% of the weakfish, and 95% of the flounder species would have been regulatory discards using today's minimum size limits [blue crab - 5" carapace length, weakfish - 12" minimum TL, flounder species - 14" minimum TL (commercial)]. Overall, the catch rates for bycatch peaked in July, elevated levels of bycatch were also observed in May and September. The highest observed mean CPUE (kg/min) for invertebrates in all regions occurred in August (0.149) and for commercial and recreational finfish in May (0.226) and July (0.273).

Brown (2009) characterized the near-shore commercial shrimp trawl fishery from Carteret County to Brunswick County from 2007 to 2008. In this study commercial fishermen were randomly selected, and observer effort was weighted by region using the NCDMF Trip Ticket Program. Over the course of the study, observations were made on 142 trips, consisting of 314 tows, achieving 5.92% coverage in number of trips. The results were stratified by net type (double seamed and tongue nets) and season (Winter: January-March, Spring: April-June, Summer: July-September, Fall: October-December). All observed trips used FFEs, Super

DRAFT

Shooter, Straight Bar and Inshore Hard TEDs were also observed on the majority of the trips; however, 4% of tongue net trips used no TED. Over 100 species were observed throughout the study in all net types; 80 in the double seamed nets and 90 in the tongue nets. Shrimp (brown, white, pink) accounted for 21% of the catch by weight in all net types. Atlantic croaker (25%) and spot (7%) were the most abundant finfish bycatch in all net types. In the double seamed fishery, the CPUE [(total weight (kg) / (headrope length* number of nets*tow time)] of Atlantic croaker was significantly higher than the other commercially important finfish bycatch in the spring and summer; in the tongue net fishery it was higher during the summer. The CPUE of spot was higher in the summer for the double seamed trawl nets. In the tongue net fishery, the spot CPUE was the highest in the fall. Overall, roughly 99% of spot by weight were classified as unmarketable bycatch. Weakfish represented the largest regulatory discard (60%) by weight in both nets. The CPUE for weakfish was the highest during the summer in the tongue net fishery; almost three times as high as what was observed in double seamed fishery during that same period. In the double seamed nets, Spanish mackerel, southern flounder, and summer flounder represented 16%, 8%, and 9% of the regulatory discards, respectively. In the tongue net fishery Spanish mackerel, southern flounder, and summer flounder represented 15%, 8%, and 8% of the regulatory discards, respectively.

In 2009, Brown (2010b) conducted another study characterizing the inshore commercial shrimp trawl fishery in the Pamlico Sound and its tributaries. Using the same protocol as used in Carteret and Brunswick County survey (Brown 2009) the catch of federally and state managed species of finfish caught in double seamed, four seamed and tongue nets was quantified. Over the course of the study, 66 commercial shrimp trawl fishing days were observed, consisting of 191 tows, achieving 1.21% coverage in fishing days. Similar to the previous study, all observed trips used the FFE. Super Shooter, Straight Bar and Inshore Hard TEDs were also observed on the majority of the trips. Sixty-nine species were observed throughout the study in all net types, 56 were observed in the double seamed shrimp trawl nets, 51 in the four seamed nets and 38 in the tongue nets. Shrimp (brown, white, pink) accounted for 23% of the catch by weight in all net types. Atlantic croaker (33%) and spot (13%) were the most abundant finfish bycatch by weight in all net types and accounted for the largest percentage of unmarketable discards. Weakfish, kingfish (*Menticirrhus spp.*), and spotted sea trout represented 6.34%, 0.79%, and 0.02% of the catch in all net types by weight, respectively. The highest observed CPUEs of Atlantic croaker and spot were observed in summer double seamed fishery. Weakfish represented the largest regulatory discards in the double seamed net (98%), four seamed nets (100%) and the tongue net fishery (86%). The highest observed CPUE for weakfish occurred in the summer four seamed fishery. Spanish mackerel, southern flounder, and summer flounder represented 0.48%, 5%, and 17% of the regulatory discards in the double seamed nets, respectively. In the four seamed net fishery, Spanish mackerel, southern flounder, and summer flounder represented 2%, 3%, and 6% of the regulatory discards, respectively. Spotted seatrout, Spanish mackerel, southern flounder, and summer flounder represented 4%, 1 %, 1%, and 9% of the regulatory discards in the tongue net fishery, respectively.

Another way of presenting and expanding bycatch data is by using the ratio of finfish to shrimp (F:S). A common method of calculating F:S ratios is to subsample the entire catch and to expand the shrimp to finfish ratio of subsample to the weight of the entire catch. Diamond (2003) cautions that F:S ratios tend to overestimate bycatch and that a CPUE estimator is the most appropriate method of scaling up individual observations to the entire fishery. The F:S ratio can be a factor of the environment and a fisherman's experience, thus if there are few shrimp in the area or a fisherman's gear is not fishing properly or he is in a poor area, a higher F:S ratio will result (Coale et al. 1994). In a study using both field data and computer simulations to compare the methods of bycatch estimation, total bycatch estimates derived with

DRAFT

the basic F:S ratio estimator by both weight and number were two to seven times higher than those based on the CPUE-mean per unit method (Diamond 2003). Both the CPUE and F:S methods tend to ignore sources of variability at several hierarchical levels, assuming that total shrimp catch and effort data are without error and that the catches are thoroughly mixed so a single sample characterizes the entire catch without variance (Diamond-Tissue 1999). Additionally, both estimation methods add a certain degree of error when they get expanded from the sampled net to the number of nets per tow. While both methods of estimating bycatch have their advantages and disadvantages, the F:S method is much easier to obtain and use than effort data and allows the use of observer data at the tow level without the additional variance caused by averaging the number of tows per trip (Vaughan and Nance 1998).

Nance (1998) reported a F:S ratio of 5.3:1 for the Gulf of Mexico, and 4.5:1 for the South Atlantic. Reported F:S ratios for North Carolina are 1.5:1 (Roelofs 1950), 1.6:1 (Diamond-Tissue 1999), 3.1:1 (Johnson 2003), 0.5:1 (Logothetis and McCuiston 2005), 1.6:1 (Johnson 2006). Using the relative biomass tables (kg) in Brown's (2010b) study characterizing the inshore commercial shrimp trawl fishery in the Pamlico Sound and its tributaries, the calculated F:S ratio (excluding sharks, and rays) was 2.7:1 (all gear types), 2.9:1 (double seamed net fishery), 2.4:1 (four seamed net fishery) and 3.4:1 (tongue net fishery). Again using Brown's (2009) relative biomass tables (kg) the calculated F:S ratio of the near-shore commercial shrimp trawl fishery from Carteret County to Brunswick County was 2.7:1 (all gears), 2.0:1 (double seamed net fishery) and 3.3:1 (tongue net fishery). Johnson (2006) notes that the F:S ratios reported in her study were highly variable (ranging from 0 to 6.9:1) and were included solely for comparison to other studies and not intended for use to estimate the total bycatch in the fishery. Diamond (2003) also cautions that due to statistically significant two- and three-way interactions among parameters (mean or variance of catch, observer coverage, correlation between the catch of fish and shrimp), bycatch estimates obtained with different methods should not be compared directly. Additionally, the methods used to analyze F:S ratios are often not well described and vary from study to study (Diamond-Tissue 1999). Table 6.18 lists the findings of various studies characterizing the commercial shrimp trawl fisheries in North Carolina and the South Atlantic.

Numerous gear evaluation studies have also been conducted in North Carolina waters (McKenna and Monaghan 1993; Coale et al. 1994; Murray et al. 1995; McKenna et al. 1996, Brown 2010a). However, this data should not be used for characterization analysis since these studies are often relegated to times of low shrimp catch rates, and as such, the bycatch data are not representative of times when shrimp catch rates are higher. For example the F:S ratio for gear studies conducted in 1994 (McKenna et al. 1996) was 5.5:1, while characterization studies conducted in 1995 by Diamond-Tissue (1999) found the F:S ratio to be 1.6:1. While these data should not be used for characterization analysis, catches can provide information on species and sizes of species vulnerable to shrimp trawls. However, it is important to note that for all discard and bycatch studies, variability exists within time periods as short as 24 hours and extends to year-to-year variability and it may not be reasonable to assume that bycatch rates in neighboring areas can give an accurate approximation of an unsampled area (Alverson et al. 1994; Alverson and Hughes 1996; Diamond-Tissue 1999). Furthermore, the ratio of discards to retained or total catch as well as raw numbers of weight are not, in themselves indicators of serious biological or ecological problems (Mangel 1993; Alverson and Hughes 1996).

DRAFT

Table 6.18 Author (year published), years sampled, area sampled, number of trips sampled, number of tows sampled, number of species observed, percent bycatch, percent finfish, percent shrimp, finfish to shrimp ratio (F:S), and bycatch to shrimp ratio (BC:S) of previous bycatch characterization work conducted in North Carolina and the South Atlantic.

Report	Years Sampled	Area Sampled	Percent Coverage	Number Trips	Number of Tows	Number of Species	Percent Bycatch [†]	Percent Finfish [†]	Percent Shrimp [†]	F:S Ratio [†]	BC:S Ratio ^{††}
Roelofs (1950)	1950	Core Sd, Pamlico Sd			17					1.5:1	
Nance (1998)	1992-96	South Atlantic (shown), Gulf of Mexico		604**	5,695**		82	54	18	4.5:1	
Diamond-Tissue (1999)	1995	Pamlico Sd, Core Sd, Cape Fear River, Ocean (off Carolina Beach)		15	52	92	69	51	31	1.6:1	
Johnson (2003)	1999-00	Core Sd, Neuse River			52		80		20	3.1:1	5.7:1
Johnson (2006)	1999-00	Back Sd, Core Sd, Southern Pamlico Sd		15	52		79	27	21	1.6:1	
Logothetis & McCuiston (2004)	2004	IWW of Brunswick Co, Onslow Co, Pender Co, Cape Fear River		64	132	87	45		55	0.5:1	0.8:1
Brown (2009)	2007-08	Brunswick Co, Cape Fear River, New Hanover Co, Onslow Co, Pender Co	5.92	142	314	110	79	56**	21	2.7:1*	3.8:1*
Brown (2010)	2009	Pamlico Sd	1.21	66	191	69	77	61**	23	2.7:1*	3.4:1*

[†]By weight (kg)

^{††}By weight (kg), includes finfish, jellyfish, rays, sharks, crabs, etc.

*Calculated using relative biomass tables (kg) from report using Logothetis and McCuiston's (2005) method that excludes sharks and rays.

**Number of trips and tows includes both South Atlantic and Gulf of Mexico

6.3.4 Biological Implications of Bycatch

Evaluating the biological impacts of bycatch is a two stage process. First, the bycatch must be characterized in both magnitude and nature. Second, information obtained from characterization efforts must be applied to population and ecosystem models to evaluate potential impacts at those levels. Although, by definition, bycatch can include both incidental and discarded catch, much of the current concern is directed toward discarded animals. This concern is largely due to a general perception that discarded bycatch is a waste of natural resources and leads to overfishing (Crowder and Murawski 1998). Beyond the obvious impacts on discarded individuals, there are also potential population and ecosystem level effects (Alverson et al. 1994; Crowder and Murawski 1998). Kept bycatch has biological impacts also, but since it is accounted for as catch such impacts are encompassed in harvest management strategies.

As previously noted, the biological significance of bycatch can be judged from a number of different perspectives, including those of the populations (e.g., of a particular species), of the fishery or fisheries that target or otherwise encounter the species, and of the general biological community (Murawski 1995). The first phase of characterization starts at the level of an individual animal. Discarded individuals suffer one of two immediate alternative fates: survival or death. Further, initial survival may still lead to chronic effects, such as delayed mortality, reduced growth, interrupted maturation, and displacement. Discarded animals are also vulnerable to increased predation, as shown by numerous observations of live discarded animals being preyed upon by birds, marine mammals, and finfishes. If this initial predation is avoided, the animals must still seek shelter and return to their normal environments, all the while exposed to the risk of predation (Murawski 1995).

In survival experiments examining the fate of discarded bycatch in the Core Sound and Neuse River shrimp trawl fisheries, 34% of the overall organisms caught were alive and healthy at the time of “discarding”, 11% were injured or non-responsive, and 56% were dead (Johnson 2003). Survival was also found to vary among species and the amount of time on deck. On average 80% of the blue crabs survived uninjured, survival did not improve with shorter tow durations or time on deck. Eleven percent of the finfish survived uninjured, 11% survived but were injured or unresponsive, and 78% were dead. Survival of croaker declined significantly with increasing time on deck; however, pinfish and spot showed no change in survival with time out of water or tow time. Spot had the lowest survival of the common species in the discards. In another survival experiment, 45% of the fish caught as bycatch were alive and healthy, 3% were alive and weak, and 52% were dead after an average holding time of 3.22 hours (Logothetis and McCuiston 2005). Logothetis and McCuiston (2005) also found that significantly more fish survived if cull times were 30 minutes or less and water temperatures were less than ~80 °F; however, the survivability of weakfish was less than 50% even when the cull times were short. Birds and blue crabs were found to be the primary scavengers on the discards, most of which were dead juvenile finfish (Johnson 2003; Logothetis and McCuiston 2005).

While discarding is generally thought of in an active sense, most fishing gears are designed to provide some degree of passive discarding. In trawling, mesh sizes are selected by choice or mandated by regulation to prevent the harvest of small sized animals and it is generally assumed that animals escaping through the mesh survive. However the possibility remains that not all animals survive, resulting in some level of unobserved mortality. This unobserved mortality is a difficult issue for both managers and scientists because when it occurs, the actual reduction in bycatch and thus mortality is lessened (Chopin and Arimoto 1995). Furthermore, since gear escapees cannot be counted by conventional fishery observer programs, they cannot

DRAFT

be monitored or included in stock assessment calculations. Chopin and Arimoto (1995) suggest that escapee mortality should be considered if gear-based measures are used as a primary management tool.

When viewed at the population level, the first instinct of many people is to assume that discarding adversely impacts populations or stocks. Such ideas lead to the widely held view that discarding, especially when the magnitude in pounds or numbers is large, contributes to overfishing and the decline of many stocks. Even if a bycatch associated fish stock is in decline, proving cause and effect is difficult because other factors such as environmental degradation may be involved (Murray et al. 1992). Unfortunately, few hypotheses about population-level impacts have been tested (Crowder and Murawski 1998). Regardless, just as large levels of discarding do not necessarily lead to significant biological impacts; it cannot be assumed that minimal discarding has only minor effects (Alverson et al. 1994). Discard impacts can only be determined through proper data collection and analytical investigations. Various studies suggest that discarding has harmed some stocks, while others seem unaffected. For example, discarding has been implicated in the decline of Gulf of Maine groundfish, Atlantic croaker in the Gulf of Mexico, and scup and black seabass in the Mid-Atlantic (Alverson et al. 1994; ASMFC 1996a; ASMFC 1996b). Conversely, sizable discarding of redfish in the Northwest Atlantic and pollock, cod, and sablefish in the Northeast Pacific represents only a fraction of the total mortality of these species and is not believed to have a significant adverse impact on population abundance (Alverson et al. 1994).

The magnitude of discarding should not be the only concern when examining population-level impacts because such effects are also related to the size or life-stage of the discarded animal. If discards are immature or below the size for optimum yield, both yield-per-recruit and spawning potential may be adversely impacted (Crowder and Murawski 1998). In other words, it is commonly known that harvesting fish before they mature and spawn can lead to recruitment overfishing and can impair a stock's ability to sustain itself. Also, harvesting a fish before it reaches some optimal size can lead to growth overfishing and reduced overall yield from the fishery. Thus, fish with slow growth rates may be more affected by bycatch mortality. These principles are unavoidable consequences of exploitation that can occur whether the fish are harvested or discarded.

In addition to impacts on individuals and populations, it is suspected that discarding can also alter entire communities. Community effects are still largely unknown, but in theory they could be significant. For instance, if an abundant species that dominates a community is removed by harvest while another species is discarded and survives, the community could eventually change to the extent that the discarded species becomes the dominant species in the ecosystem (Murawski 1995). If the newly dominant species is of less value, either ecologically or economically, both the ecosystem and the fishing economy could suffer. It is thought that such species-specific exploitation could be more damaging to the productivity of an ecosystem than exploitation of the entire community. However, such effects remain largely speculative as there has been little research on community-level effects.

Deehr (2012) investigated and modeled the impacts of shrimp trawling on the estuarine ecosystem in Core Sound, North Carolina. Using field collections, fisheries data from the NC Trip Ticket program, and Ecopath network modeling software, she created four network models of areas open and closed to shrimp trawling during spring (2007) and fall (2006 and 2007). Each model consisted of 65 compartments (including non-living detritus, bycatch, producers, and various invertebrate and vertebrate consumers), and harvests by different types of fishery

DRAFT

gears (crab pots, gill nets, haul seines, and pound nets in closed areas; shrimp trawls, skimmer trawls were added to the models in areas open to trawling).

Based on the benthic sampling, shrimp trawling had a major impact on the Core Sound ecosystem. Contrary to expectation, biomass (g C/m²) of infaunal benthic invertebrates, especially deposit-feeding polychaetes, was significantly greater in areas open to trawling. Meiofaunal biomass was significantly greater in the closed areas. Field collections of fish and invertebrates revealed that there was more biomass (g C/m²) of benthic-invertebrate feeders (such as blue crabs, spot, and pinfish) in areas closed to trawling. These results suggest a trophic cascade due to trawling may have occurred in the open areas, whereby trawls removed benthic-feeding fishes and blue crabs, released their prey (benthic polychaetes) from predation pressure, and lowered the abundance of meiofauna (prey of the polychaetes). Alternatively, the dead biomass from by-catch could fuel the growth in polychaetes and other benthos due to a direct subsidy from trawling. Further experimental work is required to test these model-derived hypotheses.

Ecopath-calculated effective trophic levels were validated using stable isotopes of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$. Trophic fractionation occurred across trophic levels, and results were comparable to published studies (for each unit effective trophic level increase there was a fractionation of +2.637‰ for $\delta^{15}\text{N}$ and +1.084‰ for $\delta^{13}\text{C}$). These results indicate that the trophic relationships established in the diet matrices reflect the observed trophic positions revealed through stable isotope analysis.

Ecopath whole-ecosystem metrics indicated that net primary productivity, trophic efficiency, ascendancy, and net primary production: respiration ratios were greater in the areas open to trawling; total system throughput and Finn Cycling Index were greater in the areas closed to trawling. Compartment-level comparisons were made using mixed trophic impacts (MTI) to determine how a small increase in the biomass of one compartment impacts all other compartments in the models. The MTI analysis for the Spring Open model indicated that a small increase in shrimp trawling in Core Sound caused large (>10%) negative impacts only on jellyfish, a bycatch species, whereas the resulting increased bycatch caused large (>10%) positive impacts on blue crabs, other smaller crabs and the crab pot fishery. The Fall Open models showed no large MTI impacts (neither positive nor negative) on any compartments with small increases in shrimp trawling or bycatch. In both Spring Open and Fall Open models, skimmer trawls caused large negative impacts on two bycatch groups: smooth dogfish and skates/rays. Thus, the greatest impacts of trawling are experienced in the spring in areas open to trawling, resulting in increases that benefit the crab pot fishery.

6.3.5 Bycatch Impacts on Stock Assessment and Prediction

Any population is a dynamic entity that will fluctuate in abundance as members enter and members leave. In a simplified example of a fish population, the entering members (or recruits) are the fish born each year and the leaving members are those removed by natural mortality and harvest (or catch). However, as indicated previously, bycatch can result in largely unknown levels of additional removals from the population. Most quantitative stock assessment techniques involve statistical analysis of catch data and, thus, require an accurate record of the entire catch to reliably estimate stock parameters such as recruitment, abundance, and selectivity. Since these parameters are crucial to forecasts of future stock conditions, any error or bias in them will lead to additional uncertainty in the predictions.

DRAFT

Before bycatch estimates can be used in stock assessments, it is necessary to convert total numbers to numbers at age and to expand estimates from known strata to unknown strata so that the entire fishing area is encompassed. Diamond-Tissue (1999) concluded that the best way to obtain unbiased estimates of bycatch is through an observer program based on randomly observed trips. A stratified random sampling design based on five geographic regions and four shrimp seasons (spring, summer, fall, winter) would be optimal. Nance (1998) notes that it is impossible to evaluate just how well a sample represents the entire fleet using objective, statistical means if vessels and trips are non-random. Based on initial estimates of variance among nets, tows, trips, months, and areas, a minimum of 60 trips per strata is needed to narrow the confidence intervals to one-half of their current range (Diamond-Tissue 1999).

Very little discard information was available in the past, so it was often assumed that discarding was a constant that could be largely ignored without causing any serious bias in assessment results (Murawski 1995). This trend is changing with the availability of additional research suggesting that while discarding may be constant in some fisheries, it is quite variable in many others; either way discards represent an unaccounted mortality in fisheries (Alverson and Hughes 1996). The challenge now lies in determining whether the additional precision gained by including discard losses justifies the expense and effort of collecting the data (Alverson et al. 1994). Since the impacts of overlooked bycatch on assessment results will vary from fishery to fishery, each case must be evaluated separately, and at least some characteristics of the bycatch must be determined.

In the most basic sense, discarded bycatch causes an underestimate of the total catch and evaluating how an assessment model responds to such an underestimate is fairly simple. It is known that responses vary among analytical techniques and depend on such factors as the age distribution of the discarded fish, the magnitude of harvest to discards, the variability and predictability of discard rates, relative year class strength, and the exploitation patterns of the involved fisheries (Alverson et al. 1994; Murawski 1995). Much of this knowledge is intuitive, and stems from understanding the interactions between input data (catch) and model outputs such as stock size and fishing mortality. Still lacking at this time are adequate bycatch estimates that could support the transition from generalized to quantitative responses. For example, if the discarded bycatch is composed of young fish and the actual removal of young fish from the population is more than that indicated by the available data then this portion of the total catch is underestimated. In generalized terms, omitting the discard data from the analysis will underestimate recruitment and, to a lesser extent, mortality rates at age. If the discarded bycatch is older fish, both numbers at age and recruitment will be underestimated and thus overall stock biomass will be underestimated as well. Quantitative responses are desirable and certainly feasible, but they require some estimate of the magnitude of the discarded bycatch.

Similar to status estimates, how discards will affect stock predictions depends on several factors, including the type of predictions being considered, variability and predictability of discard characteristics, and fishery selectivity (Alverson et al. 1994). In all situations, if discard rates cannot be predicted, then the fishery predictions will contain additional error. Short-term yield forecasts are robust if discarding and fishery selectivity are constant and predictable, but if discarding represents varying proportions of the total catch, these predictions may be impacted significantly. The impact will likely be expressed as additional uncertainty rather than as a bias (Alverson et al. 1994, Murawski 1995). According to Alverson (1994) and Murawski (1995), long-term forecasts such as equilibrium yield and spawning biomass per recruit analyses require inclusion of all sources of mortality and thus are very sensitive to discard effects. Even constant discard rates influence long-term predictions when the exploitation pattern of a fishery

DRAFT

changes, a point that can have important consequences when contemplating changes in size or mesh restrictions.

The lack of reliable discard estimates has not stopped researchers from investigating assessment impacts, but it has prevented increases in precision. Exploration of such sources as the SEAMAP database and the NMFS vessel logbook entries has provided a wide range of discard estimates for a number of fish stocks. Most assessments address the range of bycatch estimates through sensitivity analyses by comparing basic assessment results over the range of bycatch estimates and assumptions. However, estimates of the weight or number of species taken as bycatch, no matter how large or small, are meaningless without an estimate of population abundance (Diamond 2003). Those preparing or reviewing the estimates must decide which scenario seems most likely. If none of the results seem plausible, the assessment may proceed without the bycatch estimates included but with the caveat that results may be biased or contain additional uncertainty due to unknown levels of missing catch.

Unlike in the past, it is no longer acceptable to assume discards represent an unimportant removal from a stock. Under certain circumstances, discarding can and does impose uncertainty and potential bias on both estimates of current stock status and predictions of future stock conditions. This bias and error can make proper management even more difficult. While qualitative analyses of discard impacts are readily available at this time, providing the quantitative estimates that are necessary to improving stock assessments will require significant additional research and monitoring. Further, due to the extreme variation of discard characteristics, such efforts must be directed to specific fisheries and areas and must represent a long-term commitment.

6.3.6 Bycatch Impacts on the Stock Assessment of Non-Target Species

Although many species are caught as bycatch in the estuarine shrimp trawl fishery, four species, blue crab, weakfish, Atlantic croaker, and spot have, since the first studies were conducted in the 1950s and continuing to the present, accounted for the bulk of the bycatch. The bycatch of southern flounder (*Paralichthys lethostigma*) is of concern due to its overfished status. Because these five species and many other species of commercially and recreationally important finfish spend a portion of their lives in estuarine waters, bycatch in North Carolina's estuarine shrimp trawl fisheries, mainly ages 0 and 1, may have the potential to impact the stocks of these species. Natural mortality at these stages is high; however, it is believed that bycatch may adversely increase overall mortality potential (Diamond et al. 2010). Possible impacts from this increased mortality include reducing spawning stock potential and reduced yields to the fisheries (West et al. 1994). Due to the magnitude of the bycatch of these species and their importance to other commercial and recreational fisheries, a brief summary of their stock status is presented below.

However, it should be noted that resource conservation issues for these species are contained and principally evaluated in species specific management plans, from either solely North Carolina jurisdictional FMPs like blue crab and southern flounder, or coast wide ASMFC plans for weakfish, spot, and Atlantic croaker. These latter three species are also species that are covered in the North Carolina Interjurisdictional (IJ) FMP that selectively adopts management measures contained in approved federal council or ASMFC FMPs by reference as minimum standards. North Carolina in the IJ FMP acknowledges the expertise and process employed in the development of the council and ASMFC plans and that a coast wide perspective and coordinated management actions are paramount for stocks that are not within a single state's jurisdiction. So while one of the stated goals of this shrimp amendment is to "minimize harvest

DRAFT

of non-target species of finfish and crustaceans and protected, threatened, and endangered species, the extent and benefit of actions to be considered should be viewed in this broader coast wide context. Due to all the aforementioned lack of and limitations with shrimp trawl discard data, none of the ASMFC FMPs have called for additional restrictions in state's shrimp trawl fisheries.

6.3.6.1 Blue Crab Status

Reduced landings of hard blue crabs during 2000-2002 and 2005-2007, following record-high landings observed during 1996-1999 have caused concern for blue crab stocks. The 2011 fishery yielded the fifth lowest landings during the 10-year period of 2002-2011. Harvest from the Pamlico/Core sounds and tributaries increased, but continue to remain significantly below historical levels. Results of the 2011 Traffic Light Stock Assessment suggest the North Carolina blue crab stock is not overfished. However, overfishing cannot be determined at this time because data are insufficient for estimating reliable fishing mortality rates. Discard reductions of blue crabs in non-target fisheries were originally reviewed in Amendment 1 of the Blue Crab FMP; however, due to the limited amount of shrimp trawl data, discard estimates were not incorporated into the stock assessment. Amendment 2 of the Blue Crab FMP indicates there is still some concern for discards in the shrimp trawl fishery and that more fishery-dependent data are needed. The extent of delayed mortality of blue crabs in the shrimp trawl fishery was of particular concern. Amendment 2 also cites that limited tow times would help reduce mortality of sublegal crab bycatch.

The bycatch of blue crabs in the shrimp trawl fishery is of concern due to the mortality (immediate and delayed) and physical injury of culled individuals. Johnson (2006) notes mortality of crabs caught in shrimp trawls is thought to primarily to occur in the nets during trawling or the hauling back of nets. In a study of post-harvest mortality and physical injury to trawl and pot-caught crabs, McKenna and Camp (1992) found the incidence of physical injury to those crabs was similar; that is, the appendages were the most frequently damaged area. The chelipeds (pincher appendages) were the most frequently damaged appendage for both gear types; crab pot-crabs showed a greater loss than did trawl-caught crabs, 52% and 33%, respectively. There were no differences between the survival rates of damaged crabs and undamaged crabs. These findings are in agreement with those of Smith and Howell (1987), who found the appendages were the most frequently damaged structure in pot and trawl-caught American lobsters in Long Island Sound, N.Y. Additionally, Wassenberg and Hill (1989) found that 99% of the trawl-induced damage to sand crabs was restricted to the appendages.

The only observed cases of immediate mortality in crab-trawl-caught crabs occurred in June (McKenna and Camp 1992). During this trip, a large number of paper shell and soft crabs were killed in the trawling process. These findings agree with those of other investigators who found that immediate mortality in trawl-caught crustaceans was almost entirely limited to soft or paper stage individuals (Smith and Howell 1987; Wassenberg and Hill 1989).

Factors affecting the level of delayed mortality in crustaceans are temperature, exposure time, amount and level of physical injury, and total catch biomass (Smith and Howell 1987; Wassenberg and Hill 1989). Overall survival rates for trawl-caught crabs was 64%, while 93% of the crab-pot crabs survived (McKenna and Camp 1992). The effects of temperature were readily apparent; survival rates for trawl-caught crabs during the winter months were 74%, while the individuals caught in June had a 20% survival rate.

DRAFT

6.3.6.2 Weakfish Status

Weakfish are managed under the ASMFC plan as a single stock throughout their coastal range. All states from Massachusetts to Florida and the Potomac River Fisheries Commission have a declared interest in the Weakfish FMP. Responsibility for the FMP is assigned to the ASMFC Weakfish Management Board, Plan Review Team, Technical Committee, Stock Assessment Sub-Committee and Advisory Panel. The FMP for weakfish was adopted in 1985 by the ASMFC. Weakfish are currently managed under Addendum IV to Amendment #4 to the ASMFC FMP, adopted in November, 2009. Due to the depleted stock size, Addendum IV requires management measures aimed at aiding in any recovery of the weakfish stock. Addendum IV recognizes that natural mortality, rather than fishing mortality, appears to be the primary culprit in the current stock decline. As a result, the ASMFC Weakfish Management Board has implemented strict coastwide harvest limits that are intended to limit fishing pressure to aid in the recovery should conditions governing the high natural mortality subside. Amendment #4 to the FMP was designed to manage the recovered fishery and similar to Amendment #3 it had specific restrictions including: Bycatch Reduction Devices for shrimp trawls and escape panels in long haul seines, 12 inch commercial minimum size limit for all fisheries but estuarine pound net and long haul seine fisheries (seasonal 10 inch size limit), minimum mesh sizes for gill nets and trawls, and a recreational bag and size limit (currently one fish at 12 inches). According to Amendment #3 to the weakfish FMP, discard losses in the South Atlantic shrimp trawl fishery significantly increased mortality of age-0 and 1 weakfish, and both yield and spawning potential could be increased if these age classes were protected (ASMFC 1996c). BRD requirements for shrimp fisheries in the South Atlantic were introduced specifically to reduce mortality of age-0 and age-1 weakfish 30% to 40%. In addition, North Carolina is still required to maintain a closure of the area south of Cape Hatteras to flynets. One major change in Amendment #4 was an increase in the bycatch allowance for commercial fisheries from 150 pounds to 300 pounds provided that there is at least equal poundage of other species on board the vessel. In North Carolina this bycatch provision applies to gears used that do not meet the minimum mesh size requirements of Amendment #4 designed to prevent weakfish bycatch.

The 1996 stock assessment for weakfish represents one of the few examples of use of specific bycatch information in the stock assessment process. Vaughan et al. (1991) ran analyses based on different multipliers (0.0, 0.25, 0.50, 1.0) of weakfish to shrimp landings. They made the assumption that bycatch was proportional to shrimp landings and that this ratio was constant over time. However, these proportions are variable depending on location and time of year. Generally, weakfish to shrimp ratios in weight appear to range from 0.1:1 to 0.5:1.

Based on Vaughan et al. (1991), VPAs for 1982-1987 with natural mortality $M=0.3$ and without bycatch estimates, fishing mortality estimates (F) for age-0 were very small (around 0.015), while those for age-1 were much larger and increasing. However, estimates of fishing mortality at age-0 and age-1 increased values with increasing bycatch multipliers. For example, at the lowest bycatch multiplier (0.25) the estimate of F_{age} ranged from 0.3 to 0.7, a much higher value than the $F=0.015$ in the initial analysis. Initial yield-per-recruit estimates without bycatch showed almost no gain from raising the age at entry from age-0.25 to age-1, but moderate gains from age-1 to age-2 and from age-2 to age-3. However, when the bycatch multipliers entered the analysis, a significant reduction in estimated yield-per-recruit was found, and a significant gain was demonstrated from raising the age at entry from age-0 to age-1 and from age-1 to age-2. There were moderate gains from raising the age at entry from age-2 to age-3. Maximum spawning stock potential (without bycatch) showed small declines, but when the

DRAFT

bycatch multipliers were introduced, significant reductions were estimated. The 0.25 multiplier showed a small but significant gain in spawning stock potential when the age at entry was raised from age-0 to age-1 and even higher gains from increasing the age at entry to age-2.

The assumptions made in Vaughan et al. (1991) created the effect that trends in weakfish discards reflected shrimp harvest, in other words, the more shrimp caught, the more weakfish discarded. Another assumption that may be applied to weakfish stock estimates is to consider bycatch of weakfish as a function of weakfish abundance and shrimp fishing effort, but not shrimp catch. Gibson (1994) used shrimp trawl effort rather than shrimp catch and produced new estimates of weakfish bycatch in the shrimp trawl fishery. Gibson (1994) found nearly 90% of weakfish discards were age-0 fish; however, these estimates were imprecise. Discard numbers were 50% higher on average compared to a later assessment by Vaughan (1994) and showed opposite trends. Gibson's (1994) fishing mortality rates were also slightly higher than Vaughan's method and agreed with the trends in spawning stock biomass and the decline in recruitment strength.

The affect of shrimp trawl discards on the stock biomass of weakfish is still uncertain. In the 2009 ASMFC Weakfish Stock Assessment Report several methods to estimate discards were investigated, including effort based estimates, regression based estimates, and ratio extrapolation (ASMFC 2009). It was determined that there was not enough effort data for all states and years to do effort based estimations. Regression analysis was also considered inappropriate for use due to the poor fit of the predictive models. Seasonal, annual, and multi-year ratio methods were investigated as well. The multi-year estimate was found to provide the most reliable discard estimate due to the high interannual variability and large standard errors associated with the short time groupings. A positive linear relationship between the response and explanatory variable are needed for ratio extrapolation methods to work best; however, there was no evidence of such a relationship in several of the gear-species combinations investigated. It is also important to note that discard data in the southern region (North Carolina to Florida) were considered insignificant and not evaluated. Commercial discards were only evaluated for the northern region and later found to be bias by two reviewers who were concerned that the methods used to estimate discards could result in substantial uncertainty. One reviewer was concerned that discards were overestimated because multiplying the discard ratio for a given target species by total harvest of species includes harvest when that species was not the target species. The second reviewer was concerned that the gear-species combinations were too limited, missing historic fisheries with large weakfish discards. The Weakfish Technical Committee is aware of these potential sources of bias with commercial discards, citing a lack of observer data. The report also indicated that fishing mortality and discard mortality have remained low in recent years and the recent drop in weakfish productivity did not coincide with rising exploitation; however, there was a strong positive correlation between the recent rise in weakfish juvenile mortality and rising striped bass and spiny dogfish abundance (ASMFC 2009). The recent emergence of a weakfish bottleneck at age-0 is thought to be largely due to enhanced predation by these two species and not due to a surge in unreported landings and discards.

6.3.6.3 Atlantic Croaker Status

A peer-reviewed stock assessment was completed by ASMFC Technical Committee and accepted by the South Atlantic State/Federal Fisheries Management Board in August 2010. The assessment used data from both Mid- Atlantic and South Atlantic regions to produce a single, coast wide assessment, indicating that Atlantic croaker is not experiencing overfishing and is likely not overfished. Trends in independent data indicate biomass has been increasing and

DRAFT

more, older fish have been observed in the catch since the late 1980s. Absolute estimates of spawning stock biomass and fishing mortality were not given because of uncertainty in the assessment resulting from inadequate data on the magnitude of croaker discards in the South Atlantic shrimp trawl fishery. The 2010 assessment also indicates that while there are no monitoring programs in place to document the annual magnitude of discards, rough estimates of shrimp trawl discards suggest a general decline since 1995. Rough estimates of shrimp trawl discards were also used to conduct sensitivity runs of the model to determine how Spawning Stock Biomass (SSB) can be apportioned to different sources of uncertainty in the model input. Sensitivity runs of the model including rough estimates of shrimp trawl discards did not change the overall trend in SSB. Overall, the Review Panel concluded that the stock is not in trouble, noting that biomass has been on the rise, commercial catches are stable, and discards from the shrimp trawl fishery have been much reduced.

The 2004 ASMFC stock assessment also determined the stock was not overfished and overfishing was not occurring in the Mid-Atlantic region (North Carolina and north). The stock assessment showed both fishing mortality and spawning stock biomass for the Mid-Atlantic region exhibiting a cyclical trend over the time series. The Atlantic croaker stock status for the South Atlantic region (South Carolina and south) was unknown at the time and thought to make up a relatively small component of the total stock biomass. However, fishing pressure was below the target MSY and the spawning stock biomass was well above the target level. Much like the 2010 assessment, shrimp trawl bycatch was not included in the final model due to the uncertainty of the bycatch data. Model runs were completed including shrimp trawl bycatch to show the effects this fishery has on the stock even with the limited data. Sensitivity analysis evaluating the inclusion and non-inclusion of shrimp bycatch estimates, indicate that SSB_{msy} estimates are sensitive to the inclusion of Atlantic croaker caught as shrimp bycatch. However, increased SSB_{msy} estimates are also accompanied by higher total SSB estimates. The ratio of $SSB_{2002}:SSB_{msy}$ when preliminary estimates of shrimp bycatch was included indicated that the stock was unlikely to be below the threshold estimates. Also, biomass reference points from the simulation runs including shrimp trawl bycatch indicated higher SSB_{msy} values and the lower estimates of $SSB_{2002}:SSB_{msy}$ than those obtained for the base model. The range of estimates for F_{msy} (~0.4) was similar to the base model (~0.39). SSB_{msy} estimates from the simulation ranged from 48,000 to 67,000 MT with a median of 56,467 MT and were much higher than those for the base run (28,932 MT).

Diamond-Tissue (1999b) showed that by separating Atlantic croaker into different life history stages, she could examine the effects on the population of mortality at different life stages. This approach provides some insight into population changes that may be caused by bycatch. She used a stage-within age based matrix model. In this type of model, a stage-based model of the first year of life was combined with an age-based model of adults. The first year (age-0) was divided into six stages separated by biologically significant events based on major changes of morphology or habitat. Within each life stage model, she examined the population growth rate, the stable age distribution, and the elasticity (sensitivity) of the population to increases and decreases of mortality in each life stage. In order to determine elasticity of the population, baseline matrices were constructed from published and unpublished data on the life history of Atlantic croaker. Of all the data examined, only late-stage juvenile and adult mortality rates were shown to be affected anthropomorphically (fishing mortality). She then examined the trade-offs between regulating directed fisheries for adults and regulating fisheries that cause mortality on late juveniles. These simulations varied mortality from the baseline values established from data in the literature.

DRAFT

In the Atlantic and the Gulf of Mexico areas, the baseline model showed 99% of the population to be in the first year of life. The elasticity analyses showed that croaker were more sensitive to survival during age-0 than other age classes. In both regions, croaker were more sensitive to changes in fertility of age-1 fish (the age of first full reproduction) than fertility in any other year. In the analyses of other life stages, the south Atlantic population was more sensitive to fecundity than the Gulf population, but both populations were most sensitive to mortality in the oceanic larval stage than in any other stage.

By altering the late stage juvenile mortality from 10% to 200% of the baseline rate while keeping adult mortality constant, Gulf population growth rates decreased. Changing the adult mortality rates yielded similar effects. If juvenile or adult mortality was decreased, population growth rates increased. In the south Atlantic, the model was much more reactive to change. As in the Gulf, changing the mortality rate from 10% to 200% of the baseline caused population growth rates to decrease. Changing the adult mortality rate had a much larger effect on population growth rates.

Diamond-Tissue's (1999b) model results indicate that bycatch mortality at the estimated levels is not the most important factor affecting Atlantic croaker populations in the Gulf of Mexico or in the south Atlantic areas, although it can have a large negative impact on population growth rates. Both populations were most sensitive to mortality during the ocean's larval stage, followed by mortality of estuarine larvae and adults in the Gulf, and by early juvenile and adult mortality in the Atlantic. Bycatch mortality would have to be 2.5 times higher in the Gulf of Mexico and about 3.5 times higher in the south Atlantic for bycatch mortality to be the most important factor affecting population growth rate. Simulations showed that reducing late juvenile mortality by 1% and adult mortality by 3% of the baseline would stabilize the Atlantic population.

6.3.6.4 Spot Status

Commercial landings and effort have generally been decreasing in the major fisheries. This decrease accelerated in 2006 and 2007 and 2010 showed a historical low. Commercial catches in 2011 increased 64% from 2010. Catch per unit effort in the inshore gill net and offshore gillnet fisheries increased in 2011 relative to 2010. The catch per unit effort in the long haul fishery decreased in 2011 relative to 2010. Recreational landings have increased 58% from a historical low in 2010 and the mean catch per angler trip also increased. Given that spot are a short-lived species; these types of fluctuations in landings are not uncommon (Mercer 1987).

The ASMFC FMP for spot, adopted in 1987 included the states from Delaware through Florida. However, a formal coastwise spot stock assessment has not been conducted. Concerns addressed in the 1987 FMP included growth overfishing, as indicated by the dominance of unmarketable fish being landed, especially in the shrimp trawl and flynet fisheries, but also in the sciaenid pound net and long haul seine fisheries. North Carolina has addressed these concerns. North Carolina has tested bycatch reduction devices in the shrimp trawl fishery and achieved finfish reductions of 50% to 70% with little loss of shrimp. Finfish reduction devices have been required in all shrimp trawls since the fall of 1992 (15A NCAC 3J.0104) and escape panels have been required (since April 1999) in the bunt nets of long haul seines in an area south and west of Bluff Shoals in the Pamlico Sound (15A NCAC 03J.0109). The North Carolina Marine Fisheries Commission modified this rule in August 2003 to include more specific wording on installation and placement of the culling panels. Additionally, in the North Carolina flynet fishery, where a large portion of the spot catch occurs, there is a requirement for a minimum tailbag mesh of 3 1/2 inch diamond or 3 inch square. Furthermore, the state of North Carolina has banned flynet fishing in waters south of Cape Hatteras.

DRAFT

The 2010 review of the spot FMP indicates that the largest bycatch component for spot comes from the South Atlantic shrimp trawl fishery (ASMFC 2011a). The review also indicates that the non-quantifiable incidental bycatch and discard mortality of small spot in non-directed fisheries is an extremely problematic issue, citing limited discard data as one of the major problems. While the magnitude of discards from the shrimp trawl fishery is still highly uncertain, Peuser (1996) indicated that spot could account for as much as 80% of the catch by weight and 60% by number and that spot landed in these trawls are generally small and represent only one or two age classes. High priority research and monitoring recommendations listed in the 2010 ASMFC FMP review include: state monitoring and reporting on the extent of unutilized bycatch and fishing mortality on age-0 fish in fisheries that take significant numbers of spot and an evaluation of the effects of mandated bycatch reduction devices on spot catch in states with significant commercial harvest (ASMFC 2011). The 2011 Omnibus Amendment to the Interstate FMPs for Spanish mackerel, spot, and spotted seatrout states that until adequate discard estimates from the South Atlantic shrimp trawl fishery, as well as several other state fisheries that incidentally catch spot are available, a stock assessment cannot be initiated (ASMFC 2011b).

6.3.6.5 Southern Flounder Status

Based on the NCDMF 2009 stock assessment, the southern flounder stock is overfished and overfishing is occurring (Takade-Heumacher and Batsavage 2009). These findings concur with those of the 2004 stock assessment indicating that the southern flounder stock has been overfished for at least the past decade if not longer. The 2004 stock status catch-at-age indicated extremely high exploitation of age-1 and age-2 southern flounder (57% and 38% respectively), that was a concern since only 59% of age-1 and 79% of age-2 female southern flounder were sexually mature. With the addition of 1.0 million age 0-2 fish from the shrimp trawl bycatch, exploitation of juvenile southern flounder was more pronounced (19%, 52%, and 26% respectively). In absence of quantifiable observer data, the Pamlico Sound trawl survey (Program 195) was used as a proxy for estimating shrimp trawl bycatch of southern flounder in the 2006 Shrimp FMP. This data was then used to conduct a catch curve analysis and a Virtual Population Analysis (VPA) to compare the original southern flounder stock assessment with the results that included the shrimp trawl bycatch-at-age. Catch curve analysis conducted for ages 1-6 revealed that when bycatch estimates were added to the original catch-at-age model the average total mortality increased from 2.30 to 2.75, fishing mortality rate increased from 1.89 to 2.35, and fishing exploitation rate increased from 85% to 90%. Again using the same data, VPA analysis revealed that when shrimp trawl bycatch catch-at-age was added to the original data set, the fishing mortality rate did not change. At the time of the 2006 Shrimp FMP catch curve analysis and VPA indicated minimal to no differences in the affects upon the high exploitation rate of southern flounder through the fishery.

While little has changed in the availability of adequate discard data since the 2004 flounder stock assessment, aging samples from the ocean, a new January 1 birth date and a new forward calculating model (ASAP2) was used to estimated mortality and abundance. While this model was configured using discard at age matrices for the commercial gill net and recreational hook and line fishery, it was not possible to calculate discards for the shrimp trawl fishery as well as the recreational gig fishery. Thus, the current assessment could not account for all sources of removals of age-0 and age-1 fish. The inability to estimate shrimp trawl bycatch, which would consist primarily of age-0 fish, could lead to a systematic overestimation of young fish, confounding estimates of total abundance. This problem could be further compounded if there have been changes in the amount of fish caught as bycatch over time. The 2009 stock

DRAFT

assessment stresses the need for more discard information to be collected from the shrimp trawl fishery as well as other fisheries.

The 2006 Shrimp FMP took measures to address the issue of discarded sublegal flounder in the shrimp trawl fishery as directed in the approved 2004 Southern Flounder FMP recommendation 10.8.4 that stated “Recommend that the Shrimp FMP address the issue of the discard of sublegal southern flounder in the shrimp trawl fishery.” The 2006 Shrimp FMP closed upper portions of the Neuse, Pamlico, and Pungo rivers to shrimp trawling to minimize southern flounder bycatch in the shrimp trawl fishery and implemented a maximum combined 90 ft. headrope length in the mouths of the Pamlico and Neuse rivers and all of Bay River.

6.3.6.6 Summary

While the bycatch of these species has been a concern to managers since the 1950s only recently has the affect of bycatch mortality been examined. This is due in large part to the lack of adequate assessment data for these and most other species. The bycatch of weakfish in the shrimp trawl fishery has been identified as a major source of mortality for this species; however, through the use of BRDs and other management measures this mortality has been reduced. The bycatch mortality of Atlantic croaker may need to be 3.5 times higher to be the most important factor affecting population growth rate for this species (Diamond et al. 2010). It is unclear what specific impacts shrimp trawl bycatch has on the overall stock status of southern flounder given this species suppressed stock scenario. These analyses show the importance of combining adequate assessment data with the appropriate management measures to insure healthy stocks.

Obtaining unbiased and precise estimates of bycatch clearly represents a significant technical and financial challenge. However, for many target and non-target trawl species, these data may be critical to determining exploitation status and the effectiveness of management measures. The importance of discard estimates to a given species will depend on the magnitude of the discards, the fraction of the total catch represented by discards, and the variability in discard losses over time (Murawski 1995). Because of the unique nature of North Carolina’s estuarine habitats and the fact that bycatch rates vary by fishery, season, and area, North Carolina cannot depend on research efforts of the NMFS or other states in addressing bycatch losses.

While the effect that shrimp trawl bycatch has on finfish stocks is unknown, the reduction or elimination of the bycatch has a number of important implications. The reduction of fishing mortality on juvenile finfish stocks might result in more individuals recruiting into the commercial and recreational fisheries. From the commercial fisherman's perspective, less time will be spent culling the catch, fuel savings might be realized due to lower biomass in the nets, and the quality of shrimp catch should be improved. Methods and management options to reduce bycatch are discussed below.

6.3.7 North Carolina Management Strategies to Reduce Bycatch

6.3.7.1 Tailbag Mesh Size

Trawl minimum mesh size regulations are the principal method used to regulate fishing mortality on fish stocks (Smolowitz 1983). The control of net selectivity is the preferred management tool in lieu of other more stringent regulations such as temporal and spatial closures, quotas, or limited entry. The underlying principle of mesh size regulations is that undersized fish will

DRAFT

escape from the tailbag, survive, and become part of the future spawning biomass. Recent studies on the survival of fish escaping from tailbags (Main and Sangster 1988; J.T. DeAlteris, Univ. Rhode Island, Personal Communication; Simpson 1990) support the use of minimum mesh sizes as a means of reducing fishing mortality on juvenile fish. In North Carolina, the current minimum mesh size for a shrimp trawl, including the tailbag, is 1.5 inches (15A NCAC 03L .0103 (1)).

In 1949 Roelofs (1950) tested three tailbag sizes (2", 2 ¼", and 2 ½") in Pamlico Sound. Reduction rates were reported for spot, Atlantic croaker, and shrimp. Reduction rates for spot were 12.2% (2"), 42.8% (2 ¼"), and 50.5% (2 ½"). Atlantic croaker reductions were 24.8% (2"), 59% (2 ¼"), and 38% (2 ½"). Overall shrimp reduction rates were 5.6% (2"), 14.9% (2 ¼"), and 9.2% (2 ½"). In all cases, reduction rates were influenced by the size of the fish and shrimp.

The DMF conducted some preliminary tests on diamond tailbag mesh size in 1991, and square mesh tailbags in 2000. The two tailbags tested in 1991 were 1 5/8" stretched mesh (13/16" bar), and 2" stretched mesh (1" bar) tested against a 1 ½" standard stretched mesh tailbag. In 2000 a 1 ½" stretched square mesh tailbag was tested against a 1 ½" stretched mesh diamond tailbag. Results of the 1991 tests indicated that there was no apparent difference between the catches in the control net and the 1 5/8" tailbag. Tests with the 2" stretched mesh tailbag did show a difference between catch rates of spot (-46%), Atlantic croaker (-22%), total fish (-37%) and total catch (-18%). However, as was the case with the 1 5/8" tailbag not enough tows were made to test for significance differences. Tests conducted in 2000 with the 1 ½" square mesh tailbag showed a significant reduction in the catch of young of the year (YOY) weakfish (-51%), and bay whiff (-32%).

Brown (2010) compared the catch rates of modified (experimental) otter trawls in the Neuse River and Pamlico Sound. Experimental otter trawls (1 ¾" stretch mesh hung on diamond, 1 ¾" stretch mesh hung on square, 2" stretch mesh hung on square) were tested against a standard (1 ½" stretch mesh hung on the diamond) tail bag; all nets were equipped with standard TEDs. Results indicated that the catch of shrimp by weight was virtually identical in both the standard net and the experimental 1 ¾" inch stretch net that was hung on the diamond. However, testing did show a difference between catch rates of croaker (-16%), spot (-50%), flounder species (-13%) and weakfish (-2%). Tests with the 1 ¾" stretch mesh, hung on square resulted in significant reductions in croaker (-76%), spot (-77%) and weakfish (-46%). Tests with the 2" stretch mesh hung on square resulted in significant reductions in croaker (-69%), spot (-82%) and weakfish (-2%). The mean weight (kg) of bycatch in the 1 ¾" stretch mesh net hung on diamond was not significantly different from the standard net, however both the 1 ¾" stretch mesh net hung on square and the 2" stretch mesh hung on square caught significantly less bycatch compared to the standard net during the trials.

6.3.7.2 Bycatch Reduction Devices

During the 1980s the DMF and NMFS conducted studies on shrimp retention rates for various [TEDs (1985 - 1986 DMF unpublished data, and 1988 - 1989 NMFS unpublished data)], and started work on identifying means to reduce finfish bycatch in the shrimp trawl fishery (Pearce et al. 1988; Holland 1988). The 1991 Weakfish FMP recommended that South Atlantic states implement programs to reduce bycatch mortality of weakfish in their shrimp trawl fisheries by 40% by January 1, 1994. Based on results obtained during development work in 1990 and 1991 on DMF research vessels and operational testing conducted aboard a commercial trawler in 1992, the DMF required all shrimp trawlers working in state waters to equip their nets with functional fish excluders in October 1992. However, North Carolina was the only state that

DRAFT

required finfish excluders. On October 20, 1994 Amendment 2 of the weakfish FMP was passed. This amendment again required all South Atlantic states (NC-FL) to implement management measures to achieve the 40% reduction in bycatch of weakfish in the shrimp trawl fisheries by the start of the 1996 shrimping season.

Starting in 1992 DMF staff has worked with fishermen and used its own research vessel to test many different BRDs in a variety of waterbodies, seasons, and under various tidal and environmental conditions. The goal of the testing was to find devices, which maximized finfish reduction, minimized shrimp loss and meet the requirements of Amendments 1 and 2 of the weakfish FMP. The effectiveness of this gear in reducing weakfish and other fish species is a function of the size of the FFE opening and the placement of the gear in the tailbag of the trawl. A minimum opening of 5 1/2" X 6 1/2" is required for the reduction of weakfish at the mandated level. Placement in the tailbag is a function of the distance the gear is placed from the tailbag tie-off and general location in the net (top, side, or bottom). The distance from the tailbag tie-off is expressed as a ratio, BRD length/tailbag length. Where BRD length is equal to the distance from the tailbag tie-off to the opening of the FFE, and tailbag length is the length of the tailbag from the tie-off rings to the beginning of the tailbag (excluding any extension). To obtain a 40% value in weakfish reduction this ratio cannot exceed 0.65:1. Regardless of the tailbag length, the maximum mesh count cannot exceed 68 meshes from the tie-off rings. Data collected during the development of FFEs indicated that maximum reduction of weakfish was obtained when the FFE was placed 15 meshes down from to the side of the tailbag (Figure 6.8).




Figure 6.8 Diagram of Florida Fish Excluder (FFE) tested in North Carolina waters.

DRAFT

The large mesh extended funnel [LMEF (Figure 6.9)] is constructed from three sections of webbing. The forward piece is 62 meshes long, 120 meshes in circumference, 1 5/8" stretch mesh, #30 nylon twine. The center is made of 8" stretched mesh, 4 mm polyethylene, hung on the square. This section is five meshes long and 23 meshes in circumference. The rear section is similar to the first section except that it's 232 meshes long. A single hoop, constructed of 1/2" diameter plastic coated towing cable is sewn into the rear section of webbing, 4 meshes aft of the 8" webbing. This hoop is 30" in diameter. An accelerator funnel, constructed of 1 1/2", #24 depth stretched and heat set polyethylene webbing is attached to the forward section of small webbing. The funnel extends back past the 8" webbing and is reattached 4 meshes behind the hoop. Only seven meshes on top and seven meshes on the bottom are attached in the rear section. This device showed good potential in its ability to retain shrimp and exclude weakfish and other fish species. Overall this gear showed a -2% reduction in shrimp weight. Significant reduction in the weight of spot (-71%), sea mullet (-45%), Atlantic croaker (-63%), bluefish (-32%), weakfish (-50%), and total finfish (-55%) was observed with this gear.




Figure 6.9 Diagram of large mesh extended funnel BRD (LMEF) tested in North Carolina.

During the summer of 1995, a series of tests with a modified large mesh funnel excluder (LMFE; Figure 6.10) was conducted using the R/V Carolina Coast. This device consists of an extension of 4" stretched mesh, #60 nylon, hung on the square (50 meshes in circumference and 12 meshes long). Hoops of 1/2" combination cable are attached to both ends of the 4" extension. An accelerator funnel made of 1 7/8" stretched mesh, #15 nylon, runs through the 4" escapement webbing into the tailbag (15 meshes beyond the escapement webbing). The aft

DRAFT

end of the funnel is pulled tight by bungee cord attached at the top and bottom of the funnels end. The accelerator funnel is constructed from two pieces of webbing, 49 meshes (points) at the large end, 30 meshes long and cut on a 2 to 1 taper. The device was installed immediately behind the TED (mini-super shooter). Shrimp catches were reduced by 12% in the MLMEF equipped net. Significant reductions in total finfish (-24%), and total catch (-23%) weight was also observed. Since there was no reduction in weakfish weight, the accelerator funnel was modified in an attempt to increase reduction rates. The original funnel was replaced with an accelerator funnel, constructed of 1/2", #24 depth stretched and heat set polyethylene webbing cut on a 1 to 1 taper. This device was tested in Brunswick County in late August 1995. Significant reductions in the weight of weakfish (-58%), spot (-71%), and Atlantic croaker (-36%) were observed in the test net.




Figure 6.10 Diagram of modified large mesh funnel excluder (LMFE) tested in North Carolina.

From 1995 through 1996 gear development work continued using state funds. New designs developed by a local fisherman were examined for their ability to reduce weakfish. Designs tested were a 6" and 8" PVC excluder ["Sea Eagle" (Figure 6.11)]. The 6" "Sea Eagle" was tested 40 meshes above the tailbag tie-off at the top of the tailbag. Since the 6" "Sea Eagle" did not meet the minimum weakfish reduction requirement, tests were conducted with an 8" version of the device. Work with the 8" "Sea Eagle" showed that the weight of shrimp (-4.77%), weakfish (-57.80%), spot (-53.39%), Atlantic croaker (-56.70%), and total finfish (-54.33%) were significantly reduced with this gear.

In 1996, the MFC approved four BRDs for use in shrimp trawls. Proclamation SH-9-97, effective September 1, 1997, required shrimp trawlers to be equipped with one of the following approved designs: (1) a FFE measuring at least 5 1/2" x 6 1/2" (inside measurement) positioned no more than 19 meshes from the top centerline of the tailbag and located no more than 65%

DRAFT

up from the tailbag tie-off; (2) a large mesh funnel [8 or 10 inches stretched mesh; (3) a modified large mesh funnel excluder; or (4) a circular excluder constructed of PVC material measuring at least eight inches in diameter, positioned no more than 15 meshes from the top centerline and located no more than 38% up from the tailbag tie-off.

Amendment 3 to the Weakfish FMP was approved in May 1996 and changed the BRD certification requirements demonstrate a 40% reduction in catch (by number) or 50% reduction in bycatch mortality of weakfish when compared to catch rates in a naked net. Amendment 4, approved in November 2002, extended these measures.

The South Atlantic Fishery Management Council also has bycatch reduction requirements in its Shrimp FMP. Shrimp Amendment 2, approved in 1996, was consistent with Weakfish Amendment 4. However, Shrimp Amendment 6, effective in 2006 altered the Shrimp Amendment 2 BRD certification requirements creating an inconsistency with Amendment 4 of the weakfish plan. Under the 2006 amendment, the certification of any new BRD now required a reduction in the total weight of finfish by at least 30%. This inconsistency was addressed in Addendum III. This change now allows more flexible testing of BRDs, and allows the South Atlantic Council to achieve an ecosystem approach in fisheries management. On May, 11 2012 NOAA certified two new BRD devices for use in the Gulf of Mexico and the South Atlantic. Both new devices are modifications to the composite panel BRD, one of the devices adds a square mesh panel in the cod-end and the other adds a “spooker” cone inside the cod-end behind the BRD. See Appendix 2 for detailed descriptions, specifications, recommended construction and installation instructions for certified BRDs.




Figure 6.11. Diagram of “Sea Eagle” fish excluder tested in North Carolina.

DRAFT

Brown (2010) compared the effectiveness of two otter trawls equipped with a Jones-Davis BRD and a skylight panel against a standard trawl (1 ½" stretch mesh tailbag, hung on the diamond) in the Neuse River and Pamlico Sound. The Jones-Davis BRD is NMFS certified design that is similar to the expanded mesh and extended funnel BRDs, however it also has fish escape openings (windows) cut around a funnel and a webbing cone fish deflector installed behind the funnel. The skylight panel tested (4 inch stretch mesh, measuring 18 inches by 50 inches) is a commonly used Fish Escape Device (FED) that allows small fish to escape through the large mesh at the top of the net. The Jones-Davis BRD significantly reduced the mean weight of bycatch with no significant difference in the mean weights of shrimp. The Jones-David BRD was found to reduce the catch of spot by 52%. The skylight panel was also found to significantly reduce the mean weight of bycatch. There were significant reduction in the weight of spot (-12%) and flounder species (-46%) in the skylight net compared to the standard net. However, there was no significant difference in the mean weight of the shrimp catch between the net tested with the skylight panel and the standard net.

When the BRD requirements were adopted by the MFC, recreational and commercial shrimpers were considered as a single group. With the passage of the Recreational Commercial Gear License (RCGL) in 1997, recreational shrimpers were limited to a single shrimp trawl with a maximum headrope length of 26 feet and were prohibited from using mechanical retrieval methods. In 2009, mechanical retrieval was allowed however a TED was required in the trawl if mechanical retrieval was used. When testing FFEs, work was conducted aboard commercial trawlers with tow times of 60 minutes or longer. Since most RCGL holders have shorter tow times (20 minutes or less) FFEs placed 65% up from the tailbag tie-off most likely do not maximize finfish reduction. Additionally, gear testing conducted by the DMF in 1986 on the effects of light vs. heavy footrope chains on 20 foot trawls showed that bycatch of flounder, and crabs was higher in a heavily chained net while there was no difference in shrimp catches. To better reduce bycatch in RCGL shrimp trawls FFEs should be tested closer to the tailbag tie-off, and specific requirements for footrope chains should be examined.

6.3.7.3 Turtle Excluder Devices

Since 1992, the NMFS has required shrimp trawls to use TEDs to reduce the number of sea turtle strandings and incidental takes (see Protected Species section 7.2 for more information on TEDs). However, TEDs are also thought to reduce substantial amounts of bycatch as well. In Australia's northern prawn fishery, TEDs were shown to reduce the bycatch of smaller fish and invertebrates as well as reduce the number of larger sharks and rays by as much as 86% and 94%, respectively (Brewer et al. 2006). Broome et al. (2011) found that TEDs with reduced grid spacing (2 inches) was extremely effective in reducing total bycatch while maintaining minimal shrimp loss. The authors also noted that there were substantially more large rays, sharks, jelly balls and horseshoe crabs in the traditional 4 inch grid TED. Current federal law mandates that the maximum spacing between grids is four inches. In another study evaluating the performance of TEDs in the southeast Atlantic and Gulf of Mexico skimmer trawl fisheries, Price and Gearhart (2011) found that bycatch was significantly reduced for skimmers with TEDs. Reductions ranged from a mean of 40% to 98% for rays (primarily cownose rays) and 10% to 47% for finfish.

DRAFT

6.3.7.4 Alternate Gears

The development of species specific gears such as shrimp pots, pounds and cast nets could reduce finfish bycatch, minimize environmental concerns and conflicts with other fisheries, and could be more cost-effective than trawling. Even if these gears are ineffective in catching commercial quantities of shrimp, their use by recreational fishermen could result in a significant decrease in finfish bycatch.

Shrimp pots have been used in Pacific Northwest to harvest the British Columbia prawn (*Pandalus platyceros*) and in Maine to harvest northern shrimp [(*P. borealis*) Boutillier and Sloan 1987]. In 2003, the DMF became aware of the emergence of a new form of shrimp pot/trap with wings. These traps were constructed of 5/8" rigid hardware cloth and have two V-shaped wings to direct the shrimp into the traps. The wings of these pots were up to 50 feet in length and the distance between the ends of the wings measured approximately 80 feet. However, by definition these "traps" resemble pound nets more so than true pots; pound nets are defined as a trap consisting of a holding pen, one or more enclosures, lead or leaders, and stakes or anchors used to support the trap (15A NCAC 03I .0101 (3) (O)). Currently, DMF regulates shrimp traps under the same rules applied to pound nets. Thus, for a commercial shrimp pound net to be set, a permit must be issued by the DMF and the Fisheries Director shall issue a public notice of intent to consider assurance of a Pound Net Set Permit. In order for a site to be deemed suitable of a pound net set, the location shall not interfere with public navigation and be set a minimum of 300 yards from the permitted location of an existing pound net (see rule 15A NCAC 03J .0502 for full pound net permit requirements).

Brown (2006) evaluated a non-baited shrimp pound consisting of two stacked pots (18" by 18" by 36" of 5/8" mesh rigid galvanized hardware cloth) and two wings (leads) from Carteret County to Brunswick County to determine its potential as a recreational fishing gear. Brown shrimp were the predominant species, representing 96% of the total weight of the flood tide sets and 99% of the ebb tide sets. Bycatch consisted of blue crabs, white shrimp, and pinfish with mortality being extremely low. The gear was relatively inexpensive gear, easy to set up and operate. Results from this study led to the development of this gear as a RCGL gear.

Following the adoption of the 2006 Shrimp FMP, actions were taken limiting those who possess a RCGL to one shrimp pound net with each lead/wing measuring 10 feet or less with a minimum lead net mesh of 1 ½ inches, and enclosures constructed of net mesh 1 ¼ inches or greater and with all dimensions being 36 inches or less. Attendance is required at all times and all gear must be removed from the water when not being fished. The traps are most successful when set during a flood tide with one of the wings against a bulkhead or marsh shoreline. The devices are staked or anchored in place. The ends of the wings face away from the direction of the tide flow when deployed.

Sessions and Thorpe (2006) conducted a study to determine the catch potential and condition of shrimp and bycatch associated with commercial and RCGL shrimp pounds in southeastern North Carolina. The average shrimp catch rate was 4.5 lb/hour with a peak catch rate of 18.6 lb/hour. Sixteen finfish and six non-fish bycatch species were caught. Pinfish (66.4%) were the dominate fish species caught in terms of abundance, followed by menhaden (8.1%) and spot (5.3%). Commercially important finfish species (spot, croaker, pigfish, southern flounder and striped mullet) accounted for 8.1% of the total finfish bycatch by number. Blue crabs (93%) were the top non-fish by number, followed by shortfin squid (3%) and stone crabs (2.4%).

DRAFT

The overall finfish to shrimp ratio was 0.31:1 for commercial shrimp traps and 0.66: for RCGL shrimp traps. Overall finfish bycatch mortality was very low at 1.7% for commercial pounds and 0% for RCGL pounds.

The use of cast nets to harvest shrimp is a popular technique used by recreational fishermen in South Carolina and Georgia (Theiling 1988; Williams 1990), and more recently in North Carolina. Georgia also has a commercial shrimp cast netting fishery solely focused on harvesting shrimp for human consumption. Shrimp harvested by cast nets in North Carolina are typically used for bait; however a moderate percentage of the shrimp landed are thought to be consumed. Cast netting is used primarily to capture white shrimp, but may also be effective in capturing brown shrimp. In South Carolina a popular method of cast netting shrimp is to bait shrimp. In shrimp baiting, a series of poles are pushed into the bottom of shallow tidal waters. Bait balls, made from fish meal and mud, are placed at a known distance around the poles. Casting with multi- or mono-filament nets begins within minutes after baiting. In South Carolina, cast nets used in conjunction with bait balls must have a mesh size of ½ inch bar mesh (one inch stretch); there are no mesh size restrictions for shrimp landed without baiting. In Georgia, recreational and commercial cast net fishermen are restricted to a minimum mesh size of 5/8 inch bar mesh.

Currently, there is not a minimum mesh size for recreational shrimp cast nets in North Carolina. In 1992, DMF tested three different sized meshes (3/8", ½", 5/8") of cast nets in conjunction with bait balls to determine their ability to capture brown shrimp in primary and secondary nursery areas bordering Pamlico Sound (Mckenna and Clark 1993). The 5/8" bar net had the highest CPUE for brown shrimp and cast made over bait balls captured more shrimp; however, cast nets were found to be an ineffective means of harvest in this area. The Pamlico Sound has a low tidal range with circulation dominated by wind-driven currents. This lack of tidal influence could affect shrimp behavior in term of movement and feeding activity, thus making them less susceptible to baiting. The lack of suitable bait was also cited as a limiting factor. Most shrimp landed by cast nets in North Carolina are not baited. Recreational cast netting occurs in the shallow, peripheral waters of the estuaries and shallow tidal creeks. Fishing effort is typically the highest at night; however it has become more prevalent during the day in deeper areas. Cast netting for white shrimp occurs in the southern portion of North Carolina and in Core and Bogue sound. During years when white shrimp are abundant, cast netting has also become more popular in the creeks and bays throughout Dare, Hyde, Beaufort, and Pamlico counties; with limits (100 shrimp) being caught consistently (G.Judy, NCDMF. personal communication). In North Carolina, recreational fishermen using cast nets to land shrimp are limited to 48 quarts (heads on) or 30 quarts (heads off) in open waters and limited to 100 shrimp per person per day while fishing in a closed area (15A NCAC 03L .0105). Proposed changes to Rule 15A NCAC 03L .0105 replace the 100-count measurement of shrimp harvested with a cast net in closed areas to a two-quart measurement, to improve Marine Patrol Officers' safety when enforcing shrimp harvest limits. These changes have an intended effective date of June 1, 2013.

6.3.7.5 Catch Restrictions

Catch restrictions have been used by fisheries managers to maintain fish stocks, extend fishing seasons, allocate resources, and reduce bycatch. In North Carolina this method is being used to reduce the targeting of marketable finfish with shrimp trawls. From December 1 through February 28, it is unlawful to use trawl nets in internal waters to take more than 500 pounds of finfish and from March 1 through November 30 no more than 1,000 pounds of finfish may be taken (15A NCAC 3J .0104 (a) (1)). Additionally, in the Atlantic Ocean it is unlawful to possess finfish caught incidental to shrimp trawling from December 1 through March 31 unless the

DRAFT

weight of the combined catch of shrimp and crabs exceeds the weight of finfish, except that 300 pounds of kingfish may be taken south of Bogue Inlet (15A NCAC 3J .0202 (5 (a) (b))).

6.3.7.6 Harvest Seasons

Harvest seasons have been used to reduce bycatch by relegating fishing activity to times of maximum target species abundance, or by limiting activity during times of high bycatch. Currently shrimp trawling is permitted all year in North Carolina. If a specific species stock assessment indicated that measures need to be taken to reduce either the incidental or discarded catch in the shrimp trawl fishery of that species the following questions should be addressed:

- 1) How will seasons be determined?
 - a) Overall?
 - b) Area?
- 2) What criteria will be used to set seasons?
 - a) Based on historic average landings?
 - b) Maximum value?
- 3) Will allowances be made for variable conditions?
 - a) Water temperature?
 - b) Salinity?

The type of information presented in Tables 6.12 through 6.17 would provide information to answer the first two questions, while environmental data collected by the various resource agencies could be used to address the third question.

6.3.7.7 Time Restrictions

Trawl time restrictions can reduce bycatch of non-target species. In North Carolina it is unlawful to trawl for shrimp in the Atlantic Ocean off Brunswick County, 9:00 p.m. to 5:00 a.m. (15A NCAC 03J .0202 (8)). This management measure was implemented in large part to reduce the bycatch of finfish in this gear. Ingraham (2003) examined this question by conducting a study of shrimp and finfish catch rates (day vs. night) in state waters from Topsail Inlet to Little River Inlet. Data from the study showed that finfish bycatch was higher at night than during the day. Of the nine commercially important finfish species caught, southern flounder, spot, Atlantic croaker, and southern kingfish catch rates were significantly higher at night. The catch of shrimp did not vary significantly between nighttime and daytime trawling, although catches were slightly higher during the day. Limiting the number of days trawlers are allowed to fish could also limit bycatch without reducing landings. Johnson (2006) noted that twice as much shrimp were caught early in the five-day trawling week than later in the week in the estuarine shrimp trawl fishery in NC, suggesting that time restrictions could further improve the efficiency of the shrimp fishery.

6.3.7.8 Area Restrictions

Area restrictions for trawling have been used to deal with allocation, resource, habitat, and safety issues in North Carolina. During the late 1980s trawling was prohibited in Albemarle Sound and its tributaries [15A NCAC 3J .0104 (b) (3)]. This action was implemented to protect

DRAFT

the flounder gill net fishery in this area (allocation issue). Since 1978 over 124,000 acres of estuarine nursery areas have been closed to trawling to protect juvenile fish and crustaceans. MFC rule 15A NCAC 3N .0102 (a) defines Nursery Areas “as those areas in which for reasons such as food, cover, bottom type, salinity, temperature and other factors, young fish and crustaceans spend the major portion of the initial growing season.” There are approximately 77,000 acres of Primary Nurseries, 47,000 acres of Secondary Nursery areas, and 37,000 of special Secondary Nursery areas. Primary and Secondary Nursery areas are permanently closed to trawling, while Special Secondary Nursery areas can only be opened to trawling by proclamation from August 16 through May 15. In the mid 90s the sea grass beds along the Outer Banks were closed to trawling to protect this critical habitat. Over 78,000 acres of military danger zones and restricted areas are also closed to trawling for safety reasons. North Carolina has 2,220,000 acres of estuarine surface waters with approximately 1,000,000 acres (45%) closed to trawling.

6.3.7.9 Limited Entry

Limited entry methods of management restrict access to a fishery. Capping and/ or reducing fishing effort can protect the biological viability of a species and the economic integrity of the fishery. The species is protected by preventing overfishing and depletion of the stocks. The fishery is enhanced by reducing costs and increasing earnings, effectively increasing efficiency. Other benefits of limited entry programs include an incentive to conserve, more efficient management, bycatch minimization, and habitat protection. However, piecemeal implementation of limited entry programs can easily displace fishing effort from one fishery to create new problems in other areas and fisheries (Buck 1995). For bycatch reduction, limited entry systems are often used in conjunction with other management measures, such as quotas or trip limits to achieve management objectives.

DRAFT

7.0 PROTECTED RESOURCES

Protected species is a broad term that encompasses a host of species that are identified by federal or state protective statutes. The federal protective authorities are paramount and are the Endangered Species Act (ESA) and the Marine Mammal Protection Act (MMPA). Protected species in FMPs are generally discussed in relation to their implication to fisheries being prosecuted for the FMP species and that these fisheries may have an incidental take of protected species. The protected species topic herein intends to identify the principal fisheries, describe the various federal and state laws that deal with protected species, and discuss the ongoing management programs and implications of protected species interactions in the shrimp fishery.

7.1 PROTECTED RESOURCES LEGISLATION

7.1.1 Federal Endangered Species Act (ESA)

The ESA was enacted in 1973, “to provide a means whereby the ecosystems upon which endangered and threatened species depend may be conserved, (and) to provide a program for the conservation of such endangered species and threatened species.” The ESA is a comprehensive act with eighteen sections that cover many aspects of endangered species protection and management (STAC 2006).

The ESA defines a species as threatened when it is likely to become an endangered species within the foreseeable future. An endangered species is defined as any species which is in danger of extinction throughout all or a significant part of its range. A take is to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture or collect, or to attempt to engage in any such conduct (STAC 2006). Candidate species are species that appear to warrant consideration for addition to the federal ESA list. They are sometimes referred to as “species of special concern”. These species receive no substantive or procedural protection under the ESA.

Section 10 of the ESA provides for exceptions to the take prohibitions in the form of permits. These permits can be for either an intentional take or for an incidental take. Intentional take permits are intended for scientific purposes or to enhance the propagation or survival of the affected species. Incidental Take Permits (ITP) are for activities that are otherwise lawful but are expected to incidentally take a listed species. Permit holders must develop and implement conservation plans that reduce and minimize the impacts of the take. When a Section 10 permit application is reviewed and deemed appropriate, a permit is granted to authorize a specified level of takes. Along with the specified take that is authorized, the permit includes reporting requirements, and often includes other conditions that must be met (tagging, handling guidelines, data analyses, conservation plans, observer coverage, etc.).

Section 7 of the ESA relates to interagency cooperation amongst federal agencies. There are two primary provisions to this section: (1) all federal agencies shall use their authorities towards the furtherance of the goals of the ESA; (2) and each federal agency must consult with the Secretary [in practice NMFS or US Fish and Wildlife Service (USFWS)] to insure that any action funded, authorized, or carried out by the agency is not likely to jeopardize the continued existence of a listed species or result in the destruction or adverse modification of its critical habitat. Although this section relates to federal agency cooperation, it can impact state projects through a federal nexus. If a project has federal authorization, funding, or other participation, it is subject to Section 7 consultation between the federal agency and NMFS or USFWS. NCDMF

DRAFT

has received biological opinions and incidental take statements in regards to Section 7 consultations on several federally funded division research projects.

Most of the species listed as endangered or threatened fall under federal jurisdiction either with the NMFS or the USFWS. The following is a list of endangered (E) or threatened (T), or federal species of concern (FSC) species that may occur in estuarine and ocean waters of North Carolina (NCDMF 2005):

Fish

- Smalltooth sawfish (*Pristis pectinata*) E
- Shortnose sturgeon (*Acipenser brevirostrum*) E
- Atlantic sturgeon (*Acipenser oxyrinchus*) E

Reptiles

- Green sea turtle (*Chelonia mydas*) T
- Kemp's ridley sea turtle (*Lepidochelys kempi*) E
- Hawksbill sea turtle (*Eretmochelys imbricate*) E
- Leatherback sea turtle (*Dermochelys coriacea*) E
- Loggerhead sea turtle (*Caretta caretta*) T
- Northern diamondback terrapin (*Malaclemys terrapin terrapin*) FSC in Dare, Pamlico, and Carteret counties in North Carolina

Mammals

- West Indian manatee (*Trichechus manatus*) E
- Fin whale (*Balaenoptera physalus*) E
- Humpback whale (*Megaptera novaeangliae*) E
- North Atlantic right whale (*Eubalaena glacialis*) E
- Sperm whale (*Physeter catodon*) E
- Sei whale (*Balaenoptera borealis*) E

Only federally endangered or threatened species are protected by federal law.

Based on a status review and all other available information on the Atlantic sturgeon (*Acipenser oxyrinchus*), NMFS designated four separate distinct population segments (DPS) as endangered and one DPS as threatened. The Carolina and South Atlantic DPSs, both prevalent in North Carolina, were listed as endangered under the ESA on April 6, 2012.

The American eel (*Anguilla rostrata*) is currently under status review for listing as threatened by the USFWS. NMFS is also reviewing the status of the scalloped hammerhead shark (*Sphyrna lewini*) to be listed as either threatened or endangered, and Alewife (*Alosa pseudoharengus*) and blueback herring (*Alosa aestivalis*) to be listed as threatened under the ESA.

7.1.2 Marine Mammal Protection Act (MMPA)

The Marine Mammal Protection Act of 1972 was enacted in response to increasing concerns by scientists and the public that significant declines in some species of marine mammals were caused by human activities. It established a national policy to prevent marine mammal species and population stocks from declining to a point where they ceased to be significant functioning elements of the ecosystem.

DRAFT

The Department of Commerce through the NMFS is charged with protecting whales, dolphins, porpoises, seals, and sea lions. Walrus, manatees, otters, and polar bears are protected by the Department of the Interior through the USFWS. The MMPA established a moratorium on the taking of marine mammals in U.S. waters. It defines “take” to mean “to hunt, harass, capture, or kill” any marine mammal or attempt to do so. Exceptions to the moratorium can be made through permitting actions for take incidental to commercial fishing and other nonfishing activities, for scientific research, and for public display at licensed institutions such as aquaria and science centers.

The MMPA requires NMFS to categorize each commercial fishery into one of three categories based upon the level of serious injury and mortality to marine mammals that occurs incidental to each fishery. Category I fisheries pose the greatest threat and Category III fisheries the least threat. The category in which a fishery is placed determines whether fishermen are subject to certain provisions of the MMPA, such as registration, observer coverage and take reduction plan (TRP) requirements. According to the 2011 List of Fisheries created by NOAA, the Atlantic southeastern shrimp trawl fishery is considered to be in Category II (occasional mortality or serious injury) due to interactions with the bottlenose dolphin (NOAA 2010).

7.1.3 North Carolina Endangered Species Act (Chapter 113 Article 25)

Listing of protected species from a state perspective lies with North Carolina Wildlife Resource Commission (NCWRC) (NC General Statutes - Chapter 113 Article 25). The NCWRC compiled state lists of animals deserving protection over 20 years ago based on guidance from Scientific Councils on mammals, birds, reptiles, amphibians, freshwater fishes, mollusks, and crustaceans. Endangered, Threatened, and Special Concern species of mammals, birds, reptiles, amphibians, freshwater fishes, freshwater and terrestrial mollusks, and crustaceans are protected by state law. Protection for crustaceans and certain venomous snakes was enacted in 2002. However, state law does not allow for protection of invertebrate groups other than mollusks and crustaceans.

Under the state Endangered Species Act the NCWRC has the following powers and duties:

- To adopt and publish an endangered species list, a threatened species list, and a list of species of special concern, as provided for in G.S. 113-334, identifying each entry by its scientific and common name.
- To reconsider and revise the lists from time to time in response to public proposals or as the Commission deems necessary.
- To coordinate development and implementation of conservation programs and plans for endangered and threatened species of wild animals and for species of special concern.
- To adopt and implement conservation programs for endangered, threatened, and special concern species and to limit, regulate, or prevent the taking, collection, or sale of protected animals.
- To conduct investigations to determine whether a wild animal should be on a protected animal list and to determine the requirements for conservation of protected wild animal species.

DRAFT

- To adopt and implement rules to limit, regulate, or prohibit the taking, possession, collection, transportation, purchase or sale of those species of wild animals in the classes Amphibia and Reptilia that do not meet the criteria for listing pursuant to G.S. 113-334 if the Commission determines that the species requires conservation measures in order to prevent the addition of the species to the protected animal lists pursuant to G.S. 113-334. This subdivision does not authorize the Commission to prohibit the taking of any species of the classes Amphibia and Reptilia solely to protect persons, property, or habitat; to prohibit possession by any person of four or fewer individual reptiles; or to prohibit possession by any person of 24 or fewer individual amphibians.

The NCWRC develops conservation plans for the recovery of protected wild animal species, using the procedures set out in Article 2A of Chapter 150B of the General Statutes. The North Carolina Natural Heritage Program inventories, catalogues, and supports conservation of the rarest and the most outstanding elements of the natural diversity of our state. These elements of natural diversity include those plants and animals which are so rare or the natural communities which are so significant that they merit special consideration as land-use decisions are made.

Species that appear on the 2010 Natural Heritage Program List of the Rare Animal Species of North Carolina that may interact with shrimp trawls, skimmer trawls and channel nets include the loggerhead sea turtle (T), leatherback sea turtle (E), hawksbill sea turtle (E), Kemp's ridley sea turtle (E), and green sea turtle (T).

7.2 SPECIES THAT MAY INTERACT WITH THE SHRIMP FISHERY

Of the federal and state protected species listed above, only bottlenose dolphins, and sea turtles interact with the shrimp fishery. Otter trawls and skimmer trawls are the predominant gear in the shrimp fishery. Both trawls are active gears that focus on the estuarine bottom, and are restricted to areas without submerged aquatic vegetation; interactions with protected species are plausible. Channel nets used less extensively in the shrimping fishery are a passive gear and use tide flow and current to fish. There is no information on interactions with protected species and channel nets.

7.2.3 Bottlenose Dolphin

The bottlenose dolphin (*Tursiops truncatus*) inhabits temperate and tropical waters throughout the world. According to the 2009 U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessment (Waring 2009) nine bottlenose dolphin stocks have been identified in the nearshore waters of the Western North Atlantic. Two of these stocks are found in North Carolina estuaries and are identified as the Northern North Carolina Estuarine System Stock and the Southern North Carolina Estuarine System Stock. Bottlenose dolphins have been observed throughout the year in North Carolina estuarine waters but will migrate offshore when water temperatures fall below 10° C.

A marine mammal species is designated as depleted if it falls below its optimum sustainable population. The MMPA requires that a Take Reduction Team (TRT) be convened for the purpose of recommending measures for inclusion in a TRP to promote recovery of a depleted stock. The Bottlenose Dolphin Take Reduction Team (BDTRT) was convened in November 2001 and was made up of fishermen, managers, scientists, and environmental groups. The BDTRT focused on reducing serious injuries and deaths of coastal bottlenose dolphins incidental to several east coast fisheries including: the North Carolina inshore gill net, Southeast

DRAFT

Atlantic gill net, Southeastern U.S. shark gill net, U.S. Mid-Atlantic coastal gill net, Atlantic blue crab trap/pot, Mid-Atlantic haul/beach seine, North Carolina long haul seine, North Carolina roe mullet stop net, and Virginia pound net. In April 2006, NMFS published a [final rule](#) implementing the BNDTRP effective May 26, 2006 (FR Doc. 06-3909 Filed 4-25-06). Shrimp trawls were not included in the final rule.

Bottlenose dolphins are occasionally captured or entangled in various kinds of fishing gear including gill nets, seines, long-lines, crab pot lines, and shrimp trawls. The NOAA List of Fisheries classifies U.S. commercial fisheries into one of three Categories based on the level of incidental mortality or serious injury of marine mammals (NOAA 2010). Category I are fisheries with frequent incidental mortality or serious injury; Category II are fisheries where occasional incidental mortality or serious injury; and Category III are fisheries with a remote likelihood of/known incidental mortality or serious injury. Several trawl fisheries, including the southeastern shrimp trawl fishery, were elevated in 2010 from a Category III fishery to a Category II fishery in the List of Fisheries (LOF). This listing is based on interactions reported through observer reports, stranding data, and fisheries research data with multiple marine mammal stocks (<http://www.nmfs.noaa.gov/pr/interactions/lof/>) and is updated annually.

In NC there has been one known take in the lazy line of a relocation trawl and several strandings with evidence indicative of a trawl interaction. There have not been any interactions with skimmer trawls or channel nets. (Barbie Byrd, NMFS, personal communication, November 2011). No further information was provided by NMFS concerning NC dolphin interaction data.

Owners of vessels or gear engaging in a Category I or II fishery, are required under 50 CFR 229.4 to obtain a marine mammal authorization by registering with the Marine Mammal Authorization Program (MMAP). Upon receipt of a completed registration, NMFS will issue vessel or gear owners a decal to display on their vessels and an authorization certificate that the operator must possess while fishing. Fishers participating in a Category I or II fishery are required to accommodate an observer onboard your vessel(s) upon request (50 CFR 229.7) and are required to comply with any applicable take reduction plans. Currently, NMFS does not have a take reduction plan for the southeastern U.S Atlantic or the Gulf of Mexico Shrimp Trawl Fishery (<http://www.nmfs.noaa.gov/pr/interactions/lof/>).

7.2.4 Sea Turtles

Sea turtles are air-breathing reptiles with streamlined bodies and large flippers which inhabit tropical and subtropical ocean waters throughout the world. Of the seven species of sea turtle worldwide, five occur in North Carolina. They include the Kemp's ridley sea turtle (*Lepidochelys kempii*), hawksbill sea turtle (*Eretmochelys imbricate*), leatherback sea turtle (*Dermochelys coriacea*), green sea turtle (*Chelonia mydas*), and the loggerhead sea turtle (*Caretta caretta*). Although sea turtles live most of their lives in the ocean, adult females must return to land to lay their eggs on sandy beaches. They often migrate long distances between foraging grounds and nesting beaches. Kemp's ridley, green, and loggerhead sea turtles are known to move into North Carolina coastal waters as large juveniles to forage on crustaceans, mollusks, or grasses (STAC 2006). The loggerhead and green sea turtles are federally listed as threatened, while the others are listed as endangered.

Hawksbill turtles have been reported off the coast of North Carolina during the months of June, July, October and November. This species of turtle prefers shallow coastal water with depths not greater than 66 feet. Preferred habitat includes coral reefs, rocky bottoms, reefs, and coastal lagoons. Adult hawksbills primary food source are sponges, but they also eat urchins,

DRAFT

algae, barnacles, mollusks, jellyfish, and fish. Hawksbills exhibit a wide tolerance for nesting substrate type and nests are typically placed under vegetation. Nesting occurs principally in Puerto Rico and the U.S. Virgin Islands but does occur in the southeast coast of Florida and the Florida Keys. The largest threat to the hawksbill is the loss of coral reef habitat. The extent to which hawksbills are killed or debilitated after becoming entangled in marine debris has not been quantified, but it is believed to be a serious and growing problem. Hawksbills (predominantly juveniles) have been reported entangled in monofilament gill nets, fishing line, and synthetic rope. Hawksbills are incidentally taken by several commercial and recreational fisheries. Fisheries known or suspected to incidentally capture hawksbills include those using trawls, gill nets, traps, driftnets, hooks, beach seines, spear guns, and nooses (NMFS 1993b). There were no strandings reported of hawksbill sea turtles in North Carolina between 1991 and 1999, but there were nine between 2001 and 2010 (NCWRC/NMFS Sea Turtle Stranding and Salvage Network [STSSN] data).

The leatherback sea turtle is the largest turtle in the world and has a worldwide distribution in tropical and temperate waters. This species is found off the coast of North Carolina from April to October with occasional sightings into the winter. The main prey species of leatherbacks are jellyfish and tunicates and occur almost exclusively in ocean waters (STAC 2006). There is one record of a NC nesting site at Cape Lookout in 1966 (Lee and Socci 1989), and an additional nesting site was reported near Cape Hatteras in 2000. Leatherbacks become entangled fairly often in longlines, fish trap, buoy anchor lines, and other ropes and cables (NMFS 1992). Prescott (1988) implicated entanglement in lobster pot lines in 51 of 57 adult leatherback strandings in Cape Cod Bay, Massachusetts from 1977 to 1987. Between 1990 and 2000, there were 12 reported leatherback strandings in North Carolina, between 2001 and 2005 there were 75, and from 2006 through 2011, there have been 23 reported strandings (www.seaturtle.org).

The Kemp's ridley sea turtle occurs primarily in the Gulf of Mexico, but they also occur along the Atlantic coast as far north as New England. Juveniles occur year-round within the sounds, bays, and coastal waters of North Carolina. Adult Kemp's ridley turtles are primarily a bottom feeder; feeding on crabs, shrimp, urchins, starfish, jellyfish, clams, snails, and squid. Incidental take by shrimp trawls has been identified as the largest source of mortality with between 500 and 5,000 killed annually (NMFS 1993a). Manzella et al. (1988) estimated that 0.2% of the juvenile Kemp's ridleys killed by fishing gear were killed as a result of interaction with crab pots. In North Carolina 17% of the sea turtle strandings between 1990 and 2000 were Kemp's ridleys (WRC/NMFS STSSN; 1990-2000). From 2001 through 2011, there have been 785 strandings in North Carolina (www.seaturtle.org).

The green sea turtle has a circumglobal distribution in tropical and subtropical waters. In U.S. Atlantic waters, it occurs around the Virgin Islands and Puerto Rico and from Texas to Massachusetts. Green turtles are sighted in oceanic waters and within the sounds of North Carolina during the period from May through October. Due to their food preference for submerged aquatic vegetation, adult green turtles are normally found in lagoons, bays, and tidal inlets. No major nesting sites are located along the U.S. coastline; however, limited annual nesting occurs in Florida from April to July. From 1979 to 1989, there were two reported (1987, Baldwin Island and 1989, Cape Hatteras) and one confirmed (1979, Camp Lejeune) nesting sites in North Carolina. In 2009, there were three nests in North Carolina and 2010; there were 18 green turtle nests (NCWRC Sea Turtle Nest Monitoring System data). In 1992, NMFS finalized regulations to require the use of Turtle Excluder Devices (TEDs) in shrimp trawl fisheries. A significant threat to the green turtle continues to be fishing gear, primarily gill nets, but also trawls, traps and pots, and dredges. Green sea turtles have been recovered entangled

DRAFT

in trap lines with the trap in tow (NMFS 1991a). Strandings have drastically increased since 2008. From 1991 to 2000, green turtles accounted for 18% of the sea turtle strandings in North Carolina and between 2001 and 2010 they make up 32% of total strandings (WRC/NMFS STSSN).

The loggerhead sea turtle has a subtropical (and occasionally tropical) distribution, including continental shelves and estuaries along the margins of the Atlantic, Pacific, and Indian oceans. It is rare or absent far from mainland shores. The loggerhead turtle is the most common sea turtle in North Carolina (STAC 2006) and is present throughout the year, with peak densities occurring from June to September. The loggerhead turtle diet includes algae, seaweeds, horseshoe crabs, barnacles, various shellfish, sponges, jellyfish, squid, urchins, and fish. Nesting occurs along the U.S. Atlantic coast from New Jersey to Florida, however, the majority of nesting activity occurs from South Carolina to Florida. In North Carolina, nesting activity has been reported from April to September. The highest nesting densities are reported south of Cape Lookout. In 2010, there were 847 loggerhead turtle nests in North Carolina (WRC Sea Turtle Nest Monitoring System data). The primary threat to loggerhead turtle populations worldwide is incidental capture in fishing gear, primarily in longlines and gill nets, but also in trawls, traps and pots, and dredges. Loggerhead turtles account for over half of the sea turtle strandings in North Carolina (WRC/NMFS STSSN).

7.2.4.1 Sea Turtles and the Shrimp Fishery

Shrimp and flounder trawlers have been required to use TEDs since 1992. Since 2007, NOAA Fisheries has required fishing vessels that are identified through an annual determination process to take observers at NOAA Fisheries request. The NMFS prepared an Environmental Impact Statement (EIS) evaluating TEDs for use in skimmer trawls due to the non-compliance of the tow time requirements in the Gulf of Mexico. NMFS is also concerned about the increase in skimmer trawl vessel and gear size and potential impacts to sea turtles. Therefore, the NOAA Fisheries is reevaluating the efficacy of turtle conservation requirements associated with the skimmer trawl fishery. In addition, NMFS is concerned about compliance issues with TED requirements in the shrimp otter trawl fishery and have noted a variety of compliance issues ranging from lack of TED use, TEDs installed incorrectly, and TEDs sewn shut. Therefore NMFS is also considering additional management measures of the shrimp trawl fishery (NOAA 2011). During the required scoping meetings, NCDMF commented that there has been a 35% decline in skimmer trawl trips and pounds of shrimp landed by skimmer trawls were down 33%. Unlike the Gulf of Mexico skimmer trawl fleet, North Carolina has not seen an increase in vessel size. Observed increases in the Gulf gave the NMFS a reason to address the problem on non-compliance and impacts of larger boats in the Gulf of Mexico, not in North Carolina where 55-minute tow times are still sufficient (David Taylor, NCDMF, personal communication). NMFS held a public hearing on TED requirements in the skimmer trawl in June 2012. It is expected that the new regulation will be in place by March 2013.

The Sea Turtle Advisory Committee (STAC) was formed in 2003 by the North Carolina Marine Fisheries Commission (NCMFC) in response to continuing problems with sea turtle interactions in fisheries throughout the North Carolina coast. Their objective was to develop solutions for the reduction of sea turtle interactions in commercial and recreational (hook and line) fishing gear, while maintaining economically viable fisheries throughout the estuarine waters of North Carolina. Over a three year effort, the STAC identified and categorized different fishing gears. Shrimp trawls were identified as gears of primary concern with relation to sea turtle incidental catch throughout North Carolina. Skimmer trawls, butterfly nets and channel nets were

DRAFT

identified as gears of other concern, while gears of no concern used in the shrimp fishery were cast nets and dip nets.

Recommendations were provided to the NCMFC to implement observer coverage for multiple fisheries of either primary or other concern was made in order to gather information where it is limited. The STAC also supported continued efforts for gear modification and testing with the objective of reducing sea turtle interactions (STAC 2006).

Shrimp Trawls:

1. Determine and enforce TED compliance throughout North Carolina through the NC MFC creating a rule change or authorizing proclamation authority to the director of NCDMF to provide state authority to enforce TED compliance. As this is a federal regulation, initially an estimate of current compliance needs to be obtained. Following this, the NCDMF may opt to increase effort to ensure compliance with TED regulations.
2. Support turtle resuscitation education and TED education.
3. Add statewide observer coverage. The level of this coverage should have a minimum goal of 2% of the total effort by area. Coverage should increase (~10%) in areas where sea turtle interactions are occurring.

Butterfly Net, Channel Net, Skimmer Trawl:

1. Implement observer coverage. The level of this coverage should have a minimum goal of 2% of the total effort by area. Coverage should increase (~10%) in areas when/where sea turtle interactions are occurring.
2. Provide educational information on sea turtle resuscitation and reporting requirements for unharmed/injured /dead turtles.

7.2.4.2 NCDMF Programs

An agreement was established in 1979 with the WRC to exercise regulatory jurisdiction over any species of sea turtle, and their eggs and nests, consistent with designation of such species as endangered or threatened by the USFWS. In 1980, the NCMFC established a Sea Turtle Sanctuary off the coast of North Carolina to protect nesting beaches (NC Fisheries Rule – 15A NCAC 03R.0101). In 1983, proclamation authority was given to the director of NCDMF by NCMFC to close areas to protect endangered/threatened species (NC Fisheries Rule-15A NCAC 03I.0107). In 1989, an addition was made to the Marine Recreational Fisheries Statistics Survey (MRFSS) program to include a sea turtle sightings query on the survey form.

In the latter part of 2010, DMF reallocated funds to establish the Protected Resources Section within the division and obtained funding to support a statewide at-sea observer program for the gill net fishery. The new Protected Resources Section will be the lead for division actions involving protected species such as at-sea observer programs, marine mammal stranding responses and marine mammal take reduction teams, and other protected species issues that may arise (Dee Lupton, NCDMF personal communication).

Marine mammal stranding response along the central North Carolina coast, transitioned from North Carolina State University Center for Marine and Science Technology to the NCDMF in

DRAFT

October of 2010. This project is funded year to year from the John H. Prescott Marine Mammal Rescue Assistance Foundation, pending successful proposal review and acceptance. A full-time stranding coordinator was hired and stranding personnel have responded to numerous marine mammal strandings. North Carolina stranding response is divided into four areas: UNC Wilmington personnel respond to all strandings in the southern part of the state up to and including Camp LeJeune; NCDMF stranding personnel respond to strandings from Hammocks Beach State Park to Cape Lookout National Seashore and in Albemarle and Pamlico sounds; Cape Hatteras (CAHA) National Seashore stranding personnel respond to strandings in CAHA National Seashore, and North Carolina Department of Environment and Natural Resources (DENR) personnel respond to strandings from CAHA north to the VA border. Stranding personnel conduct outreach by giving public seminars at marine mammal meetings, local museums, Universities, and classrooms. Stranding personnel disseminate results and tissue samples from stranded animals to collaborating researchers and agencies.

The NCDMF observer program began in 1999 when the sea turtle stranding network noted significant increases in sea turtle strandings in the southeastern portion of Pamlico Sound. The purpose of these observations was to begin the process of characterizing effort, catch, and bycatch by area and season in various fisheries. In addition, this program was established to monitor fisheries for the potential of protected species bycatch. The data collected is used for fisheries management decisions, stock assessments, and conservation efforts for protected species. Currently, the observer program primarily focuses on large mesh gill nets but data are also being collected in small mesh gill nets and recreational hook and line. In addition sampling has just begun in long haul seines and channel nets. Data collections from observer trips includes: date, location, unit, time, season, gill net description (net length, number of net shots, mesh size, presence/absence of tie downs, vertical mesh height, hang ratio), soak time and water depth. Additionally, environmental parameters (wind, tide stage and water quality data) are collected when feasible. Total catches of target species are estimated and final disposition (kept or discarded) is recorded. Sea turtle interaction information includes species, condition, tag numbers, and final disposition. All interactions involving protected species are documented. All observers are required to adhere to these data collection parameters.

The NCDMF gear development program has provided observation data from shrimp trawls through some limited characterization work of shrimp trawlers. A study from 2009 was on the near-shore commercial shrimp trawl fishery off the southern shores of North Carolina from Carteret County to Brunswick County. This study collected relative effort and discard information on weakfish (*Cynoscion regalis*), spotted sea trout (*Cynoscion nebulosus*), spot (*Leiostomus xanthurus*), Atlantic croaker (*Micropogonias undulatus*), bluefish (*Pomatomus saltatrix*), Atlantic menhaden (*Brevoortia tyrannus*), southern flounder (*Paralichthys lethostigma*), and striped mullet (*Mugil cephalus*) as well as other species of federally and state managed species of finfish. There were three observed interactions with loggerhead sea turtles in this study. All three sea turtles were taken in the try net which are small trawls equipped with small doors, no TEDs nor BRDs, and is used to sample areas prior to setting the main trawls and to monitor the catch rates during tows. The try nets had tow times of approximately 10 minutes. All three sea turtles were released in good condition. There were no observed sea turtle interactions observed in the main nets.

Another characterization study of the shrimp trawl fishery was in Pamlico Sound in 2010 and also provided observation data. Similar information was collected regarding effort and discard of recreationally and commercially important finfish. There were no observed sea turtle interactions observed during this study.

DRAFT

In the fall of 2010, the MFC reestablished the STAC (Sea Turtle Advisory Committee) to address sea turtle bycatch. The duties of the reestablished STAC include but are not limited to: reviewing observer reports, devising means for fishermen to report turtle interactions, assisting with fishermen education, determining measures to reduce the incidental take of sea turtles, monitor observer program issues, and reviewing all future ITP provisions and take calculations prior to formal application to NMFS. The STAC will provide recommendations and guidance to the NCMFC and NCDMF in addressing protection of sea turtles in North Carolina.

Since the 1970s, the NCDMF has been proactive in developing ways to minimize impacts to threatened and endangered marine species. The NCDMF works closely with NMFS and other state and federal agencies to develop regulations that minimize impacts to protected species while trying to allow the prosecution of many economically important fisheries.

8.0 ECONOMIC STATUS

8.1 COMMERCIAL FISHERY

8.1.1 Harvesting sector

8.1.1.1 Ex-vessel value and price

The state’s trip ticket program began in 1994 when it was mandated that all commercial landings be reported to DMF. Prior to this time, landings were reported through a sampling program. Reporting the ex-vessel value of the landings continues to remain optional. It is useful in economic analyses to tie the value of annual landings back to an established baseline to control for the effects of inflation. Changes in landings values from year to year can be more clearly understood after removing the influence of changing dollar values over time. To do so, nominal values are adjusted by the Consumer Price Index (CPI) in an attempt to remove the effects of inflation over time.

The annual nominal ex-vessel value of shrimp landings typically has been volatile with sizable changes between years. The lowest nominal value of the catch was \$3.5 million in 1972. The highest nominal value for shrimp landings was \$25.4 million in 2000. Relatively speaking, 1981 represented a 69% drop in the value of landings from 1980. However, the fishery rebounded in 1982 with a 210% increase in the nominal value of landings over 1981. The value of the fishery dropped by 53% in 2001 from the record high value observed in 2000. In 2002, the value increased 54% over the 2001 value, but it remained considerably lower than the 2000 value. The nominal value hit a 20 year low in 2005 (\$4.4 million), dropping 50% over the previous year’s value; however, the fishery recovered to over \$19.2 million in 2008. The nominal ex-vessel value of landings in 2010 was \$10.7 million (Figure 8.1, Table 8.1).




Figure 8.1. Value of shrimp landings in North Carolina, 1972 – 2010 (DMF Trip Ticket Program).

DRAFT

The CPI inflation-adjusted figures (deflated to the value of a dollar in 1972) typically show less volatility. Nonetheless, significant volatility from year to year can be seen in the landings values from 1978 to 1987. With a few exceptions, the total inflation adjusted value of landings hovered around the \$2 – \$8 million mark from 1972 until 2000. The inflation adjusted value of annual landings has been in an overall downward trend in recent years. The inflation adjusted value of landings in 2005 was less than \$1million, lower than any year in over 30 years. The inflation adjusted ex-vessel value of shrimp landings in 2010 was approximately \$2 million (Figure 8.1, Table 8.1).

Changes in annual values can largely be attributed to three major causes; the number of pounds landed, price per pound received by fishermen, and in recent years, the impacts of imports. The recent history of imports and their impact on the price of shrimp is further discussed in section 8.1.1.6 of this document.

The average nominal price per pound paid to the fisherman generally rose between 1972 and 1982 (Figure 8.2, Table 8.1), rising from a low of \$0.64 in 1972 to \$2.34 in 1982. From 1983 through 1994, the price per pound fluctuated between a high of \$2.61 in 1994 and a low of \$1.73 in 1991. From 1994 to 2000, the price per pound averaged just below \$2.50 per pound. However, since 2000, the nominal price per pound paid to fishermen exhibited a decreasing trend and dropped to a low of \$1.58 in 2009.




Figure 8.2. Average price per pound of shrimp landings in North Carolina, 1972 – 2010 (DMF Trip Ticket Program).

The trend in price per pound received by fishermen becomes clearer when one takes into account the impact of inflation. The inflation adjusted prices show that with the exception of 1979, the average inflation adjusted price of shrimp was under \$1.00 per pound until 1982. Since 1983 there has been a declining trend in the average price per pound. The lowest inflation adjusted price of \$0.31 in 2009 is 52% lower than the price received in 1972, and represents the lowest inflation adjusted price received per pound in over 30 years (Figure 8.2, Table 8.1).

DRAFT

Table 8.1 shows a summary of the data presented in section 8.1.1.1 indicating by year, the number of pounds of shrimp landed, nominal values, the inflation adjusted value, nominal price per pound, inflation adjusted price per pound, and the rate of change from one year to the next for all years in which data were available since 1972.

DRAFT

Table 8.1. Detail values of pounds landed, total value, inflation adjusted value, price per pound, and percent change from year to year for shrimp landed in North Carolina, 1972 – 2010 (DMF Trip Ticket Program).

Year	Pounds Landed	% Change in Pounds	Nominal Value	% Change in Nominal Value	Inflation Adjusted Value	% Change in Inflation Adjusted Value	Nominal Price per Pound	% Change in Nominal Price per Pound	Inflation Adjusted Price per Pound	% Change in Inflation Adjusted Price Per Pound
1972	5,563,261		\$3,549,492		\$3,549,492		\$0.64		\$0.64	-
1973	5,003,417	-10%	\$4,738,223	33%	\$4,460,759	26%	\$0.95	48%	\$0.89	40%
1974	8,440,203	69%	\$4,606,363	-3%	\$3,905,598	-12%	\$0.55	-42%	\$0.46	-48%
1975	5,163,610	-39%	\$5,053,944	10%	\$3,926,670	1%	\$0.98	79%	\$0.76	64%
1976	6,642,713	29%	\$8,171,394	62%	\$6,002,887	53%	\$1.23	26%	\$0.90	19%
1977	5,600,329	-16%	\$7,239,080	-11%	\$4,993,293	-17%	\$1.29	5%	\$0.89	-1%
1978	2,960,762	-47%	\$3,883,836	-46%	\$2,489,944	-50%	\$1.31	1%	\$0.84	-6%
1979	4,941,240	67%	\$9,728,917	150%	\$5,601,498	125%	\$1.97	50%	\$1.13	35%
1980	9,823,490	99%	\$17,184,994	77%	\$8,717,630	56%	\$1.75	-11%	\$0.89	-22%
1981	2,557,426	-74%	\$5,295,209	-69%	\$2,434,981	-72%	\$2.07	18%	\$0.95	7%
1982	7,027,164	175%	\$16,411,472	210%	\$7,108,803	192%	\$2.34	13%	\$1.01	6%
1983	6,115,278	-13%	\$13,564,846	-17%	\$5,692,877	-20%	\$2.22	-5%	\$0.93	-8%
1984	5,046,163	-17%	\$10,482,761	-23%	\$4,217,319	-26%	\$2.08	-6%	\$0.84	-10%
1985	11,683,427	132%	\$21,130,303	102%	\$8,208,612	95%	\$1.81	-13%	\$0.70	-16%
1986	6,162,438	-47%	\$13,934,191	-34%	\$5,314,317	-35%	\$2.26	25%	\$0.86	23%
1987	4,416,636	-28%	\$8,178,180	-41%	\$3,009,225	-43%	\$1.85	-18%	\$0.68	-21%
1988	8,139,190	84%	\$16,509,108	102%	\$5,833,311	94%	\$2.03	10%	\$0.72	5%
1989	8,922,932	10%	\$15,620,436	-5%	\$5,265,599	-10%	\$1.75	-14%	\$0.59	-18%
1990	7,839,457	-12%	\$15,885,027	2%	\$5,080,292	-4%	\$2.03	16%	\$0.65	10%
1991	10,740,936	37%	\$18,586,613	17%	\$5,704,262	12%	\$1.73	-15%	\$0.53	-18%
1992	5,496,019	-49%	\$10,859,283	-42%	\$3,235,339	-43%	\$1.98	14%	\$0.59	11%

DRAFT

Table 8.1. (continued)

Year	Pounds Landed	% Change in Pounds	Nominal Value	% Change in Nominal Value	Inflation Adjusted Value	% Change in Inflation Adjusted Value	Nominal Price per Pound	% Change in Nominal Price per Pound	Inflation Adjusted Price per Pound	% Change in Inflation Adjusted Price Per Pound
1993	6,778,999	23%	\$13,590,604	25%	\$3,931,400	22%	\$2.00	1%	\$0.58	-1%
1994	7,292,489	8%	\$18,996,565	40%	\$5,358,005	36%	\$2.60	30%	\$0.73	27%
1995	8,669,100	19%	\$20,317,986	7%	\$5,572,781	4%	\$2.34	-10%	\$0.64	-13%
1996	5,271,273	-39%	\$13,373,962	-34%	\$3,562,980	-36%	\$2.54	8%	\$0.68	5%
1997	6,988,825	33%	\$18,204,849	36%	\$4,741,201	33%	\$2.60	3%	\$0.68	0%
1998	4,636,343	-34%	\$10,856,450	-40%	\$2,784,047	-41%	\$2.34	-10%	\$0.60	-11%
1999	9,004,535	94%	\$22,094,489	104%	\$5,543,515	99%	\$2.45	5%	\$0.62	3%
2000	10,334,915	15%	\$25,405,916	15%	\$6,167,057	11%	\$2.46	0%	\$0.60	-3%
2001	5,254,214	-49%	\$11,911,070	-53%	\$2,811,309	-54%	\$2.27	-8%	\$0.54	-10%
2002	9,969,026	90%	\$18,364,776	54%	\$4,267,080	52%	\$1.84	-19%	\$0.43	-20%
2003	6,167,371	-38%	\$10,939,078	-40%	\$2,485,073	-42%	\$1.77	-4%	\$0.40	-6%
2004	4,880,817	-21%	\$9,462,853	-13%	\$2,093,951	-16%	\$1.94	9%	\$0.43	6%
2005	2,357,516	-52%	\$4,409,124	-53%	\$943,683	-55%	\$1.87	-4%	\$0.40	-7%
2006	5,736,649	143%	\$9,141,435	107%	\$1,895,397	101%	\$1.59	-15%	\$0.33	-17%
2007	9,537,230	66%	\$17,905,334	96%	\$3,609,703	90%	\$1.88	18%	\$0.38	15%
2008	9,424,168	-1%	\$19,245,921	7%	\$3,736,499	4%	\$2.04	9%	\$0.40	5%
2009	5,407,708	-43%	\$8,527,714	-56%	\$1,661,524	-56%	\$1.58	-23%	\$0.31	-23%
2010	5,955,335	10%	\$10,691,399	25%	\$2,049,476	23%	\$1.80	14%	\$0.34	12%

8.1.1.2 Gear

From 1994 through 2010, 97% of all shrimp were caught using trawls. An additional 3% were caught using channel nets and less than 1% in other gears (Figure 8.3). Table 8.2 shows the number of pounds landed, the total value, and the price per pound for each of the gears listed in Figure 8.3 by year from 1994 – 2010.




Figure 8.3. Percent of landings by gear used to harvest shrimp in all North Carolina waters, 1994 – 2010 (DMF Trip Ticket Program).

Trawls were the primary gears used to land shrimp in every year (Table 8.2). Trawls brought in the highest price per pound in most years. The price per pound for shrimp landed in trawls ranged from \$1.59 (2009) to \$2.62 (1994), averaging \$2.12 from 1994 to 2010. The price per pound for shrimp landed in channel nets ranged from \$1.08 (2009) to \$2.41 (1997), averaging \$1.81. The price per pound for shrimp landed in other gears ranged from \$1.38 (1999) to \$6.27 (2003), averaging \$2.84.

DRAFT

Table 8.2. Gear type, pounds, price per pound, and total value of shrimp landings by gear in all North Carolina waters, 1994 – 2010 (DMF Trip Ticket Program).

Year	Gear Type	Pounds	Nominal Value	Nominal Price per Pound
1994	Trawl	7,099,215	\$18,575,429	\$2.62
	Channel Net	186,029	\$403,636	\$2.17
	Other	7,245	\$17,499	\$2.42
1995	Trawl	8,361,435	\$19,688,121	\$2.35
	Channel Net	273,092	\$568,870	\$2.29
	Other	34,573	\$60,995	\$1.76
1996	Trawl	5,068,715	\$12,910,323	\$2.55
	Channel Net	199,915	\$457,195	\$2.08
	Other	2,643	\$6,444	\$2.44
1997	Trawl	6,795,437	\$17,739,453	\$2.61
	Channel Net	191,188	\$459,963	\$2.41
	Other	2,200	\$5,433	\$2.47
1998	Trawl	4,451,934	\$10,451,410	\$2.35
	Channel Net	181,917	\$399,731	\$2.20
	Other	2,493	\$5,309	\$2.13
1999	Trawl	8,712,050	\$21,511,886	\$2.47
	Channel Net	284,443	\$571,531	\$2.01
	Other	8,042	\$11,072	\$1.38
2000	Trawl	10,070,979	\$24,775,580	\$2.46
	Channel Net	260,321	\$621,181	\$2.39
	Other	3,615	\$9,156	\$2.53
2001	Trawl	5,066,909	\$11,510,270	\$2.27
	Channel Net	185,567	\$395,426	\$2.13
	Other	1,737	\$5,375	\$3.09
2002	Trawl	9,713,978	\$17,914,977	\$1.84
	Channel Net	250,656	\$436,803	\$1.74
	Other	4,391	\$12,997	\$2.96
2003	Trawl	5,909,728	\$10,508,015	\$1.78
	Channel Net	255,892	\$420,083	\$1.64
	Other	1,751	\$10,980	\$6.27
2004	Trawl	4,730,255	\$9,230,605	\$1.95
	Channel Net	149,933	\$228,586	\$1.52
	Other	628	\$3,662	\$5.83

DRAFT

Table 8.2 continued

Year	Gear Type	Pounds	Nominal Value	Nominal Price per Pound
2005	Trawl	2,223,994	\$4,216,906	\$1.90
	Channel Net	130,710	\$187,292	\$1.43
	Other	2,813	\$4,927	\$1.75
2006	Trawl	5,549,686	\$8,902,927	\$1.60
	Channel Net	181,102	\$227,972	\$1.26
	Other	5,861	\$10,535	\$1.80
2007	Trawl	9,367,837	\$17,625,282	\$1.88
	Channel Net	165,729	\$272,177	\$1.64
	Other	3,664	\$7,876	\$2.15
2008	Trawl	9,167,896	\$18,892,082	\$2.06
	Channel Net	253,530	\$336,822	\$1.33
	Other	2,742	\$17,026	\$6.21
2009	Trawl	5,221,528	\$8,323,544	\$1.59
	Channel Net	180,704	\$195,984	\$1.08
	Other	5,476	\$8,186	\$1.50
2010	Trawl	5,819,567	\$10,499,213	\$1.80
	Channel Net	129,865	\$182,808	\$1.41
	Other	5,903	\$9,378	\$1.59

8.1.1.3 Water bodies

The majority of inshore shrimp are landed from the Pamlico and Core sounds. In every year since 1994 with the exception of 1998 and 2005, the greatest amount of shrimp in terms of pounds and value came from the Pamlico Sound compared to all other trip ticket water bodies. On average from 1994 through 2010, 50% of all shrimp landed in North Carolina came from Pamlico Sound alone. Pamlico Sound and ocean landings south of Cape Hatteras each account for over \$1 million in landings each year.

Table 8.3 shows shrimp landings for all water bodies. Some waterbodies are listed but show no landings for a given year. This does not mean those waterbodies did not have landings, rather these landings were confidential due to the low number of fishermen reporting landings. Landings that were confidential were replaced with an asterisk (*). Also, the names of water bodies and how they are used has changed over time. For example, "Inland Waterway" was separated into "Inland Waterway – Brunswick" and "Inland Waterway – Onslow" in 2003. Ocean landings were separated into landings north and south of Cape Hatteras as well within state waters (0-3 miles) and federal waters (beyond 3 miles).

DRAFT

Table 8.3. Pounds and value of shrimp landed from North Carolina water bodies from 1994 – 2010 (DMF Trip Ticket Program).

Waterbody	1994		Waterbody	1995		Waterbody	1996	
	Pounds	Value		Pounds	Value		Pounds	Value
Bay River	20,051	\$54,588	Bay River	10,021	\$19,981	Bay River	6,052	\$14,227
Bogue Sound	23,344	\$49,666	Bogue Sound	34,345	\$65,670	Bogue Sound	45,689	\$92,839
Cape Fear River	149,791	\$302,735	Cape Fear River	114,261	\$186,101	Cape Fear River	80,380	\$189,547
Core Sound	863,245	\$1,833,609	Core Sound	1,069,213	\$2,272,343	Core Sound	738,052	\$1,689,450
Croatan Sound	7,701	\$17,963	Croatan Sound	13,768	\$36,115	Croatan Sound	6,590	\$18,233
Inland Waterway	50,936	\$105,136	Inland Waterway	110,410	\$191,049	Currituck Sound	*	*
Lockwood Folly	426	\$769	Lockwood Folly	477	\$747	Inland Waterway	84,630	\$171,418
Masonboro Sound	4,638	\$9,160	Masonboro Sound	1,952	\$3,007	Lockwood Folly	*	*
Neuse River	115,689	\$320,348	Neuse River	114,705	\$284,780	Masonboro Sound	5,973	\$12,693
New River	103,078	\$284,059	New River	274,212	\$689,719	Neuse River	111,098	\$311,191
Newport River	166,828	\$311,459	Newport River	275,058	\$386,857	New River	148,264	\$420,006
North River/Back Sound	127,327	\$257,580	North River/Back Sound	196,322	\$417,171	Newport River	125,092	\$270,421
Ocean 0-3 mi, N of Cape Hatt	*	*	Ocean 0-3 mi, N of Cape Hatt	55,686	\$168,765	North River/Back Sound	56,511	\$132,244
Ocean less than 3 miles	1,372,958	\$3,621,439	Ocean 0-3 mi, S of Cape Hatt	337,606	\$862,314	Ocean 0-3 mi, N of Cape Hatt	13,318	\$31,020
Ocean more than 3 miles	277,855	\$763,765	Ocean >3 mi, S of Cape Hatt	17,649	\$48,568	Ocean 0-3 mi, S of Cape Hatt	1,232,910	\$3,176,947
Pamlico River	46,107	\$129,203	Ocean less than 3 miles	1,478,122	\$3,190,104	Ocean >3 mi, N of Cape Hatt	17,361	\$47,795
Pamlico Sound	3,861,546	\$10,720,745	Ocean more than 3 miles	303,217	\$846,978	Ocean >3 mi, S of Cape Hatt	180,351	\$475,093
Roanoke Sound	14,776	\$30,690	Pamlico River	34,756	\$86,079	Ocean less than 3 miles	329,751	\$788,277
Shalotte River	1,807	\$3,550	Pamlico Sound	4,096,435	\$10,313,455	Ocean more than 3 miles	49,752	\$139,211
Stump Sound	8,553	\$21,719	Roanoke Sound	5,632	\$12,482	Pamlico River	23,078	\$64,409
Topsail Sound	29,485	\$71,714	Shalotte River	1,491	\$2,127	Pamlico Sound	1,934,399	\$5,147,444
White Oak River	44,995	\$82,782	Stump Sound	25,546	\$47,594	Pungo River	*	*
			Topsail Sound	59,202	\$139,389	Roanoke Sound	7,896	\$19,462
			White Oak River	39,013	\$46,591	Shalotte River	394	\$907
						Stump Sound	27,088	\$65,601
						Topsail Sound	21,898	\$47,878
						White Oak River	23,825	\$45,008

*Confidential data

DRAFT

Table 8.3 (continued).

	1997			1998			1999	
	Pounds	Value		Pounds	Value		Pounds	Value
Waterbody			Waterbody			Waterbody		
Bay River	16,409	\$40,241	Bay River	1,358	\$2,709	Bay River	27,913	\$69,034
Bogue Sound	17,009	\$33,188	Bogue Sound	41,849	\$70,974	Bogue Sound	48,220	\$94,783
Cape Fear River	138,424	\$273,933	Cape Fear River	82,592	\$150,208	Cape Fear River	118,742	\$214,347
Core Sound	636,805	\$1,423,124	Core Sound	547,488	\$991,584	Core Sound	884,330	\$1,598,475
Croatan Sound	12,539	\$32,250	Croatan Sound	1,389	\$3,541	Croatan Sound	3,793	\$8,370
Inland Waterway	66,675	\$132,363	Inland Waterway	54,768	\$103,877	Inland Waterway	66,506	\$118,763
Lockwood Folly	*	*	Lockwood Folly	*	*	Lockwood Folly	*	*
Masonboro Sound	5,715	\$10,681	Masonboro Sound	4,961	\$8,371	Masonboro Sound	2,266	\$3,359
Neuse River	164,538	\$441,246	Neuse River	83,765	\$177,286	Neuse River	216,933	\$485,133
New River	244,360	\$637,018	New River	259,274	\$661,359	New River	271,883	\$626,671
Newport River	213,818	\$424,734	Newport River	71,793	\$126,734	Newport River	307,504	\$456,164
North River/Back Sound	92,489	\$224,603	North River/Back Sound	27,391	\$53,066	North River/Back Sound	160,649	\$193,871
Ocean 0-3 mi, N of Cape Hatt	21,710	\$66,135	Ocean 0-3 mi, N of Cape Hatt	*	*	Ocean 0-3 mi, N of Cape Hatt	6,638	\$21,241
Ocean 0-3 mi, S of Cape Hatt	1,030,217	\$2,618,449	Ocean 0-3 mi, S of Cape Hatt	1,493,238	\$3,695,714	Ocean 0-3 mi, S of Cape Hatt	2,468,260	\$6,668,902
Ocean >3 mi, N of Cape Hatt	*	*	Ocean >3 mi, N of Cape Hatt	14,516	\$42,554	Ocean >3 mi, N of Cape Hatt	51,502	\$174,186
Ocean >3 mi, S of Cape Hatt	205,008	\$571,483	Ocean >3 mi, S of Cape Hatt	380,907	\$1,002,254	Ocean >3 mi, S of Cape Hatt	236,725	\$584,197
Ocean less than 3 miles	243,964	\$643,232	Ocean less than 3 miles	344,408	\$810,808	Ocean less than 3 miles	67,420	\$214,004
Ocean more than 3 miles	32,609	\$89,485	Ocean more than 3 miles	18,602	\$47,936	Ocean more than 3 miles	5,007	\$17,816
Pamlico River	39,793	\$116,916	Pamlico River	14,664	\$37,008	Pamlico River	43,794	\$120,732
Pamlico Sound	3,722,785	\$10,231,549	Pamlico Sound	1,115,961	\$2,720,014	Pamlico Sound	3,876,433	\$10,191,283
Pungo River	1,303	\$3,186	Roanoke Sound	188	\$432	Pungo River	*	*
Roanoke Sound	8,568	\$21,610	Shallotte River	*	*	Roanoke Sound	1,488	\$3,130
Shallotte River	2,413	\$4,423	Stump Sound	16,038	\$36,091	Shallotte River	423	\$1,067
Stump Sound	29,139	\$65,977	Topsail Sound	36,579	\$73,690	Stump Sound	20,522	\$38,276
Topsail Sound	22,508	\$54,235	White Oak River	23,582	\$37,858	Topsail Sound	72,561	\$134,762
White Oak River	12,986	\$24,579				White Oak River	37,984	\$36,346

*Confidential data

DRAFT

Table 8.3 (continued).

Waterbody	2000		Waterbody	2001		Waterbody	2002	
	Pounds	Value		Pounds	Value		Pounds	Value
Albemarle Sound	*	*	Bay River	5,935	\$13,385	Alligator River	*	*
Bay River	35,348	\$78,560	Bogue Sound	9,906	\$13,484	Bay River	14,070	\$19,787
Bogue Sound	23,875	\$38,291	Cape Fear River	17,850	\$51,779	Bogue Sound	31,389	\$55,013
Cape Fear River	46,058	\$79,380	Core Sound	431,489	\$840,078	Cape Fear River	82,868	\$109,384
Core Sound	464,916	\$901,327	Croatan Sound	*	*	Core Sound	783,852	\$1,235,756
Croatan Sound	40,989	\$96,578	Inland Waterway	51,538	\$91,228	Croatan Sound	10,010	\$18,063
Inland Waterway	79,462	\$148,373	Lockwood Folly	*	*	Inland Waterway	55,313	\$88,650
Lockwood Folly	*	*	Masonboro Sound	1,514	\$3,014	Inland Waterway (Onslow)	2,966	\$3,316
Masonboro Sound	4,212	\$6,594	Neuse River	19,942	\$43,989	Lockwood Folly	*	*
Neuse River	210,970	\$471,504	New River	189,084	\$430,819	Masonboro Sound	3,373	\$5,116
New River	483,739	\$1,350,697	Newport River	176,502	\$241,348	Neuse River	213,697	\$373,058
Newport River	240,583	\$304,680	North River/Back Sound	71,739	\$133,593	New River	428,783	\$871,912
North River/Back Sound	216,045	\$309,372	Ocean 0-3 mi, S of Cape Hatt	1,157,075	\$2,297,258	Newport River	292,696	\$289,219
Ocean 0-3 mi, N of Cape Hatt	36,319	\$98,898	Ocean >3 mi, N of Cape Hatt	*	*	North River/Back Sound	186,314	\$212,358
Ocean 0-3 mi, S of Cape Hatt	1,397,962	\$3,565,804	Ocean >3 mi, S of Cape Hatt	100,069	\$207,035	Ocean 0-3 mi, N of Cape Hatt	*	*
Ocean >3 mi, N of Cape Hatt	29,942	\$84,146	Pamlico River	20,203	\$43,506	Ocean 0-3 mi, S of Cape Hatt	1,288,291	\$2,438,720
Ocean >3 mi, S of Cape Hatt	133,048	\$349,195	Pamlico Sound	2,890,943	\$7,337,235	Ocean >3 mi, N of Cape Hatt	*	*
Pamlico River	44,710	\$109,896	Pasquotank River	*	*	Ocean >3 mi, S of Cape Hatt	60,109	\$137,491
Pamlico Sound	6,708,334	\$17,192,339	Pungo River	*	*	Pamlico River	102,459	\$176,545
Pungo River	6,926	\$17,492	Roanoke Sound	*	*	Pamlico Sound	6,147,806	\$11,977,356
Roanoke Sound	7,298	\$15,750	Shalotte River	6,123	\$11,175	Pungo River	7,870	\$14,036
Shalotte River	896	\$916	Stump Sound	11,795	\$26,157	Roanoke Sound	32,080	\$58,859
Stump Sound	21,888	\$45,115	Topsail Sound	21,888	\$35,865	Shalotte River	*	*
Topsail Sound	39,152	\$84,948	White Oak River	62,361	\$75,401	Stump Sound	48,099	\$84,230
White Oak River	62,164	\$55,872				Topsail Sound	14,383	\$22,975
						White Oak River	137,397	\$128,142

*Confidential data

DRAFT

Table 8.3 (continued).

Waterbody	2003		Waterbody	2004		Waterbody	2005	
	Pounds	Value		Pounds	Value		Pounds	Value
Bay River	2,010	\$3,220	Bay River	*	*	Bay River	1,915	\$4,151
Bogue Sound	127,781	\$155,164	Bogue Sound	18,624	\$31,116	Bogue Sound	12,729	\$21,281
Cape Fear River	101,424	\$162,463	Cape Fear River	32,730	\$37,576	Cape Fear River	46,241	\$66,025
Core Sound	821,174	\$1,390,897	Core Sound	252,813	\$432,071	Core Sound	317,370	\$478,582
Croatan Sound	*	*	Croatan Sound	6,856	\$13,185	Croatan Sound	*	*
Inland Waterway	47,487	\$68,150	Inland Waterway	14,381	\$16,956	Inland Waterway	13,018	\$17,853
Inland Waterway (Brunswick)	18,404	\$28,735	Inland Waterway (Brunswick)	8,633	\$14,820	Inland Waterway (Brunswick)	16,746	\$20,983
Inland Waterway (Onslow)	31,972	\$48,327	Inland Waterway (Onslow)	27,523	\$35,308	Inland Waterway (Onslow)	45,855	\$68,190
Lockwood Folly	*	*	Masonboro Sound	17,722	\$18,722	Masonboro Sound	4,745	\$5,675
Masonboro Sound	6,561	\$7,470	Neuse River	87,384	\$175,348	Neuse River	110,286	\$198,067
Neuse River	102,366	\$166,540	New River	174,901	\$307,111	New River	49,506	\$88,770
New River	230,381	\$454,157	Newport River	125,039	\$139,232	Newport River	70,030	\$95,927
Newport River	142,654	\$190,650	North River/Back Sound	126,873	\$189,306	North River/Back Sound	84,838	\$116,287
North River/Back Sound	117,353	\$175,658	Ocean 0-3 mi, N of Cape Hatt	1,753	\$3,486	Ocean 0-3 mi, S of Cape Hatt	910,709	\$1,835,281
Ocean 0-3 mi, N of Cape Hatt	*	*	Ocean 0-3 mi, S of Cape Hatt	1,569,215	\$2,885,008	Ocean >3 mi, S of Cape Hatt	58,395	\$101,993
Ocean 0-3 mi, S of Cape Hatt	2,008,508	\$3,363,342	Ocean >3 mi, S of Cape Hatt	199,207	\$286,687	Pamlico River	3,903	\$9,235
Ocean >3 mi, N of Cape Hatt	*	*	Pamlico River	6,546	\$18,035	Pamlico Sound	558,104	\$1,204,022
Ocean >3 mi, S of Cape Hatt	242,477	\$413,318	Pamlico Sound	2,104,690	\$4,744,780	Pungo River	*	*
Pamlico River	11,934	\$25,109	Roanoke Sound	6,646	\$11,952	Roanoke Sound	907	\$2,226
Pamlico Sound	2,023,826	\$4,112,575	Shalotte River	*	*	Shalotte River	*	*
Pungo River	*	*	Stump Sound	9,840	\$16,378	Stump Sound	17,202	\$26,420
Roanoke Sound	2,415	\$3,978	Topsail Sound	28,312	\$35,279	Topsail Sound	26,535	\$37,665
Shalotte River	4,333	\$6,063	White Oak River	60,283	\$49,103	White Oak River	6,655	\$8,276
Stump Sound	25,010	\$37,379						
Topsail Sound	43,141	\$69,252						
White Oak River	52,052	\$49,936						

*Confidential data

DRAFT

Table 8.3 (continued).

Waterbody	2006		Waterbody	2007		Waterbody	2008	
	Pounds	Value		Pounds	Value		Pounds	Value
Bay River	*	*	Bay River	858	\$1,655	Bay River	7,144	\$12,386
Bogue Sound	70,432	\$71,766	Bogue Sound	39,385	\$52,532	Bogue Sound	57,928	\$52,670
Cape Fear River	35,843	\$48,556	Cape Fear River	46,124	\$88,767	Cape Fear River	47,264	\$83,755
Core Sound	260,588	\$359,865	Core Sound	241,093	\$361,895	Core Sound	434,900	\$661,196
Croatan Sound	2,421	\$4,639	Croatan Sound	23,961	\$51,981	Croatan Sound	4,761	\$9,375
Inland Waterway (Brunswick)	8,380	\$11,268	Inland Waterway	*	*	Inland Waterway (Brunswick)	19,944	\$39,565
Inland Waterway (Onslow)	57,007	\$60,737	Inland Waterway (Brunswick)	11,512	\$12,372	Inland Waterway (Onslow)	29,588	\$51,941
Masonboro Sound	7,603	\$5,440	Inland Waterway (Onslow)	25,631	\$37,125	Masonboro Sound	*	*
Neuse River	125,952	\$204,414	Lockwood Folly	*	*	Neuse River	391,739	\$666,697
New River	164,411	\$207,266	Masonboro Sound	335	\$413	New River	101,554	\$230,990
Newport River	199,986	\$123,387	Neuse River	139,720	\$207,794	Newport River	118,998	\$101,344
North River/Back Sound	258,670	\$237,022	New River	151,743	\$217,145	North River/Back Sound	145,782	\$138,949
Ocean 0-3 mi, N of Cape Hatt	3,331	\$6,022	Newport River	170,636	\$113,937	Ocean 0-3 mi, N of Cape Hatt	*	*
Ocean 0-3 mi, S of Cape Hatt	1,843,020	\$3,076,473	North River/Back Sound	179,602	\$213,658	Ocean 0-3 mi, S of Cape Hatt	1,787,589	\$4,298,190
Ocean >3 mi, S of Cape Hatt	125,500	\$169,992	Ocean 0-3 mi, N of Cape Hatt	32,734	\$58,075	Ocean >3 mi, N of Cape Hatt	*	*
Pamlico River	3,648	\$6,357	Ocean 0-3 mi, S of Cape Hatt	1,557,680	\$3,201,450	Ocean >3 mi, S of Cape Hatt	183,968	\$496,726
Pamlico Sound	2,477,858	\$4,473,267	Ocean >3 mi, N of Cape Hatt	*	*	Pamlico River	21,779	\$47,761
Pungo River	*	*	Ocean >3 mi, S of Cape Hatt	49,978	\$62,205	Pamlico Sound	5,944,307	\$12,125,633
Roanoke Sound	642	\$1,328	Pamlico River	30,015	\$53,571	Roanoke Sound	2,189	\$3,488
Stump Sound	11,655	\$15,775	Pamlico Sound	6,761,768	\$13,061,121	Stump Sound	31,862	\$53,968
Topsail Sound	18,925	\$22,768	Roanoke Sound	6,059	\$14,006	Topsail Sound	5,435	\$7,306
White Oak River	58,950	\$31,449	Stump Sound	16,497	\$23,204	White Oak River	20,282	\$14,570
			Topsail Sound	10,657	\$12,965			
			White Oak River	24,277	\$23,386			

*Confidential data

DRAFT

Table 8.3 (continued).

Waterbody	2009		Waterbody	2010	
	Pounds	Value		Pounds	Value
Albemarle Sound	*	*	Bay River	2,405	\$4,482
Bay River	4,192	\$6,108	Bogue Sound	34,534	\$47,578
Bogue Sound	31,643	\$38,675	Cape Fear River	137,009	\$179,609
Cape Fear River	44,658	\$71,234	Core Sound	119,470	\$190,405
Core Sound	191,151	\$247,872	Croatan Sound	1,075	\$2,121
Croatan Sound	*	*	Inland Waterway (Brunswick)	30,935	\$36,596
Inland Waterway (Brunswick)	15,873	\$15,426	Inland Waterway (Onslow)	47,345	\$69,708
Inland Waterway (Onslow)	53,465	\$56,075	Masonboro Sound	5,918	\$7,631
Masonboro Sound	*	*	Neuse River	116,953	\$187,205
Neuse River	116,298	\$167,095	New River	144,919	\$222,679
New River	22,552	\$26,134	Newport River	91,966	\$101,949
Newport River	73,951	\$58,068	North River/Back Sound	55,370	\$76,309
North River/Back Sound	65,725	\$80,887	Ocean 0-3 mi, N of Cape Hatt	*	*
Ocean 0-3 mi, N of Cape Hatt	*	*	Ocean 0-3 mi, S of Cape Hatt	1,130,146	\$2,195,822
Ocean 0-3 mi, S of Cape Hatt	860,971	\$1,468,493	Ocean >3 mi, S of Cape Hatt	103,846	\$226,230
Ocean >3 mi, N of Cape Hatt	*	*	Pamlico River	12,813	\$24,691
Ocean >3 mi, S of Cape Hatt	56,211	\$103,912	Pamlico Sound	3,837,536	\$6,988,818
Pamlico River	18,710	\$28,514	Roanoke Sound	429	\$666
Pamlico Sound	3,686,102	\$5,942,139	Shalotte River	*	*
Roanoke Sound	2,607	\$4,134	Stump Sound	19,360	\$28,561
Stump Sound	20,612	\$23,188	Topsail Sound	27,903	\$39,445
Topsail Sound	24,652	\$21,512	White Oak River	15,457	\$16,694
White Oak River	36,720	\$29,610			

*Confidential data

DRAFT

8.1.1.4 Participants and trips

DMF began a new licensing system in 1999. This new system allows for easier identification of specific fishermen with their individual landings by species and the number of trips taken where a given species was landed. Table 8.4 shows the number of participants in the shrimp fishery by year and the ex-vessel value of their landings.

The number of fishermen who participate in the fishery seems to follow the abundance of shrimp or when fishermen are receiving a good price. This indicates that some fishermen are able to rely on other species or other work when shrimping is not as lucrative. The years 1999 and 2000 saw the greatest number of participants in the fishery at over 800, while 2005 had the least number of participants at 400.

On average, from 1999 through 2010, nearly half of all fishermen who caught shrimp had ex-vessel landings values of \$10,000 or less. The percentage of fishermen who landed ex-vessel values of between \$35,000 and \$50,000 remained fairly constant at about 4-7% of all participants. The percentage of fishermen who had ex-vessel landings values between \$50,001 and \$75,000 varied in a given year; however, the general trend was downward across the time frame.

Table 8.4. Number of participants in the shrimp fishery by value of landings and year in North Carolina, 1999 – 2010 (DMF Trip Ticket Program).

	Year											
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
\$1-\$1,000	245	280	203	193	115	158	109	105	117	117	117	140
% within year	29%	30%	28%	24%	19%	27%	27%	24%	24%	23%	25%	29%
\$1,001-\$5,000	191	173	158	181	148	140	117	83	92	104	124	102
% within year	23%	18%	22%	23%	25%	24%	29%	19%	19%	20%	27%	21%
\$5,001-\$10,000	85	96	86	78	77	69	52	73	58	46	58	44
% within year	10%	10%	12%	10%	13%	12%	13%	17%	12%	9%	13%	9%
\$10,001-\$20,000	108	118	98	92	90	77	53	52	58	58	47	58
% within year	13%	13%	14%	12%	15%	13%	13%	12%	12%	11%	10%	12%
\$20,001-\$35,000	53	75	60	91	62	53	40	33	21	39	42	36
% within year	6%	8%	8%	11%	10%	9%	10%	8%	4%	8%	9%	7%
\$35,001-\$50,000	30	39	38	39	41	32	12	31	27	38	26	30
% within year	4%	4%	5%	5%	7%	5%	3%	7%	6%	7%	6%	6%
\$50,001-\$75,000	41	42	47	48	44	32	10	33	28	31	12	29
% within year	5%	4%	7%	6%	7%	5%	3%	8%	6%	6%	3%	6%
>\$75,000	80	113	32	76	22	23	7	30	81	82	33	45
% within year	10%	12%	4%	10%	4%	4%	2%	7%	17%	16%	7%	9%
Total Participants	833	936	722	798	599	584	400	440	482	515	459	484
Percent Change		12%	-23%	11%	-25%	-3%	-32%	10%	10%	7%	-11%	5%

DRAFT

Table 8.5 shows the number of fisherman and the number of trips they took in which they landed and sold shrimp for the years 1999 through 2010. From 1999 through 2010 an average of 12% of all participants only had one trip with shrimp landings. An average of 69% of all persons reporting shrimp landings had 20 or fewer trips in a given year. An average of 13% of all fishermen reported taking 41 or more trips per year. Again, abundance of shrimp, prices received for the catch, and weather events such as hurricanes greatly affect the number of trips a fisherman might make for shrimp.

Table 8.5. Number of participants and the number of trips taken that landed shrimp in North Carolina, 1999 – 2010 (DMF Trip Ticket Program).

	Year											
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
1 Trip	114	125	93	78	57	62	46	45	66	67	48	64
% within year	14%	13%	13%	10%	10%	11%	12%	10%	14%	13%	10%	13%
2-10 Trips	281	313	251	271	182	199	166	161	151	197	171	197
% within year	34%	33%	35%	34%	30%	34%	42%	37%	31%	38%	37%	41%
11-20 Trips	187	178	160	163	124	113	76	94	97	123	120	107
% within year	22%	19%	22%	20%	21%	19%	19%	21%	20%	24%	26%	22%
21-30 Trips	92	125	81	96	67	84	43	57	70	47	48	40
% within year	11%	13%	11%	12%	11%	14%	11%	13%	15%	9%	10%	8%
31-40 Trips	45	59	38	51	47	38	30	29	36	28	22	26
% within year	5%	6%	5%	6%	8%	7%	8%	7%	7%	5%	5%	5%
41-50 Trips	30	39	28	35	35	26	13	21	20	22	18	17
% within year	4%	4%	4%	4%	6%	4%	3%	5%	4%	4%	4%	4%
51-60 Trips	25	24	19	20	26	21	12	12	15	10	16	13
% within year	3%	3%	3%	3%	4%	4%	3%	3%	3%	2%	3%	3%
61-70 Trips	17	25	18	18	20	13	3	4	11	8	5	8
% within year	2%	3%	2%	2%	3%	2%	0.80%	0.90%	2%	2%	1%	2%
71-80 Trips	13	13	10	23	16	9	1	5	5	6	4	2
% within year	2%	1%	1%	3%	3%	2%	0.30%	1.10%	1%	1.20%	0.90%	0.40%
81-90 Trips	13	17	7	14	11	7	4	4	2	3	2	3
% within year	2%	2%	1%	2%	2%	1%	1%	0.90%	0.40%	0.60%	0.40%	0.60%
91-100 Trips	6	7	9	10	6	4	4	2	1	3	1	2
% within year	0.70%	0.70%	1%	1%	1%	0.70%	1%	0.50%	0.20%	0.60%	0.20%	0.40%
> 100 Trips	10	11	8	19	8	8	2	6	8	1	4	5
% within year	1%	1%	1%	2%	1%	1%	0.50%	1%	2%	0.20%	0.90%	1%
Total	833	936	722	798	599	584	400	440	482	515	459	484

DRAFT

In North Carolina, licensed commercial fishermen are legally obligated to only sell their catch to licensed seafood dealers. Figure 8.4 shows the number of North Carolina seafood dealers who purchased shrimp from commercial fishermen each year from 1994 through 2010. There is a variation in the number of seafood dealers purchasing shrimp from year to year with a low of 208 in 2008 to a high of 284 in 2002. The annual differences are due largely to availability of local shrimp as well as availability and price of imported shrimp.




Figure 8.4. Number of seafood dealers who purchased shrimp from 1994 to 2010 (DMF Trip Ticket Program).

Table 8.6 shows the number of fish dealers who purchased specific amounts of shrimp in a given year. An average of 9% of dealers purchased fewer than 100 pounds of shrimp in a given year. About 35% of dealers purchased 1,000 or fewer pounds of shrimp a year. Approximately 23% of dealers purchased more than 20,000 pounds of shrimp from fishermen. Only 9% purchased more than 100,000 pounds of shrimp.

DRAFT

Table 8.6. Number of seafood dealer and pounds of shrimp purchased by North Carolina fish dealers from North Carolina fishermen, 1994 – 2010 (DMF Trip Ticket Program).

Pounds	Year																
	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
100 Pounds or less	18	27	22	22	26	28	28	23	23	20	13	17	24	28	14	21	21
% within year	8%	10%	9%	9%	11%	10%	11%	10%	8%	8%	5%	8%	11%	13%	7%	9%	9%
101-500 Pounds	37	35	50	30	36	56	47	40	47	25	44	39	30	30	29	37	40
% within year	16%	13%	19%	12%	15%	21%	19%	18%	17%	10%	18%	19%	14%	14%	14%	15%	17%
501-1,000 Pounds	26	25	21	30	30	23	21	24	31	23	22	29	24	16	16	21	28
% within year	11%	10%	8%	12%	13%	8%	8%	11%	11%	9%	9%	14%	11%	7%	8%	9%	12%
1,001-2,000 Pounds	19	31	28	32	26	31	26	22	41	39	24	25	21	24	26	39	23
% within year	8%	12%	11%	13%	11%	11%	10%	10%	14%	16%	10%	12%	10%	11%	13%	16%	10%
2,001-5,000 Pounds	31	35	43	36	32	35	36	34	39	36	38	33	35	36	38	42	39
% within year	13%	13%	17%	15%	14%	13%	14%	15%	14%	15%	15%	16%	16%	16%	18%	17%	16%
5,001-10,000 Pounds	19	21	19	17	16	17	16	19	23	29	37	20	26	29	20	31	29
% within year	8%	8%	7%	7%	7%	6%	6%	8%	8%	12%	15%	10%	12%	13%	10%	13%	12%
10,001-20,000 Pounds	17	21	17	22	14	15	18	13	21	18	17	14	20	17	20	14	17
% within year	7%	8%	7%	9%	6%	6%	7%	6%	7%	7%	7%	7%	9%	8%	10%	6%	7%
20,001-35,000 Pounds	12	8	11	8	12	11	11	10	12	13	14	10	9	9	10	9	11
% within year	5%	3%	4%	3%	5%	4%	4%	4%	4%	5%	6%	5%	4%	4%	5%	4%	5%
35,001-50,000 Pounds	9	7	18	11	13	6	5	5	5	10	7	5	4	4	2	2	5
% within year	4%	3%	7%	4%	6%	2%	2%	2%	2%	4%	3%	2%	2%	2%	1%	1%	2%
50,001-75,000 Pounds	13	9	6	12	8	7	6	9	6	7	8	8	4	2	5	5	8
% within year	6%	3%	2%	5%	3%	3%	2%	4%	2%	3%	3%	4%	2%	1%	2%	2%	3%
75,001-100,000 Pounds	9	18	9	6	8	8	9	8	3	5	5	6	7	5	5	6	5
% within year	4%	7%	3%	2%	3%	3%	4%	4%	1%	2%	2%	3%	3%	2%	2%	2%	2%
100,001-150,000 Pounds	10	9	5	10	6	17	10	11	11	8	9	3	4	3	7	8	4
% within year	4%	3%	2%	4%	3%	6%	4%	5%	4%	3%	4%	1%	2%	1%	3%	3%	2%
150,001-200,000 Pounds	3	7	5	5	2	10	5	4	8	8	5	0	6	4	3	2	3
% within year	1%	3%	2%	2%	1%	4%	2%	2%	3%	3%	2%	0%	3%	2%	1%	1%	1%

DRAFT

Table 8.6. (continued).

Pounds	Year																
	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
More than 200,000 Pounds	9	10	4	7	5	8	16	3	14	6	3	0	6	14	13	6	6
% within year	4%	4%	2%	3%	2%	3%	6%	1%	5%	2%	1%	0%	3%	6%	6%	2%	3%
Total	232	263	258	248	234	272	254	225	284	247	246	209	220	221	208	243	239
Percent change		13%	-2%	-4%	-6%	16%	-7%	-11%	26%	-13%	0%	-15%	5%	0%	-6%	17%	-2%

8.1.1.5 Processing

Some dealers will go so far as to head shrimp for customers, but a large portion of the commercial shrimp catch is sold heads-on. Shrimp that cannot be sold fresh are frozen. A few dealers sell shrimp to be processed into other consumable products such as frozen breaded shrimp; however, there are no known shrimp processors currently operating in North Carolina.

8.1.1.6 Marketing and distribution

Seafood dealers sell shrimp to other dealers, restaurateurs, retail outlets, and directly to the consumer. There is no specific information available as to how much North Carolina shrimp is sold through each of these venues.

According to the US Department of Commerce (2011) there were four pounds of shrimp consumed per capita by Americans in 2010. According to the U.S. Census Bureau, there were 9,535,483 residents of North Carolina in 2010.

Using these two statistics provides an estimate that approximately 38 million pounds of shrimp were consumed by North Carolina residents in 2010, over six times the total weight of shrimp caught from North Carolina waters. If all shrimp caught in North Carolina remained in the state, it would only supply one sixth of the state's consumption needs. There is a large reliance on shrimp imported into North Carolina from other states and foreign countries. Thus, it appears imports represent a double-edged sword. On the one hand, they have increased the supply, and also demand due to the downward pressure on price. On the other hand, the effect on price has greatly diminished the economic returns to domestic fishermen. One result is that some fishermen have looked for more land-based work. In other cases, wives have taken full time jobs to supplement their family's income (Maiolo 2004). In some cases fishermen are becoming dealers to sell their catches to niche markets, thereby obtaining higher prices.

Supply and demand largely determine the price per pound paid to shrimp fishermen. Farm raised imports, mostly from Asia and Latin America, have expanded to meet increasing demand and, indeed, appear to have fostered it. Following the poor domestic harvest years of the late seventies and early eighties, imports had increased from fairly modest levels to 341 million pounds in 1983, 500 million in 1989 and 1990, and 759 million pounds in 2000 (Maiolo 2004). The impact of imports has been especially hard on shrimp fishermen since 2001. In that year, price per pound dropped 24% over the previous year. In 2002, the price dropped an additional 32% over the 2001 price received by fishermen. A small gain was realized in 2003 when the price per pound increased by about 13% over the 2002 price. Nonetheless, with the exception of 2008, the price per pound has remained under \$2.00. The price per pound reduction received by fishermen since 2000 can largely be attributed to the impact of imports. In recent years, imports of shrimp have remained above 1 billion pounds per year, with approximately 1.6 billion pounds of shrimp imported into the US in 2010 (NOAA 2011).

Concerned about the rising tide of imports, a group of shrimp industry individuals from the Gulf and South Atlantic formed the Southern Shrimp Alliance (SSA), made up of the shrimp producing states from North Carolina through Texas. The SSA hired two firms: one to do research for possible trade actions, and the other for lobbying.

The SSA decided to file petitions with the Federal government alleging several countries had been dumping shrimp on the US market at below cost. SSA filed trade action against six countries, and the petition was filed on December 31, 2003. Preliminary anti-dumping duties were imposed by the US Department of Commerce in July of 2004. The duties ranged from 3.4% to 67.8% on companies from the countries of Brazil, Ecuador, India, and Thailand. Furthermore, The Commerce Department found that shrimp from China and Vietnam were dumped on US markets at a rate of up to 113% below cost (NCFA 2004).

Additionally, the lobbying efforts of SSA helped to persuade the United States Congress in 2003 to set aside \$35 million to offset the economic losses suffered by shrimp fishermen from southeastern states. North Carolina received \$4.9 million of the total. Of the total, \$4.1 million was sent directly to fishermen based on their trip ticket receipts from licensed dealers that reported landings during the 2002 calendar year. Six hundred and eighty-two checks were mailed out ranging from \$7 to \$64,206; the average amount of assistance per vessel was \$5,906. Approximately \$42,000 was used by DMF to cover the costs of administering the program. Another \$160,000 of unclaimed fisherman disaster assistance funds and unused administrative services funds were channeled to the SSA on behalf of NC shrimp fishermen for legal efforts used to convince the federal government to impose tariffs on countries convicted of illegally dumping shrimp onto US markets.

Approximately \$600,000 (13.3%) of the Federal shrimp economic assistance program of 2003 was given to the North Carolina Department of Agriculture (NCDA) to develop a three-year marketing program for marketing wild-caught North Carolina shrimp. The money was used to market North Carolina wild-caught shrimp in trade and consumer publications, billboards, statewide radio and television promotions, in-store consumer awareness, recipe cards, and trade show participation. However, no additional Federal aid has been given to the NCDA to market wild-caught shrimp since 2003. The NCDA continues to market North Carolina wild-caught shrimp through the "Freshness from North Carolina Waters" seafood promotion program in various consumer and wholesaler publications and radio promotional campaigns during shrimp season (J. Aydlett, NCDA. pers. com. 2012). There are also several local programs such as, Brunswick Catch, Ocracoke Fresh, Carteret Catch, and Outer Banks Catch that promote North Carolina caught shrimp and seafood.

According to the SSA, the benefits of trade relief have not been apparent to many in the industry. Although there is increased stability in the market and declining shrimp prices have slowed, the amount shrimpers receive for their catch continues to be low. The millions collected in anti-dumping duties has been distributed to the domestic industry have overwhelmingly benefited shrimp purchasers and not fishermen. Thus, despite the influx of substantial funds into the hands of purchasers of shrimp, what shrimpers receive for their catch has continued to decline.

8.1.1.7 Economic impact of commercial fishery

In 2010, commercial shrimp landings accounted for about 15% of all the total weight and 23% of the total value of commercial shellfish landed in North Carolina. When finfish are included, shrimp accounted for 8% of the total weight and 13% of the total value of commercial seafood landings in North Carolina. The expenditures and income within the commercial fishing industry in North Carolina produce ripple effects in the state's economy. Each dollar earned and spent within the industry generates additional economic impacts by stimulating further activity in other industries which fosters jobs, income, and economic output. These impacts are calculated using IMPLAN, an economic modeling software. This software uses an input-output model to

estimate economic impacts as dollars are spent and re-spent in the state economy. In 2010, the commercial shrimp fishery in North Carolina contributed, directly and indirectly, approximately \$17.7 million to the state's economy (Table 8.7). These estimates are limited and must be viewed as conservatively low, as they do not include the economic impacts of the wholesale (seafood dealers and distributors), retail, and foodservice sectors due to lack of specific economic data for those sectors in North Carolina.

Table 8.7. Economic impact of the commercial shrimp fishery in North Carolina, 2010.

Economic inputs	\$10,691,399
Additional economic activity	\$6,964,822
Additional jobs supported	56
Total economic impact	\$17,656,221

8.1.2 Recreational fishery economics

There are two survey programs in North Carolina that collect economic data from coastal recreational fishermen. The Marine Recreational Information Program (MRIP) collects data from anglers and includes ocean landings from the coast and inside waters from the Virginia state line to the South Carolina border. Additionally, the DMF conducts creel surveys of anglers in the Cape Fear, Neuse, Pamlico, Tar, and Pungo Rivers. In the past, the DMF also collected data from recreational fishermen who are licensed to use limited amounts of commercial gear through the Recreational Commercial Gear License (RCGL). However, the RCGL survey was discontinued in 2008 due to budget cuts. See the Recreational Section (6.2) for additional information.

8.1.2.1 Marine Recreational Information Program (MRIP)

MRIP captures catch and angler participation data for finfish only. No data from any shellfish species are collected. However, some anglers may catch limited amounts of shrimp for bait using a cast net. There are no data on the economic value of this practice. Additionally, there are several live shrimp bait dealers, as live shrimp have become a popular bait among spotted sea trout fishermen. This live bait market has grown considerably, as a result, over the past 10 years.

8.1.2.2 Recreational use of commercial gear (RCGL)

Along with the heavy participation of part time commercial fishermen in the shrimp industry, the recreational use of commercial gear has had a long and contentious history. Prior to the Fisheries Reform Act of 1997, there was a growing number of participants in both user categories resulting in increased competition in the shrimp fishery. In 2002, the DMF began interviewing recreational fishermen who had purchased a license that allows them to use limited amounts of commercial gear (RCGL). These fishermen were and still are prohibited from selling their catch as it is intended solely for personal use. The RCGL holder surveys did not specifically determine the final disposition of the shrimp landed by these anglers. However, it is presumed that they use the shrimp primarily for personal consumption. This survey program ended in 2008 due to loss of funding.

Table 8.8 gives an indication of the direct economic impact of the recreational shrimp fishery by RCGL fishermen in 2007. The data are separated by those who made overnight trips as opposed to those who made day trips. In the case of the shrimp trawl fishery, the majority of fishing does occur at night. A day trip is one in which a person left their home specifically for one fishing trip and then returned to their regular residence once the fishing activity was completed. An overnight trip is defined as one in which the fishermen spent a longer period of time away from home.

Table 8.8. Economic impact of RCGL fishing trips for shrimp in 2007 (DMF RCGL Survey).

	Overnight Trips	Day Trips
Avg. # of nights	3.98	
Avg. # of miles traveled	133.19	40.3
Avg. # of people on the trip	2.54	2.25
Avg. cost of lodging/night	\$45.78	
Avg. cost of food/trip	\$83.11	\$16.49
Avg. cost of ice/trip	\$10.34	\$4.93
Avg. cost of fuel & oil/trip	\$85.90	\$39.98

The economic figures are based on an expansion of the actual values reported by RCGL fishermen and are considered the best available estimates. The direct economic impacts described below are those that can be attributed only to shrimp landings by these fishermen. In some instances, the fishermen and the non-fishers who accompanied them, engaged in other, non-fishing activities. The total expenditures were adjusted based on the average proportion of people on the trip who actually engaged in fishing activity.

The expenditures shown in Table 8.8 relate to the overall proportion of shrimp landed. Other species were typically caught and kept along with the shrimp. The economic impact was based on the percent of shrimp in the total pounds of all species kept by the fishermen on any given trip where shrimp were landed. Shrimp accounted for 84% of the total catch on trips in which shrimp were landed. Expenditures by those who made overnight trips tended to be greater when compared to day trips because of the increased costs of lodging and meals. An average overnight trip lasted approximately 4 days and resulted in total expenditures of \$362.47 attributable to shrimp landings. Day trip RCGL anglers targeting shrimp had an average trip expenditure of \$61.40. RCGL anglers targeting shrimp took 2,096 trips in 2007. The total combined expenditures were approximately \$202,861. Lodging expenditures were left out of this estimate, as the sample size within the survey was too low (9 observations). The total combined economic impact of all RCGL trips for shrimp in 2007 was \$250,583.

8.1.2.3 Other Recreational Fisheries

Some people use cast nets to catch shrimp for personal consumption in addition to those who use cast nets to land shrimp for bait. Currently there are no data on these landings or their economic impacts, however there are ongoing attempts to survey recreational anglers who use cast nets.

9.0 SOCIOECONOMIC CHARACTERISTICS

9.1 SOCIAL IMPORTANCE OF THE FISHERY

9.1.1 Commercial fishermen

There are two primary sources of recent data or accounts available that help to explain the social importance of the commercial fishery. First is a book published on the shrimp industry in North Carolina, *Hard Times and a Nickel a Bucket: Struggle and survival in North Carolina's Shrimp Industry* (Maiolo 2004). Secondly, researchers at the DMF have been conducting in-depth socioeconomic interviews with commercial fishermen since 1999. More than 1,000 fishermen have been interviewed to date. In these nearly identical surveys, 175 fishermen within the most current dataset identified themselves as shrimp fishermen.

9.1.1.1 Historical importance

Elsewhere in this document is a history of the commercial shrimp fishery in North Carolina. The DMF surveys asked the fishermen for their opinion as to how historically important they think commercial fishing is to their community. On a scale of one to ten, with one being not at all important to ten being extremely important, the average rating across all 175 persons interviewed was 9.7, indicating almost universal agreement that fishing has been historically important to their community. When asked how much does their community support commercial fishing now (using the same 10-point scale), the rating was 8.1, indicating they largely feel supported.

9.1.1.2 Community reliance on the commercial fishery

North Carolina coastal communities rely significantly less on commercial fishing now than in the past (Maiolo 2004). This is the result of the development of the communities as multiple use zones, with retirement, light industry, recreation, and tourism becoming the dominant domains of the local economies. Fewer and fewer native born residents make a full time living as fishermen like those in previous generations. DMF studies found that among commercial shrimp fishermen, the average fisherman earned about 76% of his or her income from commercial fishing. More specifically the studies found that just over half (51%) were totally reliant on fishing for their incomes. This compares with data gathered in the late 1980s where nearly all full time fishermen were committed to fishing for nearly all (95%) of their income (Maiolo 2004).

The 175 shrimp fishermen in the DMF surveys came from 47 separate coastal communities. Table 9.1 shows the communities that had the greatest number of shrimp fishermen who participated in the survey. The largest number of fishermen in the surveys who fished for shrimp came from Sneads Ferry, followed by Beaufort, Supply, Atlantic, and Belhaven, all communities known to have sizable shrimp fleets.

Table 9.1. Most frequently cited communities where shrimp fishermen live (DMF Socioeconomics Program).

Community	Percent of Respondents
Sneads Ferry	11.60%
Beaufort	9.20%
Supply	9.20%
Atlantic	5.20%
Belhaven	4.60%
Wilmington	4.60%
Cedar Island	4.00%
Morehead City	3.50%
Wanchese	3.50%
Engelhard	3.50%
Harkers Island	3.50%
Sea Level	2.90%
Kill Devil Hills	2.30%
Marshallberg	2.30%
Newport	2.30%
Other	27.70%

Studies in the 1970s and 1980s revealed that shrimp fishermen engage in a variety of both land and water based activities. Fishing activities required moving from one target species to another as opportunities prevailed, even though shrimping involved most of the effort throughout the year (Maiolo 2004). DMF found that shrimp fishermen continue to engage in a variety of capture activities throughout the year. Like most of North Carolina's commercial fishermen, these fishermen tend to diversify the species they target, gears they use, and water bodies they fish. Shrimp constituted an average of 59% of the fishing income earned by these fishermen. Table 9.2 shows the preference for other species targeted and the average percent of fishing income earned by the 175 shrimp fishermen who participated in the surveys. Other species frequently targeted by these fishermen included blue crabs (*Callinectes sapidus*), clams (*Mercenaria mercenaria*), oysters (*Crassostrea virginica*), flounder (*Paralichthys spp.*), spot (*Leiostomus xanthurus*), and striped mullet (*Mugil cephalus*).

Table 9.2. Other prevalent species targeted by shrimp fishermen and average percent of fishing income made from non-target species (DMF Socioeconomics Program).

Species	Percent who land
Shrimp	100%
Blue Crabs	35%
Clams	29%
Oysters	29%
Flounder	28%

Table 9.2 (continued).

Species	Percent who land
Spot	24%
Striped Mullet	11%
Speckled Trout	10%
Croaker	8%
Scallops	5%
Whiting	4%
Bluefish	3%
Weakfish	3%
Striped Bass	3%

9.1.1.3 Perceived conflicts

There are largely two kinds of conflicts that have been measured, those between commercial fishermen and those between commercial fishermen and others who use the water. Conflicts between the users of the public resource are not uncommon, as no one individual owns the water, yet all citizens own the water and its resources. Conflicts tend to be reported more frequently as the demand for use of the resource increases.

Extensive competition, and often ill will between the full time fishermen, part time fishermen, and recreational fishermen, characterized the shrimp fishery according to research conducted in the seventies and eighties. At that time the competition was most intense in the estuaries in July, when shrimping was at its peak. The part timers and recreational users viewed ownership of the resource as much theirs as that of the full timers (Maiolo 2004).

Maiolo (2004) reported that because many commercial fishermen feel it is their inherent right to sell what they can catch, they frequently are in favor of regulatory actions that limit the activities of fishermen other than themselves. However, not all conflicts are resolved by relying on governmental regulatory agencies. In some areas of the state there is potential for conflicts between shrimp channel net fishermen and shrimp trawl fishermen. It is worth noting that off of Harkers Island in Carteret County, the channel net and trawl fishermen have a solution that works for all concerned. Channel netters stay far enough away from each other so that each is still able to land a reasonable catch. Channel netters carry lights that allow them to signal trawlers as to where they are working. In Sneads Ferry, fishermen commonly leave their net anchors out all season, often marked by cans. Although not mandated by outside regulatory authorities, actions such as these allow the channel netters and trawl fishermen to work in the same area and minimize conflicts within and between user groups (Maiolo 2004).

One of the purposes of the Fishery Reform Act was to address the intense and often uncontrolled competition and conflict between and among the user groups, and recent data indicate there has been some success in this area. The majority of the shrimp fishermen interviewed by DMF reported not having any conflicts at all within the past year (63%). The most common conflict reported was regarding state regulations (26%), followed by federal regulations (21%), and conflicts with other recreational (13%) and commercial (13%) fishermen (Figure 9.1). Several fishermen reported more than one type of conflict, therefore the percentages do not add up to 100%.




Figure 9.1. Reported conflicts of North Carolina Shrimp Fishermen (DMF Socioeconomics Program).

9.1.1.4 Perception of important issues

Shrimp fishermen interviewed by DMF were asked to state the business issues they found to be the most important. The most important issue to these fishermen was competition from imported seafood. Not surprising, these fishermen stated that low prices for seafood was a very important issue as well, followed closely by a feeling that state regulations were a major issue. Table 9.3 lists the 12 most commonly cited issues facing shrimp fishermen at the time of the survey. While the rising cost of fuel did not make this list, the rising cost of fuel has been cited as a major concern of commercial fishermen (all fisheries included) in past surveys. In 2007, “fuel price” was first added to the survey questionnaire as a possible issue of concern and was found to be the top concern of commercial fishermen in Core Sound (Crosson 2007b). Fuel prices were also found to be the top concern of commercial fishermen in 2008 and 2009 (Crosson 2009). The influence of rising fuel prices and cheap imported shrimp are major contributors to the decline in effort seen in the fishery since 2005.

Table 9.3. Fishing related issues considered most important to shrimp fishermen (DMF Socioeconomics Program).

Ranking	Issue
1	Imported seafood
2	Low prices for seafood

Table 9.3 (continued).

Ranking	Issue
3	State regulations
4	Weather
5	Federal regulations
6	Gear restrictions
7	Keeping up with rules and proclamations
8	Closed seasons
9	Bag limits
10	Quotas
11	Size limits
12	Overfishing

9.2 RECREATIONAL FISHERY

All data regarding the social importance of the fishery come from the last annual socioeconomic survey of RCGL fishermen conducted in 2007.

9.2.1 Historical importance

North Carolina has a long history of fishermen using commercial gear for recreational purposes. The RCGL license was put into effect in 1999 as a result of the Fisheries Reform Act of 1997. Prior to that, recreational fishermen who wished to use commercial gear purchased a commercial vessel license, but did not sell their catch. The RCGL fishermen who reported landing shrimp stated that they had been fishing commercial gear on average for 20 years. It is likely that using shrimp trawls for personal harvest has been occurring ever since commercial fishermen have been harvesting shrimp using trawls.

9.2.2 Community reliance on the recreational fishery

There are no data available to indicate the level of community reliance on the recreational shrimp fishery.

9.2.3 Perceived conflicts

Twenty eight percent of the RCGL fishermen felt that there was too much fishing gear in the water where they fish. An additional 19% weren't sure if there was too much gear in the areas where they fish. The remaining 53% felt that there wasn't too much gear in the water. Over 73% of all RCGL fishermen who land shrimp say they do not have any conflicts with commercial fishermen. Nearly 90% of them stated they do not have conflicts with recreational anglers.

9.2.4 Perception of important issues

RCGL fishermen were asked for their opinions about two issues they find to be important. Of those who land shrimp, 72% agreed with the statement that they ought to be allowed to use more commercial gear. An additional 6% disagreed indicating they felt they were allowed to use plenty of gear, while 22% indicated they were not sure whether they should be allowed to use more gear.

9.3 DEMOGRAPHIC CHARACTERISTICS

9.3.1 Commercial fishermen

Table 9.4 shows a summary of the demographic characteristics of the 175 shrimp fishermen interviewed by NCDMF. Nearly all of the shrimp fishermen were white males. They averaged 50 years old and had over 30 years fishing experience. The average shrimp fisherman was currently married and had at least a high school education. Approximately 29% of the fishermen had incomes of \$15,000 to \$30,001. Another 27% had total household incomes of \$30,001 to \$50,000.

Approximately 39% of the fishermen interviewed said they fished all year long. Of those who didn't fish all year, fishing activity was lowest from January through March. The peak fishing participation months for these fishermen were May through November. Fifty one percent of the fishermen indicated that fishing was their sole source of income. Of those who had other sources of income, the most frequently cited sources of additional income included carpentry, machinery mechanic, government, and retirement pensions.

Table 9.4. Demographic characteristics of commercial shrimp fishermen (DMF Socioeconomics Program).

Variable	Category Values	Average or Percent
Years Fishing		30.17 years
Age		50.46 years
Gender	Male	97%
	Female	3%
Race	White	96%
	Black	2.90%
	Hispanic	1.10%
Education Level	Less than HS	31.79%
	HS Grad	49.71%
	Some College	12.72%
	College Graduate	5.78%

Table 9.4. (continued).

Variable	Category Values	Average or Percent
Marital Status	Married	72.70%
	Divorced	15.10%
	Widowed	3.50%
	Never Married	6.40%
	Separated	2.30%
Total Household Income	Less than \$15,000	16.10%
	\$15,001 - \$30,000	28.70%
	\$30,001 - \$50,000	27.00%
	\$50,001 - \$75,000	14.90%
	More than \$75,000	6.32%
	Refused to answer	6.90%

9.3.2 Recreational fishermen

The average RCGL holder who targeted shrimp was 52.57 years old and 75% were born in North Carolina (Table 9.5). The vast majority were males. Most of these fishermen had at least some college education and had total household incomes of greater than \$30,000 per year. On average they had been using commercial gear for nearly 20 years.

Table 9.5. Demographic characteristics of RCGL holders who targeted shrimp in 2007. (DMF RCGL Survey Program).

Variable	Category Values	Average / Percent
Years of Experience Fishing		
Commercial Gear		19.8
Born in NC		75%
Age	<16 years	1%
	17 to 25	5%
	26 to 40	18%
	41 to 60	43%
	>60 years	33%
Marital Status	Married	72%
	Divorced	13%
	Widowed	5%
	Separated	2%
	Never Married	8%

Table 9.5 (continued).

Variable	Category Values	Average / Percent
Ethnic Group	Caucasian/White	98%
	Native American	2%
Gender	Male	94%
	Female	6%
Education	< High School	17%
	High School Diploma	25%
	Some College	33%
	College Diploma	25%
Total Household Income	< \$5,000	2%
	\$5,000 to \$15,000	6%
	\$15,001 to \$30,000	15%
	\$30,001 to \$50,000	24%
	\$50,001 to \$75,000	27%
	\$75,001 to \$100,000	15%
	> \$100,000	11%

9.4 DEFINITIONS

Commercial fishing – Fishing in which fish harvested, either in whole or in part, are intended to enter commerce or enter commerce through sale, barter or trade. Since 1994, a commercial fisherman in North Carolina is required to have a license issued by the North Carolina Division of Marine Fisheries (DMF) and is allowed only to sell to a licensed dealer.

Fishing trip – A period of time over which fishing occurs. The time spent fishing includes configuring, deploying, and retrieving gear, clearing animals from the gear, and storing, releasing or discarding catch. When watercraft are used, a fishing trip also includes the time spent traveling to and from fishing areas or locales and ends when the vessel offloads product at sea or returns to the shore. When fishing from shore or man-made structures, a fishing trip may include travel between different fishing sites within a 24-hour period.

Inflation-adjusted price and value – Inflation is a general upward price movement of goods and services in an economy, usually as measured by the Consumer Price Index (CPI). Ex-vessel prices and values can be adjusted (deflated) according to the CPI to remove the effects of inflation so that the value of a dollar remains the same across years. Inflation adjusted values allow for easier comparison and analysis of changes in values over time.

Nominal (Ex-Vessel) Value and Price- The total landed dollar amount of a species (or species landing condition and market category) in a given year. Example: 100 lbs. of shrimp at a PRICE of \$.50 per pound will have a VALUE of \$50 in the year the catch was landed. These values represent the amounts paid to a fisherman by a seafood dealer.

Recreational fishing – A recreational fishing trip is any trip for the purpose of recreation from which none of the catch is sold or bartered. This includes trips with effort but no catch. Fishermen who wish to use limited amounts of commercial fishing gear to harvest fish for personal consumption in joint and coastal waters under DMF jurisdiction are required to have a RCGL.

10.0 ENVIRONMENTAL FACTORS

10.1 HABITAT

Penaeid shrimp utilize a variety of estuarine and coastal ocean habitats as described in the life history section with variations in habitat preference due to location, season, and ontogenetic stage. Penaeid shrimp are found in most habitats identified by the North Carolina Coastal Habitat Protection Plan (CHPP) including: water column, wetlands, submerged aquatic vegetation (SAV), soft bottom, and shell bottom (Deaton et al. 2010). Each habitat is part of a larger habitat mosaic, which plays a vital role in the overall productivity and health of the coastal ecosystem. The CHPP focuses on the overall fish habitat and threats to the habitat while this FMP section describes habitat conditions, threats, or needs for the various life stages of penaeid shrimp. Although penaeid shrimp are found in all of these habitats, except for hard bottom, the usage varies by habitat. Additionally, these habitats provide the appropriate physicochemical and biological conditions necessary to maintain and enhance the penaeid shrimp population. The environmental preferences (salinity, temperature, oxygen, and substrate) were described in the Life History section. The slightly different preferences in bottom substrate and salinity affect their general position in the estuary and ocean. Each habitat provides ecological services that aid in maintaining and enhancing shrimp stock sustainability, and also influences the functioning of the ecosystem overall. Protecting the integrity of the entire system is therefore necessary to manage this species. Although ecosystem protection is of vital importance to penaeid shrimp, it may be difficult to detect a cause and effect relationship between habitat protection and enhancement and shrimp stock condition due to the large natural variation in environmental conditions in North Carolina, and the relatively short life cycle of penaeid shrimp.

The SAFMC designated inshore estuarine nursery areas, offshore marine habitats used for spawning and growth to maturity, and all connecting waterbodies as Essential Fish Habitat for penaeid shrimp (SAFMC 1998). Inshore nursery areas listed by SAFMC included wetlands, SAV, and subtidal and intertidal non-vegetated bottom (soft bottom). Designated Habitat Areas of Particular Concern (HAPC) for penaeid shrimp include all coastal inlets, all state designated nursery habitats used by shrimp, and state-identified overwintering areas. In North Carolina, specific HAPC includes SAV and estuarine shorelines. In areas lacking SAV, marsh with shell hash and mud bottoms and adjoining bottoms are of particular concern (SAFMC 1998).

10.1.1 Water column

Water column habitat is defined as “the water covering a submerged surface and its physical, chemical, and biological characteristics” (Deaton et al. 2010). Adult shrimp spawn offshore in ocean waters. Brown and pink shrimp spawn in deep water over the continental shelf, while white shrimp remain nearshore in relatively shallow water (SAFMC 1993). Adult shrimp are demersal oriented in all life stages, except as larvae and post-larvae. Larvae and post-larvae depend on ocean currents to be transported through inlets into estuarine nursery grounds. Inlets are critical bottlenecks through which shrimp and many other ocean-spawned larvae must pass to complete their life cycle (Hettler and Barker 1993). Inlets accessing Pamlico Sound are limited in number and therefore are particularly important to recruitment into Pamlico Sound and its tributaries. The time of spawning varies with species, with brown shrimp spawning earliest in winter and early spring, and white and pink shrimp spawning in late spring and early summer (Table 10.1). Shrimp are transported by water circulation throughout the estuary and back into the ocean. Water quality in estuarine waters affects the viability of shrimp populations.

Table 10.1 Spawning seasons for Penaeid shrimp species in North Carolina (Pattilo et al. 1997).

Species	Spawning season
Brown shrimp	Feb-Apr
Pink shrimp	Apr-Jul
White shrimp	May-Jul

10.1.2 Wetlands

Wetlands are defined as “areas that are inundated or saturated by an accumulation of surface or groundwater at a frequency and duration sufficient to support, and under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions” (federal regulations [40 CFR 230.3(t)]; Environmental Management Commission (EMC) rules [15A NCAC 2B .0202(71)], and Deaton et al. 2010). Estuarine wetlands, which include salt and brackish marsh and estuarine shrub/scrub, generally occur along the edge of estuaries and sounds. Riverine wetlands, which include freshwater marshes, bottomlands, hardwood forest, and swamp forest, generally occur in low-salinity to fresh water along streams, creeks, and rivers. It is estimated that over 95% of commercially harvested finfish and invertebrates in the United States are wetland dependent, a strong indication of their high habitat value (Feierabend and Zelanzy 1987).

The combination of shallow water and thick vegetation provides excellent nursery and foraging habitat for juvenile shrimp and many other fish species (Graff and Middleton 2003). Shallow wetlands also provide refuge from large fish predators and a safe corridor for migration to other habitats within the system (Rozas and Odum 1987; Mitsch and Gosselink 1993). Riparian wetlands are also highly effective and well recognized for their ability to trap and filter pollutants from upland runoff, and store, spread, and slow stormwater runoff prior to entering surface waters (Mitsch and Gosselink 1993).

Primary production in salt/brackish marshes is converted into shrimp production in two ways. Wetland plants decay into detritus, which accumulates in the wetlands and adjacent soft bottom areas and is a food source for shrimp and other small organisms. Also, nutrients from the broken down organic matter support growth of benthic microalgae on, between, and near wetland vegetation (Peterson and Howarth 1987). Productivity in riverine forested wetlands in North Carolina is reported to be lower than in estuarine marsh (Brinson 1977). It is estimated that 45% of salt marsh production is exported to the estuarine system in the form of detritus, dissolved organic matter, and transient fish, including shrimp (Teal 1962).

Shrimp are considered critically linked to marsh edge habitat (SAFMC 1998; Clark et al. 2004). Studies in Texas estuaries have documented that juvenile brown shrimp and white shrimp were more abundant along the salt/brackish marsh edge than in shell bottom, SAV, soft bottom, or inner marsh (Minello 1999; Rozas and Zimmerman 2000). Turner (1977) found a positive relationship between commercial yields of penaeid shrimp and the area of intertidal vegetation present at multiple estuarine locations. This suggests that preserving existing coastal wetlands and restoring former wetlands, where possible, would be directly beneficial to shrimp populations and harvest.

Riparian wetlands covered 7% of the land in coastal river basins, and riverine forested wetlands were the most abundant type. The Cape Fear, Neuse, and Albemarle river basins have the largest acreage of riparian wetlands, primarily riverine wetlands. Pamlico, Core, and Bogue

sounds, and estuaries south of Bogue Sound, have the highest percentages of estuarine wetlands. The largest acreage of salt/brackish marsh is in the Pamlico Sound region.

Distribution, size, and abundance of shrimp are monitored in the juvenile fish sampling program (Program 120). The majority of shrimp that are collected in this program are in close proximity to shallow wetland systems. Brown shrimp are widely distributed throughout North Carolina's estuaries in both low and high salinity areas, and support relatively higher concentrations in the Neuse tributaries, Core Sound, Stump Sound, and Intracoastal Waterway in Brunswick County. White shrimp abundance is most concentrated in the Cape Fear River estuary, Brunswick County estuaries, New River, and tributaries along the western shoreline of Pamlico Sound, north of the Tar-Pamlico River. Pink shrimp occur in relatively lower concentrations along the western shoreline of Pamlico Sound, Bogue Sound, New River, lower Cape Fear River, and Intracoastal Waterway in Brunswick County. However, current DMF sampling locations do not target the primary nursery grounds of pink shrimp, and therefore, may not accurately represent juvenile pink shrimp distribution and abundance. In addition to Program 120 shrimp is sampled in program 510 for management purposes. Due to the way this data is collected it cannot be used for determining distribution or abundance indices but it can be use to determine when to open areas to trawling.

10.1.3 Soft bottom

Soft bottom habitat is defined as “unconsolidated, unvegetated sediment that occurs in freshwater, estuarine, and marine systems” (Deaton et al. 2010). Sediment composition varies with geomorphology and location within the system and may be a factor in juvenile shrimp distribution. Juvenile white shrimp prefer shallow muddy substrate. In contrast, juvenile brown shrimp prefer peat and muddy bottoms but also occur where the bottom is composed of sand, silt, clay, or shell fragments (SAFMC 1993). Although soft bottom habitat is defined as “unvegetated” and lacks visible structural habitat, the surface sediments support an abundance of microscopic plants (benthic microalgae) and numerous burrowing animals hidden below the surface.

Soft bottom plays a very important role in the ecology of estuarine ecosystems as a storage reservoir of nutrients, chemicals and microbes. Biogeochemical processing and recycling establishes a filter to trap and reprocess natural and human-induced nutrients and toxic substances or release them into the water column (Matoura and Woodward 1983), allowing chemicals to pass quickly or over several seasonal cycles through the estuary (Uncles et al. 1988). Soft bottom also provides a rich food base for juvenile and adult shrimp due to the numerous plants and animals living on and in the sediment (Peterson and Peterson 1979; Currin et al. 1995). At different life stages, shrimp feed on various organisms in bottom sediments, including microfauna such as protozoans, meiofauna, such as nematodes and copepods, and macrofauna such as amphipods, polychaetes, and other crustaceans (Peterson and Peterson 1979). Once shrimp enter ocean waters, they continue foraging on subtidal bottom, particularly on muddier bottom. Although there is little structure to hide behind, shrimp can find refuge from predators by remaining on very shallow flats that predators cannot access or by burrowing beneath soft bottom during the day, and actively foraging and moving at night (Peterson and Peterson 1979; Ross and Epperly 1985).

Soft bottom also plays a key role as a nursery area for shrimp. Primary nursery areas for juvenile brown, white, and to a lesser extent pink shrimp, include shallow soft bottom habitat, usually adjacent to wetlands (Noble and Monroe 1991). Most larval settlement occurs in the uppermost portion of shallow creek systems. Areas that have been documented to consistently

support large numbers of juvenile shrimp and other species have been designated by the MFC as Primary Nursery Areas (PNAs) (Figure 10.1a-d). For 1990-2010, data from DMF's ongoing juvenile fish monitoring program indicate that brown shrimp is one of the most abundant species found along the entire coast, along with spot, Atlantic croaker, pinfish, bay anchovy, blue crab, silver perch, and Atlantic menhaden. Brown shrimp were the fourth most abundant species in the northern region while white shrimp were nineteenth. In the southern portion of the coast, brown shrimp were the third most abundant and white shrimp the thirteenth most abundant species (DMF, unpub. data). During 1990-2010, a total of 184 species were collected from the northern juvenile sampling stations and 144 species was collected from the south juvenile sampling stations (DMF, unpub. data). Consequently, protection of these areas is a high priority for shrimp management, as well as other species.

The loss of structured habitats, such as SAV and shell bottom, over time, has most likely led to gains in the amount of soft bottom habitat, but it may be of lower quality in some areas if toxins have accumulated in the sediment. Activities that lead to the deepening, loss, or chemical contamination of shallow and intertidal habitat are the greatest threat to this habitat.




Figure 10.1a Shrimp nursery areas, including MFC designated nursery areas and SAV beds, in Pamlico Sound NC.




Figure 10.1b Shrimp nursery areas, including MFC designated nursery areas and SAV beds, for the Tar/Pamlico and Neuse Rivers.




Figure 10.1c Shrimp nursery areas, including MFC designated nursery areas and SAV beds, for the Core Banks to Topsail Island.




Figure 10.1d Shrimp nursery areas, including MFC designated nursery areas and SAV beds, for the Topsail Island to South Carolina.

10.1.4 Submerged Aquatic Vegetation (SAV)

Submerged aquatic vegetation (SAV) is a fish habitat dominated by one or more species of underwater vascular plants. The NCMFC define SAV habitat as submerged lands that:

- “(i) are vegetated with one or more species of submerged aquatic vegetation including bushy pondweed or southern naiad (*Najas guadalupensis*), coontail (*Ceratophyllum demersum*), eelgrass (*Zostera marina*), horned pondweed (*Zannichellia palustris*), naiads (*Najas* spp.), redhead grass (*Potamogeton perfoliatus*), sago pondweed (*Stuckenia pectinata*, formerly *Potamogeton pectinatus*), shoalgrass (*Halodule wrightii*), slender pondweed (*Potamogeton pusillus*), water stargrass (*Heteranthera dubia*), water starwort (*Callitriche heterophylla*), waterweeds (*Elodea* spp.), widgeongrass (*Ruppia maritima*) and wild celery (*Vallisneria americana*). These areas may be identified by the presence of above-ground leaves, below-ground rhizomes, or reproductive structures associated with one or more SAV species and include the sediment within these areas; or
- (ii) have been vegetated by one or more of the species identified in Sub-item (4)(i)(i) of this Rule within the past 10 annual growing seasons and that meet the average physical requirements of water depth (six feet or less), average light availability (secchi depth of one foot or more), and limited wave exposure that characterize the environment suitable for growth of SAV. The past presence of SAV may be demonstrated by aerial photography, SAV survey, map, or other documentation. An extension of the past 10 annual growing season criteria may be considered when average environmental conditions are altered by drought, rainfall, or storm force winds.” [2009 MFC rule 15A NCAC 03I .0101 (4)(i)].

Under NCMFC rules, SAV is a Critical Habitat Area [MFC rule 15A NCAC 03I .0100 (b)(20)].

SAV enhances the ecosystem by stabilizing and trapping sediment, reducing wave energy and cycling nutrients within the system (Thayer et al. 1984). The three-dimensional structure provides a surface for small plants and animals to attach to and provides a safe refuge and foraging area for a large number of juvenile fish and invertebrates (SAFMC 1998). Beds of SAV also produce large quantities of organic matter, which supports a complex food base for numerous fish and other organisms (Thayer et al. 1984). Similar to wetlands, the structure of SAV grass blades provides an excellent nursery area and enhances safe corridor between habitats, reducing predation (Micheli and Peterson 1999). While white shrimp may utilize freshwater SAV to some extent, brown and pink shrimp primarily utilize estuarine SAV because of salinity preferences.

Many important commercial and recreational fishery species use SAV as a nursery (Thayer et al. 1984). The blades of SAV provide protection and food for post-larvae and juvenile shrimp. Of the three penaeid shrimp species, SAV is particularly critical as a nursery area for pink shrimp (Murphey and Fonseca 1995). Juvenile pink shrimp abundance was greater in estuarine SAV beds compared to soft bottom, marsh edge, or shell bottom (Minello 1999). Brown shrimp also utilize SAV to some extent. Data from Texas estuaries suggest that brown shrimp show greater preference for SAV rather than marsh edge where both habitats occur (Clark et al. 2004). The configuration of a grass bed may also be a factor in juvenile and adult shrimp distribution (Murphey and Fonseca 1995). Additional sampling in SAV is needed to better assess the relationship of SAV condition and spatial changes to shrimp use of SAV habitat. Research is currently underway at UNC-IMS examining SAV, shellfish, and wetland utilization by fishes.

Several studies in North Carolina have shown that shrimp abundance was greater on SAV beds than on oyster beds (Ellis et al. 1996) or unvegetated soft bottom (Murphey and Fonseca 1995). These studies showed similar trends for other species as well. In Florida Bay, changes in animal abundances were compared between the 1980s and 1990s when significant loss of SAV occurred (Matheson et al. 1999). A decrease in SAV coverage appeared to result in a decrease in abundance of small fish and invertebrates that live within the seagrass canopy (such as shrimp and pipefish), while larger demersal predatory fish (such as toadfish and sharks) increased. Similarly, increases in SAV density were characterized by significant increases in crustaceans. In another study in Florida Bay, reductions in pink shrimp abundance were greater in seagrass die-off areas than in nearby undamaged or recovering areas (Roblee and DiDomenico 1992).

The presence of SAV may be the reason pink shrimp can overwinter in temperate North Carolina and thus supports North Carolina's spring pink shrimp harvest (T. Murphey, DMF, pers. com. 2012). Similarly, south Florida's seagrass and mangrove dominated estuaries also play a role as nursery areas for pink shrimp in both the Tortugas and Sanibel fisheries. In contrast, in South Carolina and Georgia, where no SAV is present, pink shrimp comprise a negligible portion of the shrimp landings (T. Murphey, DMF, pers. com. 2012).

From 1978-1993 the average pink shrimp landings comprised 27% of North Carolina's total shrimp landings, average pink shrimp landings from 1999-2010 declined to only 3% of total shrimp landing. Florida is also experiencing declines in its pink shrimp fishery as well. The reason for the decline in pink shrimp landings is thought to be due to higher fuel prices not allowing trawlers to perform more experimental trawls (A. Bianchi, NCDMF, pers. com. 2012) or colder winters may have decreased pink shrimp growth or increased mortality (T. Murphey, NCDMF, personal communication). The location of SAV beds in North Carolina is shown in Figure 10.1a-d, along with the MFC designated nursery areas.

10.1.5 Shell bottom

Shell bottom is defined in the CHPP as "estuarine intertidal or subtidal bottom composed of surface shell concentrations of living or dead oysters (*Crassostrea virginica*), hard clams (*Merceneria merceneria*), and other shellfish" (Deaton et al. 2010). In the 1990s, fisheries management agencies began to formally recognize shell bottom habitat as critical to fisheries production. The NCMFC, SAFMC, and ASMFC all recognize the importance of shell bottom.

Common terms used to describe shell bottom habitats in North Carolina are "oyster beds," "oyster rocks," "oyster reefs," "oyster bars," and "shell hash." Shell hash is a mixture of sand or mud with gravel and/or unconsolidated broken shell (clam, oyster, scallop, and/or other shellfish). Shell bottom is enhanced in some areas by the addition of cultch material. Cultch material (hard material to which oysters attach) can consist of oyster, clam, or scallop shells; gravel or marl; or other hard materials. Cultch exists naturally, as shell hash and oyster rocks. DMF's Shellfish Rehabilitation Program staff also plant cultch to enhance and restore estuarine shell bottom for oyster and hard clams.

Shell bottom is both intertidal and subtidal, and can consist of fringing or patch reefs (Coen et al. 1999). Intertidal oyster reefs in the central and southern estuarine systems may only be a few oysters thick. However, subtidal oyster mounds in Pamlico Sound may have been several meters tall (Lenihan and Peterson 1998). In North Carolina, oysters attach to and accumulate on existing oyster beds, other shell, outcropping of fossil shell beds, exposed *Spartina* roots, pilings, and rip-rap (DMF 2001). Intertidal oyster reefs in North Carolina may occur along the

edges and points of salt marsh, between salt marsh and seagrass beds, or as isolated reef features, away from other structure (Grabowski et al. 2000).

Shell bottom provides many important functions that enhance the health of the entire ecosystem for fishery and non-fishery species. Oysters filter sediment and pollutants from the water column, enhancing water quality and improving conditions for SAV growth (Coen and Luckenbach 1998). The hard multi-faceted shell structure aids in reducing wave energy, stabilizing sediment, and reducing shoreline erosion (Lowery and Paynter 2002). Oysters, like SAV and benthic microalgae, facilitate storage and cycling of nutrients. This process reduces the likelihood of coastal eutrophication and its detrimental effects on fish and fisheries. Oyster beds also increase shoreline complexity, modify circulation patterns, and enhance fish use of marsh edge habitat (Grabowski et al. 2000).

The complex three-dimensional structure of shell bottom provides protective cover for juvenile and adult shrimp. The shell structure also provides an area for small plant and invertebrate attachment, which shrimp may feed on or hide among (Meyer et al. 1996; Lenihan and Peterson 1998; Coen et al 1999). However, predatory finfish around the reefs feed, in part, on penaeid shrimp (Grabowski et al. 2000). Fringing shell bottom or shell hash also serves as a nearshore corridor between habitats such as salt marsh and SAV, which shrimp also utilize (Coen et al. 1999; Micheli and Peterson 1999).

Brown, white, and pink shrimp have been documented to utilize shell bottom habitat in South Carolina and Texas estuaries (Coen and Luckenbach 1998; Zimmerman et al. 1989), although shell bottom does not appear to be the preferred habitat, compared to salt marsh edge or SAV (Minello 1999). In North Carolina, some studies indicate use of oyster beds by pink, white, and brown shrimp (Meyer et al. 1996; Grabowski et al. 2000; Lenihan et al. 2001). Analysis of these studies in Peterson et al. (2003) concluded that pink, white, and brown shrimp were not recruitment or growth enhanced by the presence of shell bottom. Shervette and Gelwick (2008) observed higher numbers of white shrimp in oyster bottom than in non-vegetated bottom. In sounds and the lower portions of estuaries where SAV is not present, shell bottom may be more critical to penaeid shrimp. In addition, the ecosystem benefits provided by the habitat would still indirectly enhance shrimp populations.

Oysters are found along a majority of the North Carolina coast from extreme southeastern Albemarle Sound to the estuaries of the southern part of the state to the South Carolina border (DMF 2001). Oyster reefs occur at varying distances up North Carolina's estuaries, depending upon salinity, substrate, and flow regimes. In the wind-driven Pamlico Sound system north of Cape Lookout, oyster reefs consist overwhelmingly of subtidal beds. South of Cape Lookout, subtidal rocks also occur in the New, Newport, and White Oak rivers (DMF 2001). Extensive intertidal oyster rocks occur in North Carolina's southern estuaries, where the lunar tidal ranges are higher. Substantial shell hash is present in New River, eastern Bogue Sound, and along the edges of many streams and channels, such as portions of the Atlantic Intracoastal Waterway (IWW) in the southern coastal area. In the Albemarle-Pamlico estuary, oysters are concentrated in the lower portion of Pamlico Sound tributaries, along the western shore of Pamlico Sound, and to a lesser extent, behind the Outer Banks (Ross and Epperly 1986).

The current distribution of shell bottom is much less than what historically occurred (Newell 1988). Mechanical harvesting of oysters (oyster dredging) was the primary and initial cause of habitat loss (DMF 2001). Most shell bottom losses have been to subtidal beds in Pamlico Sound, where DMF has also found declines in oyster recruitment. Although mechanical harvesting of oysters has been greatly restricted, reefs have not recovered, possibly due to

stress from water quality degradation and increased occurrence of disease (Dermo, MSX) (DMF 2001). Oyster dredging removes oysters and reduces the vertical profile of oyster rocks, increasing the susceptibility of remaining shell bottom at that location to low DO and possible mortality (Lenihan and Peterson 1998; Lenihan et al. 1999). Although commercial oyster dredging has been greatly reduced, current activities continue to reduce and degrade a habitat that is utilized by shrimp. Hand harvest methods for oysters and clams can also be destructive, but on a much smaller scale. Other bottom disturbing fishing gears, such as trawls, prevent the re-establishment of oyster reefs within their historic range.

To offset some of these reductions, the NCDMF began an oyster sanctuary program in 1996. As of 2012, the DMF has established and developed 10 Oyster sanctuaries with a total 198 acres of permitted sanctuary bottom (P. Holmlund. NCDMF. personal communication). Certain bottom disturbing fishing gears such as trawls, long haul seines and swipe nets are prohibited within these sanctuaries. The harvest of shellfish by any means is also prohibited within these sanctuaries. These sanctuaries are located around Pamlico Sound and constructed of multiple, high profile mounds using mostly Class B Riprap (fossil stone) and the use of shell and seeded shell as part of the research needs. The Nature Conservancy (TNC), the National Marine Fisheries Service (NMFS) Hurricane grant 2001-2006, state appropriations through DMF, CRFL grants, and other mitigation sources provided funding. The DMF has also partnered with NCCF in several oyster restoration projects, including the large federal stimulus project. Oyster Sanctuaries are designated and delineated under North Carolina Marine Fisheries Rule 15A NCAC 03R .0117 and are protected from damaging harvest practices under rule 15A NCAC 03K .0209. The oyster sanctuary program should continue to be a high priority for funding.

Other causes of shell bottom losses include dredging for navigation channels or marina basins. These activities can physically remove or damage existing shell bottom or result in turbidity that clogs oyster gills or covers sediment completely. Hydrologic modifications in the Neuse and Pamlico rivers have decreased salinity in the downstream portions of those rivers and resulted in a downstream displacement of oysters since the 1940s (Jones and Sholar 1981). While drainage for agriculture has changed little in recent years, drainage for urban/suburban development is increasing steadily.

10.1.6 Hard Bottom

Hard bottom habitat is defined in the CHPP as “exposed areas of rock or consolidated sediments, usually colonized by a thin veneer of live or dead biota, and generally located in the ocean rather than in the estuarine system” (Deaton et al. 2010). At this time there is no documented evidence of penaeid shrimp using hard bottom.

10.2 HABITAT CONCERNS

10.2.1 Wetland Loss

It is estimated that as much as 34-50% of North Carolina’s original wetland coverage has been lost, primarily due to ditching, channelization, and filling for agriculture and development (Dahl 1990; DWQ 2000). According to the North Carolina Division of Water Quality (DWQ 2000), approximately 88% of salt/brackish marsh, 81-88% of riverine forested wetlands, and 48% of pocosins remain. From the early 1800s to the early 1900s, ditching and draining for agriculture accounted for the majority of wetland losses (Heath 1975). From 1950 to the 1990s, conversion of wetlands to managed forest and agriculture accounted for 53% and 42%, respectively, with commercial and residential development activities responsible for the remaining 5% (Bales and

Newcomb 1996). Since 1990, losses from agriculture and forestry decreased, but losses from development increased. The primary threats to wetland habitat today are dredging, filling, and hydrological alterations associated with development. Although the rate of wetland loss has slowed, losses continue to occur. Mitigation for permitted losses and voluntary restoration efforts in some areas has partially offset some of the recent losses.

One activity that has a cumulative impact on wetlands is estuarine shoreline stabilization. Hardened estuarine shorelines cause gradual, long-term wetland loss by limiting sediment inputs needed for maintenance and expansion of wetlands, and by blocking landward migration as sea level rises. Garbisch et al. (1973) showed that marsh vegetation waterward of bulkheads experienced a 63% post-construction mortality due to stress from increased turbulence and scour resulting from vertical hardened structures. Scouring action at the toe of bulkheads also deepened the adjacent water, thus reducing or eliminating intertidal habitat. The added turbulence at the base of bulkheads and deepened water depth prevents vegetation from reestablishing after construction (Knutson 1977). Several studies have found that abundance of shrimp and other organisms adjacent to bulkheaded shorelines was much less than what occurred adjacent to unaltered naturally vegetated shorelines (80-300% less) (Mock 1966; Gilmore and Trent 1974; Peterson et al. 2000). The difference was attributed to lower abundance of organic detritus and small benthic invertebrates, deeper water, and less intertidal vegetation.

Ongoing initiatives such as wetland restoration, land acquisition and preservation, and agricultural cost-share BMPs (Best Management Practices) need to be enhanced. These initiatives should be made available through continued education about the use of alternatives to vertical bulkheads. These alternatives include marsh sills which prevent erosion by allowing wetlands to dissipate the energy and still provide ecosystem functions that are lost through other means of shoreline stabilization. There should also be additional initiatives implemented to protect and enhance wetland habitat. The many fishery and water quality functions provided by wetlands make their preservation and restoration along North Carolina's coast a high priority for protection of all coastal fish habitats.

10.2.2 Soft Bottom Impacts

Soft bottom habitat may be affected by marina and dock facilities through alteration of the shoreline configuration, circulation patterns, and subsequently, changes in bottom sediment characteristics (Wendt et al. 1990). Because benthic microalgae, an important component of primary production in soft bottom habitat, are light-dependent, bottom sediments in dredged marinas will have reduced light availability due to the deeper water depth and shading from docking structures. A study estimating macroalgae and microalgae productivity before and after construction of a marina in Long Island Sound found that microalgae production on soft bottom would decline by 48% post-construction and macroalgae production would decline by 17% (Ianzuzzi et al. 1996). However, the authors concluded that some of this loss would be offset by additional microalgal production on hard structures in the marina. Operation of a marina can also affect productivity of the soft bottom community due to introduction of heavy metals, hydrocarbons, and bacteria (Chmura and Ross 1978; Marcus and Stokes 1985; Voudrias and Smith 1986). Heavy metals and hydrocarbons are toxic to many soft bottom dwelling invertebrates and benthic feeding fish (Weis and Weis 1989). Dissolved oxygen (DO) may become depleted or below optimum thresholds in dredged marina basins and channels. A North Carolina marina study found significantly lower DO concentrations (less than 5.0 mg/l) inside some marinas compared to samples from outside marinas (DEHNR 1990). Cumulatively,

docks may also negatively impact shrimp populations (Sanger and Holland 2002). Research is needed to better assess the impacts of multiple docks on shrimp and other species. In addition to impacting wetlands, estuarine shoreline stabilization can degrade soft bottom habitat by reducing or eliminating the intertidal zone, deepening shallow soft bottom habitat, or contaminating sediment from leaching of toxic preservatives from wood structures (Weis et al. 1998). Multiple studies have shown that the diversity and abundance of invertebrates and juvenile fish over soft bottom are reduced adjacent to bulkheaded areas (Mock 1966; Ellifrit et al. 1972; Gilmore and Trent 1974; O'Rear 1983; Byrne 1995; Peterson et al. 2000; Waters and Thomas 2001). Beach nourishment along ocean shorelines can alter the sediment composition of nearshore soft bottom to a condition less favorable for shrimp or result in a temporary reduction in food availability (Hackney et al. 1996). Local fishermen have noted a shift in shrimp distribution to waters further offshore at Carolina Beach and Wrightsville Beach, where storm damage reduction projects have been ongoing for many years. This change may be associated with a shift in sediment composition from muddy to sandy substrate.

While MFC rules are designed to minimize commercial fishing gear impacts to fisheries habitat, these restrictions primarily focus on restricting the use of highly destructive bottom disturbing gear from most structural habitats such as oyster or SAV beds. Soft bottom habitat, because of its low structure and dynamic nature, has historically been considered the most appropriate location to use bottom disturbing gear. Existing fishery rules that restrict bottom disturbing gears in soft bottom habitat include prohibition of trawls, dredges, and long haul seines in PNAs, [15A NCAC 3N .0104] and prohibition of trawls, or mechanical shellfish gear in crab spawning sanctuaries [15A NCAC 3L .0205] in the five northern-most inlets of North Carolina during the blue crab spawning season (March-August).

Fishing gears documented to have the greatest potential to damage or degrade soft bottom or other habitats are dredges, followed by trawls (DeAlteris et al. 1999; Collie et al. 2000). Bottom trawling is used more extensively than dredging on soft bottom habitat in both estuarine and coastal ocean waters. Shrimp trawling accounts for the majority of bottom trawling effort in North Carolina and was addressed in the 2006 Shrimp FMP. Management strategies in the 2006 Shrimp FMP that addressed both habitat impacts as well as bycatch concerns were area specific and included trawl size restrictions, a phase in period for otter trawls to be converted to skimmer trawls, designation of a special secondary nursery area, and the establishment of additional no trawling areas.

Various types of dredges used on soft bottom habitat in North Carolina cause similar bottom disturbance: crab dredges, oyster dredges, and hydraulic clam dredges. Because of the gears' teeth, crab and oyster dredges can dig deep into the sediment and cause extensive sediment disturbance. Mechanical methods for the taking of crabs is prohibited in designated Crab Spawning Sanctuaries from March through August. Although the amount of fishing effort is low, this gear is documented to cause significant damage (DeAlteris et al. 1999; Collie et al. 2000).

Hydraulic clam dredging, as well as clam "kicking", a specialized type of trawl, creates trenches and mounds of discarded material in soft bottom habitat, redistributing and resuspending sediment (Adkins et al. 1983). Water jets from the hydraulic dredge can penetrate 18 inches into bottom sediments, and uproot any biotic structure present (Godcharles 1971). Dredge tracks can remain present from a few days to more than one year, and recolonization by vegetation can take months to begin. Recruitment of clams and other benthic invertebrates does not appear to be affected by hydraulic dredging (Godcharles 1971). Because of the severe impacts to habitats, both hydraulic clam dredging and clam kicking are restricted to open sand and mud bottoms, usually deeper waters, including areas frequently dredged as

navigational channels. Overwintering pink or white shrimp could potentially be affected by this activity, although they usually overwinter in shallow vegetated areas. However, Freeman (1988) examined the effects of clam kicking on pink shrimp in Core Sound and found no significant differences in mean CPUE between an area opened to mechanical harvest and an area closed to mechanical harvest.

10.2.3 Submerged Aquatic Vegetation Impacts

The amount of SAV in North Carolina was estimated to be between 134,000 and 200,000 acres around 1990 (Orth et al. 1990; Ferguson and Wood 1994). However the current spatial distribution and acreage of SAV may be somewhat different since some areas that historically supported SAV were not mapped, and changes may have occurred since the original mapping. Along the Atlantic coast, North Carolina supports more SAV than any other state, except for Florida. The majority of SAV occurs in eastern Pamlico Sound and Core Sound in high salinity waters (Figure 9.1a-d). Because light is the primary limiting factor affecting its distribution, SAV is restricted to relatively shallow waters, usually less than 1 m in depth.

Historical accounts indicate that there have been large-scale losses of SAV in North Carolina's low salinity tributaries on the mainland side of Pamlico Sound and along much of the shoreline of western Albemarle Sound (North Carolina Sea Grant 1997; J. Hawkins, DMF, personal communication) while the high salinity grass beds to the east appear relatively stable (Ferguson and Wood 1994). Loss of low salinity SAV habitat could negatively affect white or brown shrimp. Impacts to high salinity SAV beds could be especially detrimental to pink shrimp. Protection, enhancement, and restoration of this habitat are high priorities for sustained shrimp populations.

The greatest threat to SAV is large-scale nutrient enrichment and sediment loading, which increases turbidity, reduces light penetration, and subsequently impacts SAV growth, survival, and productivity (Goldsborough and Kemp 1988; Kenworthy and Haunert 1991; Funderburk et al. 1991; Stevenson et al. 1993). Catastrophic losses of seagrass beds have been correlated with these water quality problems in other states in the past (Twilley et al. 1985; Orth et al. 1986; Durako 1994). Nutrient enrichment and/or increased sediment loads impact SAV growth, survival, and productivity by increasing chronic turbidity in the water column from suspended sediment or phytoplankton associated with algal blooms. Also, sediment, epiphytes, or drift algae can cover the surface of blades (Dennison et al. 1993; SAFMC 1998; Fonseca et al. 1998). Elevated nitrogen concentrations have also been shown to be toxic to eelgrass (Burkholder et al. 1992). In North Carolina, most of the low salinity areas that have experienced large reductions in SAV coverage (Tar-Pamlico River and Neuse River) are also designated Nutrient Sensitive Waters. Once SAV is lost, increased turbidity and sediment destabilization can result in accelerated shoreline erosion and make recolonization more difficult (Durako 1994; Fonseca 1996). Therefore prevention of any additional SAV loss through water quality maintenance and improvement is a high priority for shrimp management.

Increased sediment and nutrient loading in the water column can enter coastal waters from point source discharges, nonpoint stormwater runoff, or resuspension of bottom sediments. Specific sources that contribute to increased sediment loading include construction activities, unpaved roads, road construction, golf courses, uncontrolled urban runoff, mining, silviculture, row crop agriculture, and livestock operations (DWQ 2000). Urbanization can increase the flow and velocity of stormwater runoff, which in turn leads to increased stream bank erosion. Stream bank erosion is a significant source of sediment loading (DWQ 2000). Specific sources that contribute to increased nutrient loading include agricultural and urban runoff, wastewater

treatment plants, forestry activities, and atmospheric deposition. Nutrients in point source discharges are primarily from human waste and industrial processes. The primary contributors of nutrients from non-point sources are fertilizer and animal wastes (DWQ 2000).

In addition to effects from water quality degradation, SAV can be removed or damaged by water-based activities. Dredging for navigational channels, marinas, or infrastructure such as bridges, submarine pipelines, or cables can result in large, direct losses of SAV. Docks constructed over SAV can cause immediate loss during construction or gradual loss due to shading effects. Several studies in Florida have shown that SAV was significantly reduced or eliminated under and around docks that were less than 5.5 ft above mean high water or where light received was less than 14% of the surface light availability (Loflin 1995; Shafer 1999). In addition to direct damage from docks and marinas, indirect damage to SAV can result from boating activity associated with these structures. Shoals and other shallow bottoms supporting SAV may become scarred as boating activity to and from the docking areas increases. Boat wakes can destabilize and erode SAV beds, or resuspend sediment, reducing light penetration. As additional docks and marinas are constructed along the coast, the potential for boating-related damage increases.

North Carolina has implemented standards for dock construction to minimize impacts to SAV, including dock height above the water and minimum water depth. In North Carolina, the depth of water at the dock end is not considered in Coastal Resource Commission (CRC) rules. To minimize shading effects to wetland plants, CRC rules require a dock height of at least three feet (0.91 m) above the wetland substrate, and a pier width of no greater than six feet (1.83 m) [CRC rule 15A NCAC 07H.0208 (6)]. However, there is no requirement for height above the water surface. Results from Connell and Murphey (2004) indicate that current dock designs over SAV beds in North Carolina result in a reduction in SAV coverage and density. The DCM rules (15A NCAC 07H .1205) allow docks to be constructed over SAV where there is at least -2' normal water level (NWL). Dock criteria should be evaluated by CRC to determine if existing requirements are adequate for SAV survival and growth and what changes would be needed to allow adequate light beneath docks. The permit requirements for docks and piers may need to be changed accordingly.

Several bottom disturbing fishing gears have the potential to destroy or damage SAV. The DMF issued a report on shrimp and crab trawling impacts (DMF 1999). Also, the Fisheries Moratorium Steering Committee's Habitat Subcommittee identified specific habitat impacts from various commercial and recreational fishing gears used in North Carolina waters, and made recommendations to minimize such impacts (MSC 1996). The Fisheries Moratorium Steering Committee presented the summary of findings to the Joint Legislative Commission on Seafood and Aquaculture of the General Assembly. Fishing gear found to be potentially damaging to SAV is listed in Table 10.2.

Table 10.2 Fishing gears used in North Carolina identified as potentially damaging to submerged aquatic vegetation habitat. (Source: MSC 1996).

Severe damage	Moderate damage	Low damage or unsure
Oyster dredge	Crab trawl	Long haul seine
Crab dredge	Clam Tongs	Otter trawl
Hydraulic clam dredge		Clam hand rake
Clam trawl (kicking)		Bay scallop dredge (very little)
Bull rake		

Damage from fishing gear varies in severity. Hand gear, such as bull rakes and large oyster tongs, can uproot SAV and cause substantial damage, but generally to smaller areas than mechanical gears (Thayer et al. 1984). Current MFC rules prohibit use of rakes more than twelve inches wide or weighing more than six pounds SAV [MFC rule 15A NCAC 03K.0304 (a) (2)]. Use of hand rakes and clamming by hand are allowed.

Mobile gear, such as long haul seines or bottom trawls, can shear or cut the blades of SAV, or uproot plants without major disruption of the sediment (ASMFC 2000). Shearing of above-ground plant biomass does not necessarily result in mortality of SAV, but productivity is reduced since energy is diverted to replace the damaged plant tissue, and the nursery and refuge functions are reduced in the absence of structure. Other fishing practices can cause severe disruption of the sediment and damage the roots of SAV. Gears that disturb the sediment and below-ground plant structures, like toothed dredges, heavy trawls, and boat propellers, may cause total loss of SAV in the affected area, requiring extensive time to recover (ASMFC 2000). SAV can also be buried by excessive sedimentation associated with trawling, dredging, and propeller wash. High turbidity from use of bottom-disturbing fishing gear can reduce water clarity, affecting SAV growth, productivity, and in some cases, survival (ASMFC 2000). Although some areas such as the soundside of the outerbanks have been closed to protect SAV, other areas should be periodically evaluated to determine if boundaries need to be adjusted to avoid SAV. At this time, most of the SAV is located in areas that have been designated as PNA, SNA, or SSNA, where there is some protection preventing trawling through SAV. For more information on PNA, SNA, or SSNA see section 11 Management of the Stock. The remaining areas that do not prevent trawling in SAV are in areas in proximity to Oregon Inlet (Figure 10.1a) and Bogue Sound (Figure 10.1b) where the depths may be shallow for trawling.

All toothed dredges can cause severe damage when pulled through SAV. Because oyster dredges, crab dredges, and hydraulic clam dredges severely impact bottom structure, there are strict limits on their use in North Carolina. Use of crab dredges is restricted to an area in northern Pamlico Sound southwest of Oregon Inlet [MFC rule 15A NCAC 03R.0109] that excludes SAV beds. Use of oyster dredges is currently restricted to parts of Pamlico Sound and its tributaries. The majority of high salinity SAV occurs in areas where mechanical methods for oyster harvesting are prohibited. Amendment I of the Oyster FMP addressed oyster dredging in the shallow bays of western Pamlico Sound where dredging is now only allowed in designated areas based on a water depth criteria of six feet or more. This management strategy provides protection to brackish and freshwater SAVs.

Clam kicking can also severely impact SAV since substrate is displaced by propeller backwash (Guthrie and Lewis 1982). Peterson and Howarth (1987) found that clam kicking significantly reduced plant biomass in eelgrass and shoalgrass beds. It is likely that SAV was damaged by kicking in the past since this technique has been used in North Carolina for over 60 years. Effort was high in areas known to support SAV (Carteret County), and kicking vessels tended to operate in shallow waters (Guthrie and Lewis 1982). Because of the severe disturbance to the bottom, clam kicking is restricted to sandy bottom in Core Sound, and Newport, North, New, and White Oak rivers from December to March. The fishery is managed intensively, with strong enforcement to prevent clam kicking outside the designated areas. Much of the designated mechanical clamming areas have SAV in close proximity to them, so vessels that fish illegally outside the open areas may severely impact SAV. Turbidity generated by clam kicking may also affect adjacent SAV beds. High salinity SAV species are more likely to be impacted by mechanical clamming practices due to the location of the fishery.

Bay scallop dredges, in contrast to oyster and crab dredges, cause less severe damage to SAV because they are smaller [not over 50 lb (22.68 kg)] and have no teeth. They are intended to glide along the substrate surface, taking bay scallops lying on the surface within SAV beds. Most damage observed by DMF staff has not been from the dredge, but from propeller scarring while pulling the dredge, particularly when the season opening coincides with low tide (T. Murphey, DMF, personal communication). Amendment I of the Bay Scallop FMP put in place an adaptive management strategy to determine harvest levels based on scallop abundance. All management triggers with the exception of the highest trigger allows hand harvest only. At the highest trigger, hand harvest is allowed at the beginning of the season (last Monday in January) with scallop dredging delayed until the first week in March. This strategy allows for removal of scallops in the shallow waters by hand harvesters first, followed by opening dredging later in the season to fish those scallops in waters too deep for hand harvest. In addition, opening day for harvest occurs on a high tide to allow dredgers to clean out any scallops in hand harvested areas. This management strategy minimizes damage to SAV from propeller scarring by dredging vessels (T. Murphey, DMF, personal communication).

Fishery restrictions already exist for most of the gears used in North Carolina that are potentially damaging to SAV. Additional law enforcement may be needed to enforce buffers around SAVs. In addition, the boundaries of areas where dredging or trawling is allowed should be evaluated and adjusted, if necessary, to adequately protect all SAV beds and provide a buffer of unvegetated area to reduce turbidity impacts. Because of the location and magnitude of fishing effort and SAV beds, it appears that trawling in Core and Bogue sounds has the greatest potential for significant fishing gear impacts on existing SAV beds.

10.3 WATER QUALITY

Adequate water quality is necessary to maintain the chemical properties of the water column that are needed by shrimp, as well as sustain SAV, shell bottom, and soft bottom habitats that support shrimp. Human activities that degrade water quality or alter water flow can negatively impact shrimp growth or survival. For example if salinity or dissolved oxygen (DO) concentrations are altered beyond the known preferences of shrimp, shrimp distribution or growth rates may be affected. Toxins can be assimilated into shrimp tissue and alter growth and reproduction. The common causes of water quality use support impairment in North Carolina's coastal river basins are excessive sediment loading and low DO (DWQ 2000). Hydrological modifications, low DO and toxin contamination are probably the greatest water quality concerns for penaeid shrimp.

10.3.1 Hydrological modifications

Hydrological modifications occur when streams and creeks are channelized (deepened and straightened), dredged, or ditched to improve drainage of adjacent lands or for navigation (North Carolina Sea Grant 1997), and often result in increased runoff. Runoff from agriculture, urban/suburban development, and transportation infrastructure carries sediment, nutrient, and toxic chemical pollutants (DWQ 2000). Sediment, the number one pollutant of waterways in the United States, clogs oyster gills and buries shells (Coen et al. 1999). Excess nutrients can fuel algal blooms and low DO events, and in turn, cause mortality of benthic organisms on deep, subtidal shell bottom (Lenihan and Peterson 1998). Heavy metals, petroleum products, pesticides, and other toxic chemicals in the runoff can kill sensitive oyster larvae (Wendt et al. 1990; Funderburk et al. 1991).

Channelized streams are often deeper, with more extreme flows, less woody debris and less variable depth than natural streams. These changes primarily affect smaller species and early life stages that use shallow stream margins, since these areas are reduced with channelization. Channelization potentially affects shrimp in several ways. By removing the meanders of the channel and increasing the slope of the shoreline, water velocities in the altered stream are higher and erosion of the shoreline and sediment loading increases. In many channelized streams, storm flows are confined primarily to the main channel rather than passing through wetlands and achieving some filtration of pollutants, deposition of sediment, and water storage. In addition, the natural woody vegetation along the sides of the stream is often removed in the process of channelization. Consequently, loading and movement of sediment and other nonpoint source pollutants are often greater in channelized sections than natural streams, which can have negative impacts on water quality and therefore fish habitat (EPA 2001). Nutrient concentrations, particularly for nitrogen and phosphorus, may increase with channelization. Elevated water velocities can also deter or prevent movement of adult and juvenile fish. In addition, spoil banks created by dredge disposal along the shoreline prevents shrimp from accessing adjacent wetlands.

Several studies have found that the size, number, and species diversity of fish in channelized streams are reduced and the fisheries associated with them are less productive than those associated with unchannelized reaches of streams (Tarplee et al. 1971; Hawkins 1980; Schoof 1980). Pate and Jones (1981) compared nursery areas that were altered and unaltered by channelization and found that brown shrimp, spot, croaker, southern flounder, and blue crab were more abundant in nursery habitats with no man-made drainage. They attributed this reduction in organisms to the unstable salinity conditions that occurred in areas adjacent to channelized systems following moderate to heavy rainfall (>1 inch/24 hr).

10.3.2 Low oxygen

Adequate supply of DO is critical to survival of benthic invertebrates and fish. Low-oxygen conditions (hypoxia) can occur naturally in a system from flushing of swamp waters, which characteristically have low DO, or from stratification of the water column due to wind, temperature, and salinity conditions. However, low-oxygen conditions can also be fueled by increased stormwater runoff carrying nutrients and oxygen-consuming wastes, which result in excessive oxygen demand in the water column or sediment. Algal blooms deplete the water column of DO as respiration from the dense concentrations of plants consumes oxygen at night (DWQ 2000). Dissolved oxygen can be further depleted as bacteria use oxygen to decompose the algae's organic material. Algal blooms may occur naturally in coastal waters or occur with greater frequency or intensity upon inputs of nutrients. Dissolved oxygen depletion in the water column occurs most often in summer. Warmer water holds less DO and increases microbial decomposition. In addition, warmer water, calm winds, and reduced freshwater inflow in the summer reduce mixing and aeration of water. The stratified bottom layer of water is prevented from receiving oxygenated surface waters and rapidly becomes depleted of oxygen. Shallow water estuaries with less frequent flushing often develop persistent stratification and bottom-water hypoxia that can last for weeks to months (Tenore 1972). Low oxygen events in coastal waters of the United States are becoming more frequent, larger in extent, and longer lasting due to increasing eutrophication (Cooper and Brush 1991; Breitberg 1992; Lenihan and Peterson 1998).

In freshwater streams, DWQ water quality (use support) data indicate low DO as a major cause of impairment in the Neuse River basin (132 mi), Chowan River basin (46 mi), Pasquotank River basin (40 mi), Roanoke River basin (24 mi), Tar-Pamlico River basin (13 mi), and White Oak

River basin (8 mi) (DWQ 2000). In estuarine waters, low DO was a major source of impairment in the Cape Fear (5,000 acres) and the Pasquotank river basins (1,125 acres). In the Neuse River, recent estimates suggest that up to 30-50% of the estuarine bottom during summer is unsuitable habitat due to hypoxia (Seldberg et al. 2001; Eby and Crowder 2002). Since shrimp live on the bottom in estuaries where hypoxia and anoxia (no oxygen) have been reported to occur, the species may be negatively affected by low oxygen events.

Brown shrimp and some other organisms are capable of detecting and avoiding waters with low oxygen concentrations (Wannamaker and Rice 2000). Where shrimp had access to water with 4 or 2 mg/l DO rather than 1 mg/l DO, shrimp strongly preferred and moved to the higher oxygenated waters. Migration of benthic organisms from hypoxic or anoxic waters can lead to high densities of organisms in oxygenated areas, increased competition, and increased predation by opportunistic predators (Eby and Crowder 2002; Seldberg et al. 2001). Although fish have the ability to migrate away from hypoxic areas and seek refuge in shallower oxygenated waters, wind-driven circulation can rapidly transport the hypoxic bottom-water into shallow waters, so that fish cannot escape (Paerl et al. 1998). Dissolved oxygen (DO) depletion has historically been the major factor driving fish kill activity in North Carolina. Low DO levels occur under a variety of conditions but are more common during the summer or following major storms and hurricanes. Consequently, in the wake of periods of hot weather and the arrival of Irene, low DO was the most frequently reported cause for fish kills during the 2011 season (DWQ 2011).

Although direct mortality does not appear to be a significant factor for shrimp, prolonged periods of hypoxia could stress and negatively impact penaeid shrimp and significantly alter the estuarine system. Studies on white shrimp found that growth rates of white shrimp were reduced in waters having less than 3.5 mg/l DO, feeding was affected in waters 2-3 mg/l DO, and oxygen uptake was reduced by 50-70% in 2 mg/l DO (Gray et al. 2002). When a benthic community is severely depleted by a low oxygen event, ecological successional patterns of the benthos are altered (Luettich et al. 1999). The various successional stages may affect or benefit different benthic feeders to differing extents. For example, early successional communities composed of very small, shallow-burrowing opportunists (capitellid worms) and meiofauna may favor small species, such as penaeid shrimp and larval and juvenile croaker and red drum, but not provide food for large adult fish species. Partially recovered benthic communities consisting of polychaetes and small juvenile clams could benefit demersal species like spot, croaker and blue crabs. A fully recovered community with deep burrowing polychaetes and large clams might benefit adult spot and hogchoker, but not shrimp (Luettich et al. 1999).

Hypoxia and anoxia can occur naturally, but can also be attributed, in part, to anthropogenic changes in the system, including excess nutrient and organic loading from waste discharges, nonpoint runoff, streambank erosion, and sedimentation (Schueler 1997). Oxygen depletion in the water column was positively correlated with accumulation of organic material in the sediments (Luettich et al. 1999). Several studies have indicated that the frequency, duration, and spatial extent of low oxygen events have increased over the years due to increasing eutrophication of coastal waters from human and animal waste discharges, greater fertilizer use, loss of wetlands, and increased atmospheric nitrogen deposition (Cooper and Brush 1991; Dyer and Orth 1994; Paerl et al. 1995; Buzzelli et al. 2002). More information is needed to fully understand consequences on the estuarine food web and to what extent anoxia affects the soft bottom community. Efforts are needed to reduce anthropogenic nutrient loading, particularly in systems that have a history of hypoxia and anoxia.

10.3.3 Toxins

While toxins can fluctuate between the sediment and water column, concentrations of toxic chemicals tend to accumulate in sediments at concentrations several orders of magnitude greater than in overlying waters (Kwon and Lee 2001). The bioavailability and transport of a chemical depends on the form of the chemical incorporated into the sediments, the feeding habits and condition of aquatic organisms, and the physical and chemical conditions of the environment. Toxic chemicals can become active in soft bottom sediment or overlying waters through several mechanisms, including resuspension from natural weather events or human activities, such as dredging and trawling.

Toxins in sediments or the water column can affect benthic invertebrates by inhibiting or altering reproduction or growth, or causing mortality in some situations (Weis and Weis 1989). Early life stages are most vulnerable to toxins (Funderburk et al. 1991). Because macroinvertebrate diversity significantly declines with increasing sediment contamination, food resources for benthic feeders, like shrimp, may be limited in highly contaminated areas (Weis et al. 1998; Brown et al. 2000; Dauer et al. 2000). While the survival of some aquatic organisms is affected by toxins, other organisms survive and bioaccumulate the chemicals to toxic levels, passing them along in the food chain. Multiple studies have shown clear connections between concentrations of toxins in sediments and those in benthic feeding fish and invertebrates (Kirby et al. 2001; Marburger et al. 2002). Heavy metal contamination of sediments has been documented to result in elevated trace metal concentrations in shrimp, striped mullet, oysters, and flounder (Kirby et al. 2001; Livingstone 2001).

There is some information available on the effect of certain toxic chemicals on different shrimp species. A study on the effect of copper, a common chemical associated with marinas, on a penaeid shrimp (*Metapenaeus dobsoni*) found that shrimp were tolerant to low concentrations of copper (0.05 mg Cu 1 super (-1)). However shrimp growth was significantly reduced when exposed to higher concentrations (0.15 mg Cu 1 super (-1)) (Manisseri and Menon 2001). Cellular damage to the hepatopancreas also occurred to shrimp exposed to 50-150 ppb Cu (Manisseri and Menon 1995). Another study examined mercury concentrations in both shrimp and blue crab, and found that blue crabs collected in the field with pink shrimp had higher mercury concentrations. The lower levels found in pink shrimp were attributed to shorter residence times in the contaminated area, differences in feeding habits, and the ability to excrete mercury somewhat faster (Evans et al. 2000).

Toxic chemicals come from localized point sources, as well as from diffuse nonpoint sources. Point sources include industrial and municipal waste discharges. Nonpoint sources of toxins include urban runoff containing household and yard chemicals, roadways, marinas and docks, boating activity, runoff from agriculture and forestry, industrial emissions, spills from industrial shipping, and dredge spoil disposal (Wilbur and Pentony 1999).

Because low concentrations of heavy metals in the water column can be easily incorporated into fine-grained sediment, chemicals can accumulate in the sediment to toxic levels and be resuspended into the water column (Riggs et al. 1991). Studies have shown that fine-grained sediments are the primary reservoir for heavy metals, particularly organic rich muds (ORM) (Riggs et al. 1991). Since ORM are the most extensive sediment type in North Carolina's estuaries, and since many primary nursery areas are composed of ORM, resuspension of contaminated ORM sediments in PNAs is of particular concern.

The extent of sediment contamination in North Carolina coastal waters is not well known. Sediment sampling is not conducted by the DWQ since there are no sediment standards in the state. Studies examining sediment contamination at sites in North Carolina soft bottom areas have found various levels of contamination. The EPA Environmental Monitoring and Assessment Program surveyed 165 sites within North Carolina's sounds and rivers during 1994-1997 to evaluate condition of bottom sediments (Hackney et al. 1998). Highest contamination levels occurred in low salinity areas with low flushing and high river discharge. Benthic populations were dominated by tolerant opportunistic species and benthic communities had low species richness. Laboratory bioassays showed that sediments from many sites were toxic to biological organisms. However, because of the low sample size, frequency of sampling, and the confounding effects of hypoxia in areas sampled, results from this study may not accurately assess the condition of North Carolina sediments (C. Currin, NOAA, personal communication).

Concentrations of heavy metals in the Neuse and Pamlico estuaries have been assessed (Riggs et al. 1989; Riggs et al. 1991). In the Neuse River, surface sediments contained elevated levels of several heavy metals, including zinc, copper, lead, and arsenic. Furthermore, 17 areas between New Bern and the mouth of the river were identified as "contaminated areas of concern". The contaminated sites were primarily attributed to permitted municipal and industrial treatment plant discharges. Marinas were also found to contribute substantial amounts of copper and variable amounts of zinc and lead. Nonpoint sources were more difficult to evaluate. In the Pamlico River, heavy metal contamination was less severe, although arsenic, cobalt, and titanium exceeded the levels found in the Neuse River. These studies suggest that sediment contamination in some estuarine areas especially those where both ORM and waste water discharges are present, may be significant and could affect fish populations and the base of their food chain. To better determine if contaminated sediment is a significant threat to shrimp habitat, the distribution and concentration of heavy metals and other toxic contaminants in freshwater and estuarine sediments need to be adequately assessed and areas of greatest concern need to be identified. Continued minimization of point and nonpoint sources of toxic contaminants is vital for protection of the entire ecosystem.

10.3.4 Tiger Shrimp

Tiger shrimp (*Penaeus monodon*), a non-native species of shrimp, have been observed in NC waters since 1988 when they were believed to have been released accidentally from an aquaculture facility in Bluffton, SC (Knott et al. 2012). Tiger shrimp have been observed from NC to TX. Although the impacts are not definitive at this time, tiger shrimp may pose a disease threat to native shrimp species. The NCDMF has been recording reported observations of tiger shrimp in NC waters since 2008. Whenever the public encounters what is believed to be a tiger shrimp it is reported to NCDMF and confirmed if possible. There have been a steady number of reports from the North Carolina ocean and estuarine waters throughout the coast since 2008 until 2011 when there was a large increase in the number of reports (Table 10.3). The reason for this increase is unclear, however the majority of those shrimp reported occurred in mostly southern shrimp catches after Hurricane Irene came through in August of 2011. This increase may be a result of local news stories after Hurricane Irene, or a potential spawning offshore community. The U.S. Geological Survey is investigating a potential community by collecting individuals and genetically testing them to determine the relationship (P. Fuller, USGS, personal communication 2012). The full impact of tiger shrimp in NC waters needs to be further investigated.

Table 10.3 Reported observations of tiger shrimp in NC since 2008.

Year	Yes ¹	No ²	Total Number of reported tiger shrimp
2008	12	4	16
2009	10	10	20
2010	1	4	5
2011	54	203	257

¹Confirmed by NCDMF and NC Coastal Federation staff.

²Reported tiger shrimp not confirmed may still be tiger shrimp.

10.4 HABITAT AND WATER QUALITY PROTECTION

10.4.1 MFC Authority

Presently, the MFC has authority to manage, restore, develop, cultivate, conserve, protect, and regulate marine and estuarine resources. Marine and estuarine resources are defined as “All fish [including marine mammals, shellfish, and crustaceans], except inland game fish, found in the Atlantic Ocean and in coastal fishing waters; all fisheries based upon such fish; all uncultivated or undomesticated plant and animal life, other than wildlife resources, inhabiting or dependent upon coastal fishing waters; and the entire ecology supporting such fish, fisheries, and plant and animal life” (G.S. 113-129).

Although MFC’s primary responsibilities are management of fisheries (seasons, size and bag limits, licensing, etc.), the MFC also has authority to comment on State permit applications that may have an effect on marine and estuarine resources or water quality, regulate placement of fishing gear, develop and improve mariculture, and regulate location and utilization of artificial reefs. MFC authority is found at G.S. 143B-289.51 and 289.52. The MFC and DMF should continue to comment on activities (state, federal, and local permits) that may impact estuarine water quality and work with permitting agencies to minimize impacts. Additionally, the MFC and DMF should solicit and support Fishery Resource Grant (FRG) and Coastal Recreational Fishing License Grant (CRFL) projects that may provide information necessary for protection, management, and restoration of water quality. Water quality standards should be based on the assimilative capacity of, and impacts to, the entire system. Several plans for water quality management have recommended strategies that need to be implemented to improve water quality. The MFC should continue to support management and research recommendations as outlined by the CHPP.

10.4.2 Authority of Other Agencies

Several divisions within the North Carolina Department of Environment and Natural Resources are responsible for providing technical and financial assistance, planning, permitting, certification, monitoring, and regulatory activities that have a direct or indirect impact on coastal water quality and habitat. The North Carolina Division of Coastal Management (DCM) is responsible for development permits along the estuarine shoreline in 20 coastal counties. Wetland development activity throughout North Carolina is permitted through the United States Army Corps of Engineers (COE) and the North Carolina Division of Water Quality (DWQ; 401-certification program). The DWQ permits and regulates discharges to surface waters, and

monitors water quality throughout the state. DWQ has established a water quality classification and standards program for “best usage” to promote protection of surface water supply watersheds, high quality waters, ecosystem functions, and the protection of unique and special pristine waters with outstanding resource values. Classifications, particularly for High Quality Waters (HQW), Outstanding Resource Waters (ORW), Nutrient Sensitive Waters (NSW) and Water Supply (WS) waters, outline protective management strategies aimed at controlling point and nonpoint source pollution. Various federal and state agencies, including DMF, evaluate projects proposed for permitting and provide comments and recommendations to the DCM, DWQ, and COE on potential habitat and water quality impacts. Various public agencies (state and federal) and private groups acquire and manage natural areas as parks, refuges, reserves, or protected lands, which helps to protect adjacent public trust estuarine water quality.

10.4.3 Nursery Area Protection

Existing rule definitions for fish habitat areas were revised by the Marine Fisheries Commission (MFC) in April 2009 in Marine Fisheries Commission Rule 15A NCAC 03I .0101 (4). The word “critical” was omitted since all fish habitats, under the ecosystem concept are critical to a properly functioning system as a whole. Regulatory protections exist for fish habitats areas such as primary nursery areas, secondary nursery areas, submerged aquatic vegetation, and shellfish producing areas.

Nursery areas are necessary for the early growth and development of virtually all of North Carolina’s important seafood species. Nursery areas need to be maintained, as much as possible, in their natural state, and the populations within them must be permitted to develop in a normal manner with as little interference from man as possible. In order to protect the integrity and resources of nursery areas, it is necessary to prohibit the use of bottom disturbing gears and severely restrict or prohibit excavation and/or filling activities.” The MFC and WRC first designated primary nursery areas in 1977 and 1990 respectively, based on field sampling. Approximately 162,000 acres of coastal fishing waters are currently designated by the MFC as Primary, Secondary and Special Secondary Nursery Areas.

There are specific protections for designated nursery areas included in the rules of three DENR commissions. For example, MFC Rule 15A NCAC 03N .0104 prohibits the use of trawls, dredges, long haul and swipe seines, and mechanical methods for oysters and clams in PNAs. Once an area has been designated as a PNA by the MFC, the area also comes under the protection of existing CRC rules (15A NCAC 07H .0208) and EMC rules [15A NCAC 02B .0301 (c)] that protect physical and water quality parameters of PNAs as a class. Various in-water work moratoria are also regularly required by state and federal agencies to protect sensitive habitat areas such as nursery and spawning areas from turbidity-related impacts. Due to the importance of that designation to DMF and the other agencies interested in protecting these nurseries, it is very important not to denigrate the integrity of the nursery area designation by naming areas that do not qualify under the rigorous biological sampling protocol and criteria established for nursery areas. .

The DMF conducted trawling and seine surveys in the early 1970s to inventory the state’s estuarine resources. The result of these surveys was the identification of estuarine areas that consistently supported populations of juvenile shrimp, crabs and finfish. Protection of these areas is imperative because they provide food, protection and proper environmental conditions (salinity and bottom type) for development and growth of young fish and crustaceans. Ninety percent of North Carolina’s commercially and recreationally important species are dependent on the estuary during some stage of their life cycle. The MFC adopted regulations in 1977 to

protect these estuarine areas known as nursery areas. Nursery areas are defined in Rule 15A NCAC 03I .0101(4)(f) as: “ Those areas in which for reasons such as food, cover, bottom type, salinity, temperature and other factors, young finfish and crustaceans spend the major portion of their initial growing season. Primary nursery areas are those areas in the estuarine system where initial post-larval development takes place. These are areas where populations are uniformly early juveniles. Secondary nursery areas are those areas in the estuarine system where later juvenile development takes place. Populations are composed of developing sub-adults of similar size which have migrated from an upstream primary nursery area to the secondary nursery area located in the middle portion of the estuarine system.”

The primary criteria for determining primary nursery areas are abundance of selected recreationally and commercially important species during recruitment periods, size composition, species diversity, bottom type and depth.

Abundance

Abundance of selected species is the primary criteria for selecting nursery areas. Selection of species for analysis is based on the area being considered, however, the species typically used in analysis are: brown shrimp, blue crab, spot, Atlantic croaker and southern flounder.

Size Composition

Another important characteristic of a nursery area is size of species in nursery area. Primary nursery areas are distinguished by the presence of finfish and crustacean populations that are uniformly very early juveniles. Size data are collected to determine the proportion of target species that are juveniles.

Species Diversity

The utilization of the site by various species is another indication of nursery area function. An index of species diversity summarizes community structure and takes into account, species richness as well as evenness of individuals among species.

Bottom Type

Bottom sediments in existing primary nursery areas are primarily coarse silt or clay with a high organic content. Most substrates are variously covered with detritus consisting of tree or shrub leaves, bark and branches, and dead marsh grasses. Numerous species of juvenile estuarine fishes in North Carolina are associated with this bottom type. Bottom type is categorized as mud, sand, or a combination of mud and sand.

Depth

Existing primary nursery areas generally have water depths of less than six feet. The most productive and valuable zone in many estuaries is the intertidal and shallow subtidal area.

The DMF's estuarine trawl sampling program (Program 120) provides data to identify nursery areas. It also provides a long-term database of annual juvenile recruitment of economically important species as provided by the core stations. This database has been used for designation of new nursery areas in the past and continues to be the main source of data and information used to designate future potential nursery areas.

Once a waterbody has been identified as a potential nursery area site, a sampling station for that area is established. Once that station has been established, it is sampled a minimum of three years. Comparison stations in approved PNAs located in close proximity to the proposed nursery area must also be established if they are not already a core station. Other PNAs

located in the same major waterbody are also included in the sampling. These PNA stations also must be sampled a minimum of three years, preferably on the same day and same tide.

Some areas that may be identified as a potential nursery area site will be unsuitable to trawling due to depth or underwater obstructions. Other gear types may need to be considered in order to properly compare those areas with nearby nursery areas. If this is the case, a gear is selected that can be used in both the potential nursery area and the comparison nursery areas. Once a sampling scheme has been determined, sampling continues for a minimum of three years.

Proposed PNA designations with supporting data are presented to the MFC. Comparisons of abundance and size are presented, along with environmental parameters including depth, sediment type, and salinity and compared to nearby designated PNAs. If the MFC approves the PNA to go to the advisory committees and the public for comment, it will then follow the normal rulemaking procedures to be designated.

Additional rules protecting PNAs were created under the authority of the Coastal Area Management Act (CAMA) of 1974. CAMA provided rules for coastal development, such as prohibiting dredging of channels, canals and boat basins in primary nursery areas. Construction of marinas that require dredging is also prohibited in PNAs. These restrictions are based on the quality of scientific analyses that goes into designation of nursery areas.

It is important to recognize the distinction between the generic term “nursery area” and the specific regulatory designations of “Nursery Area”. For example, Pamlico Sound maintains a diversity of habitat functions. Its abundance of young finfish as well as shrimp and crabs is well documented and therefore is often termed a nursery area. However, the regulatory designation of a nursery area is specific to the MFC’s intent of balancing competing public trust uses with the goal of habitat protection.

10.4.4 Coastal Habitat Protection Plan

The Fisheries Reform Act of 1997 (FRA 1997) mandated the Department of Environment and Natural Resources (DENR) to prepare a Coastal Habitat Protection Plan (CHPP -- G. S. 143B-279.8). The legislative goal for the CHPP is long-term enhancement of the coastal fisheries associated with coastal habitats and provides a framework for management actions to protect and restore habitats critical to North Carolina’s coastal fishery resources. The Coastal Resources Commission, Environmental Management Commission, and the Marine Fisheries Commission must each approve and implement the plan for it to be effective. These three Commissions have regulatory jurisdiction over the coastal resources, water, and marine fishery resources. The CHPP was initially approved in December 2004, updated in 2010 (Deaton et al. 2010), and implementation plans are developed every two years. Actions taken by all three commissions pertaining to the coastal area, including rule making, are to comply, “to the maximum extent practicable” with the plans. The CHPP will help to ensure consistent actions among these three commissions as well as their supporting Department of Environment and Natural Resources agencies and will be reviewed every five years.

The CHPP describes and documents the use of habitats by species supporting coastal fisheries, status of these habitats, and the impacts of human activities and natural events on those habitats. Habitats are categorized as wetlands, submerged aquatic vegetation (SAV), soft bottom, shell bottom, ocean hard bottom, and water column. The plan explains the environmental requirements, ecological value, status, and threats of the six fish habitats and

includes management recommendations to protect and enhance the entire coastal ecosystem. Much of the information provided in this section of the FMP came from the CHPP.

The CHPP recommends that some areas of fish habitat be designated as “Strategic Habitat Areas” (SHAs). Strategic Habitat Areas are defined as specific locations of individual fish habitat or systems of habitat that have been identified to provide critical habitat functions or that are particularly at risk due to imminent threats, vulnerability or rarity. While all fish habitats are necessary for sustaining viable fish populations, some areas may be especially important to fish viability and productivity. Protection of these areas would therefore be a high priority (Deaton et al. 2010). Habitat mapping and change over time is the foundation for SHA identification. DMF has an ongoing bottom mapping program, and other agencies and universities provide habitat distribution information as well. The process of identifying and designating SHAs was initiated in 2005. The coastal regions of NC have been divided into regions for the SHA process. SHAs for the Albemarle Sound and Northeast Coastal Ocean were completed in 2009 and the SHAs for the Pamlico Sound area and Central Coastal Ocean were completed in 2011. The remaining portions of NC should be completed within the next couple of years. SHAs should be identified and delineated that will enhance protection of penaeid shrimp. See Section 10 for recommended habitat and water quality actions.

Habitat protection, conservation, and restoration are essential to accomplish the goal and objectives of this plan. The FRA gives precedent to the CHPP and stipulates that habitat and water quality considerations in the fishery management plan be consistent with CHPP. Management actions recommended in this plan that are under MFC authority will be acted upon directly, while those management actions under other DENR authorities will be considered and acted upon through the CHPP implementation process and the appropriate agencies. Through that process, the MFC, Coastal Resources Commission (CRC), and Environmental Management Commission (EMC), and WRC should adopt rules to protect critical habitats for shrimp as outlined in the Coastal Habitat Protection Plan (CHPP). The DENR should develop a strategy to fully support the CHPPs process with additional staff and funding. The MFC and DMF should continue to comment on activities that may impact aquatic habitats and work with permitting agencies to minimize impacts and promote restoration and research. Research needs to be conducted to investigate the impacts of trawling on various habitats.

10.5 STATUS OF 2006 SHRIMP FMP ENVIRONMENTAL FACTORS RECOMMENDATION

Since the 2006 Shrimp FMP, habitat and water quality conditions appear to be the same or in some cases, somewhat better. The area of submerged aquatic vegetation coverage appears to be expanding in estuaries south of New River and in the lower salinity estuaries of the Neuse and Tar-Pamlico. The latter increase could be related to nutrient reduction efforts in those river basins, but may also be a result of several years of drought. Wetland acreage continues to decline from permitted losses and natural erosion associated with storms and rising sea level. Efforts have increased to restore more subtidal oyster beds in Pamlico Sound through DMF’s oyster sanctuary program and partnerships with non-profit organizations. Water quality degradation, in terms of aquatic life use support impairment, is greatest in freshwater streams in the Neuse and Cape Fear River basins and in estuarine creeks in the Neuse River basin. Fish kill events have declined in number but have been increasing in size.

In reviewing the 2006 Shrimp FMP habitat and water quality management recommendations, many have been implemented or are substantially underway. Many of these were also components of the CHPP implementation plan. They include:

Habitat

1. Coast-wide imagery of SAV was taken in 2007/2008 and has been mapped.
2. Identification and designation of Strategic Habitat Areas has been completed for the Albemarle and Pamlico Sounds along with their associated rivers.
3. Dredging of PNA, SAV, and shellfish habitat is avoided through DMF's permit review process.
4. CRC has revised dock rules to require review by resource agencies for General Permit dock applications located over SAV, shell bottom, or PNAs, and where water depth is less than 2 ft MLW to avoid boating related impacts.
5. Additional bottom disturbing gear restrictions have been implemented through the bay scallop and oyster fishery management plans to avoid damage to SAV and oysters.
6. Additional funding has supported expansion of oyster sanctuaries, development of a shell recycling program to supplement cultch planting, and acceleration of shell bottom mapping.
7. Ecosystem Enhancement Program is in the process of evaluating non-traditional but effective mitigation techniques for wetland, oyster, and SAV impacts, and improving the mitigation process.
8. Neuse and Tar-Pamlico NSW nutrient reduction measures have successfully reduced nutrient loading by more than their 30% reduction goals for point source dischargers and agriculture.
9. DWQ revised coastal stormwater rules that limit impervious surface and run-off in coastal areas.
10. Loss of additional riparian wetlands has been minimized through the permitting process, land acquisition, and land use planning.
11. Resource and regulatory agencies have been cooperating to promote alternative shoreline hardening measures. These measures include marsh sills.
12. The MFC has created an informational paper on the effects of contaminants. DMF has worked with partners to collect unused medicines as part of Operation Medicine Drop.

Water Quality

1. DMF staff continues to work with the permitting and commenting agencies to enhance protection of water quality. The MFC utilizes its permit commenting authority outlined in G.S. 143B-289.52 as needed.
2. DCM has created a clean marina program to promote environmentally friendly marinas.
3. Wetland buffers along coastal streams and rivers have been used to enhance wetlands and improve water quality.

10.5.1 Coastal Habitat Protection Plan Actions

There are many actions that natural resource managers can take to sustain and enhance habitat and water quality conditions for shrimp. High priority needs include:

- Preserving existing coastal wetlands and restoring wetlands
- Protecting PNAs from dredging and water quality degradation
- Protecting and enhancing SAV habitat
- Assessing sediment contamination in NC estuaries and effects on shrimp
- Reducing pollutant loading from point and non-point sources

These management needs are currently being addressed through several existing CHPP recommendations (Deaton et al. 2010) and implementation actions (DENR 2011) that were approved by the CHPP Steering Committee. Listed below are those CHPP recommendations

and implementation actions that could be beneficial for protecting and improving habitat and water quality issues affecting shrimp. Numbering refers to the CHPP recommendations. Implementation actions are denoted by (I) following the recommendation number.

- 2.1 Support Strategic Habitat Area assessments by:
 - a) Coordinating, completing, and maintaining baseline habitat mapping (including seagrass, shell bottom, shoreline, and other bottom types) using the most appropriate technology
 - b) Selective monitoring of the status of those habitats

Of specific importance for shrimp is:

- remapping and monitoring SAV in North Carolina to assess change in distribution
- assessing the distribution, concentration, and threat of heavy metals and other toxic contaminants in freshwater and estuarine sediments and identify the areas of greatest concern to focus water quality improvement efforts
- monitoring to determine if additional areas should be designated as Primary Nursery Areas due to their nursery importance to shrimp

2.2 Identify, designate, and protect Strategic Habitat Areas.

- 3.1 Expand habitat restoration in accordance with restoration plan goals, including:
 - a) Creation of subtidal oyster reef no-take sanctuaries
 - b) Re-establishment of riparian wetlands and stream hydrology
 - c) Restoration of SAV habitat and shallow soft bottom nurseries

Of specific importance for shrimp is protection and restoration of coastal wetlands and SAV.

- 3.3 Protect habitat from fishing gear effects through improved enforcement, establishment of protective buffers around habitats, modified rules, and further restriction of fishing gear where necessary.

Of specific importance for shrimp is periodic re-examination of areas where trawling, oyster dredging or mechanical harvest is currently allowed to determine if conflicts with habitat protection exist.

- 3.4 Protect estuarine and public trust shorelines and shallow water habitats by revising shoreline stabilization rules to include consideration of erosion rates and prefer alternatives to vertical shoreline stabilization measures that maintain shallow nursery habitat.

- 3.7 (I) Develop an interagency policy for marina siting to minimize impacts to ecologically important shallow habitats such as Primary Nursery Areas (PNA), Anadromous Fish Spawning Areas (AFSA), and SAV.

- 4.1 Reduce point source pollution discharges by:
 - a) Increasing inspections of wastewater treatment facilities, collection infrastructure, and disposal sites
 - b) Providing incentives for upgrading all types of discharge treatment systems

- c) Developing standards and treatment methods that minimize the threat of endocrine disrupting chemicals on aquatic life.
- 4.5 Improve strategies throughout the river basins to reduce non-point pollution and minimize cumulative losses of fish habitat through voluntary actions, assistance, and incentives, including:
- a) Improved methods to reduce pollution from construction sites, agriculture, and forestry
 - b) Increased on-site infiltration of stormwater
 - c) Encouraging and providing incentives for low-impact development
- 4.6 Improve strategies throughout the river basins to reduce non-point pollution and minimize cumulative losses of fish habitat through rule making, including:
- a) Increased use of effective vegetated buffers
 - b) Implementing and assessing coastal stormwater rules and modify if justified
 - c) Modified water quality standards that are adequate to support SAV habitat
- 4.8 Reduce non-point source pollution from large-scale animal operations

11.0 MANAGEMENT STRATEGIES

11.1 HISTORY OF SHRIMP MANAGEMENT

Shrimp management in North Carolina has evolved from early biological work done in the mid 1960s. At that time, studies were conducted on the behavior of our three species of shrimp (pink, brown and white), their growth rates, mortality and migration, habitat preferences, and salinity and temperature tolerances.

A major step in the evolution of management came in 1978 with the designation of PNA and SNAs. These are the shallow bays and tributaries with low salinities, muddy bottoms and detritus where the shrimp spend their post-larval and juvenile development. In these shallow waters, food is abundant, salinities and temperatures are optimal, and there are few predators. No trawling is allowed in PNAs and SNAs to allow the shrimp to grow to harvestable size with as little man-made disturbance as possible. A SSNA designation originated in the 1980s to protect the shrimp during the majority of the season and allow harvest toward the end of the season, when shrimp are of harvestable size and juvenile fish have migrated out of the bays. The Fisheries Director may, by proclamation, open any portion of SSNAs to shrimp or crab trawling from August 16 through May 14. Management rationale included minimizing waste by delaying the trawl opening date to reduce the finfish bycatch and to ensure catches of larger shrimp. North Carolina was the first state to require the use of BRDs in shrimp trawls and requires them to be installed in specific tailbag locations in order to reduce the incidental catch of juvenile finfish. The strategy of DMF has been to protect the sensitive nursery areas and critical habitats while working to reduce bycatch as much as possible (see section 6.3 Bycatch).

Other management measures that have been implemented include the 1 ½ inch minimum mesh size in shrimp trawls, no trawling areas in the Outer Banks sea grass beds, military restricted areas, and weekend closures in internal waters from 9:00 p.m. on Friday nights to 5:00 p.m. on Sunday nights, among others. The Director has proclamation authority to open and close waters within the estuaries and the Atlantic Ocean based on shrimp size and environmental conditions. This flexibility in opening and closing shrimping areas is a valuable management tool, but it makes managers subject to the lobbying efforts of the various user groups.

Shrimp management varies from the southern portion of the state to the northern part because of species behavior and differences among geographic areas as well as preferences of the user groups. In the Roanoke Island area, which is the northernmost range for NC shrimp, the management of SSNAs is based more on the protection of juvenile finfish than on the harvest of shrimp. Sampling is conducted to insure that the small fish have left the bays and, if shrimp are present, the area is opened. Abundant shrimp in the northern part of the state is such a rare occurrence that nearly any size is considered harvestable, and by August 16, they are usually of sufficient size.

Before the implementation of the 2006 Shrimp FMP, attempts were made to limit the frequent movement of shrimp lines by meeting with the fishermen, discussing the problems, and seeking answers acceptable to the majority, while offering reasonable protection for the small shrimp. For example, a meeting was held at Harkers Island in 1997 about a possible solution to North River shrimp line and by unanimous choice, a permanent line was agreed to and implemented. The line works well, unless there are tremendous numbers of shrimp or significant rainfall, which cause smaller ones to spill over into the open area. Still the shrimp are marketable and provide income to the fishermen, particularly the early summer pink shrimp.

Closing an area in mid-season may result in a “grand opening” later. Areas like Adams Creek, Newport or North River may have up to 200 boats, regardless of the abundance of shrimp. This large number of vessels operating in confined waterbodies results in dangerous navigational situations. Fish kills following shrimp openings in New River and Bay River in the past have brought attention to trawling impacts. The detrimental effects of these openings to the bottom and juvenile fish in the area make it very desirable to avoid them whenever possible. The implementation of the 2006 Shrimp FMP led to the development of other management strategies to protect habitat, reduce bycatch, minimize user conflict and bring consistency to the management of the shrimp fishery. The following sections outline these management strategies as developed in the 2006 plan. Details of the development of these strategies may be found in the appendices of the 2006 Shrimp FMP (NCDMF 2006).

11.2 SHRIMP MANAGEMENT BY SIZE

Shrimp grow at different rates depending on water temperature and salinity. As growth increases, shrimp migrate to deeper, saltier waters of the sound and eventually to the ocean. As shrimp migrate to the ocean, they enter areas that are open or may be opened by the DMF to the harvest of shrimp. Sampling is conducted by the DMF staff to determine if an area should be opened or closed, based primarily on size and count. Over time, target sizes for opening different waterbodies have evolved and allow for better flexibility of management for both recreational and commercial shrimping.

Although highly variable, the density of shrimp in the nursery areas during the spring as well as weather conditions in the critical spring nursery months determine the number and size of shrimp in the different waterbodies. Overcrowding and its associated competition for food and space cause the shrimp to migrate downstream earlier than normal with wind and rainfall compounding the problem. At times when this occurs, the event is over before a closure can take effect or the shrimp have crossed the line established by consensus, which the Division will honor.

Shrimp in the Southern District, with no extensive bays and sounds to grow and develop begin to migrate at a smaller size. The waters of Onslow, Pender, New Hanover, and Brunswick counties that are available for opening to trawling are typically located either in or landward of the Intracoastal Waterway (IWW), which runs the entire length of all four counties' coastlines. Portions of these narrow waters may remain closed during part of the shrimping season or not open at all, depending on the size of the shrimp observed in the DMFs samples. Target opening size in Brunswick and portions of New Hanover counties is 40-50 count (heads on). In Onslow and parts of Pender counties, sampling has shown that a 20-30 count target size can be achieved before migration occurs. Channels that connect the IWW with the Atlantic Ocean have been left open to allow some harvest of shrimp as they migrate from closed areas to the ocean. Trawling in these migration routes has become the subject of discussion among shrimpers as well as the public because of concerns about bycatch of other species as well as interference with navigation. One migration route that has been the subject of recent controversy is the channel leading to Blue Water Point Marina in Brunswick County.

The target size of shrimp in the majority of the Central District and Pamlico District is 26-30 or 27-35 count (per pound heads-on) although White Oak River shrimp tend to be smaller with a 45-50 count (heads-on) targeted size.

Consideration must be given to the entire range of users, from the 15' outboard in the shallow water sounds and river tributaries to the 85' ocean trawler. In most cases, 100 pounds of 45 count (heads-on) shrimp would be much more valuable if permitted to grow to 16-20s, even factoring in the mortality suffered in the meantime. Even this statement has its exception in the spring pink shrimp fishery in the North River area of Carteret County when 45 count shrimp bring up to \$2.50 per pound. Managing for 16-20 count shrimp would eliminate the majority of the shrimp fleet and leave the catch to larger trawlers in Pamlico Sound and the ocean and to some channel netters. The current management strategy is to allocate some of the public resource to all groups.

Unusual weather events or the occurrence of unusually high numbers of small shrimp may occasionally force closures of normally opened areas like a portion of Neuse River or in the ocean south of Cape Fear.

Target sizes for opening have evolved: 26-30 count from Pamlico Sound to White Oak River; 45-50 count in the White Oak River; 20-30 count in New River and parts of Pender County; and 40-50 count in Brunswick and parts of New Hanover counties. At the present time modal groups are used and some shrimp are larger and some smaller than our target. Openings based on these target sizes have addressed the variability within the state of boat sizes and size preferences of the user groups, geographical differences in the shrimp size at migration, weather events, and socio-economic conditions.

11.3 SHRIMP MANAGEMENT BY AREA

Historically, the DMF has used a number of criteria to determine if trawling should be allowed in estuarine waters. These criteria include habitat issues such as aquatic vegetation, water depth and bottom types; shrimp size and abundance; economic and social factors; user conflicts; and bycatch issues.

DMF has utilized rules and proclamations to manage trawling in ocean and internal coastal waters. The intention of these rules and proclamations has been to allow the harvest of shrimp and crabs in estuarine waters but prohibit directed finfish trawling. Openings and closings of specific areas are based primarily on the size of the shrimp.

The closure of nursery areas and the protection of sea grass beds through rules, and proclamations are designed to minimize the bottom-disturbing effects of trawling (see Section 10. Environmental Factors). Trawling is limited primarily to the large bodies of water, such as the rivers, sounds and ocean. Shoals, wrecks, obstructions, oyster rocks, and algal and bryozoan growth make some of this open water area inaccessible to trawls. There are also areas opened to shrimping that receive very little effort because shrimp abundance is low.

Shellfish management areas (SMAs) are another critical habitat where trawling is prohibited (15A NCAC 03N.0104 and 0105.03J.0103). While these regulations protect the substrate from physical damage by trawls, bottom-disturbing gear used adjacent to the SMAs impacts oyster reefs indirectly by re-suspending sediment. As sediment disperses away from the disturbance and settles to the bottom, it can bury oyster larvae, adults, or shell, deterring successful recruitment of larvae due to lack of an exposed hard substrate (Coen et al. 1999). Additionally, excessive sedimentation can also harm shellfish by clogging gills, increasing survival time of pathogenic bacteria, or increasing ingestion of non-food particles (SAMFC 1998).

DMF conducts regular sampling to monitor shrimp size and abundance and takes appropriate action based on the samples. Waters eligible to be opened to trawling may also be closed if the size of the shrimp is too small. Closures of this nature are primarily influenced by economics since small shrimp have little value and if there is no market, the resource is wasted. Affected areas include those where shrimp size changes predictably based on annual cycles and environmental conditions as well as those areas where the habitat has changed in response to physical changes such as inlet closures and shoaling. Waters have also been closed in order to reduce or eliminate conflicts with other users and traditional uses such as navigation. These would include closure of crab pot areas and navigation channels where shrimping activity has been problematic.

11.3.1 Shrimp Management in the Southern District

DMF has been managing the shrimp harvest since the early 1970s. In 1977, based on sampling conducted over a number of years the DMF designated nursery areas (both primary and secondary) throughout the State that were closed to all bottom disturbing gear, including shrimp trawls. Many of these nursery areas are in the southern part of the State and include those areas that are most biologically sensitive to trawling. Additional areas were closed in the 1980s in reaction to an increase in fishing effort. Time and area closures were the only tools available to deal with the increase in effort. The net result of all these closures is that approximately one-third of the waters in Onslow, Pender, New Hanover and Brunswick counties can be opened to shrimp trawling. However, portions of these waters may remain closed or not be open at all depending on the size of shrimp observed in DMF's samples. While this strategy helps protect these areas, it forces the fishery to operate in a smaller area thus increasing user conflicts.

The areas that can be opened to shrimping are typically located either in or landward of the IWW which runs the entire length of the Onslow, Pender, New Hanover and Brunswick counties coastline. The heads-on counts used to determine whether to open an area vary by area based on historical sampling which indicates at what size shrimp tend to migrate from different water bodies. In Brunswick and portions of New Hanover counties, where shrimp migrate at smaller sizes, DMF attempts to open on a 40-50 count shrimp. In Onslow and parts of Pender counties, sampling has shown that a 20-30 count can be attained before migration occurs. Channels that connect the IWW with the Atlantic Ocean are normally left open at all times to allow some harvest of shrimp as they migrate from closed areas to the ocean. Trawling in these migration routes has become the subject of discussion amongst shrimpers as well as the public because of concerns about bycatch as well as interference with navigation.

11.3.1.1 Brunswick County

The Brunswick County coastline stretches for approximately 33 miles and is bound by the Cape Fear River Inlet on the east end and by the Little River Inlet on the west end. Four barrier islands, all of which are densely developed, are separated by five inlets along the coastline.

The IWW in Brunswick County is managed based on the size and abundance of the shrimp taken in the DMF's samples. The area is usually open until the beginning of June when it is closed because of small brown shrimp. In most years, portions may be opened in late June or early July to allow harvest of brown shrimp and then closed in late July or early August when small white shrimp recruit to the area. Occasionally, small white shrimp may appear before the brown shrimp reach a harvestable size, thus delaying an opening until the whites are harvestable, usually in September but sometimes as late as November. Principle harvest areas




Figure 11.1 Map of shrimp management areas in Brunswick County.

are behind Oak Island, from the Holden Beach Bridge to Shallotte River and from the Ocean Isle beach bridge to the Sunset Beach Bridge (Figure 11.1).

The IWW channel from the Sunset Beach Bridge to the South Carolina State Line and the Calabash River are rarely opened to trawling because of the abundance of small shrimp. The area from Sunset Beach Bridge to Calabash River is usually opened toward the end of the season so that the shrimp won't be "lost" to South Carolina.

The channels that connect the IWW with the Atlantic Ocean usually remain open during the entire year to allow harvest of shrimp that are migrating to the ocean. In rare instances of very heavy rainfall, these channels may be closed. The areas include Elizabeth River, Dutchman Creek, Montgomery Slough, Jink's Creek and Bonaparte Creek. Trawling in Montgomery Slough and the Elizabeth River has become the subject of discussion amongst shrimpers as well as the public because of concerns about bycatch as well as interference with navigation. Eastern Channel, located behind Ocean Isle Beach, is a shallow channel (less than one meter at mean low tide) that connects the IWW at Marker 93 to Jink's Creek. These waters have not been opened to harvest in over 20 years.

The Shallotte River was opened and closed to shrimp trawling based on size and abundance until 1998. However, DMF sampling has shown that these shrimp rarely reach large sizes with the heads-on count remaining greater than 60 during most of the season. Consequently, the last time DMF opened Shallotte River was a span of time in 1998 between July 8 and September 9.

11.3.1.2 Cape Fear River Complex

The waters of the Cape Fear River, the Basin, Second Bay, Buzzard's Bay (the Bays) and Bald Head, Cape and Bay Creeks (the Creeks) are part of the Cape Fear estuarine system (Figure 11.2). Bottom types range from sand near the inlet and creek mouths to mud in some of the bays and channels near Snow's Cut. There are active clam and oyster fisheries in the bays, creeks and the river upstream to the Fort Fisher Ferry Terminal. These fisheries occur primarily by hand and in shallow water though there are tong and bull rake clam fisheries in the deeper areas. In some of the deeper areas of the Cape Fear, clam and shrimp fisheries co-exist. There are active crab pot and gillnet fisheries throughout the entire estuary. There are several Primary, Secondary and Special Secondary Nursery Areas located in the Cape Fear River.

The area in the Cape Fear that is open to shrimping is dredged on a regular basis for navigation purposes. The river is managed on the size of shrimp and various parts of the river are opened and closed based on the DMF's samples. The upstream line was placed at Snow's Cut for many years because of the abundance of small shrimp above this line. The line was moved upstream in 2003 based on larger shrimp being present at that location. The river has not been closed in recent years because when small shrimp were in the open areas the participants have chosen not to harvest them.

The bays south of Fort Fisher known as the Basin or First Bay, Second Bay and Buzzard's Bay have been managed in the past as a unit with openings and closings based on the DMF's samples. New Inlet drained these areas but closed after a series of hurricanes in the late 1990s and circulation is now through the Cape Fear. Since the inlet closed, DMF has observed a shift in the biological characteristics of these waters towards more of a nursery area. Consequently, the size of the shrimp tends to remain small in this area and have remained closed since 2002.




Figure 11.2 Map of shrimp management areas in the Cape Fear River Complex.

Historically, the Bald Head Creeks were usually opened in late June or early July based on the size of shrimp. Areas opened included the lower portions of the Creeks. However, following the implementation of the 2006 Shrimp FMP, no trawling areas were established in the bays south of Fort Fisher and Bald Head Creeks. The main river has remained open to shrimping with potential opening dates set by proclamation and determined by DMF sampling.

Trawling trips in the main part of the Cape Fear are usually day trips and fishery operations are performed primarily from small boats using otter trawls, although vessels up to 50 feet may work in the channels of the Cape Fear. The 2006 Shrimp FMP restricted the total headrope length for otter trawls to 90 feet in the Cape Fear River. There are no other mobile gears used but there has been some use of channel nets in the past.

11.3.1.3 Intracoastal Waterway and Sounds from Carolina Beach to Rich's Inlet

The estuarine waters of the IWW channel and adjacent sounds between Carolina Beach and Rich's Inlet stretch over 21 miles and include four inlets separating four barrier islands, three of which (Figure Eight, Wrightsville, Carolina Beach) are heavily developed (Figure 11.3). These waters are bordered on the south by the Carolina Beach Yacht Basin (CBYB) and to the north by Rich's Inlet. The largest inlet is Masonboro Inlet and it is located approximately in the center of these estuaries where it separates Wrightsville Beach from Masonboro Island.

Bottom types are primarily sand throughout the area with the exception of more soft muddy substrates in the sounds and portions of the IWW. Submerged aquatic vegetation (SAV) is limited to a few patches in the shallow sound areas. There are active oyster, clam, and crab fisheries throughout the area. These fisheries are prosecuted in the sounds and along the edges of the IWW. The waters contain a few shellfish leases and DMF maintains six SMAs from Hewlett's Creek north to Rich's Inlet. In addition, DMF and the Coastal Federation a non-governmental organization located in Newport, NC, have collaborated on construction of an oyster sanctuary in the mouth of Hewlett's Creek. Areas closed to the harvest of shellfish due to pollution are abundant and include all or portions of creeks on the mainland side of the IWW as well as most of the Wrightsville Beach area and buffers around numerous marinas.

Most all of these areas receive very minimum shrimping effort with little or no impact on shellfish resources. Exceptions are a section of the IWW in Myrtle Grove Sound (William's landing) and the CBYB. Additionally, some of the channels around Wrightsville Beach also receive shrimping effort at various times during a typical year. Both commercial and recreational shrimpers utilize these waters.

The William's Landing area has been difficult to manage because the shrimp often migrate before reaching larger sizes (30-40 count, heads-on) except in the fall. In some years, large concentrations of algae (*Grassilaria* and *Ulva spp.*) prevent the use of trawls until the shrimp grow to an acceptable count while in other years there has been harvest of small shrimp. The CBYB is opened and closed based on the size of shrimp present. Channels around Wrightsville Beach remain open to allow harvest of shrimp migrating to the ocean. Historically, the area of the IWW from the Wrightsville Beach drawbridge to Marker #105 at Green's Channel has always remained open to shrimping but received little effort from commercial or recreational fisherman. However, the IWW was closed to trawling from Marker #105 to the Wrightsville Beach drawbridge following the adoption of the 2006 Shrimp FMP. Actions were also taken to manage the IWW from Marker #139 to Marker #146 as a SSNA, opening by proclamation from August 16 through May 14. However due to the abundance of small shrimp and large concentrations of algae, this area is rarely opened during this time period.




Figure 11.3 Map of shrimp management areas in the Intracoastal Waterway and sounds from Carolina Beach to Rich's Inlet.

11.3.1.4 Intracoastal Waterway and sounds from Rich's Inlet to New River

The estuarine waters of the IWW channel and the adjacent sounds and bays between Rich's Inlet and New River Inlet are managed as a single waterbody by the DMF (Figure 11.4). A section of this waterbody bounded by Marker #17 to the north and the Surf City swing bridge to the south is designated as SSNA. Historical data (since 1972) collected by DMF indicates these waters support large aggregations of commercially important finfish as well as shellfish and crustaceans.

Bottom types range from mud and muddy/sand in the IWW to mostly sand near the inlets. The shallow waters of Topsail Sound and some of the estuarine areas around New River Inlet contain patches of SAV.

There are active clam and oyster fisheries in the entire area. Hand harvest for oysters and clams take place in the shallow areas throughout these waters on both public bottom and leased areas, while mechanical harvest of clams is allowed in the IWW from New River to south of the Surf City bridge ("BC" Marker). DMF maintains Shellfish Management Areas throughout the area, all of which are located in waters closed to shrimping with mobile gears. DMF and the Coastal Federation have collaborated to begin construction of oyster sanctuaries in Stump Sound.

The typical management cycle for these waters is; the IWW north of Marker #17, the IWW south of the Surf City swing bridge and Banks Channel in Topsail Sound remain open during the entire year unless unusually high rainfall amounts or overcrowded nursery areas force large numbers of small shrimp into them prematurely. Waters in the SSNA, with the exception of the middle portion of the SSNA, are typically opened sometime after August 15. The middle portion of the SSNA from Marker #45 to the Highway 210-50 high-rise bridge usually remains closed until late in the season because of the abundance of small white shrimp.

The fishing is dominated by small boats that trawl, float net and skim in the main channel of the IWW and in a 100-foot strip on the side of the IWW that is open from Marker #49 to Marker #105. Channel nets are set outside of the marked channel from Marker #15 at New River to just south of the Surf City Bridge and in Topsail Sound. Banks Channel serves as a migration route for emigrating shrimp; gears used include trawls, skimmers and most recently shrimp traps and shrimp pounds.

11.3.1.5 New River

The DMF manages the New River based on nursery area classification (Figure 11.4). The waters upstream of the Highway 172 Bridge were designated by rule as a SSNA in 1996. The areas of the SSNA that are impacted by the opening include the river above the bridge up to the marked closure line running from Grey's Point to the opposite side of the river. Trawling in any of the tributary creeks is prohibited. The river consists mostly of shallow bays with the exception of the marked navigation channel. Bottom types range from sand and sand/mud to live shell bottom. The DMF actively manages seven SMAs in this portion of New River.

The use of otter trawls upstream of the Highway 172 Bridge was phased out in 2010 following the adoption of the 2006 Shrimp FMP. Those who wished to continue to harvest shrimp in the waters above the 172 Bridge were allowed a four year grace period to convert to skimmers.




Figure 11.4 Map of shrimp management areas in the Intracoastal Waterway and sounds from Rich's Inlet to New River.

Subsequently, crab trawls were also phased out of this area as part of the 2006 Shrimp FMP. Prior to the 2006 Shrimp FMP, crab trawlers would often fish above the Highway 172 Bridge to target flounder more so than crab; however, stricter minimum size limits for flounder made it economically unfeasible for crab trawlers to harvest only crabs in this area. Currently, the waters upstream of the Highway 172 Bridge are only open to boats equipped with skimmer rigs. During the revision of this plan, it was requested by several crab trawlers to reconsider allowing trawlers to fish above the Highway 172 Bridge; however this management strategy will remain in place (see appendix 1).

The DMF typically issues a proclamation opening the waters above the Highway 172 Bridge around the middle portion of August. Once a proclamation has been issued, these waters remain open until May 14th. Initial sampling of core and optional stations in the recruitment or nursery areas starts in August and is completed prior to August 16th. The waters below the 172 Bridge are open to trawling year round; however, over the past few years there has been very little effort in this part of the river.

11.3.1.6 Chadwick Bay

Chadwick Bay is a small high salinity waterbody encompassing 841 acres located just south of the mouth of New River and adjacent to the IWW and the New River Inlet (Figure 11.4). The southern portion of the bay is classified as a Primary Nursery Area (PNA) characterized by shallow water depth (< 5 feet) and a sandy mud substrate with patches of SAV. Fullard Creek is the major tributary of Chadwick Bay and minor tributaries include Charles Creek and Bumps Creek. The upper portion of Fullard and all of Charles Creek and Bumps Creek are designated by DMF as PNAs. Although the lower portion of Fullard Creek is not currently classified as a nursery area, it is not opened to shrimping because of the abundance of juvenile finfish. Prior to April 1, 2011 the remainder of Chadwick Bay was opened by proclamation to shrimping when the shrimp reached a harvestable size (30-40 heads-on count).

In the past the DMF utilized two different strategies in managing Chadwick Bay. In years when brown shrimp were abundant and large, the bay was opened in July along with the White Oak River, Queen's Creek and Bear Creek. In other years when brown shrimp were less abundant, a Chadwick Bay shrimp opening on white shrimp may have occurred in August or September in conjunction with the openings in New River and/or Stump Sound. However, the 2006 Shrimp FMP requested that a trawl survey be initiated to determine if Chadwick Bay functioned as a SSNA. Based on the species diversity, habitat, and size of shrimps, crabs, and fishes caught in the bay during the survey, Chadwick Bay was found to function as a SSNA. Under its new classification, trawling is permitted by proclamation from August 16th to May 14th. By managing Chadwick Bay as a SSNA, the harvest of juvenile shrimp, finfish, and crustaceans in areas where they spend a major portion of their initial growing season will be reduced by eliminating any openings before August 16th. This management strategy allows for larger, more valuable shrimp to be harvested before they move out into open waters. Additionally, the potential negative effects of trawling on the shallow soft bottom habitat and SAVs of the bay is reduced by decreasing the time when trawling is permitted.

The Chadwick Bay shrimp fishery is primarily conducted with trawls, although, in recent years, the use of skimmers has increased in the commercial portion of the fishery. RCGL holders frequently shrimp in the bay, especially on opening days.

11.3.2 Shrimp Management in the Central District

Management of shrimping in the Central District takes place from the White Oak River on the Onslow/Carteret County line to Core Sound in Carteret County. The Central District also manages the south side of the Neuse River in Craven County. Areas that are open and closed to shrimping through proclamation include: the, West Bay/Long Bay, Thorofare Bay, several tributaries in Core Sound and Adams Creek, located on the south side of the Neuse River.

The DMF issues a proclamation during the first week of June showing shrimp lines for the beginning of the season. This proclamation establishes closures in, Jarrett Bay and the West Bay-Long Bay and Thorofare Bay areas. This proclamation also designates closures of the SSNAs located in Core Sound. The DMF conducts nighttime sampling in both the closed portion and the open portion of a waterbody with a small 20-foot otter trawl with ½ inch bar mesh in the body and ¼ inch bar mesh in the tailbag. Tow times are between 5 minutes and 20 minutes. Shrimp are counted and a subset of the sample is measured to determine sizes or counts. Salinities and water temperatures are also recorded. Target counts vary dependent on the waterbody and range from 26-30 count to 31-35 count (heads-on). In an area like the White Oak River, where shrimp do not grow very large, the count is around 45-55. Based on this sampling, lines may be moved by proclamation to protect small shrimp until they are large enough to harvest.

11.3.2.1 White Oak River

White Oak River is located on the Onslow/Carteret County line and has the town of Swansboro at its mouth (Figure 11.5). Due to the presence of oyster rocks and shoals, there are only a few places that are able to be trawled in the river. They are Hills Bay below Jones Island, the mouth of Pettiford Creek, the Turnstake, and Cahoon's Slough above Jones Island. Recreational shrimpers as well as a few commercial shrimpers use the White Oak River.

Before the 2006 Shrimp FMP, the river was closed at the Highway 24 Bridge with the issuance of the first shrimp proclamation in early June. Sampling for opening White Oak River generally began around the end of June because of the tendency for shrimp to migrate early. Historically, the DMF opened White Oak between July 10 and July 20 to the Gator Gap where the river widens near Bluff Point. Small shrimp were often forced across that line and the DMF tried alternative line locations with varying success that allow for shrimping in the lower portion of the river while protecting small brown and white shrimp upstream. Adjusting the line was difficult due to the amount of oyster rocks in the river. Shrimpers like to tow on the line, therefore placement of the line over oyster rock lead to habitat destruction of those rocks.

Issues that had to be considered in the previous management of this river besides shrimp size were weather conditions and lunar stage. Early northerly winds with a lot of rain or a hurricane can force the small shrimp to migrate before the normal opening dates. A full or new moon on top of that may also cause the DMF to open on a smaller count so they can be caught.

When the bridge was the closure line, there was no shrimp trawling allowed in White Oak River. If the shrimp move out before the river was opened, then the only fishermen who benefited were a few channel net fishermen and maybe ocean trawlers. Over the years options were considered to leave the river closed at all times to protect the oyster rocks, but that was inconsistent with permitting mechanical clam harvest up to the Turnstake and did not allow trawlers to catch the shrimp at all. Therefore a permanent line was established after the




Figure 11.5 Shrimp management areas in the White Oak River and Bogue Sound.

adoption of the 2006 Shrimp FMP in the area of Cahoon Slough to Hancock Point..

11.3.2.2 Bogue Sound

Bogue Sound for the most part has permanent closure lines (Figure 11.5). The sound is closed to trawling on the north or mainland side of the IWW and in a portion of the western part of the sound. These areas remain closed because of the nature of the bottom. The area along the northern side of the IWW acts as a nursery area and also borders several primary nursery areas. SAVs with bay scallops are located in the closed portion of the western part of Bogue Sound. This SAV area was examined in 2008 for changes in SAV habitat per the 2007 Bay Scallop FMP. Minor changes to this line will be incorporated into the shrimp proclamation that is issued during the first week in June beginning in 2012. There have been requests made to open the western side of the IWW, particularly around Broad Creek. These requests usually come from skimmer trawl fishermen who have problems fishing in the waterway. The DMF has not opened this area because it functions as a nursery area for shrimp and other species.

11.3.2.3 Newport River

The Newport River is a relatively small estuary of about 63 square miles located north of Morehead City in Carteret County (Figure 11.6). Average depth is less than three feet with a maximum depth in natural channels of six feet and 40 feet in the dredged channels near the State Port. The western portion of the Newport River has bottoms composed of silts, clays and oyster rocks and the eastern part is composed of a firm sand bottom. There is a PNA and a SSNA located in the western portion as well.

Before the 2006 FMP, the Newport River had a long history of disagreements concerning the best location of a shrimping closure line. Lines used in the past were the Hardesty Farm line, the White Rock line (SSNA line) and the Turtle Rock line (PNA line). During this long period of conflict that peaked in the mid-1980s, the line would move several times during a season in response to requests by fishermen and the variation in shrimp size. By October of each year the river would open to the PNA line with the opening of the SSNA by proclamation. Shrimp harvest generally begins in June with the presence of brown shrimp and can continue into November and sometimes as late as December if white shrimp are abundant. The primary conflict had historically occurred in the fall, between two groups of fishermen. One group wanted the Hardesty Farm line established because shrimp that have migrated downstream to that line are a more marketable size and that line provided more towing room for their larger vessels. The other group of fishermen with smaller vessels preferred the White Rock line (SSNA) in order to access the shrimp before they moved down to the Hardesty Farm line. The White Rock line is located in shallow water, where the larger boats are unable to work because only a small portion of the White Rock line is deep enough for trawling.

Juvenile spot, croaker, brown shrimp, blue crab and southern flounder utilize the PNA and SSNA habitats in Newport River. Shellfish leases, DMF cultch plantings and natural oyster rock are also located in the SSNA. Through the 2006 Shrimp FMP, the Hardesty Farm line became a permanent line by rule; therefore protecting leases, cultch plantings and other oyster resources from being trawled over or covered in sediment. However with the implementation of the Hardesty farm line, the Fisheries Director no longer has the authority to open the Newport River SSNA.




Figure 11.6 Map of the shrimp management areas in the Newport River, North River, Ward Creek, The Straits, and Back Sound.

The implementation of the line has been successful because it protects oyster habitats, leases and cultch plantings as well as small shrimp that move out of Harlowe Creek in the early summer and provides a buffer when the abundance of juvenile shrimp, heavy rainfall or strong northerly winds pushes the shrimp downstream of their normal location. During the revision of this plan, it was requested by several fishermen to reconsider management of Newport River shrimping lines; however this management strategy will remain in place (see appendix 1).

11.3.2.4 North River

North River also has a long and interesting shrimp line history. This river was managed with two lines for years. These were the Long Point line and the Oyster House line. Both lines were established to protect small brown shrimp in the early summer (Long Point line) and small white shrimp in the fall (Oyster House line). The point of contention with these lines was when to open to the Oyster House line. Concerns with opening the area too late included the shrimp moving on a northeast wind as well as on rain and/or full or new moon. In June of 1997 a public meeting was held to discuss permanent lines in North River. It was agreed to move the Wards Creek line downstream to the mouth of the creek and move the Long Point line upstream to the next point north (Figure 11.6). These lines offered deeper water, more shelter to work in a northeast wind and provided an adequate buffer for both brown and white shrimp. The locations of these lines do allow for small brown shrimp to be caught at the beginning of the season. The old line at Long Point has been used a couple of times since the implementation of the permanent line concept because of pressure to close by fishermen because of the small brown shrimp in the area. However once the proclamation was issued, there was pressure from fishermen to honor the new permanent line. Currently, the DMF continues to keep this line as a permanent closure line unless unusual conditions such as in 2003 where high amounts of rainfall displaced small shrimp into open areas causing the DMF to close all of North River as well as the Straits. Opening dates are determined by shrimp size based on DMF sampling.

11.3.2.5 Jarrett Bay

The DMF also manages Jarrett Bay under different strategies. Since 2001, Jarrett Bay is closed to the range markers in early June by proclamation and is opened to the chimney line in July. This is to protect small shrimp in the bay until they are big enough for harvest. In the past, the DMF has opened Jarrett Bay to the chimney line in June because of pressure from fishermen out of the Marshallberg area. These fishermen say this line is easier to tow and they can keep the shrimp from moving out of the bay. Only half of the range marker line can be towed and there is more algae outside of the bay creating a lot of fouling of nets. Jarrett Bay also has a special secondary nursery area that allows it to be opened to the bridge after August 15.

11.3.2.6 Core Sound

The banks side of Core Sound from Wainwright Island to a portion of Back Sound is a shallow sand bottom area with SAV and SAV habitat was protected from shrimp trawling and mechanical clam harvest by a mix of proclamations and rule. The implementation of the 2006 Shrimp FMP placed the entire banks side of Core Sound and the eastern portion of Back Sound into rule (Figure 11.7).

The tributaries of Core Sound on the mainland side are designated as SSNA. They include Jarrett Bay, Brett Bay, Nelson Bay, Thorofare Bay-Barry Bay and Cedar Island Bay. In the




Figure 11.7 Map of shrimp management areas in Core Sound and its surround estuaries.

northwestern portion of Core Sound, bordering parts of the southern portion of the Pamlico Sound, Thorofare Bay, Long Bay-Ditch Bay and Turnagain Bay are also designated as SSNA. Prior to August 1, 2004 these SSNAs would be opened to trawling no earlier than October 15th because they were also trawl prohibited areas and coordinated whenever possible with the opening of the Newport River to diffuse effort. A rule change removing these areas from the Trawl Nets Prohibited Rule now allow these areas to be opened between August 16th and May 14th when shrimp reach a harvestable size and fish abundance is at relatively lower levels.

11.3.3 Shrimp Management in Pamlico District

Typically, as with the Central District, the annual shrimp management process begins when the DMF issues a proclamation during the first week of June that shows the location of shrimp closures lines that the season begins with. As sampling dictates, lines may be moved downstream by proclamation to protect small shrimp until they are large enough to harvest. The DMF uses a small 20 foot otter trawl with 3/4 inch mesh in the body and 1/4 inch mesh in the tailbag. This small trawl is used to determine the size structure of all the shrimp and fish in the waterbody, so that the impacts will be known. The target count size is in the neighborhood of 26-30 count or 31-35 count (heads-on). When sampling indicates that the majority of the shrimp in a closed area have reached this target size, the area is opened by proclamation.

In years when shrimp occur in great numbers, they compete for space and food and spill out into the open trawl areas because the closed nursery areas cannot contain them. Also, heavy rainfall and strong northerly winds during the month of June will cause the shrimp to move out of the closed areas seeking higher salinity. The DMF's response to finding the small shrimp in these open trawling areas has been to close them by proclamation to protect the shrimp until they reach harvestable size. This harvestable size has been the source of controversy for over twenty years.

Before implementation of the 2006 Shrimp FMP, the DMF was reluctant to close larger bodies of water like Neuse and Bay rivers or migration routes like Adams Creek. Occasionally, shrimp will be driven out of the creeks from Oriental to the mouth of the Neuse River, and from the tributaries of Bay River. When shrimp size dictated that these areas, particularly Neuse River, be closed, the closure line itself was an issue. Closing the entire river, or placing a line following channel markers running from offshore Oriental to Maw Point was used with mixed success. This enabled the larger boats to run along that line and catch small shrimp to the exclusion of the smaller boats. Smaller recreational boats were not able to work in more open and unsheltered waters and the harvestable shrimp size desired by recreational fishermen before opening is smaller than the size desired by commercial interests. For example, a 41-45 count shrimp may be more suitable to some and they want to see areas opened when that size is achieved.

"Grand openings" were also a problem with area closures. They result in a massive concentration of all types and sizes of boats in a very confined area like Adams Creek or Bay River. This increased finfish bycatch and discards because of the increased effort, increased conflict between vessels, and decreased the amount of shrimp available after the opening as opposed to a gradual migration out of a closed area over time when the shrimp themselves are ready to run. Opening times were sometimes at issue. A Sunday evening opening is convenient for Marine Patrol as far as marking the area. More odd times such as Monday at noon tended to diffuse the number of boats present at once for a "grand opening" as they gradually show up to fish that night.

An issue with the dynamic nature of the opening and closing of intensively managed areas was keeping the public informed. Immediately after an area was closed, calls by fishermen would begin, asking when the area would re-open. Proclamations require 48 hours notice and fishermen need more time than that to plan their activities.

11.3.3.1 Neuse River

The Neuse River is one of the state's larger rivers and separates Pamlico County to the north from Craven and Carteret counties to the south (Figure 11.8). The river is one mile wide at New Bern and five miles wide near its mouth, with depths ranging from 12 to 23 feet. Although shrimp and crab trawling are technically permitted from New Bern downstream to the Pamlico Sound (except when closed due to small shrimp size), shrimp are only found as far upstream as Slocum Creek. The majority of the Neuse tributaries are designated primary, secondary, or special secondary nursery areas. Shrimp generally grow in these nursery areas during the early spring and begin migrating out of them and into the river proper in July. Once in the river, they migrate around Cedar Island into Core Sound, or down Adams and Clubfoot creeks toward Beaufort Inlet to the ocean.

Before implementation of the 2006 Shrimp FMP, the management of the Neuse River had included opening the river in early June and leaving Adams Creek and West Bay opened. At that time, the river was opened to shrimp and crab trawling up to the joint-coastal line adjacent to New Bern and could be closed by proclamation due to the presence of small shrimp. In years when shrimp were scarce or of average abundance, the closure lines remained the same. When there were great numbers of juvenile shrimp in the tributaries or heavy rainfall in the critical weeks prior to reaching harvestable size, causing early movement, closures were implemented to protect the small shrimp until they reached harvestable size.

South River is currently left opened to trawling. It rarely contains shrimp, but is trawled regularly during the summer months for crabs. Most of Turnagain Bay is a SSNA, which opens with the other SSNAs in mid October.

The line that protected small shrimp on the north side of the river ran along the channel markers from Dawson Creek to the mouth of Neuse River. This line was first used in 1999 and again in 2000 when overcrowding, weather, or both forced small shrimp out of the Oriental area creeks and complaints began about catching small shrimp. The line along the channel markers was difficult to enforce and often the same size shrimp were found on the open side of the line as in the closed area. Once closed, either at the channel markers or at the river's mouth, there was always a considerable difference of opinion among the public as to the appropriate opening size, with larger commercial boats wanting a larger count and RCGL fishermen being satisfied with 40 or 50 (heads-on) count. Based on DMF sampling, the river would open on approximately 30-35 count shrimp in mid-July. When the river, creeks and bays are opened, even though there is a conscious effort to open as many areas as possible together to distribute the fishermen, there is always the grand opening aspect to contend with. For example, as many as 200 boats have been present for past opening days in Adams Creek.

Several changes in the 2004 Blue Crab FMP update, effective September 1, 2005, had indirect benefits to the Neuse River shrimp fishery. The change in designated crab pot areas in most areas of the Neuse River from a distance offshore to the six-foot depth contour and prohibiting trawling within that contour from June through November greatly decreased shrimp trawling effort in the river, particularly by the smaller commercial vessels and the RCGL fishermen. The




Figure 11.8 Map of the shrimp management areas in The Neuse River and its tributaries.

plan also set a minimum mesh size, four inch stretched mesh, for crab trawls in the western half of the Pamlico Sound and its tributaries, including Neuse River.

Management strategies in the Neuse were further modified in the 2006 Shrimp FMP restricting total headrope lengths to 90 feet upstream of the northeast line from Winthrop Point on the eastern shore of Adams Creek to Windmill Point at the entrance of Greens Creek at Oriental, effective July 1, 2006. The 2006 Shrimp FMP also established a no trawling line from Wilkinson Point to Cherry Point at the entrance of Pierce Creek. These management strategies were established to minimize juvenile southern flounder bycatch while still achieving the overall goal and objectives of the 2006 Shrimp FMP.

11.3.3.2 South Side of Neuse River

South River, Adams Creek and the outer portion of Clubfoot Creek typically stay open unless all of Neuse River closed (Figure 11.8). Adams Creek and Clubfoot are popular areas for the recreational shrimper to fish because they are small waterbodies with protection from bad weather. South River typically has very few shrimp but is a popular crab trawl area. The DMF tries not to close these areas because of concerns of grand openings. These result in a large number of small and large boats in a small waterbody. This concentration of effort on opening day increases finfish bycatch and discards, vessel conflict and decreases the amount of shrimp available after the opening.

11.3.3.3 Bay River

Bay River is a tributary of Pamlico Sound, located in Pamlico County, between the Pamlico and Neuse rivers (Figure 11.8). Trawling (shrimp and crab) is only allowed in the main stem of the river. All feeder creeks and bays are classified as either Nursery Areas (Primary or Secondary) or no trawl areas. A majority of the shrimp landed from Bay River are caught by shrimp trawls by vessels less than 40 feet in length, with a small percentage of the landings reported from skimmer trawls and channel nets. Other commercial fisheries in Bay River include crab pot, crab trawl, gill net, oyster, and long-haul.

Historically, Bay River may or may not have closed to protect small shrimp. Most closures typically occurred in mid to late June with openings in mid-July. However, following the adoption of the 2006 Shrimp FMP, actions were taken to modify openings and closures based on count size (31-35 count) and abundance. Actions were also taken in the 2006 Shrimp FMP restricting total headrope lengths to 90 feet upstream of the closure line. These management strategies were put in place to further minimize juvenile southern flounder bycatch while still achieving the overall goal and objectives of this FMP.

11.3.3.4 Pamlico River

The Pamlico River is a tributary of Pamlico Sound (Figure 11.9). Prior to the 2006 Shrimp FMP, trawling (shrimp and crab) was allowed in the main stem of the river. All feeder creeks and bays are classified as either Nursery Areas (Primary, Secondary, Special Secondary) or Inland waters all of which are closed to trawling. Overall this system is approximately 82,705 acres in size of which 76,516 acres (93%) are under DMF jurisdiction. The majority of the Pamlico tributaries are classified as Primary Nursery areas, Secondary Nursery areas, Special Secondary Nursery areas, or no trawl areas. Restrictions were put in place following the




Figure 11.9 Map of shrimp management areas in the Pamlico and Pungo Rivers.

adoption of the 2006 Shrimp FMP making it unlawful to take shrimp with trawls with a combined head rope greater than 90 feet in the waters upstream of a line between Pamlico Point and Willow Point. Further actions were taken to close the waters to trawling upstream of a line between Wades Point and Goose Creek. These management strategies were also established to minimize juvenile southern flounder bycatch while still achieving the overall goal and objectives of the 2006 Shrimp FMP. Shrimp openings typically occur in June and may or may not close due to the presence of small shrimp. Over the last 16 years the Pamlico River has not been closed to shrimp trawling. Other commercial fisheries in the Pamlico River include crab pot, crab trawl, gill net, eel potting, pound netting, and long-haul.

11.3.3.5 Pungo River

The Pungo River is a tributary of Pamlico Sound (Figure 11.9). Overall, the Pungo River is approximately 32,741 acres in size. Before the implementation of the 2006 Shrimp FMP, trawling (shrimp and crab) was allowed in the main stem of the river. All feeder creeks are classified as either Nursery Areas (Primary, Secondary, Special Secondary) or Inland waters all of which are closed to trawling. Historically, the main stem of the river would open to trawling in June. However, with the adoption of the 2006 Shrimp FMP, actions were taken prohibiting the use of shrimp trawls upstream of a line from Wades Point to Abels Bay. Other commercial fisheries in the Pungo River include crab pot, crab trawl, gill net, eel potting, pound netting, and long-haul.

11.3.3.6 Pamlico Sound

Pamlico Sound system extends from Oregon Inlet south to Core Sound (Figure 11.10). Salinity varies from 25- 30 ppt near the three inlets to near zero in the upper tributaries. Two large river systems (Neuse and Tar-Pamlico) provide the major fresh water inputs. The average depth of the sound is 16 ft. Numerous small creeks and bays surround Pamlico Sound. The Sound is divided into two basins east and west of Bluff Shoal. Extensive low salinity *Juncus* marshes border the sound and many of the tributary bays and creeks. Significant SAV beds occur in the sound, with high salinity species (e.g., eel grass) along the shoals behind the Outer Banks in the east and low salinity species (e.g., widgeon grass, wild celery) along some of the western shores. There are diurnal tides of two to three feet near the three inlets, but virtually no lunar tides away from the inlet areas. However, wind tides exceeding two feet regularly occur during storms.

Trawling (shrimp and crab) is only allowed in the main portion of the sound. All feeder creeks and bays are classified as either Nursery Areas (Primary, Secondary) or no trawl areas all of which are closed to trawling. The Pamlico Sound has the potential to close and open when shrimp are of sufficient size. Over the last 16 years portions of western Pamlico Sound have been closed six times to shrimp trawling. Other commercial fisheries in Pamlico Sound include crab pot, crab trawl, crab dredging, oyster dredging, clam kicking, gill net, pound netting, and long-haul.

11.3.3 Shrimp Management in the Northern District

Species specific shrimp sampling programs do not exist in the Northern District since shrimp management is only necessary during banner shrimp years and since ongoing shrimp data are available within the DMF juvenile trawl program data base (Program 120). An exception is data




Figure 11.10 Map of shrimp management areas in the Pamlico Sound and its tributaries.

collected in Stumpy Point Bay. Juvenile brown shrimp stations within Stumpy Point Bay were sampled by the Pamlico District staff from 1975-1986. In order to maintain this historical data

base, the Manteo office continues to sample Stumpy Point Bay for juvenile brown shrimp recruitment from 1989-present. Brown shrimp samples are typically taken during the last week of May and the first week of June using a 10.5 ft trawl with 1/4" mesh in the body and 1/8" mesh in the cod end. Thereafter, brown shrimp data are collected in the juvenile trawl survey (Program 120) from which management decisions may be made based on shrimp size and abundance.

The low relative abundance of white and pink shrimp in the northern district requires minimal sampling effort except during times of extreme environmental conditions. In such cases, sampling efforts may be initiated on demand in order to provide the foundation for shrimp management decisions of the Division.

11.3.3.1 Croatan Sound

Croatan Sound is bound by Pamlico Sound to the south, extends along the west side of Roanoke Island, to Albemarle Sound to the North. This system is approximately 26,272 acres in size. Spencers Creek is the only nursery area located in Croatan Sound and is closed to trawling. Additionally, one hundred and thirty five acres are classified as inland areas and are also closed to trawling. The majority of the shrimp trawling in Croatan Sound occurs in deep holes and sloughs. The Croatan Sound has the potential to close and open when shrimp are of sufficient size. Other commercial fisheries in Croatan Sound include crab pot, crab trawl, gill net, and pound netting.




Figure 11.11 Map of shrimp management areas in the Croatan and Roanoke Sounds.

11.3.3.2 Roanoke Sound

The Roanoke Sound system extends from Oregon Inlet north, along the east side of Roanoke Island to Albemarle Sound. This system is approximately 21,168 acres in size. Trawling (shrimp and crab) is only allowed in the main portion of the sound. With the exception of Outer Broad Creek, all feeder creeks and bays are classified as either PNAs, SNAs, SSNAs or no trawl areas. The majority of the shrimp trawling in Roanoke Sound occurs in Roanoke channel, Outer Broad Creek, and the Wanchese Channel. The SSNAs of Outer Shallowbag Bay, and Kitty Hawk Bay-Buzzard Bay, are also popular, when open. The Roanoke Sound has the potential to close and open when shrimp are sufficient size. Other commercial fisheries in Roanoke Sound include crab pot, crab trawl, gill net, pound netting, fyke net and long-haul.

11.3.4 Atlantic Ocean

North Carolina's coastline on the Atlantic Ocean is comprised of barrier islands that stretch approximately 300 miles. Shoals extending perpendicular from shore accompany capes and inlets along North Carolina's coastal ocean. On average, 22% of shrimp landed in North Carolina are harvested from these nearshore (0-3 miles) ocean waters. Near-shore hardbottom areas, dense concentrations of marine algae, artificial reefs and shipwrecks limit the amount of trawlable bottom available to commercial fishers. On average, only 3% of shrimp landed in North Carolina are harvested from offshore (>3 miles) ocean waters.

Since shrimp that migrate from the estuaries are usually large, the DMF does not actively manage the ocean waters. However, in the past and exclusively off the Brunswick county coast, DMF has been requested by the fishermen to take a more active role in the management of the ocean shrimp fishery. These requests were precipitated as result of the heavy hurricane or tropical storm induced rains that have impacted southeastern North Carolina with regularity since the mid 1990s. Fresh water from these heavy rains dramatically reduces salinities in the estuaries causing the shrimp to prematurely migrate from the estuaries into the ocean. When this occurs, DMF generally closes the impacted ocean and estuarine waters to shrimp trawling. During the revision of this plan, it was requested that trawling be allowed within a closed area located off Bogue Banks. This area is closed from Beaufort Inlet to Salter Path, NC and extends one half mile off shore and was put in place due to conflict with beach users and pier users. It was requested that the line be moved to within one quarter mile of shore, however this management strategy will remain in place (see appendix 1).

11.4 SUMMARY OF MANAGEMENT ACTIONS

11.4.1 Rules (new, modifications, or technical changes)

No rule changes required.

11.4.2 Legislative Action (new, modifications, or technical changes)

No legislative action is required.

11.4.3 Actions by Other Agencies

Various sections of state government will need to implement these actions to accomplish the processes outlined below:

1. Support Strategic Habitat Area assessments

2. Remap and monitor SAV in North Carolina to assess change in distribution
3. Assess the distribution, concentration, and threat of heavy metals and other toxic contaminants in freshwater and estuarine sediments and identify the areas of greatest concern to focus water quality improvement efforts
4. Monitor to determine if additional areas should be designated as Primary Nursery Areas due to their nursery importance to shrimp
5. Identify, designate, and protect Strategic Habitat Areas.
6. Expand habitat restoration in accordance with restoration plan goals of coastal wetlands.
7. Protect habitat from trawling and mechanical harvest gear effects through improved enforcement, establishment of protective buffers around habitats, modified rules, and further restriction of fishing gear where necessary.
8. Protect estuarine and public trust shorelines and shallow water habitats by revising shoreline stabilization rules to include consideration of erosion rates and prefer alternatives to vertical shoreline stabilization measures that maintain shallow nursery habitat.
9. Develop an interagency policy for marina siting to minimize impacts to ecologically important shallow habitats such as Primary Nursery Areas, Anadromous Fish Spawning Areas , and SAV.
10. Reduce point source pollution discharges by:
 - a) Increasing inspections of wastewater treatment facilities, collection infrastructure, and disposal sites. Providing incentives for upgrading all types of discharge treatment systems
 - b) Developing standards and treatment methods that minimize the threat of endocrine disrupting chemicals on aquatic life.
11. Improve strategies throughout the river basins to reduce non-point pollution and minimize cumulative losses of fish habitat through voluntary actions, assistance, and incentives, including:
 - a) Improved methods to reduce pollution from construction sites, agriculture, and forestry
 - b) Increased on-site infiltration of storm water
 - c) Encouraging and providing incentives for low-impact development
12. Improve strategies throughout the river basins to reduce non-point pollution and minimize cumulative losses of fish habitat through rule making, including:
 - a) Increased use of effective vegetated buffers
 - b) Implementing and assessing coastal storm water rules and modify if justified
 - c) Modified water quality standards that are adequate to support SAV habitat

13. Reduce non-point source pollution from large-scale animal operations

11.4.4 Management Related Research Needs

High Priority

- Continue to conduct bycatch characterization work across all strata (for example: dominant species, season, areas, vessel type, number of nets/rigs, headrope length).
- Initiate/increase state monitoring and reporting on the extent of unutilized bycatch and fishing mortality on fish less than age-1 in the shrimp trawl fishery.
- Continue to develop and test methods to reduce bycatch in the commercial and recreational shrimp trawl fisheries.
- Obtain mortality (immediate and post harvest) estimates of culled (active and passive) bycatch from gears used in the recreational and commercial shrimp fisheries.
- Continue to develop standard protocol for bycatch estimations.

Medium Priority

- Conduct research to quantify the number of protected species interactions with the shrimp fishery.
- Continue to develop and test methods to reduce interactions with protected species in the commercial and recreational shrimp trawl fisheries.
- Initiate sampling to investigate if additional areas currently open to shrimping need changes to their habitat designations
- Evaluate the effectiveness and efficiency of the current sampling protocol used to manage shrimp.

Low Priority

- Continue to support research to determine the status of protected species along the NC coast to better anticipate and prevent interactions (for example: migration patterns and habitat utilization).

11.4.5 Biological Research Needs

High Priority

- Continue to define and quantify the intensity, duration and spatial scale of trawling effort in NC estuaries.
- Determine species interactions and predator/prey relationships for prominent shrimp trawl bycatch.

- Determine how the resuspension of sediment, siltation, and non-point source pollution from adjacent land use practices impacts trends in shrimp abundance and habitat degradation.
- Determine the spatial and biological characteristics of submerged aquatic vegetation that maximize their ecological value to shrimp for restoration and conservation purposes.

Medium Priority

- Continue to map and quantify the habitat structure and sediment types in North Carolina estuaries.
- Continue to measure the effects of trawling on sediment size distribution and organic carbon content.

Low Priority

- Continue to investigate the impact of tiger shrimp in NC waters.
- Initiate research to determine the impacts of endocrine disrupting chemicals (EDCs) on the various life stages of shrimp.

11.4.6 Social and Economic Research Needs

Medium Priority

- Expand current social and economic surveys to specifically collect information on shrimp fishermen.
- Continue to determine the extent of recreational shrimp harvest that is occurring. This group primarily use cast nets to take shrimp either for bait or personal consumption.

11.4.7 Data Needs

High Priority

- Effort data needs to be collected to provide estimates based on actual time fished (or number of tows), rather than number of trips.
- Improve accuracy of self-reported license gear survey data, or investigate other means of accurately obtaining shrimp fleet characteristic.

11.4.8 Education

High Priority

- Encourage research and education to improve the understanding of new innovative BRDs and TEDs.
- Encourage research and education to improve the understanding and management of the shrimp resource as well as the fishery.

12.0 BYCATCH IN THE SHRIMP FISHERY AND MANAGEMENT OPTIONS

The DMF, at the direction of the MFC, presented the 2012 Shrimp FMP revision to the MFC Southern Regional AC, the MFC Northern Regional AC, the MFC Habitat and Water Quality AC and the MFC Shellfish/Crustacean AC and also took public comment at each of these committees. With the exception of the Southern AC, all of the committees voted to revise the Shrimp FMP. However, due to the overwhelming public comment concerning the issue of bycatch in the shrimp trawl fishery and the acknowledgement that bycatch is an issue in the shrimp fishery; the DMF changed its initial recommendation to the MFC from a revision to move forward with amending the Shrimp FMP. The MFC approved the plan amendment but limited the scope of the amendment to bycatch issues in the commercial and recreational fisheries.

A Shrimp FMP AC was formed in January 2013 and met over a period of eight months to become familiar with the content of the revision text in general and the bycatch issue specifically (see Section 6.3) and to review different bycatch management options. The division proposed a holistic approach to review the numerous options under consideration and directed the AC to assess the different management options through a series of evaluation matrices. Each evaluation matrix listed management options along with an initial list of potential impacts discussed by the Plan Development Team (PDT). Quantifying the potential biological gain to affected bycatch species populations was not possible with existing data; therefore it was important for the AC to consider reasonable and practicable management strategies to reduce bycatch while balancing the economic and social value of the shrimp fishery. The AC was directed to the following two FMP objectives during their deliberations:

- Minimize waste and enhance economic value of the shrimp resource by promoting more effective harvesting practices.
- Minimize harvest of non-target species of finfish, and crustaceans, and protected, threatened, and endangered species

The AC assessed bycatch reduction, economic impacts, social impacts, and inter-fishery impacts for each management option to the shrimp fishery. Enforcement and authority/administration was only assessed by the PDT. These evaluation matrices provided focused deliberations and provided a starting point for thorough and meaningful discussions in determining the best approaches for reducing bycatch in the shrimp trawl fishery. The AC was able to add options and remove options as well as change or rephrase the initial impacts as contemplated for each management option.

Twenty-nine different management options were brought forward to address eight different issues during monthly meetings from May through August 2013. Each of these issue papers follows in this Section 12, including both sets (AC and PDT) of evaluation matrices. The AC voted to remove four of those options from the evaluation process. After all options were evaluated, the members of the AC were sent an option selection package and asked to select what he or she considered to be the five best options to reduce bycatch. This enabled discussion to be focused on the best options and combinations of those options and to discuss the details needed to develop management recommendations. The AC deliberated and recommended actions for the MFC to consider (Section 12.10) to address bycatch in the shrimp fisheries.

The division also assessed management options in a similar manner. The PDT provided assessed matrices to the Management Review Team (MRT) who finalized each matrix for the DMF. Each PDT member also selected what he or she considered to be the best five options to reduce bycatch. The PDT then developed management recommendations which were then

sent to the MRT. The MRT reviewed and modified the PDT recommendations into the DMF recommendations listed in section 12.11.

12.1 TRAWLING IN THE NEW RIVER ABOVE THE HIGHWAY 172 BRIDGE

I. ISSUE

Request to reexamine the provision in the 2006 Shrimp Fishery Management Plan (FMP) which prohibits the use of otter trawls upstream of the Highway 172 Bridge over the New River.

II. ORIGINATION

Request by the Shrimp Advisory Committee (AC)

III. BACKGROUND

The use of otter trawls upstream of the Highway 172 Bridge was phased out in 2010 following the adoption of the 2006 Shrimp FMP. Those who wished to continue to harvest shrimp in the waters above the Highway 172 Bridge were allowed a four year grace period to convert to skimmers. Subsequently, crab trawls were also phased out of this area as part of the 2006 Shrimp FMP. Prior to the 2006 Shrimp FMP, crab trawlers would often fish above the Highway 172 Bridge to target flounder more so than crab; however, stricter minimum size limits for flounder made it economically unfeasible for crab trawlers to harvest only crabs in this area. In 2011, a request was made by the New River shrimp and crab trawlers to reexamine this provision. Following this request an issue paper was written for the 2011 Shrimp FMP revision. The findings of that issue paper revealed that skimmers were more effective at catching the target species than conventional otter trawls, otter trawl bycatch had been significantly reduced while the mean catch per trip (lb) for shrimp remained fairly high for the rest of the river, and the highest mean catch per trip (lb) for crab trawlers occurred in 2010 when trawlers were not allowed access to the New River Special Secondary Nursery Area (SSNA). The Plan Development Team's (PDT) recommendation was to continue to prohibit otter and crab trawlers in the New River SSNA. Currently, the waters upstream of the Highway 172 Bridge are only open to boats equipped with skimmer rigs.

The waters upstream of the Highway 172 Bridge (Figure 12.1) were designated by rule as a Special Secondary Nursery Area (SSNA) in 1996. The areas of the SSNA impacted by trawl openings include the river above the bridge up to the marked closure line running from Grey's Point to the opposite side of the river. Trawling in any of the tributary creeks is prohibited. The river consists mostly of shallow bays with the exception of the marked navigation channel. Bottom types range from sand and sand/mud to live shell bottom. The Division of Marine Fisheries (DMF) actively manages seven Shellfish Management Areas (SMAs) in this portion of New River.

Data from the DMF Trip Ticket Program were used to describe the commercial shrimp fishery in the New River from 1994 to 2011 (Tables 12.1-12.5). Landed bycatch by gear was calculated and ratios (in pounds) of marketable bycatch relative to shrimp catch were also calculated for the four main gears: channel nets, otter trawls, skimmer trawls, and various miscellaneous gears (cast nets, gill nets, etc). Marketable bycatch from skimmers was consistently lower than with the other gears. Marketable bycatch landings in channel nets were also low, with the exception of 2000-2002 when significant amounts of blue crabs were landed in this fishery. In 2005, trip limits were put in place to restrict harvest of crabs in channel nets in the first Blue Crab FMP [15A NCAC O3J.0106 (h)]. During this three-year period, ratios of pounds of shrimp

per pound of marketable bycatch in the channel nets were 4:1, 2:1, and 3:1 respectively. These bycatch ratios apply only to the portion of bycatch retained and sold.

The number of trips made by the major shrimp gears indicates a decrease in effort for all gears from 1994 to 2011 (Figure 12.2). Prior to the 2006 Shrimp FMP, channel nets were fished in the waters above and below the Highway 172 Bridge. Currently, channel nets are only allowed to be set above the 172 Bridge Channel when the river opens to skimmers by proclamation. Effort has remained low since the 2006 FMP, with only a slight increase in the number of participants (19) and trips (322) occurring in 2010 (Table 12.1). However, overall effort has not increased since the 2010 otter trawl ban. An average of 31 participants made an average of 715 channel net trips prior to the otter trawl ban and an average of 16 participants made 264 trips following the ban. While channel net landings were down as a whole as compared to the average for the years prior to the ban, the mean catch per trip was up 32.9% following the ban on otter trawls (Figure 12.3).

The numbers of otter trawl trips and participants dropped significantly in the New River following the ban of trawls above the Highway 172 Bridge (Table 12.2). Prior to the ban an average of 411 trips was made by 79 participants from 1994 to 2009 and an average of 13 trips was made by eight participants from 2010 to 2011. Overall, there was 97.0% reduction in the number of trips and a 90.5% reduction in the number of participants. Otter trawl landings in the New River dropped 95.3% after the ban (Figure 12.4). Prior to the ban, an average of 58,034 lb of shrimp were landed by otter trawls; following the ban an average of 2,749 lb were landed below the bridge. On average 174 lb of shrimp was landed per trip from 1994 to 2009; however, following the ban the mean catch per trip increased 25.2% to 218 lb per trip during 2010-2011 (Figure 12.3). Sold bycatch was reduced by 96.2%, dropping from 4,231 lb to 161 lb annually.

Much of the skimmer trawl effort in the New River has occurred in the Special Secondary Nursery (SSNA) located above the Highway 172 Bridge. Immediately following the otter trawl ban in 2010 there was a slight increase in the number of skimmer trawl trips and participants; however, these increases were not seen in 2011 (Table 12.3). Prior to the ban an average of 35 participants made 309 skimmer trawl trips annually from 1994 to 2009. Following the otter trawl ban, 13 participants made 214 skimmer trawl trips. Much like the other fisheries, skimmer trawl landings have fluctuated in response to year class strength (Figure 12.4). Prior to the otter trawl ban, skimmer trawl landings averaged 66,331 lb annually. In 2010, during the first year of the ban, skimmers landed 102,032 lb and 18,729 lb in 2011. Despite the fact that skimmer trawl landings did not increase after the otter trawl ban, the mean catch per trip increased by 11.7% (Figure 12.3).

While the number of fishermen using miscellaneous gears such as cast nets, shrimp pounds and gill nets to land shrimp in the New River have been historically low, the number of participants and trips has not increased after the otter trawl ban (Table 12.4). Prior to the ban, an average of 1,660 lb of shrimp were landed annually by seven participants. Following the otter trawl ban the landings dropped substantially (landings confidential) and the mean catch per trip declined 57.7% compared to the years prior to the ban (Figure 12.3).

The number of trips made by crab trawls also indicates a decrease in effort from 1994 to 2011 (Table 12.5). Following the adoption of the 2006 Shrimp FMP, there were no reported trips from 2007 to 2009 (12.5). In 2010, 32 trips were made by nine participants below the 172 bridge, landing 23,383 lb of crab. In 2011, 23 trips were made by five participants landing 5,680 lb of crab. Average landings of crabs following the trawl ban have been reduced by 24.2% (Figure 12.6). Prior to the trawl ban above the Highway 172 Bridge, mean catch per trip ranged from 64 to 725 pounds from 1994 to 2006 (Figure 12.6). In 2010, an average of 731 lb of crab per trip

was landed below the Highway 172 Bridge and an average of 247 lb in 2011. Overall, 489 lb of crab were caught per trip following the ban, well above the 262 lb per trip average observed from 1994 to 2006 when trawls were allowed above the bridge. Following the ban, sold bycatch was reduced by 96.2%.

IV. AUTHORITY

§ 113-134. Rules.

§ 113-182. Regulation of fishing and fisheries

§ 143B-289.2. Marine Fisheries Commission – powers and duties.

V. DISCUSSION

As part of the 2006 Shrimp FMP, otter trawls were prohibited by proclamation upstream of the Highway 172 Bridge in the New River beginning in 2010. Subsequently, this also prohibited the use of crab trawls, eliminating a traditional Sneads Ferry fishery, prompting the remaining members of that fishery to question the prohibition. In response to this query, trip ticket harvest data was used to investigate if the prohibition of all trawls, including crab trawls, above the Highway 172 Bridge was successful. As part of the 2011 Shrimp FMP revision, the PDT recommended that the Division continue to prohibit the use of otter and crab trawls above the Highway 172 Bridge to minimize waste/bycatch and disturbance to the bottom (*status quo*). The area above the Highway 172 Bridge is still designated as a SSNA (Figure 12.1).

For all gears, shrimp landings and effort in the New River were reduced substantially during 2010-2011 (Tables 12.1-12.4). Trip ticket harvest data indicates that following the prohibition of otter trawls in the New River SSNA, average shrimp landings declined for channel nets, otter trawls, skimmer trawls, and miscellaneous gears (cast nets, gill nets, etc) by 46.9%, 95.3%, 9%, and 95.6%, respectively. The average number of trips declined for these gears by 63.1%, 97.0%, 97.0%, and 89.9%, respectively. Statewide reductions in shrimp landings and effort were also observed for all gears during 2010-2011 (Tables 12.6-12.9). During this time period average statewide otter trawl landings declined 10.2% and effort (trips) declined 57.9%. Trip ticket data also indicated that the average statewide skimmer trawl landings declined by 58.4% and effort (trips) declined by 58.4% as compared to the pre-ban period (1994-2009). Also following the prohibition of otter trawls in the New River SSNA, average crab trawl landing declined in the New River by 24.2% and statewide by 82.6% (Tables 12.5 and 12.10). During the post-ban period effort (trips) declined by 66.6% in the New River crab trawl fishery; statewide the number of trips declined by 88.7%. It is important to note that not all of the reductions in landings and effort in New River are a result of the prohibition of otter and crab trawls above the Highway 172 Bridge. Declines in the average landings (shrimp, crab, sold bycatch) observed following the ban of trawling above the Highway 172 may be more of a result of environmental conditions and year-class strength. Economic hardships, increased supply of imports and rising fuel prices may have also contributed to the observed reductions in effort (participants, trips) and landings.

While landings declined for all gears targeting shrimp in the New River, the mean catch per trip (lb) increased with the exception of miscellaneous gears during 2010-2011 (Tables 12.1-12.4). The mean catch rate (lb/trip) of miscellaneous gear in the New River decreased 57.7% and increased only 3.9 % statewide (12.4 and 12.9). On average, the annual mean catch of shrimp per trip (lb) increased for otter trawls by 25.2% following the proclamation prohibiting their use above the Highway 172 Bridge in the New River. The mean catch rate (lb/trip) of skimmer trawls and channel nets in the New River increased by 11.7% and 32.9%, respectively. Statewide, the mean catch per trip (lb) of skimmer trawls declined by 12.3% and the mean catch

per trip (lb) of channel nets increased by 11.7% (12.6 and 12.8). By reducing otter trawl effort in the New River it appears that additional shrimp, once landed by otter trawls, may now be available to other gears, such as skimmer trawls and channel nets. However, increases in the mean catch per trip (lb) may also be a result of fishermen trying to circumvent rising operating costs by limiting the number of trips made until shrimp are more abundant or at a more marketable size. Since the prohibition of otter and crab trawls in the New River SSNA the mean catch per trip (lb) for crab trawls increased by 86.3% (Table 12.5). Crab trawls operating below the Highway 172 Bridge caught an average of 227 lb more crab per trip than when they were allowed in the SSNA. During this same time period (2010-2011), the statewide mean catch per trip (lb) for crab trawlers only increased 3.2% (Table 12.10). Much like the gains observed in the catch rates of the New River shrimp fishery, increases in the mean catch per trip (lb) of the crab trawlers may be the result of economic and environmental factors.

In addition to increased mean catch rates (lb/trip) for otter and crab trawls, sold bycatch has been reduced in the New River following the closure of the waters above the Highway 172 Bridge. In the years following the closure, sold bycatch from otter trawls decreased 96.2% in the New River (Table 12.2). This is slightly higher than the reductions observed statewide (-84.3%) during that same time period (Table 12.7). Sold bycatch from the New River crab trawl fishery also declined 96.2% following the closure (Table 12.5); markedly higher than 63.1% decline observed statewide (Table 12.10). As with the otter trawl fishery, reductions in the number of trips (-66.6%) and participants (-45.2%) most likely led to the observed reductions in sold bycatch in the New River crab trawl fishery. While overall reductions in bycatch (sold and discarded) cannot be quantified using trip ticket data, the declines in sold bycatch observed in the New River may suggest that overall bycatch has declined as effort has declined as a result of stricter regulations and higher operating cost.

Continuing to prohibit the use of all trawls, including crab trawls, above the Highway 172 Bridge protects the New River SSNA from bottom disturbing activities. Trip ticket data also indicates that since the closure, the mean catch rates (lb/trip) of otter and crab trawls have increased, as well as those of channel nets. Prohibiting otter and crab trawls above the Highway 172 Bridge appears to have been successful at reducing bycatch while maintaining catches of target species; however, additional data over a longer time series is needed to account for the influence of environmental conditions, year-class strength and abundance. Opening the waters to crab and shrimp trawls would only reverse the progress made in the 2006 Shrimp FMP.

VII. EVALUATION MATRICES

Advisory Committee Evaluation Matrix for Re-examination of Trawling in the New River Above the HWY 172 Bridge

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability Impact	Authority/Administrative	Other Impacts
1. <i>Status quo</i>	Limits bycatch in the Special Secondary Nursery Area. Reduces waste/fish kills on opening day. Encourages the use of non-bottom-disturbing gears. +	Will not create shifts in effort to other fisheries +/-	Continues loss of traditional otter trawl fishery in Special Secondary Nursery Area. +/-	Commercial and recreational fishing will continue with no changes in gear use or conflict. Not evaluated	Same level of enforcement. Not evaluated	Continued proclamation authority. No rule change needed. Not evaluated	Benefit to existing Shellfish Management Areas. Not evaluated
2. Allow all trawlers in the New River SSNA	Increases bycatch in the Special Secondary Nursery Area -	Increases harvest on opening day. Possible increase in pay to shrimpers +?	Re-establishes traditional otter trawl fishery in Special Secondary Nursery Area (+). Public will view as a step back in management (-). +/-	May increase conflict between otter and skimmer trawlers, as well as other user groups. Not evaluated	Same level of enforcement. Not evaluated	Implemented by proclamation authority. Not evaluated	No benefit to existing Shellfish Management Areas. Not evaluated
3. Prohibit all trawlers and skimmers in the New River Special Secondary Nursery Areas AC elected to not consider this option	Eliminates all bycatch in the Special Secondary Nursery Area. Eliminates waste/fish kills on opening day. Encourages the use of other gears.	Eliminates traditional Sneads Ferry fisheries in Special Secondary Nursery Area. Loss of income for fishermen and fish houses. Additional income from other gears may be marginal due to limited bottom space and efficiency.	Loss of traditional shrimp fisheries in Special Secondary Nursery Area.	Reduced conflict between recreational and commercial fishermen. Potential to increase the catch of bycatch species in the Special Secondary Nursery Area by other fisheries. Potential to increase competition among channel netters (limited areas in SSNA).	Same level of enforcement	Implemented by proclamation authority. Eliminates sampling associated with opening and closing Special Secondary Nursery Area.	Benefit to existing Shellfish Management Areas.

DMF Evaluation Matrix for Re-examination of Trawling in the New River Above the HWY 172 Bridge							
Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability Impact	Authority/Administrative	Other Impacts
1. <i>Status quo</i>	Limits bycatch in the SSNA. Reduces waste/fish kills on opening day. Encourages the use of non-bottom-disturbing gears. +	Will not create shifts in effort to other fisheries +	Continues loss of traditional otter trawl fishery in Special Secondary Nursery Area -	Commercial and recreational fishing will continue with no changes in gear use or conflict +	Same level of enforcement +	Continued proclamation authority. No rule change needed. +	Benefit to existing Shellfish Management Areas +
2. Allow all trawlers in the New River Special Secondary Nursery Area	Increases bycatch in the Special Secondary Nursery Area -	Increases harvest on opening day, potentially affecting market price. -	Re-establishes traditional otter trawl fishery in Special Secondary Nursery Area. May increase conflict between otter and skimmer trawlers. -	May increase conflict between otter and skimmer trawlers, as well as other user groups. -	Same level of enforcement +	Implemented by proclamation authority +	Potential increase of siltation on Shellfish Management Area -
3. Prohibit all trawlers and skimmers in the New River Special Secondary Nursery Area	Eliminates all bycatch in the Special Secondary Nursery Area. Eliminates waste/fish kills on opening day. Encourages the use of other gears. +	Eliminates traditional Sneads Ferry fisheries in SSNA. Loss of income for fishermen and fish houses. Additional income from other gears may be marginal due to limited bottom space and efficiency -	Loss of traditional shrimp fisheries in Special Secondary Nursery Area. -	Reduced conflict between recreational and commercial fishermen. Potential to increase the catch of bycatch species in the SSNA by other fisheries. Potential to increase competition among channel netters (limited areas in Special Secondary Nursery Area). +/-	Same level of enforcement +	Implemented by proclamation authority. Eliminates sampling associated with opening and closing SSNAs. +	Benefit to existing Shellfish Management Areas +

Table 12.1 Catch and effort data on shrimp and landed bycatch for channel nets in New River, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lb)	Sold bycatch(lb)	Mean catch per trip (lb)	Ratio of shrimp to sold bycatch
1994	37	544	47,556	747	87	64
1995	39	850	87,536	1,435	103	61
1996	36	585	62,590	1,894	107	33
1997	44	1,122	86,610	3,065	77	28
1998	29	856	80,714	428	94	189
1999	40	1,453	124,727	4,444	86	28
2000	45	1,380	163,109	38,998	118	4
2001	41	1,112	137,595	79,793	124	2
2002	38	1,257	163,831	61,907	130	3
2003	33	835	100,667	1,685	121	60
2004	32	570	59,799	4,370	105	14
2005	19	126	15,379	886	122	17
2006	18	206	57,011	240	277	238
2007	15	255	36,742	1,043	144	35
2008	14	168	40,892	750	243	55
2009	10	118	16,558	259	140	64
2010†	19	322	39,297	1,279	122	31
2011†	12	205	45,803	862	223	53
Pre-Ban Avg. (1994-2009)	31	715	80,082	12,621	130	56
Post-Ban Avg. (2010-2011)	16	264	42,550	1,070	173	42
% change Pre vs. Post-Ban Avg.	-49.4	-63.1	-46.9	-91.5	+32.9	-25.0

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.2 Catch and effort data on shrimp and landed bycatch for otter trawls in New River, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lbs)	Sold bycatch(lbs)	Mean catch per trip (lbs)	Ratio of shrimp to sold bycatch
1994	120	807	53,787	7,115	67	8
1995	152	1,186	152,285	12,142	128	13
1996	96	508	42,113	3,941	83	11
1997	109	828	79,788	3,721	96	21
1998	109	569	109,034	4,875	192	22
1999	141	755	77,956	4,537	103	17
2000	157	614	163,640	7,479	267	22
2001	70	186	14,926	4,389	80	3
2002	76	445	91,652	4,710	206	19
2003	67	247	39,264	5,612	159	7
2004	62	174	32,618	4,085	187	8
2005	26	58	11,820	1,528	204	8
2006	21	88	26,029	666	296	39
2007	36	71	21,117	1,735	297	12
2008	19	36	11,499	1,127	319	10
2009	7	10	1,016	30	102	34
2010†	10	13	3,450	5	265	690
2011†	5	12	2,048	317	171	6
Pre-Ban Avg. (1994-2009)	79	411	58,034	4,231	174	16
Post-Ban Avg. (2010-2011)	8	13	2,749	161	218	348
% change Pre vs. Post-Ban Avg.	-90.5	-97.0	-95.3	-96.2	+25.2	+2,088.8

Table 12.3 Catch and effort data on shrimp and landed bycatch for skimmer trawls in New River, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of shrimp to sold bycatch
1994	5	12	1,468	7	122	226
1995	25	85	21,554	0	254	0
1996	34	224	42,677	267	191	160
1997	41	341	75,029	188	220	400
1998	43	302	69,396	13	230	5,338
1999	49	449	68,813	222	153	310
2000	77	615	155,949	2,508	254	62
2001	44	306	36,043	1,879	118	19
2002	51	832	173,091	1,701	208	102
2003	55	564	89,780	1,356	159	66
2004	37	432	82,384	385	191	214
2005	24	155	21,714	307	140	71
2006	15	169	76,501	121	453	632
2007	27	265	93,094	152	351	611
2008	20	148	48,834	12	330	4,246
2009	9	42	4,973	3	118	1,658
2010†	16	297	102,032	330	344	309
2011†	10	130	18,729	401	144	47
Pre-Ban Avg. (1994-2009)	35	309	66,331	570	218	882
Post-Ban Avg. (2010-2011)	13	214	60,380	366	244	178
% change Pre vs. Post-Ban Avg.	-62.6	-30.9	-9.0	-35.9	+11.7	-79.8

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.4 Catch and effort data on shrimp and landed bycatch for miscellaneous gear (cast nets, gill nets, etc.) in New River, 1994-2011 (courtesy of DMF trip ticket program). *Data confidential due to less than three participants reporting landings.

Year	Participants	Trips	Shrimp (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of shrimp to sold bycatch
1994	*	*	*	*	*	*
1995	24	162	12,837	11,043	79	1
1996	12	20	884	1,528	44	1
1997	11	53	2,934	4,394	55	1
1998	3	6	130	442	22	<1
1999	5	10	387	553	39	1
2000	11	18	1,041	827	58	1
2001	7	9	519	819	58	1
2002	5	5	209	184	42	1
2003	5	16	670	27	42	25
2004	6	5	100	710	20	<1
2005	4	4	594	1,039	149	1
2006	10	64	4,870	349	76	14
2007	6	16	790	2,100	49	<1
2008	3	6	329	631	55	1
2009	*	*	*	*	*	*
2010†	3	3	140	104	47	1
2011†	*	*	*	*	*	*

Table 12.4 continued.

Year	Participants	Trips	Shrimp (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of shrimp to sold bycatch
Pre-Ban Avg. (1994-2009)	7	25	1,660	1,568	58	3
Post-Ban Avg. (2010-2011)	2	3	72	52	24	1
% change Pre vs. Post-Ban Avg.	-72.2	-89.9	-95.6	-96.7	-57.7	-77.4

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.5 Catch and effort data on crab and landed bycatch for crab trawls in New River, 1994-2010 (courtesy of DMF trip ticket program). *Data confidential due to less than three participants reporting landings.

Year	Participants	Trips	Crab (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of crab to sold bycatch
1994	7	35	10,848	492	310	22
1995	15	94	33,616	3,512	358	10
1996	14	47	8,284	519	176	16
1997	14	187	33,196	2,777	178	12
1998	10	62	3,988	373	64	11
1999	12	32	23,214	489	725	48
2000	11	42	17,643	555	420	32
2001	16	103	17,476	446	170	39
2002	13	77	12,190	183	158	67
2003	15	101	18,732	459	185	41
2004	23	159	41,192	863	259	48
2005	14	125	28,060	113	224	248
2006	*	*	*	*	*	*
2007	-	-	-	-	-	-
2008	-	-	-	-	-	-
2009	-	-	-	-	-	-
2010†	9	32	23,383	61	731	386
2011†	5	23	5,680	3	247	1893
Pre-Ban Avg. (1994-2009)	13	82	19,181	833	262	47
Post-Ban Avg. (2010-2011)	7	28	14,532	32	489	1,140
% change Pre vs. Post-Ban Avg.	-45.2	-66.6	-24.2	-96.2	+86.3	+2,332.2

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.6 Statewide catch and effort data on shrimp and landed bycatch for channel nets, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of shrimp sold to bycatch
1994	148	2,109	185,585	2,350	88.0	79
1995	176	2,279	272,892	3,701	119.7	74
1996	126	1,473	198,653	3,585	134.9	55
1997	136	2,088	191,188	6,404	91.6	30
1998	113	1,864	181,915	3,043	97.6	60
1999	120	2,589	284,257	17,187	109.8	17
2000	122	2,167	260,321	41,280	120.1	6

Table 12.6 continued.

Year	Participants	Trips	Shrimp (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of shrimp sold to bycatch
2001	97	1,623	185,277	80,288	114.2	2
2002	88	1,865	250,656	62,513	134.4	4
2003	86	1,697	255,892	3,523	150.8	73
2004	83	1,351	149,933	5,553	111.0	27
2005	57	864	130,710	2,138	151.3	61
2006	60	896	181,102	2,131	202.1	85
2007	67	954	165,729	7,521	173.7	22
2008	66	1,101	253,530	7,903	230.3	32
2009	60	1,084	180,704	4,199	166.7	43
2010†	57	1,063	129,865	3,303	122.2	39
2011†	40	531	97,908	2,433	184.4	40
Pre-Ban Avg. (1994-2009)	100	1,625	208,022	15,832	137	42
Post-Ban Avg. (2010-2011)	49	797	113,887	2,868	153	40
% change Pre vs. Post-Ban Avg.	-51.7	-51.0	-45.3	-81.9	+11.7	-5.0

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.7 Statewide catch and effort data on shrimp and landed bycatch for otter trawls, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lb)	Sold bycatch (lb)	Mean catch per trip (lb)	Ratio of shrimp sold to bycatch
1994	845	14,583	5,240,153	666,665	359.3	8
1995	888	15,481	5,729,152	478,805	370.1	12
1996	705	11,007	3,055,860	428,639	277.6	7
1997	722	12,702	4,911,799	448,060	386.7	11
1998	513	8,297	2,019,600	577,421	243.4	3
1999	667	10,817	5,275,158	392,835	487.7	13
2000	793	10,521	7,847,702	299,773	745.9	26
2001	553	7,734	3,493,218	235,398	451.7	15
2002	639	10,030	7,511,154	270,553	748.9	28
2003	439	6,682	3,179,629	315,436	475.8	10
2004	421	5,358	2,581,743	217,756	481.8	12
2005	272	2,890	1,078,088	67,411	373.0	16
2006	297	3,252	2,891,435	84,524	889.1	34
2007	338	4,464	7,123,976	138,746	1,595.9	51
2008	364	4,204	6,764,108	161,531	1,609.0	42
2009	340	3,890	4,049,599	123,416	1,041.0	33
2010†	355	3,943	4,280,703	68,106	1,085.6	63
2011†	301	3,003	3,889,637	27,984	1,295.3	139
Pre-Ban Avg (1994-2009)	550	8,245	4,547,023	306,685	659	20
Post-Ban Avg (2010-2011)	328	3,473	4,085,170	48,045	1,190	101
% change Pre vs. Post-Ban Avg.	-40.3	-57.9	-10.2	-84.3	+80.8	+401.8

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.8 Statewide catch and effort data on shrimp and landed bycatch for skimmer trawls, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lb)	Sold bycatch(lb)	Mean catch per trip (lb)	Ratio of shrimp sold to bycatch
1994	79	1,118	203,866	678	182.3	301
1995	128	1,563	424,181	1,636	271.4	259
1996	102	1,179	188,666	4,824	160.0	39
1997	143	2,203	339,056	1,828	153.9	186
1998	92	1,058	179,387	786	169.6	228
1999	155	2,080	599,465	1,666	288.2	360
2000	180	2,429	624,010	3,671	256.9	170
2001	135	1,765	314,994	5,262	178.5	60
2002	158	3,565	831,511	3,919	233.2	212
2003	130	2,535	475,582	8,004	187.6	59
2004	101	2,097	377,173	1,537	179.9	245
2005	72	1,101	176,928	719	160.7	246
2006	87	1,344	686,475	436	510.8	1576
2007	84	1,556	586,700	2,891	377.1	203
2008	92	935	365,331	234	390.7	1558
2009	60	807	181,458	189	224.9	960
2010†	64	1,095	284,972	381	260.2	748
2011†	31	327	55,576	404	170.0	138
Pre-Ban Avg. (1994-2009)	112	1,708	409,674	2,392	245	416
Post-Ban Avg. (2010-2011)	48	711	170,274	392	215	443
% change Pre vs. Post-Ban Avg.	-57.7	-58.4	-58.4	-83.6	-12.3	+6.4

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.9 Statewide catch and effort data on shrimp and landed bycatch for miscellaneous gear (cast nets, gill nets, etc.), 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lb)	Sold bycatch(lb)	Mean catch per trip (lb)	Ratio of shrimp sold to bycatch
1994	49	185	10,719	53,426	57.9	<1
1995	106	557	50,594	55,855	90.8	1
1996	39	186	4,766	22,234	25.6	<1
1997	51	241	6,247	17,448	25.9	<1
1998	37	167	3,576	31,368	21.4	<1
1999	37	144	9,999	48,305	69.4	<1
2000	63	171	5,611	21,818	32.8	<1
2001	40	213	3,511	14,937	16.5	<1
2002	52	233	5,053	17,030	21.7	<1
2003	25	148	1,826	15,496	12.3	<1
2004	19	154	1,792	5,308	11.6	<1
2005	16	118	2,687	3,297	22.8	1
2006	21	111	5,785	3,110	52.1	2
2007	23	62	3,473	8,646	56.0	<1
2008	20	65	2,653	4,764	40.8	1
2009	14	152	5,468	2,137	36.0	3

Table 12.9 continued.

Year	Participants	Trips	Shrimp (lb)	Sold bycatch(lb)	Mean catch per trip (lb)	Ratio of shrimp sold to bycatch
2010†	17	136	5,984	2,186	44.0	3
2011†	17	163	5,404	5,598	33.2	1
Pre-Ban Avg (1994-2009)	38	182	7,735	20,324	37	1
Post-Ban Avg (2010-2011)	17	150	5,694	3,892	39	2
% change Pre vs. Post-Ban Avg.	-55.6	-17.7	-26.4	-80.9	3.9	213.9

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

Table 12.10 Statewide catch and effort data on crab and landed bycatch for crab trawls, 1994-2011 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Crab (lb)	Sold bycatch(lb)	Mean catch per trip (lb)	Ratio of crab sold to bycatch
1994	239	3,394	1,858,304	153,728	547.5	12
1995	213	1,918	1,045,927	98,997	545.3	11
1996	285	4,051	3,075,373	150,693	759.2	20
1997	293	4,595	3,268,736	152,629	711.4	21
1998	258	5,303	3,065,385	161,243	578.0	19
1999	200	3,246	1,799,454	126,029	554.4	14
2000	167	2,051	922,254	105,831	449.7	9
2001	194	2,332	984,162	107,758	422.0	9
2002	126	958	1,113,491	78,914	1,162.3	14
2003	131	1,605	1,252,366	135,128	780.3	9
2004	170	1,670	886,719	90,318	531.0	10
2005	94	1,027	378,714	47,897	368.8	8
2006	34	243	129,312	19,650	532.1	7
2007	27	115	25,839	6,352	224.7	4
2008	42	278	1,555,327	49,946	5,594.7	31
2009	57	436	911,907	79,812	2,091.5	11
2010†	51	261	286,359	18,693	1,097.2	15
2011†	39	210	199,181	53,462	948.5	4
Pre-Ban Avg (1994-2009)	158	2,076	1,392,079	97,808	991	13
Post-Ban Avg (2010-2011)	45	236	242,770	36,077	1,023	10
% change Pre vs. Post-Ban Avg.	-71.5	-88.7	-82.6	-63.1	+3.2	-27.4

†Otter and crab trawls not allowed above Highway 172 Bridge in the New River

New River Trawl Areas




Figure 12.1 Map of the New River showing the areas for the PNAs and SSNA as well as the other trawl closure line.




Figure 12.2 Shrimp trips by gear in New River, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.




Figure 12.3 Mean catch of shrimp in New River, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.




Figure 12.4 Total catch of shrimp in pounds by gear in the New River, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.




Figure 12.5 Number of trips and participants in the New River crab trawl fishery, 1994-2011. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.




Figure 12.6 Pounds of crab landed and number of pounds of crab landed per trip in the New River crab trawl fishery, 1994-2010. Dotted line represents the average pounds per trip landed from 1994-2010. †Otter and crab trawls not allowed above Highway 172 Bridge in the New River.

12.2 EVALUATION OF THE SKIMMER TRAWL AND OTHER GEARS USED FOR SHRIMPING IN NORTH CAROLINA

I. ISSUE

The exploration of the skimmer trawl along with other gears to harvest shrimp and reduce bycatch in estuarine waters

II. ORIGINATION

The Marine Fisheries Commission, the public and Division staff

III. BACKGROUND

The management of shrimp occurs by district (Figure 12.7). The majority of estuarine shrimping occurs in the Pamlico Sound (81%) which is co-managed by the Pamlico District and Northern District. The Central District and Southern District make up 11% and 4% respectively. The Pamlico and Northern Districts make up approximately 4% of total estuarine shrimp landings excluding Pamlico Sound.

Effort has decreased over time in otter trawls from a high of 15,482 trips in 1995 to a low of 3,004 trips in 2011. Skimmer trawl effort peaked in 2002 at 3,565 trips but has since decreased to 327 trips in 2011 (Table 12.11). Channel nets show a similar decrease from 2,589 trips in 1999 to a low for 531 trips in 2011. Compared to the overall number of otter trawl trips, skimmer trawls, channel nets, cast net and pound net trip numbers are low (Table 12.11). This is

probably due to the overall effectiveness of otter trawls compared to the limited effectiveness of these other gears, based on where they are able to be fished. There was a slight increase in the number of cast net and pound net trips occurring from 2007 through 2011 (Table 12.11). The average number of otter trawl trips is five times the average number of trips made with skimmer trawls with average catch per unit effort (CPUE) 583 lb of shrimp per trip. CPUE for skimmer trawls is 240 lb of shrimp per trip while CPUE for channel nets is 129 lb per trip (Table 12.11). There is generally low effort in the use of cast nets and shrimp pound nets. Pound net data availability is limited because of confidentiality, due to less than three dealers reporting landings (Table 12.11).

The shrimp fishery is the second most valuable fishery in North Carolina. The otter trawl portion of the shrimp fishery annually contributes an average of \$9,776,788 dockside value to the economy with an average value per trip of \$1,267. Skimmer trawls and channel nets also make large annual contributions to the fishery with values of \$566,512 and \$359,367 respectively (Table 12.1). The value per trip is considerably lower for skimmer trawls (\$355) and channel nets (\$234) than otter trawls (Table 12.1). This may be attributed to differences in the amount of gear fished by each type of vessel or overall gear effectiveness.

Participation in the estuarine shrimp fishery has decreased in the otter trawl, skimmer trawl and channel net fisheries. Cast net and shrimp pound net participation has been variable over time with no apparent trends. Otter trawl participation was highest with 888 participants in the fishery in 1995. The fishery has experienced low participation for the last several years with a 66% decline to 301 participants in 2011. Skimmer trawl participation was highest in 2000 with 180 participants but similar to otter trawls, has dropped 83% to 31 participants in 2011. Channel nets have also shown a steady decline in participation dropping 77% from 176 participants in 1995 to 40 participants in 2011.

Factors to be considered when evaluating the effectiveness of skimmer trawls are targeted shrimp species, seasons, water depths, clear bottom, tidal current and tidal range and economic potential. Skimmer trawls are most effective on white shrimp in shallow water depths over bottom clear of structure so that the rigid frames do not foul on the bottom.

The effectiveness of other gears such as channel nets, cast nets and shrimp pounds is determined by what shrimp species is being targeted, seasons, water depths, bottom type, tidal current and tidal range and economic potential. Similar to skimmer trawls, channel nets require specific areas to fish based on bottom contours and water depth. Channel nets are fished in the evening, during a falling tide, in areas that are so specific, they are named by the fishermen. Effective water depths range from 12 to 20 feet deep and tend to be located in deep holes where tidal flow is strong enough to hold the net open.

Cast netting for shrimp has been a popular method to catch bait for hook and line fishing, and in more recent years, for food. In addition, fishermen have the ability to use cast nets in areas closed to shrimping, such as nursery areas, areas closed due to small shrimp and areas closed due to habitat concerns. The limit for cast netting shrimp will be changed to two quarts June 1, 2013 to enable law enforcement to more safely and efficiently enforce this rule. Cast netting requires the skill to effectively throw a round cast into the water and works best when the cast netter is near marsh edge, along a river, off a dock, bridge or boat, over clear non-structured bottom. Although cast netting generally targets white shrimp, brown shrimp may also be harvested with cast nets.

Shrimp pounds require high tidal flow and concentrations of shrimp moving toward channels and inlets. These are best for brown shrimp in shallow water at night; however, they are not as effective for white shrimp that move in deeper water. Shrimp pounds appear to be more functional in Topsail Sound compared to other regions because of the high tidal current and soft bottom habitat (Session and Thorpe 2006).

Southern District

The shrimp fishery within the estuarine waters of the Southern District makes up a large number of small boats fishing in the Intracoastal Waterway, New River and Cape Fear River. Otter trawl effort has declined over time with a maximum of 3,330 trips in 1995 and the lowest number of trips (599) taken in 2006 (Table 12.12). A management strategy that was put in place through the 2006 Shrimp Fishery Management Plan in New River limits otter trawls to the waters below the Highway 172 Bridge, only allowing skimmer trawls to work the waters above the bridge. As a result of this management strategy, marketable bycatch from otter trawls has declined and otter trawl effort has been reduced. While it has only been two years since otter trawls were phased out, preliminary data indicates this has been a viable management strategy; potential lower bycatch, lowering discard mortality and reducing bottom disturbing activities while maintaining the catches of shrimp.

Skimmer trawls are typically fished in shallow waters, no deeper than 15 feet and are used to target white shrimp. In the Southern District, skimmer trawls work the IWW in Onslow and Pender counties up and into New River, Bear Creek and Queens Creek (Figure 12.8 - Figure 12.11). There is very little, if any, effort in New Hanover and Brunswick counties due to the tidal range and bottom contour. Overall, 21% of skimmer trawl landings come from the Southern District from 1994-2011. Skimmer trawl trip number has been variable through the time series and effort in the fishery is most likely based on the year to year availability of white shrimp. The highest number of trips were made in 2002 (1,073 trips) and the lowest number occurred in 1994 (13 trips) (Table 12.12). Recreational Commercial Gear License (RCGL) trawl trips in the southern area decreased from 1,100 in 2002 to 500 in 2008 (otter trawls and skimmer trawls were not differentiated).

Effort in the Southern District for channel nets is highest in New River with fishing also occurring in the IWW, Topsail and Stump Sound (Table 12.12). The majority of cast netting occurs in the IWW with some effort occurring in the Cape Fear River and Lockwood Folly targeting white shrimp in the late summer and early fall. Channel net effort has decreased to a low of 282 trips in 2011 since 1999 when 1,967 trips were recorded (Table 12.12). Cast net trips and landings have remained variable ranging from zero trips made in 2006 to 122 trips made in 2002. Based on a 2012 recreational cast net survey, 37% of recreational cast netters fished in the Southern District. While overall landings from shrimp pounds are low, effort is increasing, especially in the Southern District where tidal flow is probably a little more consistent for its use. However, data are considered confidential. No data were collected on shrimp pounds during RCGL surveys.

Central District

The shrimp fishery in the Central District occurs in Core Sound, North River, Newport River, Bogue Sound and White Oak River. In the Central District, shrimp fishermen tend to fish both otter and skimmer trawls. Fishermen will start out the shrimp season using otter trawls in the spring to target pink shrimp and in the summer to target brown shrimp. The fishermen will then switch over to skimmer trawls in the fall to fish for white shrimp. They may switch back to using otter trawls in the late fall to catch those shrimp that are migrating through deeper channels as

they make a seaward migration. Otter trawl activity has steadily declined since 1995 where 7,150 trips were made to a low of 173 in 2011. Similar to the Southern District, Central District skimmer trawl effort has been variable over time with a high of 2,391 trips in 2002 and a low of 105 trips in 2011 (Table 12.13). Skimmer trawl activity occurs in White Oak River, part of the IWW in Bogue Sound and on the banks side between Hoop Hole Creek and Salter Path. Newport River, North River, and areas around Harkers Island are also popular areas for skimmer trawls (Figure 12.11 - Figure 12.13). The mainland side of Core Sound and its tributaries are also worked by skimmer trawls. Similar to the Southern District, RCGL trawl trips decreased from 1,000 trips in 2002 to 130 trips in 2008.

In the Central District the majority of channel nets are fished in Newport River, North River, and Core Sound (Table 12.3). Channel net effort has been stable for most of the time series; however the highest effort occurred during 1994 with 1,296 trips made. Channel net effort in 2010 and 2011 showed a decline from 476 trips to 249 trips in 2011 (Table 12.13). Most cast netting occurs in the White Oak River, Newport River and North River with variable effort. Forty-seven percent of recreational cast netters surveyed in 2012 were in the Central District of which 26.5% of those were in Bogue Sound. Based on Trip Ticket data, there has been no shrimp pound effort or landings since 1997 (Table 12.13).

Pamlico District

In the Pamlico District, the majority of the fishing effort is from otter trawls in the rivers. Though effort has declined in this district, the declines do not appear to be as steep as in the Southern and Central Districts (Table 12.14). The highest effort was in 1997 with 966 trips and lowest effort occurring in 2009 with 224 trips (Table 12.14). Skimmer trawl activity is low and variable, but effort has increased over time in areas in the Neuse River along the shore between Dawson Creek and the town of Oriental and in Adams Creek (Figure 12.13). The RCGL survey shows that most recreational trawling occurred in the Pamlico District and effort remained steady throughout the time period. An average of 1,600 trips was made from 2002 to 2008.

Channel nets and shrimp pounds are used very little in the Pamlico District and in recent years there have been no reported landings from these gears. In the Pamlico District, the predominant tidal currents and range are driven by the wind compared to the Southern and Central Districts where tides are more lunar driven, thus explaining the limited use of this gear in these areas (Table 12.14). Very little cast netting is done in the Pamlico District with 0% of recreational cast netters surveyed in 2012.

Northern District and Pamlico Sound

The otter trawl is the predominant gear used in the Northern District as well as in Pamlico Sound (Table 12.15 and Table 12.16). Otter trawl effort has decreased in the Pamlico Sound, with only 1,452 trips occurring in 2011, down from peak effort of 4,117 trips during this time series (Table 12.16). There appears to be some use of skimmer trawls within Pamlico Sound along the mainland side from Wysocking Bay to Stumpy Point Bay when shrimp abundances are high (Figure 12.14). RCGL data indicates lower but somewhat steady trip numbers in the Northern District; decreasing from 742 trips in 2002 to 337 trips in 2008.

There is sporadic but limited use of channel nets and cast nets in Pamlico Sound. Only 1.5% of the cast netters surveyed fished in the northern district. In recent years, no shrimp pounds have been fished in Pamlico Sound. On average 12 skimmer trawl trips were made from 1994 to 2011. The Pamlico Sound can be too rough and is generally too deep for skimmer trawls,

therefore making the otter trawl the most efficient gear to be used in Pamlico Sound (Table 12.16).

IV. AUTHORITY

§ 113-134. Rules.

§ 113-173. Recreational Commercial Gear License.

§ 113-182. Regulation of fishing and fisheries.

§ 143B-289.2. Marine Fisheries Commission – powers and duties.

V. DISCUSSION

The majority (89%) of the estuarine shrimp harvest in North Carolina comes from otter trawls. However, major concerns of otter trawls is the capture and discard of various amounts of other non-target species and discard mortality associated with otter trawls. Other concerns include any impacts that may have an influence on the amount of resources available to recreational and commercial fishing. Commercially and recreationally valuable finfish species such as southern flounder, weakfish, spot, and Atlantic croaker are of special concern to fishermen in North Carolina.

McKenna and Clark (1991) explored ways to reduce bycatch in the shrimp fishery through the development and testing of shrimp pot designs and the feasibility of using cast nets to harvest brown shrimp in the Pamlico Sound complex. They found that the limiting factors of shrimp pot development and use of cast nets was shrimp behavior and the lack of suitable bait.

There are also areas (15A NCAC 03R .0106) closed to trawling because of the presence of juvenile finfish and shrimp and habitat concerns but may be conducive to passive type gears and cast nets at certain times of the year. Because of these concerns, this paper explores the use of gears other than otter trawls to harvest shrimp and the bycatch that occurs in these gears. The more popular gears used include skimmer trawls, channel nets, cast nets, and shrimp pounds. These other gears combined make up approximately 11% of the average total estuarine shrimp landings from 1994 to 2011 compared to 89% from the otter trawl.

Otter Trawl

An otter trawl net is dragged along the bottom behind a towing vessel. The mouth of the net is held open by two large "doors" which are attached to either side of the net. The net is dragged behind the boat or on either side of a vessel attached to outriggers. Once the cod end of the net is filled or hauled for a certain time, it is hauled back aboard the vessel where the catch is spilled from the cod end and the net redeployed.

Commercial shrimp trawling in North Carolina began in 1916 in the Southport area. The practice spread throughout the rest of North Carolina over the next couple of decades. Following World War II, there was a considerable increase in effort. Technological advances in the shrimp industry have increased the catching efficiency of larger boats. In the 1940s and early 1950s, a 45-60 foot vessel pulled a single trawl with a head rope length of 60-65 feet. Due to improvements in engine design, the same sized vessel, using four-barreled rigs, can now pull four nets with a combined head rope length of 120-160 feet. Four-barreled rigs allow fishermen to pull two nets from each outrigger.

Skimmer Trawl

A skimmer trawl consists of two nets, mounted on both sides of a vessel on a rigid frame and pushed through the water column. Skimmer trawls were introduced in North Carolina by two Carteret County fishermen who had read about them in a 1989 National Fishermen magazine. They approached North Carolina State University Sea Grant Program to investigate the effectiveness of skimmer trawls in North Carolina waters. Following the request, a series of Sea Grant papers were later published (Coale et al. 1994; Rudershausen and Weeks 1999, Hines et al. 1999). The Rudershausen and Weeks (1999) study examined the mechanical efficiency of experimental skimmer trawl frames constructed of different materials and the Hines et al. (1999) study examined the catch composition of high (12 ft) and low profile (3 ft) skimmer trawls.

Coale et al. (1994) compared shrimp catch and bycatch between a skimmer trawl and a four-seamed balloon otter trawl fishing in close proximity in Straits Channel and North River in Carteret County from June through August of 1991. The skimmer trawl was found to be very effective for catching white shrimp while the otter trawl was more effective at catching brown shrimp. On average, the skimmer trawl caught less bycatch (mean, 0.47 kg/min) than the otter trawl (mean, 0.66 kg/min). It is important to note that the otter trawl tested in this study was not equipped with a turtle excluder device (TED) or a bycatch reduction device (BRD); BRDs were not required in North Carolina until 1992. Bycatch was thought to be reduced as a result of the skimmer's tailbag being fished more frequently. The tailbag of a skimmer can be dumped while the gear continues to fish, thus reducing culling times per tow allowing more fish to be returned to the water alive. Based on live well experiments, 12 finfish species exhibited lower mortality when caught by the skimmer trawl.

Scott-Denton et al. (2006) also concluded that bycatch rates in skimmer trawls are lower compared with historical and current estimates of bycatch associated with otter trawls in Louisiana. In their study, the discard to landing ratio was much lower for skimmer gear (0.63:1) compared to that of the otter trawls in the Gulf of Mexico (4.56:1; Harrington et al. 2005). While bycatch associated with skimmers appears to be lower as a whole compared to otter trawls, Hines et al. (1999) found that low profile net (3 ft) was not effective in reducing bycatch as compared to the high profile net (12 ft) and actually was less effective at capturing brown shrimp.

The benefits of the skimmer trawl include the reduction of finfish bycatch, less bottom disturbance, less fuel consumption, more effective fishing time, and less culling time (Coale et al. 1994; Rudershausen and Weeks 1999; Scott-Denton 2006). The down side of skimmer trawls is that while they are more efficient at catching white shrimp as compared to an otter trawl, they are less efficient at catching brown and pink shrimp (Coale et al. 1994). The white shrimp season can be very short in duration (6 to 8 weeks) and only occurs in the summer and early fall. Skimmer trawls are less likely to be used during a poor white shrimp season which is dependent on prevailing winter water temperatures. The bottom where skimmer trawls work must be free of obstructions due to the rigidity of the gear mouth opening. The vertical height of the skimmer frame also limits the depth by which it can fish; typically working best in shallow water less than 10 to 12 ft deep. Hines et al. (1999) noted that deeper depths substantially increased steering problems due to drag. In contrast, otter trawls are more versatile allowing fishermen to follow shrimp to deeper waters or channels (Hines et al. 1999).

Channel Nets

Channel nets are stationary nets that fish the surface and middle depths on an outgoing tide. They resemble a trawl anchored and staked to the bottom to keep it open. Channel nets offer the advantages of less fuel consumption although there is very little information about bycatch in the channel net. Channel net caught shrimp are considered by the public as a higher quality shrimp. Since these shrimp are migrating out of the estuaries when caught, they tend to be harder shelled, cleaner and bigger than trawled shrimp. Channel nets are set at night on an ebb tide with mouth oriented toward the direction of the oncoming current. The tailbag of the channel net is emptied into a skiff every 15 to 30 minutes. The net is retrieved from the water before the tide changes to prevent it from being turned inside out. There are a limited number of areas that channel nets can be set, with the majority of the effort concentrated in the estuarine waters from Beaufort Inlet to Rich's Inlet. Channel nets must be set near inlets where the current is strong and shrimp have concentrated to move out to sea. Channel nets account for 5% of the average annual shrimp landings.

Cast Nets

The cast net is another type of gear used to harvest shrimp. It consists of a circular net weighted around the perimeter that is thrown out over the shrimp. The weighted edges of the cast net sink to the bottom enveloping the shrimp which is then pulled into the catcher by a line attached to the top of the net. The cast net is most successful on white shrimp in the fall as they school in large concentrations and leave the creeks and tributaries and head for the sounds and, eventually, the ocean. A few pink and brown shrimp are captured around the marshes and shallows during the summer. Throwing from boats or bridges over creeks is productive when they are migrating.

There are little data on catch rates and shrimp size in cast nets. Whitaker et al. (1991) examined catch rates and size of white shrimp caught with cast nets of different mesh sizes over bait. They looked at 3/8 inch, 1/2 inch and 5/8 inch mesh sizes in the Ashley River in South Carolina at night. Overall catch rates of all mesh sizes combined were 37 shrimp per cast with 3/8 inch mesh averaging 55 shrimp per cast, 1/2 inch averaging 36 shrimp per cast and 5/8 inch averaging 21 shrimp per cast. Heads on count estimates were 71, 59, and 41 for 3/8, 1/2, and 5/8 inch mesh respectively. Bycatch from all three mesh sizes were low and included blue crab, catfish, mullet, silver perch, menhaden and brief squid. This study resulted in a recommendation for a minimum mesh size of 1/2 inch mesh cast net for the South Carolina shrimp baiting fishery.

McKenna and Clark (1993) tested the feasibility of using cast nets to harvest brown shrimp in primary and secondary nursery areas bordering Pamlico Sound. They evaluated three different sized meshes of cast nets (3/8 inch, 1/2 inch, and 5/8 inch) over bait balls made of menhaden meal and mud. No shrimp were captured, although there were high densities of shrimp in the area. They modified the sampling methodology by adding a heavier weighted 1/2 inch mesh cast net over bait and also included additional bait types such as canned dog and cat food, bricks soaked in menhaden oil and areas of no bait. They sampled at sunrise, sunset, in the middle of the day and during the night. Nineteen brown shrimp and one white shrimp were captured in 139 throws. The 5/8 mesh net had the highest CPUE for brown shrimp at 0.44 shrimp/cast and over bait balls at 0.18 brown shrimp/cast. Bycatch was low and included Atlantic menhaden, silver perch, blue crab, spot, southern flounder, Atlantic croaker and southern kingfish. They concluded that a lack of tidal influence could affect shrimp behavior in terms of movement and feeding activity and the lack of suitable bait.

Shrimp Pounds

Although not a significant contributor to shrimp landings, shrimp pounds have recently been developed and employed in the taking of brown shrimp. Shrimp pounds are semi-permanent or non-permanent static pounds with a V-shaped lead that directs a shrimp to a funnel connected to a box-shaped pound in shallow waters of the Intracoastal Waterway (IWW) and beach areas in the evenings. One of the leads extends to the shoreline and the other extends out towards a channel or deeper water. Shrimp enter the nets at night as they migrate. The larger shrimp are penned in the pound while the smaller ones are allowed to pass through. These shrimp pounds are used in the southern region of North Carolina and have been around since the mid 1990s. There is minimal disturbance to bottom habitat, and low finfish bycatch. Shrimp and bycatch remain mobile within the pounds thus increasing survivability of finfish bycatch. Sessions and Thorpe (2006) reported average catch rates of shrimp to be 4.5 lb//hour with larger shrimp occurring in June and July. Finfish bycatch consisted of 16 species and were caught between June and November. Catch rates for finfish bycatch averaged 0.1 lb//hour. The average finfish to shrimp ratio was 0.31:1. Pinfish were the most abundant finfish caught making up 66% of the total bycatch by number; menhaden and spot made up 8% and 5% respectively. Spot, croaker, pigfish, southern flounder and striped mullet cumulatively accounted for 8.5% of the total bycatch. Blue crab, shortfin squid, and stone crab were the top ranked invertebrate bycatch (Sessions and Thorpe 2006).

Mortality rates were low with a total of 98% of the finfish species released alive and 95% released alive in good condition. Only menhaden (17%) and one filefish experienced mortality within the gear. Other finfish that were released in fair or poor condition included menhaden, pinfish, white perch, and spot.

Brown and Price (2006) investigated a shrimp pound consisting of two stacked pots and two leads to determine its potential as a recreational gear. Average catch rates of shrimp were 2.7 lb of shrimp and 0.24 lb of bycatch on flood tide while average catch rates on an ebbing tide was 2.3 lb of shrimp and 0.11 lb of bycatch. Brown shrimp made up the majority of the shrimp catch at 96% and 99% of total weight from flood tide sets and ebb tide sets, respectively. Bycatch consisted of blue crabs and pinfish. Only one spot suffered mortality throughout this study. Moon phase, current speed, salinity, and temperature were found to effect catch per unit effort. Shrimp pounds are easy to set up and operate, inexpensive, and easily transported as compared to trawls. However, the pounds only work in certain habitats and are most successful when set during a flood tide with one wing against a bulkhead or marsh shoreline. Results from this study led to the development of shrimp pounds as a Recreational Commercial Gear License (RCGL) gear and another choice for the RCGL holder because of low bycatch and low impacts to the habitat as compared to the otter trawl. Currently, recreational shrimp pounds must be attended, properly marked, and set a minimum of 100 yards from another RCGL shrimp pound or 300 yards from a permitted commercial shrimp pound.

The estuarine otter trawl fishery is the most effective way to harvest shrimp in North Carolina and makes the shrimp fishery the second most valuable fishery in the state behind the blue crab fishery. Skimmer trawls, when used in specific areas during white shrimp season have been shown to be effective in catching commercial quantities of shrimp with less bycatch. However skimmer trawls are limited by water depth and bottom type, and almost exclusively operate in the late summer/early fall white shrimp fishery. Currently skimmer trawls are being fished in the areas they are most effective. Further skimmer trawl characterization studies are also needed.

Other gears specific to harvesting shrimp may have less bycatch than the otter trawl, but are less effective at catching commercially viable market quantities of shrimp. Each gear requires a specific environmental condition to operate efficiently such as depth, bottom profile, and/or current or tidal flow requirements. The seasonality of brown and white shrimp is also important to consider when and where other gears can be used. All of the gears discussed are currently being fished in areas where they are most effective. The use of cast nets in areas otherwise closed to shrimping could be considered based on shrimp availability and size; still allowing fishermen access to the resource while limiting bycatch.

VII. EVALUATION MATRICES

Advisory Committee Evaluation of Skimmers and Other Gears for Shrimping in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. Effort reduction has resulted in reduced bycatch and will continue. Gear is more effective, even if effort is reduced +/-	Will not create shifts in effort to other fisheries. Maintains present market value of fishery. +/-	Allows flexibility of use of gears in the fishery. +	Commercial and recreational fishing will continue with no changes in gear use or conflict. not evaluated	Same level of enforcement. not evaluated	Continued proclamation authority. No rule change needed. not evaluated	Allows for further characterization and bycatch reduction studies to fill data gaps prior to new regulations. not evaluated
2. Designate skimmer trawl areas/seasons. Consider an August-November skimmer trawl season in Newport River, North River, Jarrett Bay, Phase in skimmer trawls	Likely decrease in the amount of bycatch in specific areas and during specific seasons. Areas where gear can be used are limited due to physical characteristics of area. +	More profitable at certain times in certain areas. Cost of re-rigging (-). Greater efficiency (+) Greater areas for skimmer trawls (+) Loss of marketable bycatch (-) ++/- -	May reduce the seasonal availability of local brown and pink shrimp. -	Likely to increase conflict among commercial fishermen. Effort shifts may impact other fisheries in same area. Potential to increase other fisheries' catches of adult bycatch species. not evaluated	Need determination of enforceable boundaries for skimmer areas. No definition of "skimmer" in rule. not evaluated	Implemented by proclamation authority or rule change. Development of criteria for designating skimmer-only areas needed. Extensive mapping of boundaries by seasons needed. not evaluated	Potential to decrease impact from other trawls. not evaluated
3. Designate channel nets, pound nets, and cast nets areas/seasons	Likely decrease in the amount of bycatch in specific areas and during specific seasons. Areas where gear can be used are limited due to physical characteristics of area. +	May reduce flexibility in landings and value of landings in specific areas and during specific seasons (channel nets cannot be set Dec-March 1). Loss of marketable bycatch. Cost of additional gear. Reallocation of resource to another user group. Based on user group, could be a + or a - . Economic impact-everyone gets a piece of the pie. +/-	May increase conflict within each fishery. May reduce the seasonal availability of local shrimp. Increased gear may restrict waterway. Pound nets eliminate areas other gear can be fished. -	May increase conflict among commercial fishermen. Effort shifts may impact other fisheries in same area. Potential to increase other fisheries' catches of adult bycatch species. not evaluated	Need determination of enforceable boundaries for each gear and area. Pound nets must be permitted and have public comment period. not evaluated	Development of designation criteria needed. Extensive mapping of boundaries by gear and seasons needed. not evaluated	Potential to decrease impact from other trawls. not evaluated

Advisory Committee Evaluation of Skimmers and Other Gears for Shrimping in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
4.Allow limited quantities of shrimp to be harvested with cast nets in closed areas except for nursery areas Consider 4 quarts as an option	Unlikely to reduce bycatch because of very low bycatch in cast nets. <p align="center">+</p>	Possible additional source of income as a bait fishery. Expands the ability to cast net to more consumers. <p align="center">+</p>	May increase animosity (-). May open up areas for recreational and commercial users (+). <p align="center">+/-</p>	May increase conflict between recreational and commercial fishermen. Encourages increased disturbance in sensitive areas (e.g., SAV). <p align="center">not evaluated</p>	Need determination of enforceable boundaries for cast net areas. Increased enforcement for harvest limits. <p align="center">not evaluated</p>	Implemented by proclamation authority and rule change. Development of criteria for designating cast net only areas needed. Extensive mapping of boundaries by gear and seasons needed. <p align="center">not evaluated</p>	
5.Eliminate Recreational Commercial Gear License otter trawls Added by AC but not evaluated							

DMF Evaluation of Skimmers and Other Gears for Shrimping in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery +/-	Will not create shifts in effort to other fisheries. Maintains present market value of fishery. +	Allows flexibility of use of gears in the fishery. +	Commercial and recreational fishing will continue with no changes in gear use or conflict +/-	Same level of enforcement +	Continued proclamation authority. No rule change needed. +	Allows for further characterization and bycatch reduction studies to fill data gaps prior to new regulations. +
2. Designate skimmer trawl areas/seasons	Likely decrease in the amount of bycatch in specific areas and during specific seasons. Areas where gear can be used are limited due to physical characteristics of area. +	May reduce flexibility, landings and value of landings in specific areas and during specific seasons. Loss of marketable bycatch. May see otter trawl effort shift to other areas. Cost of re-rigging. -	Likely to increase conflict between skimmer and otter trawlers. May reduce the seasonal availability of local brown and pink shrimp. -	Likely to increase conflict among commercial fishermen. Effort shifts may impact other fisheries in same area. Potential to increase other fisheries' catches of adult bycatch species. -	Need determination of enforceable boundaries for skimmer areas. No definition of "skimmer" in rule. -	Implemented by proclamation authority or rule change. Development of criteria for designating skimmer-only areas needed. Extensive mapping of boundaries by seasons needed. -	Potential to decrease habitat impact from otter trawls. +
3. Designate channel nets, pound nets, and cast nets areas/seasons	Likely decrease in the amount of bycatch in specific areas and during specific seasons. Areas where gear can be used are limited due to physical characteristics of area +	May reduce flexibility in landings and value of landings in specific areas and during specific seasons (channel nets cannot be set Dec-March 1). Loss of marketable bycatch. Cost of additional gear. -	May increase conflict within each fishery. May reduce the seasonal availability of local shrimp. Increased gear may restrict waterway. Pound nets eliminate areas other gear can be fished. -	May increase conflict among commercial fishermen. Effort shifts may impact other fisheries in same area. Potential to increase other fisheries' catches of adult bycatch species. -	Need determination of enforceable boundaries for each gear and area. Pound nets must be permitted and have public comment period. -	Development of designation criteria needed. Extensive mapping of boundaries by gear and seasons needed. -	Potential to decrease habitat impact from otter trawls. +
4. Allow limited quantities of shrimp to be harvested with cast nets in closed areas except for nursery areas	Unlikely to reduce bycatch because of very low bycatch in cast nets -	Possible additional source of income as a bait fishery. Potential harvest of small shrimp before reaching marketable size. +/-	May increase animosity with non-cast net fishermen. May open up areas for recreational and commercial users. -	May increase conflict between recreational and commercial fishermen. Encourages increased disturbance in sensitive areas (e.g., SAV). -	Same level of enforcement +	Implemented by proclamation authority and rule change. Development of criteria for designating cast net only areas needed. Extensive mapping of boundaries by gear and seasons needed. -	




Figure 12.7 NCDMF Fishery Management Districts




Figure 12.8 Skimmer Trawl Locations in Pender County.




Figure 12.9 Skimmer Trawl Locations in Onslow County.




Figure 12.10 Skimmer Trawl Locations in New River.




Figure 12.11 Skimmer Trawl Locations in Bogue Sound, White Oak River and Queens Creek.




Figure 12.12 Skimmer Trawl Locations in Newport River, North River, Back Sound and Lower Core Sound.




Figure 12.13 Skimmer Trawl Locations in Neuse River and Upper Core Sound.




Figure 12.14 Skimmer Trawl Locations in Pamlico Sound.

Table 12.11 Number and average pounds, trips, and value in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in state estuarine waters, 1994-2011.

Year	Shrimp Trawl			Skimmer Trawl			Channel Net			Cast Net			Shrimp Pound		
	Pounds	Trips	Value	Pounds	Trips	Value	Pounds	Trips	Value	Pounds	Trips	Value	Pounds	Trips	Value
1994	5,240,153	14,585	\$13,797,757	203,866	1,118	\$382,118	185,585	2,109	\$402,539	236	15	\$566	0	0	\$0
1995	5,729,152	15,482	\$13,759,068	424,181	1,563	\$760,945	272,892	2,279	\$568,260	1,266	36	\$2,645	1,680	13	\$4,226
1996	3,055,860	11,008	\$7,809,425	188,666	1,179	\$439,670	198,653	1,473	\$454,963	637	51	\$1,769	**	**	**
1997	4,911,799	12,702	\$12,958,128	339,056	2,203	\$763,231	191,188	2,088	\$459,963	70	36	\$380	**	**	**
1998	2,019,600	8,297	\$4,473,965	179,387	1,058	\$375,854	181,915	1,864	\$399,726	620	50	\$1,587	0	0	\$0
1999	5,275,158	10,817	\$12,928,539	599,465	2,080	\$899,582	284,257	2,589	\$571,077	4,936	63	\$5,600	0	0	\$0
2000	7,847,702	10,521	\$19,585,614	624,010	2,429	\$1,087,923	260,321	2,168	\$621,181	928	71	\$2,582	0	0	\$0
2001	3,493,218	7,734	\$8,506,491	314,994	1,765	\$497,427	185,277	1,623	\$394,717	289	140	\$2,316	0	0	\$0
2002	7,511,154	10,030	\$14,159,626	831,511	3,565	\$1,136,668	250,656	1,865	\$436,803	386	161	\$5,131	**	**	**
2003	3,179,629	6,682	\$6,011,535	475,582	2,535	\$714,348	255,892	1,697	\$420,083	271	105	\$7,822	0	0	\$0
2004	2,581,743	5,358	\$5,523,421	377,173	2,097	\$529,413	149,933	1,351	\$228,586	142	115	\$2,334	0	0	
2005	1,078,088	2,890	\$2,016,414	176,928	1,101	\$263,381	130,710	865	\$187,292	116	82	\$1,087	**	**	**
2006	2,891,435	3,255	\$5,059,891	686,475	1,344	\$590,720	181,102	897	\$227,972	41	20	\$635	637	9	\$907
2007	7,123,976	4,465	\$13,595,395	586,700	1,556	\$672,596	165,729	954	\$272,177	740	11	\$1,398	**	**	**
2008	6,764,108	4,206	\$13,516,404	365,331	935	\$432,017	253,530	1,101	\$336,822	531	16	\$1,184	**	**	**
2009	4,049,599	3,890	\$6,452,588	181,458	807	\$173,889	180,704	1,084	\$195,984	218	65	\$1,837	**	**	**
2010	4,280,703	3,946	\$7,649,074	284,972	1,095	\$384,020	129,865	1,063	\$182,808	126	37	\$1,321	**	**	**
2011	3,889,637	3,004	\$8,178,854	55,576	327	\$93,420	97,908	531	\$107,651	231	64	\$3,187	**	**	**
Average	4,495,706	7,715	\$9,776,788	383,074	1,598	\$566,512	197,562	1,533	\$359,367	655	63	\$2,410	257	2	\$642
lb/trip	583			240			129			10			105		
value/trip	\$1,267			\$355			\$234			\$38			\$263		

Table 12.12 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Southern District*, 1994-2011.

Year	Otter Trawl		Skimmer Trawl		Channel Net		Cast Net		Shrimp Pound	
	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips
1994	278,655	3,031	1,493	13	66,714	812	194	4	0	0
1995	422,595	3,330	21,812	88	124,951	1,160	943	21	**	**
1996	226,007	2,334	43,398	231	97,579	781	83	33	0	0
1997	315,710	2,683	77,508	370	112,073	1,374	56	33	0	0
1998	261,126	2,032	77,837	368	114,942	1,161	578	46	0	0
1999	274,212	2,132	84,770	558	188,513	1,967	4,252	58	0	0
2000	299,827	1,878	173,429	727	199,071	1,697	905	67	0	0
2001	90,356	1,197	48,532	414	158,568	1,340	125	114	0	0
2002	237,973	1,603	210,461	1,073	188,847	1,461	129	122	**	**
2003	227,572	1,492	123,395	821	156,717	1,132	112	67	0	0
2004	124,393	903	100,636	574	89,094	844	39	39	0	0
2005	108,779	745	45,773	288	64,263	451	84	53	**	**
2006	96,497	599	99,271	264	102,498	383	0	0	**	**
2007	104,801	785	101,744	323	54,615	420	**	**	**	**
2008	105,572	830	54,910	183	73,126	403	323	10	**	**
2009	91,048	945	25,180	135	61,509	421	**	**	**	**
2010	213,305	1,358	123,349	376	71,148	586	62	25	**	**
2011	111,845	909	22,270	145	69,090	282	127	59	**	**

*Cape Fear, Inland IWW, Inland IWW Brunswick, Inland IWW Onslow, Lockwood Folly, Masonboro Sound, New River, Shallotte River, Stump Sound, Topsail Sound

** Confidential

Table 12.13 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Central District*, 1994-2011.

Year	Otter Trawl		Skimmer Trawl		Channel Net		Cast Net		Shrimp Pound	
	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips
1994	902,950	6,989	197,467	1,098	118,507	1,296	42	11	0	0
1995	1,051,793	7,150	399,169	1,456	144,641	1,113	133	9	914	8
1996	744,692	5,939	142,843	920	98,885	651	554	18	**	**
1997	642,743	5,624	251,558	1,756	77,654	710	**	**	**	**
1998	542,580	4,531	101,550	690	66,973	703	**	**	0	0
1999	837,823	4,795	502,839	1,491	95,509	621	654	4	0	0
2000	515,230	3,395	431,102	1,610	61,021	469	**	**	0	0
2001	459,414	3,448	265,594	1,340	26,709	283	162	25	0	0
2002	762,620	3,734	606,769	2,391	61,810	404	257	39	0	0
2003	813,464	3,510	348,207	1,688	99,175	565	160	38	0	0
2004	260,366	1,778	262,269	1,464	60,839	507	103	76	0	0
2005	296,469	1,313	128,569	795	66,319	413	32	29	0	0
2006	213,278	879	556,690	970	78,535	513	37	19	0	0
2007	184,556	774	358,624	1,029	111,114	534	0	0	0	0
2008	344,098	672	253,178	619	180,404	698	**	**	0	0
2009	148,416	763	131,497	608	119,195	663	80	29	0	0
2010	111,363	559	146,517	644	58,693	476	61	11	0	0
2011	35,417	173	15,415	105	28,818	249	15	4	0	0

*Bogue Sound, Core Sound, Newport River, North River/Back Sound, White Oak River

** Confidential

Table 12.14 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Pamlico District*, 1994-2011.

Year	Otter Trawl		Skimmer Trawl		Channel Net		Cast Net		Shrimp Pound	
	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips
1994	177,999	734	3,794	3	0	0	0	0	0	0
1995	152,274	706	3,200	19	801	5	**	**	704	4
1996	136,442	498	2,425	28	2,189	41	0	0	0	0
1997	215,821	966	5,849	57	**	**	0	0	0	0
1998	99,774	443	0	0	0	0	0	0	0	0
1999	285,904	649	8,513	30	**	**	**	**	0	0
2000	280,918	674	17,009	83	**	**	0	0	0	0
2001	51,999	284	867	11	0	0	**	**	0	0
2002	324,068	658	13,807	100	0	0	0	0	0	0
2003	112,105	367	3,980	26	0	0	0	0	0	0
2004	83,030	511	11,425	53	0	0	0	0	0	0
2005	113,982	336	2,586	18	0	0	0	0	0	0
2006	116,879	271	14,545	66	0	0	**	**	0	0
2007	139,692	286	30,795	116	0	0	72	4	0	0
2008	372,697	616	47,833	120	0	0	**	**	0	0
2009	121,901	224	17,254	58	0	0	**	**	0	0
2010	117,346	380	14,771	73	**	**	**	**	0	0
2011	104,863	448	17,191	73	0	0	**	**	0	0

*Bay River, Neuse River, Pamlico River, Pungo River

**Confidential

Table 12.15 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in the Northern District*, 1994-2011.

Year	Otter Trawl		Skimmer Trawl		Channel Net		Cast Net		Shrimp Pound	
	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips
1994	21,251	330	0	0	0	0	0	0	0	0
1995	19,230	179	0	0	0	0	0	0	0	0
1996	14,323	287	0	0	0	0	0	0	0	0
1997	20,339	323	0	0	0	0	0	0	0	0
1998	1,540	30	0	0	0	0	0	0	0	0
1999	5,128	132	0	0	0	0	0	0	0	0
2000	47,281	588	0	0	0	0	0	0	0	0
2001	845	21	0	0	0	0	0	0	0	0
2002	40,557	472	0	0	0	0	0	0	0	0
2003	2,747	43	0	0	0	0	0	0	0	0
2004	13,149	229	0	0	0	0	0	0	0	0
2005	881	28	0	0	0	0	0	0	0	0
2006	3,063	43	0	0	0	0	0	0	0	0
2007	28,998	88	0	0	0	0	**	**	0	0
2008	6,904	82	0	0	0	0	**	**	0	0
2009	10,746	99	**	**	0	0	0	0	0	0
2010	1,488	26	0	0	0	0	0	0	0	0
2011	2,051	22	0	0	0	0	0	0	0	0

*Albemarle Sound, Alligator River, Croatan Sound, Currituck Sound, Pasquotank River, Roanoke Sound

**Confidential

Table 12.16 Number of pounds and trips in shrimp trawls, skimmer trawls, channel nets, cast nets and pound nets in Pamlico Sound, 1994-2011.

Year	Otter Trawl		Skimmer Trawl		Channel Net		Cast Net		Shrimp Pound	
	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips
1994	3,859,298	3,501	1,112	4	**	**	0	0	0	0
1995	4,083,261	4,117	0	0	**	**	44	5	0	0
1996	1,934,396	1,950	0	0	0	0	0	0	0	0
1997	3,717,187	3,106	4,141	20	**	**	**	**	0	0
1998	1,114,581	1,261	0	0	0	0	0	0	0	0
1999	3,872,092	3,109	**	**	0	0	0	0	0	0
2000	6,704,446	3,986	2,470	9	**	**	0	0	0	0
2001	2,890,604	2,784	0	0	0	0	0	0	0	0
2002	6,145,936	3,563	**	**	0	0	0	0	0	0
2003	2,023,741	1,270	0	0	0	0	0	0	0	0
2004	2,100,805	1,937	2,843	6	0	0	0	0	0	0
2005	557,977	468	0	0	**	**	0	0	0	0
2006	2,461,717	1,463	15,970	44	**	**	0	0	**	**
2007	6,665,929	2,532	95,538	88	0	0	300	3	0	0
2008	5,934,836	2,006	9,410	13	0	0	**	**	0	0
2009	3,677,487	1,859	7,514	5	0	0	**	**	0	0
2010	3,837,201	1,623	**	**	0	0	0	0	0	0
2011	3,635,461	1,452	699	4	0	0	**	**	0	0

** Confidential

12.3 RESEARCH RESULTS – THE USE OF TURTLE EXCLUDER DEVICES (TEDS) IN COMMERCIAL SKIMMER TRAWL OPERATIONS

ISSUE

Results from testing the use of a Turtle Excluder Device (TED) in commercial skimmer trawl fisheries.

ORIGINATION

The North Carolina Marine Fisheries Commission voted to amend the Shrimp Fishery Management Plan for the purpose of examining ways to address finfish bycatch in North Carolina shrimp fisheries. While TEDs are used in shrimp otter trawl fisheries to exclude protected species bycatch, they also reduce unwanted finfish bycatch. This paper will present findings of TED testing in commercial skimmer trawl fisheries to date.

BACKGROUND

Skimmer trawl operations consist of two rigid “L” shaped frames attached to each side of the vessel forward of the midline with nets attached along the two sides of the frame (Figure 12.15). The frames are lowered into the water perpendicular to the gunwale of the vessel with the outer portion of the frame, which is affixed with a skid, resting on the sea floor. The lead line of the trawl is attached to the skid on the outer portion of the frame and a bullet weight along the inner portion, which spreads the net horizontally and vertically. A tickler chain shorter than the lead line is attached at the same locations as the lead line. The nets are pushed through the water, and at certain intervals, the “lazylines” or “easy lines”, which are attached just ahead of the tail bags, are retrieved and the catch is dumped on deck for culling while the mouth of the net continues to fish. Frames, bullet weights, and lazylines are all typically retrieved with winches.

Skimmer trawls are used in North Carolina, Florida, Alabama, Mississippi, and Louisiana. Skimmer trawls operate in shallow water bodies (< 12 ft) as the nets hang from frames on each side of the vessel and are pushed through the water column. Skimmer trawls can also operate in greater depths (~20 ft) in some areas in Louisiana, but this is not typical and concentrated in areas with a high tidal volume. In North Carolina, skimmer trawls became prevalent in the early 1990s as technology was transferred from Louisiana fishermen (Hines et al. 1999). Skimmer vessels in North Carolina are typically 30 ft long with crews of one or two fishermen, and operate in estuarine waters in late summer/fall (August – October) when white shrimp (*Litopenaeus setiferus*) are most prevalent. Some fishermen may also target brown shrimp (*Penaeus aztecus*) in the early summer when concentrations are high, but this is not common practice. During the 2011 fishing season, only 327 skimmer trawl trips were reported to the North Carolina Division of Marine Fisheries (NCDMF) trip ticket database. These trips landed about 55,550 lb of shrimp representing less than 2% of the total commercial shrimp landings throughout the state in 2011 (NCDMF Trip Ticket Database 2012).

Turtle Excluder Devices (TEDs) have been required throughout the southeast Atlantic and Gulf of Mexico otter trawl shrimp fisheries since the early 1990s. Skimmer trawls have remained exempt from TED requirements in lieu of tow time restrictions (55 min and 75 min, seasonally), except in Florida, where state management requires the use of approved TEDs in skimmer trawls. However, tow times may often be exceeded, which poses a threat to endangered or threatened species (Scott-Denton et al. 2007). To examine the effectiveness of TED use in skimmer trawl fisheries, the National Marine Fisheries Service (NMFS) – Harvesting Systems

Unit has been conducting comparative testing in skimmer trawls since 2008 in MS, AL, LA, and NC. The purpose of this research has been to test and develop a TED configuration that will minimize shrimp loss and function effectively for commercial fishermen should TED use in skimmer trawls be made a requirement.

Prior to the third year of testing, a mass sea turtle stranding event occurred along the MS coastline in the late spring 2010, which prompted a draft emergency rule to require TEDs in skimmer trawls for the southeastern Atlantic. However, the rule was not enacted because of the Deepwater Horizon oil spill in the Gulf of Mexico and the subsequent closure of commercial fishing activities in this area. Coincidentally, the commercial shrimp fishery had opened in the area adjacent to the strandings just prior to the stranding event.

On June 24, 2011, NMFS published a notice of intent to prepare an Environmental Impact Statement and hold scoping meetings to address incidental bycatch and mortality of sea turtles in the southeastern shrimp fishery (76 FR 37050). One management option presented at the scoping meetings was the implementation of TEDs in commercial skimmer trawls. Since that time, public hearings have been held throughout the southeast and Gulf of Mexico to obtain public comments on proposed skimmer trawl TED regulations.

During the summer and fall 2012, NMFS obtained multiple observations aboard commercial skimmer trawl vessels in the Gulf of Mexico. Sea turtle interactions ($n = 24$) were observed during many of these trips, and it was determined that 58% of these captures were small individuals that could potentially pass through the approved maximum 4 inch bar spacing on TED grids. Due to the size of the sea turtles captured and frequency of interactions, the proposed rule to require TEDs in skimmer trawl operations was withdrawn in November 2012. Further characterization work throughout the Gulf of Mexico and North Carolina is scheduled for 2013. In addition, the Harvesting Systems Unit is currently seeking funding to conduct reduced bar spacing TED testing in skimmer trawls.

Methodology

Skimmer trawl TED testing was conducted aboard contracted commercial fishing vessels with a TED installed on one side of the vessel in a twin trawl operation while the other side has served as a control net. Tows were limited to a 55 minute maximum for all testing. Prior to TED testing, several tows were conducted to assess potential side bias and conduct any necessary gear tuning. To reduce side bias, TEDs were switched between each side of the vessel on a daily basis.

NMFS Observers (Galveston, TX) manned and recorded data on all trips. For each tow, observers recorded the total catch and total shrimp weight for both the port and starboard nets. Sample baskets were selected from each trawl and examined for species composition and weights. Weights and counts of all marketable shrimp from the sample basket were recorded. The remainder of the sub-sample was separated and weighed by species group: finfish, non-shrimp crustaceans, invertebrates other than crustaceans, and debris. Other select species (e.g., skates, rays, sharks) were also separated, counted and weighed.

The catch was analyzed by catch categories of total catch, shrimp catch, and bycatch to determine reduction rates as measured by weight of catch (kg) with the use of a TED. Reduction rates were also calculated for some select species and species groups of finfish (teleost fish), crustaceans (non-shrimp), invertebrates, and debris when sample sizes of each

group were sufficient. It is important to note that finfish species were not sampled individually for these studies.

The objectives of these studies were to: quantify the difference in shrimp catch associated with TED use in skimmer trawls; quantify reductions in bycatch (species groups) associated with TED use in skimmer trawls; identify handling problems or specialized handling techniques required when using TEDs in skimmer trawls; and determine optimal TED configurations by area.

Research Results

2008 – Mississippi and Alabama Skimmer Trawl TED Testing

In 2008, two vessels conducted testing in MS and AL (Figure 12.16). The two TED configurations tested in 2008 consisted of a large (50”) and mid-size (40”) grid. Both TEDs were installed in a top-opening configuration and were fitted with double-cover flaps. Total catch was significantly reduced with means ranging from a 14% to 51% reduction on both vessels (Table 12.17). Bycatch (all species groups combined) was reduced significantly and ranged from 16% to 55%. Shrimp reductions ranged from 4% to 8% (Table 12.17).

Significant reductions with the use of a TED were observed on both vessels in 2008 for teleost fish ranging from 10% to 47% (Table 12.18). Rays were also reduced significantly in this testing on both vessels with ray reduction rates ranging from 80% to 84%. Crustaceans and invertebrate (non-shrimp) reductions were not found to be significantly reduced on either vessel (Table 12.18).

2009 - Mississippi and Alabama Skimmer Trawl TED Testing

In 2009, testing was also conducted in MS and AL with two contracted commercial vessels (Figure 12.17). Both TED configurations used during this study were top-opening, mid-sized (40”) grid TEDs with double cover escape openings. Percent reductions were only significant on one vessel for total catch and bycatch categories (Table 12.19). Mean total catch reductions ranged from 3% to 19%, while bycatch was reduced on average from 3% to 25% (Table 12.19). Shrimp reductions ranged from 1% to 3%, but these were not statistically significant values (Table 12.19).

Species group reductions (crustaceans, invertebrates, teleost fish, and rays) were not found to be significant barring the teleost fish group on one vessel (Table 12.20), where teleost fish were significantly reduced by a mean of 32% (Table 12.20). Reductions in crustaceans, invertebrates, and rays were not found to be significant with the two vessels tested in 2009 (Table 12.20).

2010 - North Carolina Skimmer Trawl TED Testing

Six contracted skimmer trawl vessels from three separate areas of North Carolina estuaries were contracted to conduct comparative TED testing (Figure 12.18). Four different TED configurations were tested during this portion of the study; top-opening double cover (TODC), bottom-opening double cover (BODC), top-opening single cover (TOSC) and bottom-opening single cover (BOSC). All inshore single flap openings stretched to 44 inches and all grids were “Super Shooter” style TEDs.

All six vessels in the 2010 skimmer TED testing in NC showed significant reductions in the total catch, and bycatch groups (Table 12.21). Reductions in total catch for TED equipped nets ranged from a mean of 18% to more than 32% (Table 12.21). Bycatch reductions ranged from 23% to 43%. Statistically significant shrimp reductions were observed for TED equipped trawls on two vessels and ranged from a mean of 9% to 11% on these two vessels. Average shrimp loss was less on the other four vessels ranging from an increase of 1% to a 7% shrimp loss, but these reductions were not significant (Table 12.21).

Four species groups (crustaceans, invertebrates, teleost fish, and rays) were analyzed for reduction rates in the 2010 NC study (Table 12.22). Significant reductions in teleost fish and rays were generally observed on all six vessels. Teleost fish reductions with the use of TEDs ranged from a mean of 10% to more than 27% (Table 12.22). Significant reduction in rays by weight ranged from a mean of 55% to 98% (Table 12.22). Crustacean and invertebrate reductions were more variable on the six vessels in the 2010 study relative to teleost fish and ray groups (Table 12.22).

2011 – North Carolina Skimmer Trawl TED Testing

Three commercial skimmer trawl vessels were contracted to conduct the 2011 comparative TED and usability testing on traditional fishing grounds in the inshore waters of North Carolina (Figure 12.19). Testing in NC during 2011 used TEDs on each side of the skimmer trawl operation with one TED serving as a control. This was done only in 2011 to address fishermen concerns over potential drag that a TED on one side of the vessel may be creating, and was also designed to potentially identify an optimal TED configuration for the fishery in this area. In the 2010 NC study, the TODC TED configuration showed an approximate 11% shrimp loss, but these results were statistically significant and this TED was chosen as the control for the 2011 study. The 2011 testing in NC compared the TODC (control) to three other experimental TED configurations; BODC, TOSC, and BOSC.

Total catch, bycatch, and shrimp reduction rates for the three TEDs tested in the 2011 NC study (BOSC, TOSC, and BODC) relative to the control TED (TODC) showed varied results (Table 12.23). Increases in total catch, bycatch and shrimp were observed with the use of both the BOSC and TOSC TED configuration. Total catch was increased by 9% and 11% with the BOSC and TOSC, respectively (Table 12.23). Bycatch rates ranged from an increase of 6% to 13% with the use of the BOSC and TOSC, respectively. Shrimp catch rates ranged from an increase of nearly 26% with the use of the BOSC, while an increase of less than 1% was observed with the use of the TOSC (TOSC). For these two TED configurations, reduction rates of total catch, bycatch and shrimp were only significant for shrimp ($p = 0.0007$) with the BOSC TED. This significant increase in shrimp may indicate that the bottom opening single cover flap was not functioning properly on this vessel. This increased catch may also have been further confounded by the relatively low catch numbers per tow (mean ~ 2.0 kg/tow) observed throughout testing. For the BODC, reductions rates relative to the control TED in both total catch (5%) and bycatch (8%) were observed. Shrimp catch rates were increased by 2% with the use of the BODC (Table 12.23). For total catch, bycatch, and shrimp groups, reductions rates were all non-significant barring the shrimp increases observed with the use of the BOSC.

Four species groups (crustaceans, debris, invertebrates and teleost fish) were analyzed for each of the three experimental TED types in the 2011 NC study (Table 12.24). Mean crustacean reductions ranged from 4% to 16% between the experimental TED configurations and the control TED, and were only significant ($p = 0.04$) with the use of the TOSC configuration. Invertebrates were reduced (mean = 11%, $p = 0.41$) with the use of the BODC,

but increased percentages were observed in the both the BOSC ($p = 0.19$) and TOSC ($p = 0.01$) relative to the control TED. Mean debris was reduced by 43% ($p = 0.06$) and 46% ($p = 0.0009$) for the BOSC and BODC, respectively. With the use of the TOSC, mean debris was increased (21%, $p = 0.30$) relative to the control TED. Teleost fish bycatch was reduced on average by 4% ($p = 0.25$) and 10% ($p = 0.09$) in the BODC and TOSC, respectively. However teleost fish bycatch increased in the BOSC by approximately 8% ($p = 0.09$) relative to the control (Table 12.24).

2012 – North Carolina Skimmer Trawl TED Testing

Two commercial skimmer trawl vessels were contracted to conduct comparative function and usability TED testing in the inshore waters of North Carolina including Queen's Creek, North River, and Core Sound (Figure 12.20 and Figure 12.21). The TED used for this study was a prototype ('D-Shaped'; smaller (33" x 33" grid), and lighter) TED. This configuration was constructed and tested in response to fishermen concerns that larger (oval) grids used in skimmer trawl operations may decrease fuel efficiency and/or have the tendency to drag the bottom in relatively shallow water fishing grounds. Two TED configurations were tested with the prototype D-shaped grid including a TODC and a BODC.

Total catch was reduced up to 26% on average with the use of either the TODC or the BODC TED configuration (Table 12.25, Figure 12.22). Reductions of total catch ranged from 14% (BODC) up to 35% (TODC), and these reductions rates were highly significant ($p = 0.0000$). Bycatch reductions with the use of a TED showed similar results and ranged from a mean of 25% (BODC) to 37% (TODC), (Table 12.25, Figure 12.22). These reductions ranged from a low of 13% to a high of 52%, and all values were significant ($p = 0.0000$).

Shrimp catches were reduced by 7% to 10% on average with the use of this TED. Shrimp reductions ranged from 2% to 17% for the BODC and TODC TED configurations, respectively. Reductions with the use of the BODC TED configuration were lower (7%) compared to the TODC TED configuration which showed a reduction of 10%. All shrimp reductions were significant ($p < 0.05$), (Table 12.25, Figure 12.22).

Four primary species groups (crustaceans, debris, invertebrates, and teleost fish) were analyzed on Vessel 1 - (Top Opening Double Cover) TED configuration, while rays and sharks were also collected during operations on Vessel 2 - (Bottom Opening Double Cover) TED configuration installed in one side (Table 12.26, Figure 12.23).

Mean percent (kg) reductions were observed with both TED types for all species groups barring an increase in the amount of debris with the use of the BODC. Mean reduction percentages were significant ($p < 0.0001$) for crustaceans and teleost fish with the use of both TED types. Teleost fish reductions ranged from 11% to 57% with a mean reduction of 37% with the TODC TED, and a mean reduction of 22% with the BODC.

Significant reductions ($p = 0.047$) in the mean percentage of invertebrates were observed with the use of the TODC TED, while a 100% reduction ($n = 7$, $p = 0.0015$) in rays were observed in Vessel 2 with the BODC (Table 10, Figure 9). Sharks ($n = 17$) were reduced by 77% on average, but these values were not statistically significant (Table 12.26, Figure 12.23).

2011 and 2012 Louisiana Skimmer Trawl TED Testing

Four commercial skimmer vessels were contracted in 2011 and four were contracted in 2012 to conduct comparative TED testing with naked nets. Smaller and lighter D-shaped TEDs were used for all testing. In 2011, TOSC, BOSC, and TODC TED configurations were tested. All single cover TEDs consisted of 44-inch inshore TED openings. Results of 2011 testing indicated that top opening TEDs outperformed bottom opening configurations with regard to shrimp retention (Figure 12.24). Bycatch, composed primarily of floating vegetation, caused severe clogging problems in bottom opening TEDs, while top opening TEDs readily excluded the vegetation. Clogging of bottom opening TEDs resulted in a significant shrimp loss that averaged greater than 21% (Figure 12.24). In contrast, use of the TOSC configuration resulted in a non-significant 1.7% average shrimp gain over a naked net. The TODC performed even better resulting in a non-significant 4.8% gain in shrimp catch (Figure 12.24). Average bycatch reduction, which includes vegetative debris, for the three configurations ranged from an average of 13.8% for the TODC TED to 40.9% for the BOSC TED (Figure 12.24). One green sea turtle was captured in a naked net during testing in 2011 and was released alive.

During 2012, only TODC TED configurations were tested against naked nets. Data are currently being analyzed but preliminary results indicate results similar to those observed during 2011 testing. Three Kemp's ridley sea turtles were captured in naked nets during 2012 testing and all were released alive.

Skimmer Trawl Characterizations – Louisiana

In 2004 and 2005, NMFS initiated observer coverage on a voluntary basis in commercial skimmer trawl operations throughout the coastal waters of Louisiana for the purpose of estimating target and bycatch rates by area and season (Scott-Denton et al., 2007). A total of 307 tows were sampled in this characterization study with vessels ($n = 3$) operating in 1.3 fathoms of water on average. Extrapolated catch estimates were dominated by kept penaeid shrimp (66%), while fish species represented 19%; discarded penaeid shrimp comprised approximately 6% of the catches, and debris and crustaceans combined represented about 10% of the catches. None of the vessels observed were equipped with Bycatch Reduction Devices (BRDs) or TEDs. No sea turtles were observed during these observations in 2004 and 2005.

Further observations in the skimmer trawl fishery were obtained in 2012 as mandatory observer coverage was established in Louisiana due to concern for potential interactions with threatened or endangered species. The objectives of this characterization were to document interactions with protected or endangered species and to quantify target and bycatch species (Pulver et al. 2012). A total of 796 tows were sampled during this time with vessels ($n = 26$) operating in 1.6 fathom depths on average. Extrapolated catch estimates from 274 nets, were dominated by finfish (47%), while penaeid shrimp represented 45% of the catch. Crustaceans, debris, invertebrates and discarded shrimp comprised approximately 10% of the catch. About 47% of the vessels observed in this study had BRDs installed in their nets, while 5% of the boats were equipped with TEDs. Twenty-four sea turtles were captured on these trips and all released alive.

I. AUTHORITY

§ 113-134. Rules.

§ 113-182. Regulation of fishing and fisheries.

§ 143B-289.2. Marine Fisheries Commission – powers and duties.

DISCUSSION

Comparative TED testing has been conducted in MS, AL, NC and LA between 2008 and 2012 (Price and Gearhart, 2011; Price and Gearhart *In press*). These studies assessed the functionality of TEDs through comparisons between nets with TEDs and naked nets (no TED). Results from the 2008 to 2010 studies indicated that standard 40" x 33" bent bar grid (super-shooter-style) TEDs perform well in shallow water skimmer trawl operations with minimal shrimp loss (~5%) and significant reductions in total bycatch (~25%). In addition, industry feedback identified several usability deficiencies that were addressed through operational and rigging solutions. Some of the problems identified were: TED twisting, TED chaffing, and bag lifting problems. Vessels used for testing all had slightly different rigging configurations, which is common throughout this diverse fishery. This required slightly different solutions for each problem identified depending on the vessel configuration. However, all of the problems that arose were solved through either adding floatation or chaffing gear or adjusting lifting points on the tailbag. Even though the TEDs worked well, a common request among all users was a request for a smaller, lighter frame. This prompted testing of a smaller D-shaped configuration during the following seasons.

In 2011, and 2012 testing was conducted in NC and LA with the use of a prototype 'D-shaped' TED. This TED was a smaller (33" x 33" grid), and lighter TED than previously tested. This design was much lighter than traditional oval grids and enabled fishing in shallow water with minimal chaffing due to the flat bottom of the D-shaped grid vs. the rounded bottom of the oval grids. The TED was also easier to handle aboard smaller vessels which are found throughout the industry. The TED also performed well during the initial 2011 LA trials with fishermen providing positive feedback about the ease of use over the oval TEDs.

In addition to usability improvements, the D-shaped TEDs also performed better than the oval TEDs in many cases. During LA testing, slight increases in shrimp catch were observed for top opening TEDs. However, bottom opening TEDs in LA performed poorly due to the composition of the bycatch which included lots of floating vegetative debris. This was not the case in NC, where bottom opening TEDs outperformed top opening TEDs. These results illustrate how TED performance can vary widely from location to location depending on local conditions and catch composition. What works well in one location may perform poorly in another, which shows how important it is to conduct testing under differing conditions to identify the best possible TED configurations for each area and situation.

Following 2012 testing, a proposed rule to require TEDs in skimmers was withdrawn, but it is likely that a rule requiring TEDs in these operations is forthcoming in the near future. Currently, NMFS is evaluating reduced bar spacing on TED grids to discern the potential for decreasing smaller (sized) sea turtle interactions or captures. This work will begin with the annual TED testing in Panama City, FL in June 2013. In addition, NMFS is trying to secure funding to conduct TED testing with reduced bar spacing in LA and NC in the late summer and early fall 2013. Collectively, TED testing to date has identified functional TED types and configurations that can work in skimmer trawl operations under various conditions. While some shrimp loss was identified in these studies, further testing is necessary to refine these estimates. In each of

these studies, TED use also resulted in significant reductions of finfish bycatch. Based on testing to date, the use of TEDs in skimmer trawls appears to be a viable management option for reducing sea turtle bycatch without significantly impacting shrimp catch.

II. EVALUATION MATRIX

AC Evaluation of Research Results – The Use of Turtle Excluder Devices (TEDs) in Commercial Skimmer Trawl Operations							
Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
1. <i>Status quo</i>	No change to current bycatch in skimmer trawl fishery. N	No added expense of reconfiguring gear to incorporate a TED. N	Allows skimmer trawl operators option of using a TED. +	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. N	No proclamation or rule change needed. N	Allows for ongoing characterization and TED testing. +
2. Require TEDs in skimmer trawl operations in North Carolina	Potential to reduce protected species and other bycatch in the skimmer trawl fishery. +	Cost increase to acquire and maintain TEDs. Potential shrimp loss with TED use in skimmer trawls. Increased cost with state/federal rule differences. Increase in cost for redesigning rigs. -	Change from historical skimmer fishery. Provides for general public support of increased sea turtle protection. +	Gear change would have no impact between commercial and recreational fishermen. Gear change remains the same for Recreational Commercial Gear License holders N	Require increased training/patrols. -	Will require rule change (03L.0103). State rule could be more restrictive than current federal regulations. -	May precede potential federal rule and require industry to reconfigure gear multiple times. Consider turtle sizes in NC relative to TED bar spacing. -

DMF Evaluation of Research Results – The Use of Turtle Excluder Devices (TEDs) in Commercial Skimmer Trawl Operations							
Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
1. <i>Status quo</i>	No change to current bycatch in skimmer trawl fishery. -	No added expense of reconfiguring gear to incorporate a TED. +	Allows skimmer trawl operators option of using a TED. +	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. N	No proclamation or rule change needed. N	Allows for ongoing characterization and TED testing. +
2. Require TEDs in skimmer trawl operations in North Carolina	Potential to reduce protected species and other bycatch in the skimmer trawl fishery. +	Cost increase to acquire and maintain TEDs. Potential shrimp loss with TED use in skimmer trawls. Increased cost with state/federal rule differences. -	Change from historical skimmer fishery (-). Provides for general public support of increased sea turtle protection (+). +/-	Gear change would have no impact between commercial and recreational fishermen. N	Require increased training/patrols. -	Will require rule change (03L.0103). State rule could be more restrictive than current federal regulations. N	May precede potential federal rule and require industry to reconfigure gear multiple times. -

Table 12.17 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch (bycatch + shrimp catch), bycatch, and shrimp catch (kg) by vessel for 2008 skimmer trawl TED testing conducted in MS and AL.

Vessel	TED Type	Species Group	Control Net			Exp Net (TED)			Reduction (wt.)			p-value
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	Top Opening	Total Catch	31	194.02	59.93	31	166.04	57.22	8.22	14.42	20.63	< 0.0001
	Double Cover	Bycatch	31	174.94	57.89	31	147.82	55.81	8.65	15.50	22.36	< 0.0001
		Shrimp	32	19.14	8.34	32	18.33	7.49	- 0.11	4.24	8.59	0.0200
2	Top Opening	Total Catch	28	87.20	41.42	28	42.76	17.15	35.30	50.96	66.61	< 0.0001
	Double Cover	Bycatch	28	79.56	41.92	28	35.76	17.49	37.49	55.05	72.61	< 0.0001
		Shrimp	30	7.81	2.13	30	7.15	2.07	4.29	8.37	12.45	0.0001

Table 12.18 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, teleost fish, and rays) (kg) by vessel for 2008 skimmer trawl TED testing conducted in MS and AL.

Vessel	TED Type	Species Group	Control Net			Exp. Net (TED)			Reduction (Wt.)			p-value
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	Top Opening Double Cover	Crustac	4	0.08	0.12	4	0.14	0.17	-919.26	- 80.65	757.97	0.339
		Inverte	31	1.09	1.25	31	1.26	1.27	- 64.90	- 15.70	33.50	0.241
		Teleost fish	31	130.83	43.35	31	118.26	52.82	1.46	9.61	17.76	0.017
		Rays	18	27.68	35.31	18	5.34	3.97	17.25	80.70	144.15	0.008
2	Top Opening Double Cover	Crustac	28	0.56	1.64	28	0.19	0.17	- 49.35	65.26	179.88	0.129
		Inverte	28	0.36	0.54	28	0.23	0.20	- 21.10	36.30	93.69	0.109
		Teleost fish	28	49.80	23.35	28	26.33	13.06	32.18	47.13	62.07	< 0.0001
		Rays	27	22.45	27.02	27	3.56	7.90	38.02	84.13	130.25	< 0.0001

Table 12.19 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch (bycatch + shrimp catch), bycatch, and shrimp catch (kg) by vessel for 2009 skimmer trawl TED testing conducted in MS and AL.

Vessel	TED Type	Species Group	Control Net			Exp. Net (TED)			Reduction (wt.)			p-value
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	Top Opening Double Cover	Total Catch	36	17.88	11.15	36	14.52	9.17	8.94	18.78	28.62	< 0.001
		Bycatch	36	12.91	9.64	36	9.60	7.27	11.97	25.68	39.39	< 0.001
		Shrimp	36	4.97	3.38	36	4.93	3.61	- 6.48	0.87	8.22	0.407
2	Top Opening Double Cover	Total Catch	39	26.22	10.91	39	25.47	10.58	- 2.26	2.85	7.96	0.134
		Bycatch	39	20.93	10.93	39	20.35	10.37	- 3.38	2.80	8.97	0.183
		Shrimp	39	5.29	3.61	39	5.12	3.54	- 2.85	3.08	9.02	0.150

Table 12.20 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, teleost fish, and rays) (kg) by vessel for 2009 skimmer trawl TED testing conducted in MS and AL.

Vessel	TED TYPE	Species Group	Control Net			Exp. Net (TED)			Reduction (Wt.)			p-value
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff.	U 95% CI	
1	Top Opening Double Cover	Crustac	35	0.07	0.08	35	0.06	0.07	- 15.97	20.84	57.64	0.143
		Inverte	35	2.72	2.58	35	2.69	2.98	- 15.76	1.25	18.25	0.442
		Teleost fish	35	10.00	8.35	35	6.78	5.32	14.49	32.18	49.87	< 0.001
		Rays
2	Top Opening Double Cover	Crustac	38	0.07	0.06	38	0.06	0.06	- 8.96	18.70	46.36	0.108
		Inverte	38	1.96	3.18	38	1.99	3.18	- 19.74	- 1.13	17.49	0.064
		Teleost fish	39	18.26	9.52	39	17.72	8.81	- 3.11	2.92	8.94	0.166
		Rays	25	0.75	0.95	25	0.95	2.11	-194.92	- 26.59	141.74	0.348

Table 12.21 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch (bycatch + shrimp catch), bycatch, and shrimp catch (kg) by vessel for 2010 skimmer trawl TED testing conducted in NC.

Vessel	TED Type	Species Group	Control Net			Exp. Net (TED)			Reduction (wt.)			p-value
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	Top Opening Dble Cover	Total Catch	64	32.43	24.25	64	21.99	10.81	14.28	32.19	50.10	<0.001
		Bycatch	64	25.36	23.25	64	15.68	9.60	15.92	38.16	60.40	<0.001
		Shrimp	64	7.07	4.07	64	6.31	4.33	4.81	10.75	16.68	<0.001
2	Btm Opening Dble Cover	Total Catch	52	13.55	7.14	52	10.93	5.27	9.21	19.34	29.46	<0.001
		Bycatch	52	6.95	3.81	52	4.91	2.94	15.53	29.39	43.25	<0.001
		Shrimp	52	6.60	5.53	52	6.03	4.35	- 0.51	8.76	18.04	0.021
3	Top Opening Single Cover	Total Catch	60	21.44	14.95	60	17.09	9.51	6.38	20.25	34.13	0.002
		Bycatch	60	13.25	14.02	60	9.44	7.35	6.58	28.74	50.90	0.004
		Shrimp	60	8.33	5.01	60	7.78	4.85	- 2.34	6.52	15.39	0.418
4	Btm Opening Dble Cover	Total Catch	61	17.40	11.27	61	14.12	10.31	9.79	18.82	27.85	<0.001
		Bycatch	60	12.92	10.58	60	9.89	9.32	11.44	23.48	35.51	<0.001
		Shrimp	60	4.57	3.77	60	4.42	3.96	- 3.90	3.19	10.28	0.192
5	Btm Opening Single Cover	Total Catch	64	20.75	11.42	64	17.01	9.06	7.56	18.00	28.45	<0.001
		Bycatch	64	12.81	10.82	64	8.98	7.01	13.00	29.96	46.91	<0.001
		Shrimp	64	7.93	6.02	64	8.04	6.29	- 6.10	- 1.30	3.49	0.291
6	Top Opening Dble Cover	Total Catch	45	13.04	7.69	45	8.86	4.42	17.90	32.05	46.21	<0.001
		Bycatch	45	9.63	5.95	45	5.46	2.54	25.86	43.33	60.79	<0.001
		Shrimp	42	3.66	3.32	42	3.65	3.20	- 5.49	0.22	5.94	0.469

Table 12.22 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, teleost fish, and rays) (kg) by vessel for 2010 skimmer trawl TED testing conducted in NC.

Vessel	TED Type	Species Group	Control Net			Exp. Net (TED)			Reduction (Wt.)					
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff.	U 95% CI	p-value		
1	Top Opening Dble Cover	Crustac	64	0.99	0.58	64	0.84	0.70	-	0.32	15.64	31.61	0.041	
		Inverte	36	0.88	1.38	36	0.67	1.17	-	0.67	23.44	47.56	0.009	
		Teleost fish	64	19.49	11.66	64	14.13	9.37	-	15.01	27.50	40.00	<0.001	
		Rays	33	8.03	27.09	33	0.31	0.75	-	23.49	96.11	215.71	0.056	
2	Btm Opening Dble Cover	Crustac	51	0.52	0.42	51	0.37	0.25	-	4.73	28.60	52.48	0.014	
		Inverte	43	0.74	1.63	43	0.54	1.13	-	23.46	27.49	78.44	0.135	
		Teleost fish	52	4.69	2.69	52	3.52	2.01	-	12.19	25.02	37.85	< 0.001	
		Rays	36	1.18	1.74	36	0.53	0.97	-	2.10	55.00	107.91	0.036	
3	Top Opening Single Flap	Crustac	60	0.78	0.56	60	0.73	0.49	-	12.74	6.06	24.85	0.265	
		Inverte	37	0.11	0.18	37	0.10	0.18	-	67.17	8.81	84.79	0.412	
		Teleost fish	60	8.77	8.21	60	7.88	6.92	-	1.90	10.10	22.10	0.044	
		Rays	43	4.03	10.71	43	0.84	1.06	-	2.64	79.17	160.98	0.030	
4	Btm Opening Dble Cover	Crustac	59	0.61	0.53	59	0.65	0.68	-	35.82	-	5.75	24.31	0.346
		Inverte		0.39	0.51		0.30	0.32	-	24.53	23.71	71.94	0.177	
		Teleost fish	61	9.90	8.73	61	8.12	7.95	-	7.38	17.98	28.58	<0.001	
		Rays	41	1.64	2.26	41	0.99	1.45	-	7.73	39.75	87.24	0.069	
5	Btm Opening Single Flap	Crustac	62	0.66	0.64	62	0.50	0.46	-	5.39	24.64	43.89	0.007	
		Inverte	22	0.08	0.24	22	0.09	0.23	-	110.66	-	23.21	64.24	0.262
		Teleost fish	64	9.22	7.75	64	7.63	5.93	-	5.37	17.28	29.18	0.001	
		Rays	41	2.82	6.30	41	0.90	1.46	-	3.29	68.27	139.84	0.037	
6	Top Opening Dble Cover	Crustac	28	0.04	0.05	28	0.02	0.06	-	45.11	33.66	112.44	0.241	
		Inverte	-	
		Teleost fish	45	6.04	3.23	45	5.36	2.48	-	0.74	11.17	23.07	0.034	
		Rays	29	5.50	5.15	29	0.11	0.41	-	62.38	98.04	133.70	< 0.001	

Table 12.23 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch, bycatch, and shrimp catch (kg) for each experimental TED type in the 2011 NC skimmer trawl TED testing.

Vessel	Exp TED Type *	Species Group	N	Control Net		Exp Net		Reduction (wt.)			p-value
				Mean	SD	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	BOSC	Total Catch	50	10.91	7.27	11.85	6.47	-20.43	- 8.64	3.14	0.0683
		Bycatch	50	9.31	7.01	9.84	6.25	-17.58	- 5.69	6.19	0.1672
		Shrimp	50	1.60	0.87	2.01	1.16	-45.13	-25.81	- 6.50	0.0007
2	TOSC	Total Catch	44	11.36	6.74	12.60	9.95	-40.95	-10.92	19.12	0.2157
		Bycatch	44	9.33	6.20	10.56	9.36	-49.26	-13.22	22.82	0.2099
		Shrimp	44	2.04	1.08	2.04	1.18	- 9.10	- 0.35	8.41	0.4683
3	BODC	Total Catch	45	21.45	8.00	20.31	10.29	- 3.17	5.33	13.82	0.1186
		Bycatch	45	15.88	6.32	14.65	8.51	- 1.63	7.74	17.12	0.0660
		Shrimp	45	5.58	3.41	5.66	3.86	-11.59	- 1.55	8.49	0.3766

* Control TEDs for all three vessels were TODC (Top Opening Double Cover)

Table 12.24 Summary statistics, results of paired t tests, power analyses, percent differences (kg) and corresponding 95% confidence intervals for groups (crustaceans, debris, invertebrates, teleost fish) for each experimental TED type in the 2011 NC skimmer trawl TED testing.

Vessel	Exp TED Type *	Species Group	N	Control Net		Exp Net		Reduction (wt.)			p-value
				Mean	SD	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	BOSC	Crustac	49	1.27	0.91	1.22	0.90	- 9.77	4.32	18.41	0.2741
		Debris	45	0.50	1.31	0.28	0.51	- 21.63	43.01	107.64	0.0637
		Inverts	43	0.27	0.43	0.34	0.39	- 84.72	- 23.81	37.11	0.1924
		Teleost fish	50	7.30	5.69	7.90	5.37	- 20.87	- 8.18	4.52	0.0932
2	TOSC	Crustac	44	1.56	0.92	1.31	0.68	- 1.78	15.66	33.10	0.0442
		Debris	43	2.14	4.24	2.64	5.30	-129.53	- 23.45	82.63	0.3048
		Inverts	43	1.13	1.63	2.62	4.55	-403.13	-132.70	137.73	0.0095
		Teleost fish	44	4.47	3.28	4.01	2.73	- 4.60	10.24	25.09	0.0874
3	BODC	Crustac	45	1.96	0.87	1.75	0.97	- 2.58	10.52	23.63	0.0678
		Debris	43	1.08	1.17	0.58	0.59	16.79	46.19	75.58	0.0009
		Inverts	21	0.09	0.17	0.08	0.11	- 82.53	10.71	103.96	0.4092
		Teleost fish	45	12.66	6.29	12.16	8.35	- 7.14	3.97	15.08	0.2472

* Control TEDs for all three vessels were TODC (Top Opening Double Cover)

Table 12.25 Summary statistics, results of paired t tests, percent differences and corresponding 95% confidence intervals for total catch, bycatch, and shrimp catch (kg) for each experimental TED type in the 2012 NC skimmer trawl TED testing.

Vessel	Exp TED Type *	Species Group	Control Net			Exp. Net (TED)			Reduction (wt.)			p-value
			N	Mean	SD	N	Mean	SD	L 95% CI	% Diff	U 95% CI	
1	TODC	Total Catch	45	17.49	8.92	45	13.01	7.03	16.60	25.59	34.58	0.0000
		Bycatch	45	10.00	8.20	45	6.27	7.08	22.68	37.28	51.89	0.0000
		Shrimp	45	7.49	3.62	45	6.74	2.74	2.60	9.99	17.39	0.0032
2	BODC	Total Catch	56	34.90	16.59	56	27.62	13.33	14.14	20.85	27.56	0.0000
		Bycatch	57	23.34	14.45	57	17.54	9.15	13.41	24.83	36.26	0.0000
		Shrimp	57	10.95	6.53	57	10.17	6.14	1.73	7.08	12.43	0.0045

Table 12.26 Summary statistics, results of paired t tests, power analyses, percent differences (kg) and corresponding 95% confidence intervals for groups (crustaceans, invertebrates, fish, debris, rays and sharks) for each experimental TED type in the 2012 NC skimmer trawl TED testing.

Vessel	Exp TED Type	Species Group	N	Control Net		Experimental Net		Reduction (wt.)			p-value
				Mean	SD	Mean	SD	L 95% CI	% Diff	U 95% CI	
Vessel 1	TODC	Crustac	47	1.05	0.52	0.73	0.50	18.19	30.76	43.33	0.0000
		Debris	47	0.91	0.94	0.71	0.88	- 15.75	22.03	59.81	0.1472
		Inverts	44	1.50	2.50	0.83	1.05	- 5.87	44.78	95.43	0.0465
		Teleost fish	47	6.40	7.62	4.01	6.77	18.07	37.38	56.69	0.0000
Vessel 2	BODC	Crustac	57	1.38	0.91	1.10	0.84	5.63	19.92	34.20	0.0055
		Debris	46	0.06	0.11	0.10	0.22	-233.32	- 68.59	96.13	0.0826
		Inverts	54	0.10	0.13	0.08	0.07	- 6.95	26.50	59.95	0.0640
		Teleost fish	57	21.03	12.08	16.48	9.60	10.99	21.65	32.30	0.0001
		Rays	7	13.10	7.23	0.00	0.00	48.98	100.00	151.02	0.0015
		Sharks	17	1.47	3.95	0.33	0.35	- 61.24	77.32	215.87	0.1318




Figure 12.15 Diagram of typical skimmer trawl operation (Source: Hein and Meier 1995).




Figure 12.16 Skimmer trawl TED testing locations during 2008 in Mississippi and Alabama.




Figure 12.17 Skimmer trawl TED testing locations during 2009 in Mississippi and Alabama.




Figure 12.18 Skimmer trawl TED testing locations in 2010 in North Carolina.




Figure 12.19 Skimmer trawl track lines in the 2011 NC skimmer trawl TED testing.




Figure 12.20 Trawl locations for Vessel 1 (TODC TED configuration) in the 2012 NC Skimmer Trawl TED testing.




Figure 12.21 Trawl locations for Vessel 2 (BODC TED configuration) in the 2012 NC Skimmer Trawl TED testing.




Figure 12.22 Percent difference for total catch, bycatch, and shrimp (kg) for each experimental TED type (TODC and BODC) in the 2012 NC skimmer trawl TED testing.




Figure 12.23 Percent difference for species groups by each experimental TED type (TODC and BODC) in the 2012 NC skimmer trawl TED testing.






Figure 12.24 Percent difference for species groups by each experimental TED type (TOSC, BOSC, and TODC) in the 2011 LA skimmer trawl TED testing.

12.4 CONSIDERATION FOR A COMMERCIAL LIVE BAIT SHRIMP FISHERY IN NORTH CAROLINA

I. ISSUE

The exploration of the option of establishing a commercial live bait shrimp fishery in North Carolina

II. ORIGINATION

MFC Southern Regional Advisory Committee and the public

III. BACKGROUND

Management of the shrimp fishery in North Carolina is based on regional management and shrimp size. Regional management allows flexibility within areas and waterbodies because of regional differences. There are several criteria that are used to determine opening areas to shrimping and shrimp trawling including habitat, economic and social factors, user conflicts, bycatch issues and shrimp size. Shrimp grow at different rates depending on water temperature and salinity. As growth increases, shrimp migrate to deeper and saltier waters, and eventually move out to the ocean. Presently, the division manages shrimp based on count size, or number of shrimp per pound. For example, thirty-count means that for one pound of shrimp, there are 30 shrimp in that pound. Shrimp count size ranges from 80+ count shrimp to 10-15 count shrimp depending on the area.

Because North Carolina shrimp management is based on larger shrimp compared to the small bait size shrimp (60-80 count), waters will close if sampling indicates that there are small shrimp in the area. This then prohibits the harvest of those small shrimp for live bait by recreational and commercial harvest. However, when areas are closed, recreational and commercial fishermen are allowed to harvest two quarts per person with a cast net. Creating a live bait shrimp fishery in North Carolina may provide another economic opportunity for shrimpers and increase the value of smaller shrimp thus improving the overall value of the fishery.

At the Southern Advisory Committee in September 2012, a shrimper requested that committee consider a regulatory process for bait shrimping. The division's management for large shrimp causes the area where he shrimps to close due to the presence of small shrimp which he is targeting. He requested access to small shrimp over the weekends and access to areas that are closed by proclamation because of small shrimp.

Bycatch does occur in the smaller bait trawls but at net bycatch mortality is generally likely low due to short tow times, and culling times. However as temperatures increase, mortality usually increases as well.

IV. AUTHORITY

§ 113-134. Rules.

§ 113-182. Regulation of fishing and fisheries

§ 143B-289.2. Marine Fisheries Commission – powers and duties.

V. DISCUSSION

Live shrimp are popular bait for recreational fishermen targeting spotted sea trout and red drum as well as other popular recreational finfish. Currently, North Carolina does not manage shrimping for bait and fishermen harvesting shrimp as live bait must comply with current rules and proclamations that are in place for shrimp harvested for consumption. The number of pounds of live bait shrimp is recorded in the trip ticket program, as “numbers” and “dozens”. The number of pounds of live bait is low, ranging from 129 lb in 1994 to 2,074 lb in 2008 but is increasing over time along with the number of dealers reporting and the number of trips (Table 12.27). Value for this fishery is high compared to food shrimp with value increasing over time (Table 12.27). Value per pound has been, on average, between \$10.00 and \$15.00 a pound with an increase in 2011 at \$27.00 a pound. There is a steady number of fishermen participating in the fishery with over half the catches made from shrimp trawls (65%) followed by cast nets (12%), skimmer trawls (10%), and channel nets (5%). Seventy-two percent of the live bait shrimp landings come from the Cape Fear River, the Intracoastal Waterway, Stump and Topsail sounds. The division is unable to account for shrimp sold as dead bait because there are no data collected on the disposition of shrimp landings. All other states in the south Atlantic and Gulf manage for shrimp bait harvest.

Table 12.27 Number of Pounds of Live Bait Shrimp (dozens, numbers) 1994 through 2011.

Year	Number of Pounds	Number of Dealers	Number of Trips	Number of Participants	Value
1994	129	5	69	4	\$1,163
1995	204	11	85	8	\$1,834
1996	242	10	118	12	\$3,657
1997	249	8	130	10	\$2,627
1998	175	14	126	16	\$1,908
1999	418	11	60	10	\$1,252
2000	469	12	88	10	\$6,684
2001	266	8	150	11	\$4,338
2002	805	11	222	16	\$12,976
2003	1,027	12	201	17	\$25,758
2004	1,154	10	218	14	\$19,210
2005	921	14	178	15	\$7,843
2006	1,349	13	142	14	\$30,132
2007	909	14	134	14	\$14,009
2008	2,074	11	133	10	\$34,572
2009	1,652	15	249	14	\$22,942
2010	1,710	16	250	14	\$30,994
2011	1,923	17	279	10	\$52,673

South Carolina

A commercial trawler can sell shrimp for both bait and consumption with a land and sell license. Cast netters are restricted to 48 quarts and are limited 12 dozen shrimp dead or live from

December 16 to April 30. Dead Shrimp can only be sold by cast netters if they are caught in legal trawling areas (with all necessary licenses). These regulations were established to prohibit the sale of shrimp caught by shrimp baiting and “deep holing” where fishermen cast net shrimp without bait (Larry DeLancey, personal communication). Bait dealers who are harvest live shrimp to be sold as bait must have a bait dealer license, live bait tanks aboard the harvesting vessel with a compatible aeration system. They may not have any dead shrimp on board. Bait dealers must also be certified as a *bona fide* bait dealer and must have that certification in hand while harvesting live shrimp for bait.

Georgia

Commercial bait shrimping is open year round. Those that fish for shrimp for live bait to be sold and/or engage in the sale of shrimp for live or dead bait must possess a bait dealer license and personal commercial fishing license. Commercial bait fishermen may pull trawl nets up to 20 feet headrope length constructed of mesh smaller than one inch or greater than one and three-eighths inches when stretched. No bycatch reduction devices or turtle excluder devices are required in bait trawls. Harvest limits are no more than 50 quarts of shrimp at any one time with less than 10 percent dead. Fishing at night is not allowed. The Georgia Department of Natural Resources established bait zones in areas of rivers and creeks that are open for bait shrimp fishing. Fishing boats must maintain bait-holding facilities on the vessel.

Florida

Bait shrimping is allowed with roller trawls only with the exception of the Northeast Region where one trawl is allowed with 5/8 inch mesh in the body and 1/2 inch in the cod end. There are also live well requirements with vessels mandated to be equipped with tanks containing a minimum of 16 cubic feet of continuously aerated saline water during harvest and transport.

Alabama

Commercial bait shrimpers are allowed one trawl not to exceed 50 ft headrope length in open shrimp areas and one trawl not to exceed 16 ft headrope length in areas closed to commercial shrimping or in exclusive bait areas. Anyone engaged in taking, catching, transporting, or selling live bait and transporting dead bait must be in possession of a live saltwater bait dealer’s license. Exclusive bait areas are special areas opened each day from 4:00 a.m. to 10:00 p.m. Seasonal bait areas are those areas open to commercial and recreational taking of live bait when adjacent waters are closed. There are live well requirements. There is a maximum tow time of 20 minutes and harvest is limited to two standard shrimp baskets of live or dead shrimp.

Mississippi

Commercial bait shrimping is open year-round. Fishermen are allowed one trawl with no more than 25 ft headrope length, however there are some water bodies that have restrictions down to 16 ft headrope length. There is a tow time restriction of 25 minutes for bait trawls. Size limits are 100 count shrimp or lower and only daytime fishing is allowed. Fishermen can possess no more than 30 lb of dead shrimp. Major bays are closed but live bait may be taken in some bays. Mississippi requires a license for live bait catcher boats; these boats must be equipped with shaded holding tanks with aeration and water circulation.

Louisiana

Bait shrimp may be harvested at any time. During the closed shrimp season, bait shrimp can be harvested with cast nets less than 8.5 feet in radius, hand operated dip nets with a diameter not to exceed 3 feet, bait traps and bait seines less than 30 ft with a maximum mesh size of ¼ inch bar mesh that are manually operated on foot. A special bait dealer's permit is required to take live bait shrimp during the closed season beginning May 1 and between the spring and fall inshore shrimp season.

Texas

Texas requires a bait shrimp boat to be licensed and must operate only under commercial bait shrimp regulations. They may only fish in places authorized for bait shrimp and sell only to a bait shrimp dealer or sportsman. Bait shrimpers can possess only one trawl net with no more than a 54 ft headrope length. Bait shrimping is open year round with a 200 lb limit. From November to August, 50% of the shrimp must be alive and all heads must be attached August through November.

VII. EVALUATION MATRIX

AC Evaluation Matrix for Consideration for a Commercial Live Bait Shrimp Fishery in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. N	No change in market value of bait shrimp. N	Continues inflexibility for fishermen fishing for bait because management is for larger shrimp. -	Supplies hook and line fishermen with bait. N	Same level of enforcement. N	No rule change needed. N	Harvest of 2 quarts of shrimp with a cast nets in closed areas is allowed. Other means of catching shrimp 7 days a week exist. + / N
2. Establish a permitted commercial live bait shrimp fishery with weekend access and access to areas closed by proclamation because of shrimp size.	Will increase the amount of bycatch in the shrimp fishery. Weekend access and areas closed to proclamation due to shrimp size (-) -	May increase economic value of smaller live shrimp used for bait. Possible cost to purchase a permit. Increase revenue to state. +	Will create conflict between food shrimp fishermen and bait shrimp fishermen. Increase in competition. Public may view as a step back in management. -	Supplies hook and line fishermen with bait. Increases accessibility to bait. +	May increase enforcement duties by increasing permit checks and inspections. Not evaluated	Will require the creation of a new permit with general and specific conditions by rule to establish a live bait shrimp fishery. Undermines current management by size in the Shrimp FMP. Not evaluated	Will require establishment of bait shrimping criteria to establish a fishery. Negates the weekend resting period. Not evaluated
3. <u>Establish a permitted commercial live bait shrimp fishery with weekend access.</u> This option added by the AC.	Will increase the amount of bycatch in the shrimp fishery. Not enough bycatch to cause an impact (N). Increase in effort could cause more bycatch (-) N/-	<u>May increase economic value of smaller live shrimp used for bait. Possible cost to purchase a permit. Increase revenue to state.</u> +	<u>Will create conflict between food shrimp fishermen and bait shrimp fishermen. Increase in competition. Public may view as a step back in management.</u> -	<u>Supplies hook and line fishermen with bait. Increases accessibility to bait</u> +	<u>May increase enforcement duties by increasing permit checks and inspections.</u> Not evaluated	<u>Will require the creation of a new permit with general and specific conditions by rule to establish a live bait shrimp fishery. Undermines current management by size in the Shrimp FMP.</u> Not evaluated	<u>Will require establishment of bait shrimping criteria to establish a fishery. Negates the weekend resting period.</u> Not evaluated

DMF Evaluation Matrix for Consideration for a Commercial Live Bait Shrimp Fishery in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. -	No change in value of bait shrimp. N	Continues inflexibility for fishermen fishing for bait because management is for larger shrimp. -	Supplies hook and line fishermen with bait. +	Same level of enforcement. +	No rule change needed. +	Harvest of 2 quarts of shrimp with a cast nets in closed areas is allowed. Other means of catching shrimp 7 days a week exist. +
2. Establish a permitted commercial live bait shrimp fishery with weekend access and access to areas closed by proclamation because of shrimp size.	Will increase the amount of bycatch in the shrimp fishery. -	May increase economic value of smaller live shrimp used for bait. Possible cost to purchase a permit. +/-	Will create conflict between food shrimp fishermen and bait shrimp fishermen. Increase in competition. Public may view as a step back in management. -	Supplies hook and line fishermen with bait. +	May increase enforcement duties by increasing permit checks and inspections. -	Will require the creation of a new permit with general and specific conditions by rule to establish a live bait shrimp fishery. Undermines current management by size in the Shrimp FMP. -	Will require establishment of bait shrimping criteria to establish a fishery. Negates the weekend resting period. -

12.5 GEAR MODIFICATIONS IN NORTH CAROLINA SHRIMP TRAWLS TO REDUCE FINFISH BYCATCH

I. ISSUE

Reduce finfish bycatch in the North Carolina shrimp trawl fishery through gear modifications and Bycatch Reduction Devices (BRDs).

II. ORIGINATION

The public and Division staff (the Shrimp Fisheries Management Plan (FMP)) Plan Development Team (PDT), Advisory Committee (AC), and the North Carolina Marine Fisheries Commission (MFC).

III. BACKGROUND

Bycatch associated with the commercial shrimp trawl fishery remains a controversial and complex issue. There are few studies that attempt to quantify the impact of shrimp trawling on finfish populations primarily because the magnitude of discards is largely unknown. Even with the lack of data necessary to evaluate the impact of shrimp trawl bycatch, a good argument can be made that there is at minimal a resource management issue because finfish bycatch in shrimp trawls is so perceived as a problem by many in the public (Murray et al. 1992).

Policies at both the state and federal level have been adopted as conservation and management measures to minimize bycatch and bycatch mortality and incorporate that goal into management considerations (DMF 2006). At the federal level the gear technology certified by the Council for use in the penaeid shrimp fishery attempts to balance biological, ecological, and economic trades-offs by reducing finfish bycatch while minimizing shrimp loss. Amendment 6 to the South Atlantic Fishery Management Council (SAFMC) Shrimp FMP reviewed the status of the five species of greatest concern in the South Atlantic (weakfish, king mackerel, Spanish mackerel, Atlantic croaker and spot) and concluded that there is no evidence to indicate that the mortality of finfish caused by the shrimp trawl fleet (with TEDs implemented) is having a significant adverse effect on finfish stocks. This practicability analysis concluded that current management measures minimize bycatch and bycatch mortality to the extent practicable in the penaeid shrimp fishery.

A Bycatch Reduction Device (BRD) is any device, trawl modification, or a combination of devices in a shrimp trawl which reduces finfish and other bycatch (NOAA 2008). Extensive research on hundreds of BRDs and gear modifications has been conducted on both the state and federal levels in pursuit of these goals (Price, personal communication).

Summary of State BRD Testing

During the 1980s the NCDMF and National Marine Fisheries Service (NMFS) conducted studies on shrimp retention rates for various Turtle Excluder Devices (TEDs) (1985 – 1986 DMF unpublished data, and 1988 – 1989 NMFS unpublished data), and started work on identifying means to reduce finfish bycatch in the shrimp trawl fishery (Pearce et al. 1988; Holland 1988).

DMF began working both independently, using its own research vessel and other resources and in cooperation with industry to test various BRDs and gear modifications in 1992. Tests have been conducted in multiple waterbodies and seasons to attempt to encompass varying

environmental conditions. The goal of the testing was to find devices that maximized finfish reduction, minimized shrimp loss and met the requirements of Amendments of the Atlantic States Marine Fisheries Commission (ASMFC) Weakfish FMP (1985-1986 DMF unpublished data; McKenna 1993; Pearce et al. 1988; Brown 2010a).

Based on results obtained during development work DMF required all shrimp trawlers working in state waters to equip their nets with functional fish excluders in 1992, becoming the first state to do so. Amendment 2 of the ASMFC Weakfish FMP required all South Atlantic states (NC-FL) to implement management measures to achieve a 40% reduction by number in bycatch of weakfish in the shrimp trawl fisheries by 1996 (ASMFC 1996).

Although BRD testing has continued sporadically in North Carolina, no new devices have been identified that meet the weakfish reduction requirements. Currently the DMF allows five BRDs for use in state waters (SH-3-2012). Both federal and state certified BRDs are summarized in the attachment.

Florida Fish Excluders (FFE) are the primary BRD employed in North Carolina state waters (NCDMF 2006; Brown 2009; and Brown 2010b). The 2006 FMP stated that over 80% of commercial and close to 100% of recreational shrimpers use the FFE. This high rate of use can be attributed to the fact that this is the only device that is certified for use in both NC state waters and the Exclusive Economic Zone (EEZ). However, the NCDMF specifications for the FFE are different than the federal specifications.

Its effectiveness in reducing weakfish and other fish species is a function of the size of the opening and placement. A minimum opening of 5 ½" X 6 ½" is required for the reduction of weakfish at the mandated level (DMF 2006). Placement in the tailbag is a function of the distance the gear is placed from the tailbag tie-off and general location in the net (top, side, or bottom). The distance from the tailbag tie-off is expressed as a ratio: BRD length/tailbag length where BRD length is equal to the distance from the tailbag tie-off to the opening of the FFE, and tailbag length is the length of the tailbag from the tie-off rings to the beginning of the tailbag (excluding any extension). To obtain a 40% value in weakfish reduction this ratio cannot exceed 0.68 (NCDMF 2006). Data collected during the development of FFEs indicated that maximum reduction of weakfish was obtained when the FFE was placed 15 meshes to the side of the tailbag (NCDMF 2006). While no data was collected, several fishermen did some preliminary testing of using two FFEs, but saw no benefit to bycatch reduction as compared to using one FFE (Mckenna 1993).

The "Sea Eagle" Fish Excluder showed 5% reduction in shrimp, 58% reduction in weakfish, 53% reduction in spot, 57% reduction in Atlantic croaker, and 54% overall finfish reduction (NCDMF 2006; Mckenna 1993).

The Large Mesh and Extended Funnel BRDs showed 2% reduction in shrimp by weight, 71% reduction in spot, 63% reduction in Atlantic croaker, 50% reduction in weakfish and an over 55% reduction in finfish (Mckenna 1993).

BRD Certification Procedures (See Attachment)

States are responsible for certifying BRDs for use in state waters. Procedures for certifying BRDs vary from state to state. Georgia is the only state that specifically addresses the use of federally certified BRDs in state waters by reference:

Note: Any BRD certified by NMFS is allowable in Georgia waters. BRDs currently certified for use in federal waters are also described in Amendment 2 to the SAFMC Shrimp Management Plan” (Georgia 2012).

NMFS requires BRDs in shrimp trawl nets working in the federal waters of the Gulf of Mexico and within the Exclusive Economic Zone of the South Atlantic regions (50CFR Part 622 Appendix D).

In 2004 the preferred alternative for the certification of new BRD's in the Final Amendment 6 (2005) to the Fishery Management Plan (FMP) for the Shrimp Fishery of the South Atlantic Region recommended that for a new BRD to be certified, it must be statistically shown that the device can reduce the total weight of finfish by at least 30%. This created an inconsistency between the SAFMC Shrimp FMP and the ASMFC Weakfish FMP. Addendum III to Amendment 4 of the ASMFC weakfish plan changed BRD requirements to be consistent with Amendment 6 of the SAFMC Shrimp FMP by changing previous requirements from the 40% reduction in weakfish by number to 30% by weight for finfish. This allowed more flexible testing of BRDs, and allowed the South Atlantic Council to achieve an ecosystem approach in fisheries management. This new requirement allows the potential for new BRDs to be developed that have greater bycatch reduction rates than those currently in use.

Gear Modifications

One available management measure to reduce bycatch are gear modification requirements. Research should be directed at webbing/net design for bycatch reduction in trawls (NOAA 2006). The control of net selectivity is a preferred management tool in lieu of other more stringent regulations such as temporal and spatial closures, quotas, or limited entry. Most fishing gears are designed to provide some degree of passive discarding. To understand gear modifications that could potentially reduce unwanted bycatch, one must first understand the gear (Figure 12.25). Many net parameters have been tested to increase net performance (Harrington et al. 1988) and some modifications to these same parameters may provide bycatch reduction.

Potential gear modifications requirements that could reduce bycatch include: trawl door modifications (e.g., beam trawls, wing trawl system); net design (e.g., topless trawls, Figure 12.26); modifications to tickler chains or footrope; mesh size, twine size/type; bycatch reduction devices (BRDs), sorting grids (e.g., nordmore grate, vonin flexi-grid); or turtle excluder device (TED) modifications to further reduce bycatch. Gear testing conducted by the DMF in 1986 on the effects of light vs. heavy footrope chains on 20 foot trawls showed that bycatch of flounder, and crabs was higher in a heavily chained net while there was no difference in shrimp catches (NCDMF 2006). Gear modifications that are easy to deploy, reduce bycatch, and maintain shrimp catch are more acceptable to the fishing industry than area or seasonal closures (Murray et al. 1992). Industry involvement in the development of these devices will most likely result in better designs and increased acceptance.

OTTER TRAWL COMPONENTS




Figure 12.25 Diagram of typical otter trawl (NOAA 2008).




Figure 12.26 Topless trawl (Northeast Fisheries Science Center)

Tailbag Mesh Size

Currently, it is unlawful to take shrimp with trawl nets with stretch mesh lengths less than 1 ½ inches in North Carolina (15A NCAC 03L .0103).

Mesh size is often mandated by regulation to prevent the harvest of small sized animals. Trawl minimum mesh size regulations are the principal method used to regulate fishing mortality on fish stocks (Smolowitz 1983). The underlying principle of mesh size regulations is that undersized fish will escape from the tailbag, survive, and become part of the future spawning biomass. Studies on the survival of fish escaping from tailbags (Main and Sangster 1988) support the use of minimum mesh sizes as a means of reducing fishing mortality on juvenile fish. The possibility remains that not all fish that escape from the tailbag survive, resulting in some level of unobserved mortality. This unobserved mortality is a difficult issue for both managers and scientists because if it occurs, the actual reduction in bycatch and mortality is lessened. Furthermore, since gear escapees cannot be counted by conventional fishery observer programs, they cannot be monitored or included in stock assessment calculations. Chopin and Arimoto (1995) suggest that escapee mortality should be considered if gear-based measures are used as a primary management tool.

In 1949, the first studies on methods to reduce juvenile finfish bycatch in Pamlico Sound, North Carolina examined three mesh sizes of tailbags (2, 2 ¼, and 2 ½ inch). Spot were reduced 12%, 43%, and 50% in the 2, 2 ¼, and 2 ½ inch nets, respectively. Reduction rates for Atlantic croaker were 25%, 59%, and 38%. Shrimp reduction rates were 6%, 15%, and 9% in the 2, 2 ¼, and 2 ½ inch nets respectively (Roelofs 1950).

During July 1991, NCDMF conducted some preliminary tests on the culling efficiency of 2 tailbag sizes (2 inch stretch mesh and 1 5/8 inch stretch mesh) vs. a standard 1 ½ inch stretch mesh net. Five tows were made with each gear type and tow time was standardized to 1 hour. The 2 inch stretch mesh net reduced total finfish weight by 37%, spot weight by 46%, Atlantic

croaker by 22%, and shrimp weight increased by 7%. The 1 5/8 inch stretch mesh net fished similar to the 1 1/2 inch stretch mesh net and there was no apparent difference between the catches of the test and control net. However, not enough tows were made with either gear to test for significance.

NCDMF tested 5 experimental otter trawls in the Neuse River and Pamlico Sound, North Carolina from July 2008 to June 2009. One of those tests compared a standard 1 1/2 inch stretch mesh tail bag to a 1 3/4 inch stretch mesh tailbag. A total of 30 tows were made during this test, with tow times standardized to 1 hour. The total catch of shrimp by weight was virtually identical in both nets while total finfish weight was reduced by 32% in the 1 3/4 inch stretch mesh net. Atlantic croaker was reduced by 16% and spot was reduced by 50% by weight (Brown 2010a).

TEDs with reduced bar spacing

Federal regulations require all shrimp otter trawls to be equipped with TEDs and require the vertical bars in the TED to be no more than 4 inches apart (CFR 223.207(a)(4)). NCDMF adopts the federal regulations for TEDs by reference.

Some observations indicate that TEDs with bars spaced less than 4 inches were also successful at excluding other mega-fauna such as sharks, rays, and large fish. Two studies have investigated the effectiveness of using reduced bar spacing in TEDs as a BRD (Broome et al. 2011; Hataway 2010).

Hataway (2010) compared bycatch and shrimp rates of nets equipped with a TED with 2 inch bar spacing vs. a TED with 4 inch bar spacing on a commercial shrimp trawler. The study was conducted in the inshore waters of Mississippi and Louisiana from Horn Island, MS to Breton Sound, LA. The gear was tested in March, April, July, August, September, and October 2010. A total of 65 tows with a mean tow time of 2 hours and 12 minutes were conducted. Results indicate that the net equipped with a 2 inch (bar spacing) TED, reduced shrimp catch by nearly 9%, Atlantic croaker by 33%, and total finfish by nearly 50% (Hataway 2010).

Broome compared nets equipped with a TED with 2 inch bar spacing vs. a TED with 4 inch bar spacing in the near shore waters of North Carolina from Carolina Beach Inlet to Lockwood Folly Inlet. The gear was tested from September to December 2010. A total of 43 tows were conducted. This study concluded that the TED with the 2 inch reduced grid spacing is an effective way to reduce bycatch with minimal shrimp loss (Broome 2011).

T90 Tailbags and Skylight Panels

T90 refers to sections of nets or entire tailbags that are hung on the square, meaning that as the net is hung vertically the meshes will form squares. T90 tailbags and skylight panels use meshes hung on the square as BRDs. The square meshes do not collapse when being towed as do standard tailbags hung on the diamond. The open square meshes reduce drag on the net, and thus increases fuel efficiency and allows low water pressure areas, both of which allow for increased finfish escapement.

In 2000, the NCDMF conducted tests comparing a standard 1 1/2 inch stretch mesh tailbag hung on the diamond to a 1 1/2 inch stretch mesh T90 (hung on the square). The tests showed a

significant reduction (51%) in the catch of young of the year weakfish in the T90 tailbag (NCDMF 2006).

DMF tested 5 experimental otter trawls in the Neuse River and Pamlico Sound, North Carolina from July 2008 to June 2009. Two of those tests compared a standard 1 ½ inch stretch mesh tail bag (hung on the diamond) to a 1 ¾ inch stretch mesh T90 tailbag (hung on the square) and a 2 inch stretch mesh T90 tailbag (hung on the square). A total of 30 tows were made during each test, with tow times standardized to 1 hour. Reductions in total weight of shrimp were 22% in the 1 ¾ inch stretch mesh T90 tailbag and 13% in the 2 inch stretch mesh T90 tailbag, however neither of these reductions were statistically significant. The reductions in total weight of bycatch in both T90 tailbags were statistically significant. Total finfish reduction was 61% and 57% in the 1 ¾ inch stretch mesh T90 tailbag and the 2 inch stretch mesh T90 tailbag, respectively. Atlantic croaker bycatch was reduced by 76% and 69% in the 1 ¾ inch T90 and 2 inch T90 tailbags. Spot was reduced by 77% and 82% in the 1 ¾ inch T90 and 2 inch T90 tailbags. Weakfish bycatch was reduced by 46% and 2% in the 1 ¾ inch T90 and 2 inch T90 tailbags (Brown 2010a).

Industry Involvement

In cooperation with personnel from the North Carolina Sea Grant, an industry gear advisory committee was established in 1989. The advisory committee was comprised of commercial and recreational fishermen, net makers, seafood dealers, and resource managers. The purpose of this committee was to act as consultants throughout the design and testing phase of a gear development project to reduce bycatch in the North Carolina trawl fisheries (Mckenna et al. 1992). The committee suggested two finfish excluding techniques: skylight panels and large mesh tailbags. Skylights were examined in a NCDMF pilot study in the summer of 1990 and two sizes of tailbags were examined during the summer of 1991.

Because of the regional importance of reducing bycatch in the shrimp trawl fishery, government agencies, private industry, and the Sea Grant College Programs, NMFS collaborated in extensive testing of trawl gear to assess optimal design and performance. Fishermen and net makers donated time and materials, NMFS divers were used to assess the gear underwater, and in the field. From this testing, a new class of trawl was developed, the tongue trawl (Harrington et al. 1988).

Industry involvement in the development of TEDs was crucial to their success. Lessons learned in the initial development of TEDs can be applied in the development of BRDs. Concerns of the environmental community were not seriously considered until the concerned groups became polarized. Much of the initial research was done without involvement and input from the industry. This lack of widespread involvement from the commercial industry resulted in fewer ideas for gear innovations and greater opposition to the program. Additionally, this resulted in site-specific problems associated with the gear not being discovered until the program was fully implemented (Murray et al. 1992).

IV. AUTHORITY

§ 113-134. Rules.

§ 113-182. Regulation of fishing and fisheries

§ 143B-289.2. Marine Fisheries Commission – powers and duties.

V. DISCUSSION

BRD Certification Procedures

State

Issues relating to modifying the current NCDMF BRD Certification Procedures, relate primarily to weakfish. In 1991 Amendment 1 to the Weakfish Fishery Management Plan (FMP) was adopted. This amendment recommended that South Atlantic states implement programs to reduce bycatch mortality of weakfish in their shrimp trawl fisheries by 40% by January 1, 1994.

Addendum III to Amendment 4 of the ASMFC weakfish plan was passed to make the weakfish plan consistent with SAFMC Shrimp Amendment 6 with regard to BRDs. Without modification to Weakfish Amendment 4, new BRDs certified for use in federal waters (using the 30% finfish definition) would not be certified for use in state waters. The weakfish plan which requires BRDs in state waters will still dictate how DMF can certify BRDs for use in state waters. North Carolina must abide by SAFMC's and ASMFC's requirements for BRD certification. In short, NMFS certification will have to be obtained for any future BRDs we approve for use in state waters (L. Paramore, personal communication).

Federal

The disconnect between the state and federally certified BRDs causes issues with shrimpers and net makers trying to comply with both sets of rules since many shrimpers fish in both state and federal waters. Allowing all federally certified BRDs to be used in North Carolina state waters would alleviate this.

DMF could by reference adopt for use in state waters any federally certified BRDs. This is done for TEDs. However, NC Marine Patrol does not support regulations that are by reference. It poses issues with officer and fishermen knowledge of the regulations, enforcement and upholding referenced regulations in a court of law (H. Knudsen, personal communication).

If it is decided to allow federally certified BRDs for use in state waters, all approved BRDs and descriptions would need to be included in the proclamation. Before updating the proclamation, DMF will have to verify which BRDs are federally certified and the requirements for use and installation in the net.

Another issue with adopting federally certified BRDs is that it is unlikely any significant reduction in bycatch would occur. The vast majority of shrimpers in North Carolina use the Florida Fish Excluder (Brown 2009; Brown 2010; NCDMF 2006). Most net makers and fishermen would likely just adapt their current FFE to federal regulations. One positive is that there would be little or no cost to do so for fishermen. There are reports of fishermen optimizing FFEs for shrimp retention reducing its effectiveness (NOAA 2004).

Gear Modifications

From the doors to the tailbag, various gear modifications can be made to potentially reduce bycatch. Nets are typically designed to keep finfish from escaping; in this case we want them to escape. Modifications to gear designed to reduce finfish bycatch and retain shrimp must take into consideration the various behaviors and characteristics of bycatch and target species. Differences in the biology and behavior of round fish, flatfish, and the three commercial shrimp

species are often related to environmental influences such as tides, wind, and bottom type. Different net types have been developed to capitalize on the different behaviors of shrimp species. Brown and pink shrimp are typically more active at night and are found in closer association with the bottom than white shrimp. When targeting pink and brown shrimp, fishermen are more concerned with the spread of the net rather than net height. Net height becomes more important when targeting white shrimp (Harrington et al. 1988).

Development of BRDs must be tested in many areas and over several seasons, since there is considerable variation in conditions both spatially and temporally. It is important to understand that the development of BRDs is a long process, and is dependent on a number of factors. The first step is to design and build the necessary modifications to the gear. Test prototypes and make adjustments until satisfied with the gear to be tested. To meet federal requirements of certifying BRDs it often takes 150 tows. That does not include calibration of the test and control nets or any tows made during initial testing. Often after testing a gear modification for bycatch reduction, alterations to the initial design are needed. Averaging three to five tows a day (practicing normal fishing operations); it would take 30 to 50 days of testing. If multiple industry boats were involved, each boat's contribution would be reduced. The boats would also be allowed to sell any marketable catch, further reducing the burden. That burden may be further mitigated on larger boats (four-barrel rigs) which would have three standard nets to the one test net with potential shrimp loss. It is possible to make a significant number of tows and realize the gear is not effective in reducing bycatch or has poor shrimp retention.

There is no one gear design or modification that will work in every situation. What works during the summer brown shrimp fishery may not be effective in the fall white shrimp fishery. The goal of gear researchers is to give the industry additional tools and techniques to use under various real life field situations.

A consideration with any state regulations concerning gear modifications is the limitation on certain environmental rules imposed by North Carolina General Statutes. Both the Department of Environment and Natural Resources (DENR) and the Marine Fisheries Commission (MFC) may not adopt a rule for the protection of the environment or natural resources that imposes a more restrictive standard, limitation, or requirement than those imposed by federal law or rule. There are exceptions to this statute, but currently none of those are met (G.S. 150B-19.3). The division is seeking clarification on the applicability of the statute to this issue.

The various gear modifications for current consideration as bycatch reduction devices discussed below are promising but all are in the early stages of testing. The division cannot recommend any of these be required until sufficient field test have been conducted.

Tailbag Mesh Size

Currently the minimum mesh size of trawl nets used to target shrimp in North Carolina is 1 ½ inch stretch mesh (15A NCAC 03L.0103). Increasing the minimum mesh size could decrease the amount of bycatch caught. Many shrimpers already use 1 ¾ inch stretch mesh (or larger) tailbags. However, there may be some shrimpers who target smaller shrimp due to market demands that would be adversely affected by regulations increasing the minimum mesh size allowed.

An individual tailbag costs approximately \$150-\$200. Shrimpers use between one and four of these per operation, depending on the rig type. Most shrimpers use two different net types throughout the season (two-seamed and tongue nets), and some fishermen may use additional

net types. Shrimpers can change tailbags between net types, but most have dedicated tailbags for each net type. This could lead to some shrimpers needing eight to 12 new tailbags to meet any new regulation of minimum mesh size. The impact of this cost could be minimized by allowing the industry a year or two to implement as gear typically requires replacement every few years.

TEDs with reduced bar spacing

In addition to reducing the incidental takes of sea turtles TEDs can also function as BRDs. Studies such as Hataway's (2010) and Broome's (2011) have shown minimal shrimp loss and significant finfish reduction using TEDs with reduced bar spacing. While the gear has been tested in the inshore waters of Mississippi and Louisiana and the near shore waters of North Carolina, it has not been tested in the inshore waters of North Carolina. NCDMF has applied for a Bycatch Reduction Engineering Program (BREP) grant that will test TEDs with both 2 inch and 3-inch spaced bars against a standard 4 inch spaced bar TED. Results from this study, if awarded will be available in late 2014.

The reduced bar spacing is effective in eliminating large sharks and rays, reducing risks of injury to deck hands. The gear is also effective at reducing jellyfish and is often referred to as "jelly ball shooters". Potential benefits of the reduced bar spaced TEDs include potential fuel savings by reducing unwanted bycatch in the tailbag, shorter culling times resulting from less bycatch, and a higher quality shrimp.

The industry has concerns with the effectiveness of this gear in the inshore waters of North Carolina. The concern relates to the weight of the TEDs causing the net to dig into the bottom (Mikey Daniels, personal communication). This concern could potentially be mitigated through additional floats on the TED or constructing TEDs out of lighter weight materials, but testing of this gear in the inshore waters of North Carolina is needed to determine that. Another concern is that the reduced bar spacing could result in the gear clogging up in areas and times of high grass and/or debris concentrations (Broome et al. 2011). There would likely be a reduction in marketable finfish from this gear.

The cost of a TED is approximately \$350. Shrimpers use between one and four of these per operation, depending on the type of rig per net type. Most shrimpers use two different net types throughout the season (two-seamed and tongue nets), and some fishermen may use additional net types. Shrimpers can change TEDs between net types, but most have dedicated TEDs for each net type. This could lead to some shrimpers needing eight to 12 new TEDs to meet any new regulation of TED bar spacing. The impact of this cost could be minimized by allowing the industry a year or two to implement as gear typically has to be replaced every few years.

A final consideration with any state regulations concerning TEDs is that currently, NCDMF adopts by reference federally certified TEDs and NCDMF's rules concerning TEDs cannot be stricter than the federal rules (G.S. 150B-19.3). Any state modification to the existing TED regulations would require a rule change (15A NCAC 03L.0103 (g)).

T90 Tailbags and Skylight Panels

T90 tailbags as well as other applications of square mesh panels (e.g., skylight panels) have been shown to be effective in reducing finfish bycatch in shrimp trawls and other trawl fisheries (Brown 2010a; Courtney et al. 2007). The meshes in T90 tailbags remain open allowing escapement of unwanted finfish, and are particularly good at allowing small fish to escape

(Walker et al. 2010). The reduction in unwanted bycatch also allows the net to spread more than a full tailbag. The T90 tailbag is less likely to wobble than a standard tailbag, reducing damage to targeted catch and allows the net to fish more efficiently (Figure 12.27). The open meshes allow water to flow through and reduce drag, increasing fuel efficiency (Knuckey et al. 2008). In contrast to other BRDs, the catch accumulates in the square mesh (T90) tailbag allowing small fish and other animals to escape after swimming a much smaller distance (Courtney et al. 2007).




Figure 12.27 Standard trawl (top) and T90 trawl (bottom) (Knuckey et al. 2008).

Due to the way the tailbags are cut, square mesh tailbags produce more waste of webbing and are somewhat more difficult to hang than the traditional tailbag hung on the diamond. T90 tailbags would likely be 50% higher in cost than a traditional tailbag. Initial costs to transition to T90 tailbags could be mitigated by allowing the industry a year or two to come into compliance with any new regulations.

Tests conducted on two variations of this gear in North Carolina showed no statistically significant loss of shrimp (Brown 2010a). The reduction in shrimp that was observed was at least in part a reduction of small shrimp that are often culled anyway. Also, any reduction in shrimp catch could likely be made up due to the efficiency of the gear. There is reduced drag and because the tailbag is not being filled with unwanted bycatch, longer tows or bigger nets are possible.

Another concern with this gear is finfish “gilling” or “marshing” themselves in the open meshes. This was not observed during testing in 2008 and 2009 (Brown 2010a). There is concern from

the industry that T90 tailbags will lose a lot of shrimp catch when the tailbag is very full. With reduced bycatch, the tailbag is not as likely to become full. This can also be mitigated with reduced tow times, which would likely lower mortality of bycatch that is caught. There is some reluctance to use T90 tailbags because some fishermen believe that they are more likely to tear or the knots slip compared to the diamond-mesh netting when under strain (Walker et al. 2010). This emphasizes the need for continued studies, as even the age and wear of gear can affect its effectiveness.

T90 selector panels and T90 lengtheners have also shown potential as gear modifications that can potentially reduce bycatch (Walker et al. 2010). The state and industry should encourage the research and development of this gear modification. There are currently two such modifications being tested by the industry. A T90 ring (Figure 12.28) in two variations is being tested by Mikey Daniels (personal communication).



Figure 12.28 T90 extension constructed by Mikey Daniels.

Industry Involvement (Form an Industry Work Group/Further Research)

Industry involvement in the development of effective BRDs and gear modifications is vital. The fishermen are the ones who best understand how nets work, modifications that may help reduce bycatch while retaining shrimp, and ways to keep the BRDs practical and not cumbersome to use. Industry involvement in the development of effective TEDs was crucial to their success, both in terms of their effectiveness and of their acceptance. When the industry is involved in the development of gear modifications to reduce bycatch, they not only bring their expertise of nets but have ownership of the solution, which aides in the acceptance of new gear regulations. The industry is more likely to get behind new technologies or methodologies if they feel they have had a role and a financial stake in its development (NOAA 2006). Cooperative research programs between investigators and the industry are becoming more common.

Fishery managers should actively seek out industry involvement in identifying, developing, evaluating, and implementing BRDs and modifications to gear. This cooperation aids in information and knowledge transfer within the industry and government organizations. The

development of an Industry Work Group that brings together net makers, fishermen, scientists, and other stakeholders could aid in the development of more effective BRDs and fishing methods or practices. The team would act as consultants throughout the research, design, and testing phases of a gear and fishing method development project to reduce bycatch in the North Carolina shrimp trawl fishery. A similar team has been used by NCDMF in the past (Mckenna et al. 1992). Through such a group, NCDMF and the industry could promote, recognize, and encourage fishermen who make significant contributions to the effort to reduce bycatch.

Beyond the Industry Work Group, industry involvement could be useful in the development and testing of BRDs. Funding is often a limiting factor for gear development programs. The NCDMF has very limited resources to conduct BRD development testing. NCDMF has and will continue to seek outside funding to conduct this type of research, but the need for these outside sources of funding could be mitigated with industry involvement in the process. If industry stakeholders agreed to provide vessels, gear to test (NCDMF does have limited gear that could be used), and crew members to assist NCDMF staff with sampling, the burden on NCDMF's resources would be reduced to personnel and travel cost (which are still limited). The more members of industry involved the fewer burdens on each individual.

VII. EVALUATION MATRIX

AC Evaluation of Gear Modifications in Shrimp Trawls to Reduce Finfish Bycatch in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. -	No change in value of shrimp fishery. N	Allows flexibility of use of BRDs currently certified through the state but not the use of federally certified BRDs in the fishery. -	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. Not evaluated	No rule change required. Not evaluated	Allows for further characterization and bycatch reduction studies prior to new regulations. Not evaluated
2. Update and certify BRDs through the state BRD certification process	May result in decreased bycatch in the shrimp fishery through development of more effective BRDs in the future. No reduction in bycatch in the short term future. +	None in short term. May result in less fuel consumption, less culling time, bigger shrimp and more shrimp in the long term. Increase in gear cost. +/-	No change in short term. Allows flexibility of fishermen use of state certified BRDs but not the use of federally certified BRDs in the fishery. May gain public and industry support. -	Commercial and recreational fishing will continue with no changes in conflict. N	Minor increase in enforcement duties by increasing number/types of BRDs to enforce. Not evaluated	Can be accomplished by proclamation authority. Not evaluated	Is a lengthy process and would tie up division's resources. There are issues with weakfish criterion. Not evaluated
3. Allow federally certified BRDs	Most fishermen would likely adopt federal fisheye and would significantly reduce bycatch. +	May result in less fuel consumption, less culling time if switch to more efficient BRDs. Initial cost for changing BRDs. +/-	Allows flexibility of fishermen to use state and/or federally certified BRDs. May gain public and industry support. +	Commercial and recreational fishing will continue with no changes in conflict. N	Will increase enforcement duties by increasing the numbers/types of BRDs to enforce. Will require training of officer on new federal regulations allowed on fisheye and others. Not evaluated	Can be accomplished by proclamation authority. DMF would have to update proclamation periodically as federal specifications change. Not evaluated	May be issues with ASMFC concerning weakfish. Negates area specific testing by NCDMF to optimize fisheye for weakfish reduction. Not evaluated
4. Increase minimum tailbag mesh size	Potential decrease in bycatch. +	Impacts fishermen targeting small shrimp and bait fishery. Some cost to fishermen, can be mitigated over time. Potential reduction in catch. -	May gain public support. Industry view may be mixed. N	May cause conflict with those currently using smaller mesh. -	Same level of enforcement. Not evaluated	Will require a rule change. (03L. 0103(a)(1)) Not evaluated	There is a need for additional research. Not evaluated

AC Evaluation of Gear Modifications in Shrimp Trawls to Reduce Finfish Bycatch in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
5.Require T90 Tailbags and/or skylight panel	Potential decrease in bycatch. +	Major cost to fishermen. Potential reduction in catch. Need replacing more often -	May gain public support. Industry view may be mixed. N	Commercial and recreational fishing will continue with no changes in conflict. N	Same level of enforcement with some additional training. Not evaluated	Will require a rule change. (03I .0101(3)(n)). Not evaluated	Is not currently federally certified or state certified. There is a need for additional research. Not evaluated
6.Require reduced bar spaced TEDs	Potential decrease in bycatch. +	Some cost to some fishermen based on bar spacing. Potential reduction in catch. Most expensive TED. -	May gain public support. Industry view may be mixed. N	Commercial and recreational fishing will continue with no changes in conflict. If RCGL required to have TEDs, may cause conflict. -	Same level of enforcement with some additional training. Not evaluated	Will require rule change (03L.0103(g)). State rule could be more restrictive than current federal regulations. Not evaluated	There is a need for additional research. Not evaluated
7. Initiate industry testing of BRDs and gear modifications and implement regulations based on findings	Potential decrease in bycatch through development of more effective BRDs and fishing methods in the future. +	Potential impact to fishermen in the long term due to costs associated with modifications and BRD requirements. -	Increased cooperation between stakeholders. Potential to encourage development. +	No initial change. Potential impacts dependent on findings. N	Same level of enforcement with potential of additional training. Not evaluated	Potential for rule changes. Not evaluated	Without a secure source of funding, progress could be minimal. Not evaluated
8.Convene Stakeholder Work Group	Potential decrease in bycatch through development of more effective BRDs and fishing methods in the future. +	Potential impact to fishermen in the long term due to costs associated with modifications and BRD requirements. -	Increased cooperation between stakeholders. Potential to encourage development. +	Commercial and recreational fishing will continue with no changes in conflict. +	No initial change. Could potentially impact future regulations. Not evaluated	Support structure of group not yet identified. Not evaluated	Opportunity to develop ongoing partnerships based on scope of the group. Not evaluated

DMF Evaluation of Gear Modifications in Shrimp Trawls to Reduce Finfish Bycatch in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/Administrative	Other Impacts
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. -	No change in value of shrimp fishery. N	Allows flexibility of use of BRDs currently certified through the state but not the use of federally certified BRDs in the fishery. -	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. N	No rule change required. +	Allows for further characterization and bycatch reduction studies prior to new regulations. +
2. Update and certify BRDs through the state BRD certification process	May result in decreased bycatch in the shrimp fishery through development of more effective BRDs in the future, if fishermen embrace the use of more effective BRDs. No reduction in bycatch in the short term future. +	None in short term. May result in less fuel consumption, less culling time, bigger shrimp and more shrimp in the long term. +	No change in short term. Allows flexibility of fishermen use of state certified BRDs but not the use of federally certified BRDs in the fishery. May gain public and industry support. +	Commercial and recreational fishing will continue with no changes in conflict. N	Minor increase in enforcement duties by increasing number/types of BRDs to enforce. N	Can be accomplished by proclamation authority. +	Is a lengthy process and would tie up division's resources. There are issues with weakfish criterion. -
3. Allow federally certified BRDs	Most fishermen would likely adopt federal fisheye and would not significantly reduce bycatch. May reduce bycatch if fishermen switch to more efficient BRDs. +/-	Little to no cost for fishermen, particularly those staying with fisheye. May result in less fuel consumption, less culling time if switch to more efficient BRDs. Initial cost for changing BRDs. +/-	Allows flexibility of fishermen to use state and/or federally certified BRDs. May gain public and industry support. +	Commercial and recreational fishing will continue with no changes in conflict. N	Will increase enforcement duties by increasing the numbers/types of BRDs to enforce. Will require training of officer on new federal regulations allowed on fisheye and others. -	Can be accomplished by proclamation authority. DMF would have to update proclamation periodically as federal specifications change. N	May be issues with ASMFC concerning weakfish. Negates area specific testing by NCDMF to optimize fisheye for weakfish reduction. -
4. Increase minimum tailbag mesh size	Potential decrease in bycatch. +	Impacts fishermen targeting small shrimp. Some cost to fishermen, can be mitigated over time. Potential reduction in catch. -	May gain public support. Industry view may be mixed. +/-	May cause conflict with those currently using smaller mesh. -	Same level of enforcement. +	Will require a rule change. (03L. 0103(a)(1)) -	There is a need for additional research. -

DMF Evaluation of Gear Modifications in Shrimp Trawls to Reduce Finfish Bycatch in North Carolina

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
5.Require T90 Tailbags and/or skylight panel	Potential decrease in bycatch. +	Some cost to fishermen, can be mitigated over time. Potential reduction in catch. /may result in less fuel consumption +/-	May gain public support. Industry view may be mixed. +/-	Commercial and recreational fishing will continue with no changes in conflict. N	Same level of enforcement with some additional training. N	Will require a rule change. (03L.0101(3)(n)). How is it measured now? -	Is not currently federally certified or state certified. There is a need for additional research. -
6.Require reduced bar spaced TEDs	Potential decrease in bycatch. +	Some cost to fishermen, can be mitigated over time. Potential reduction in catch. -	May gain public support. Industry view may be mixed. +/-	Commercial and recreational fishing will continue with no changes in conflict. N	Same level of enforcement with some additional training. N	Will require rule change (03L.0103(g)). State rule could be more restrictive than current federal regulations. -	There is a need for additional research. -
7. Initiate industry testing of BRDs and gear modifications and implement regulations based on findings	Potential decrease in bycatch through development of more effective BRDs and fishing methods in the future +	Potential impact to fishermen in the long term due to costs associated with modifications and BRD requirements -	Increased cooperation between stakeholders. Potential to encourage development. +	No initial change. Potential impacts dependent on findings. N	Same level of enforcement with potential of additional training. N	Potential for rule changes. SEC Permit structure in place. +/-	Without a secure source of funding, progress could be minimal. -
8.Convvene Bycatch Reduction Workgroup	Potential decrease in bycatch through development of more effective BRDs and fishing methods in the future. +	Potential impact to fishermen in the long term due to costs associated with modifications and BRD requirements. -	Increased cooperation between stakeholders. Potential to encourage development. +	Commercial and recreational fishing will continue with no changes in conflict. N	No initial change. Could potentially impact future regulations. N	Support structure of group not yet identified. -	Opportunity to develop ongoing partnerships based on scope of the group. +

ATTACHMENT

Summary of BRD Certification Procedures

State

The DMF developed BRD certification procedures in the 1990s and Proclamation SH-3-2012 has a special provision that allows persons to test BRD designs.

The following summarizes the current DMF BRD certification procedures (DMF procedures):

The fisherman must submit a design that to DMF for initial evaluation by a gear review panel. This panel is made up of one net maker, a Sea Grant representative, and the DMF gear biologist. If the design is approved, the fisherman will be issued a permit valid for two weeks to test the device. After the two week trial, the fisherman can request a two week extension for additional tuning and testing, request DMF staff to accompany them for final testing, or inform DMF that the device was ineffective.

Certification testing must meet minimum requirements for methodology including net calibration, and the presence of an observer for and data collection. Nets must be switched between sides after every day. At least 30 tows are required for final certification. When the fisherman requests that DMF personnel accompany him/her for final BRD certification then the fisherman is responsible for all costs (except DMF time), and must help in sampling the catches.

For a new BRD to be certified by the DMF it must reduce the bycatch of weakfish by 50% by numbers, reduce total finfish bycatch weight by 45%, and show minimal shrimp loss. For the device to be certified in the EEZ the device must also show a 40% reduction in the number of Spanish mackerel. The data collected will be reviewed by the gear review panel, and their recommendation will be forwarded to the Director of the DMF. If approved by the Director, the recommendations will be forwarded to the ASMFC Weakfish Management Board for their approval. If the ASMFC approves the BRD a new proclamation will be issued allowing the use of the new gear.

Federal

The procedures for federal certification of BRDs are similar to that of DMF, with some notable differences (NOAA 2008).

Fishermen may need to test BRDs for use in state waters with state officials, but for the data collected in such evaluations to be considered by National Oceanic and Atmospheric Administration (NOAA) Fisheries Service for certifications, the operations plan and data collection procedures must meet criteria found in NOAA's Bycatch Reduction Device Testing Manual (NOAA 2008).

A BRD candidate must apply for a Letter of Authorization (LOA), which includes a brief statement of the purpose and goal of activity, diagrams and descriptions of the gear to be tested, and an operations plan describing the scope, duration, methods, and locations of the test. Pre-certification tests can be conducted to assess the preliminary effectiveness of a prototype. No observer is required during pre-certification. For certification testing, testing must meet minimum requirements of methods, calibration of nets, data collection conducted by an observer. A minimum sample size of 30 tows is required for statistical analysis, using

appropriate statistical procedures such as Bayesian analyses (NOAA 2008). However in practice, the standards for certification often take between 100-150 tows (Hataway, pers comm). It is the responsibility of the applicant to ensure that a qualified observer is on board during certification tests.

To be certified for use in the southeastern shrimp fisheries, data collected under a standardized sampling procedure must demonstrate a 30% reduction in finfish biomass (Federal Register 50 CFR Part 622). It should be noted that this criterion was recently changed; previously there was a requirement to reduce red snapper by 50% and Spanish mackerel by 40%, but citing difficulties in obtaining significant results on specific species that may be seen in very low abundances, these criterion were removed (Hataway, pers comm).

The BRD candidate must also meet to the following conditions to be certified:

1. There is at least a 50-percent probability that the true reduction rate of the BRD candidate meets the bycatch reduction criterion.
2. There is no more than a 10-percent probability that the true reduction rate of the BRD candidate is more than 5 percentage points less than the bycatch reduction criterion.

There is also a provisional certification (allowing further tuning and testing) that is effective for 2 years and is based on the following condition:

1. There is at least a 50-percent probability that the true reduction rate is no more than 5 percentage points less than the bycatch reduction criterion.

Certified BRDs

State

The DMF currently allows the following BRDs to be used in state waters (SH-3-2012):

- Florida Fish Excluders (FFE) can be either diamond shaped measuring at least 5 ½ inches X 6 ½ inches or 6 inches X 6 inches (inside measurement) or oval shaped measuring 9 inches by 5 inches and must be positioned no more than 19 meshes from the top centerline of the tailbag and located no more that 65% up from the tailbag tie-off rings
- Eight inch PVC “Sea Eagle” Fish Excluder is a circular excluder constructed of PVC positioned no more than 15 meshes from the top centerline and no more than 38% up from the tailbag tie-off.
- Large Mesh and Extended Funnel BRDs: These devices consist of a funnel of small mesh netting within a cylinder of large mesh netting, held open by one semi-rigid hoop.
 - General ten inch and eight inch large mesh and extended mesh funnel BRD
 - Eight inch and ten inch inshore large mesh and extended funnel BRD
- Large Mesh Funnel Excluder (LMFE): This device consists of a funnel of small mesh netting within a cylinder of larger mesh netting, held open by two semi-rigid hoops.

A single test trawl (try net), channel nets, float nets, fixed nets, and butterfly nets are exempted from required BRDs.

Federal

BRDs currently certified for use in the federal waters of the Gulf of Mexico and within the Exclusive Economic Zone (EEZ) of the South Atlantic regions (50CFR Part 622 Appendix D) include:

- **Extended Funnel:** The extended funnel BRD consists of an extension with large mesh webbing in the center and small mesh webbing on each end held open by a semi-rigid hoop. A funnel of small mesh webbing is placed inside the extension to form a passage for shrimp to the cod end. It also creates an area of reduced water flow to allow for fish escapement through the large mesh. One side of the funnel is extended vertically to form a lead panel and area of reduced water flow. There are two sizes of extended funnels BRDs, a standard size and an inshore size for small trawls.
- **Expanded Mesh.** The expanded mesh BRD is constructed and installed exactly the same as the standard size extended funnel BRD, except that one side of the funnel is not extended to form a lead panel.
- **Fisheye.** The fisheye BRD is a cone-shaped rigid frame constructed from aluminum or steel rod of at least ¼ inch diameter, which is inserted into the cod end to form an escape opening. Minimum escape opening of 5 inches. In gulf EEZ it must be installed top center no further forward than 9 feet from the cod end drawstring (tie-off rings), in South Atlantic EEZ must be installed top center no further forward than 11 feet from the tie-off rings.
- **The Gulf fisheye BRD** is a cone-shaped rigid frame constructed from aluminum or steel rod of at least ¼ inch diameter, which is inserted into the top center of the cod end, and is, offset not more than 15 meshes perpendicular to the top center of the cod end to form an escape opening.
- **The Jones-Davis BRD** is similar to the expanded mesh and the extended funnel BRDs except that the fish escape openings are windows cut around the funnel rather than large mesh sections. In addition, a webbing cone fish deflector is installed behind the funnel.
- **Modified Jones-Davis BRD** is a variation to the alternative funnel construction method of the Jones-Davis BRD except the funnel is assembled by using depth-stretched and heat-set polyethylene webbing instead of the flaps formed from the extension webbing. In addition, no hoops are used to hold the BRD open.
- **Cone Fish Deflector Composite Panel BRD** is a variation to the alternative funnel construction method of the Jones-Davis BRD, except the funnel is assembled by using depth-stretched and heat-set polyethylene webbing with square mesh panels on the inside instead of the flaps formed from the extension webbing. In addition, no hoops are used to hold the BRD open.
- **Square Mesh Panel (SMP) Composite Panel** is a panel of square mesh webbing placed in the top of the cod end to provide finfish escape openings.

12.6 EFFORT MANAGEMENT FOR BYCATCH REDUCTION IN THE NORTH CAROLINA SHRIMP TRAWL FISHERY

I. ISSUE

An examination of using time and seasonal restrictions in the shrimp trawl fishery to reduce bycatch in coastal fishing waters

II. ORIGINATION

The public and Division staff

III. BACKGROUND

A primary component of the management in the 2006 Shrimp Fishery Management Plan is the management of the shrimp fishery by waterbody and by size. Typically, creeks and tributaries of the larger rivers and sounds are closed at the beginning of the year to protect small shrimp and finfish and are opened up during July when they are larger and more valuable. However in the Southern District, areas such as the Intracoastal Waterway (IWW) may remain open until June when small brown shrimp arrive. Special Secondary Nursery Areas (SSNA) open in August and October (Core Sound) when shrimp are larger and the majority of small finfish have migrated out. There are four existing time restrictions used in current shrimp management. There is the weekend closure to shrimp trawling except in the ocean, a nighttime closure to trawling in the ocean off Brunswick County, a nighttime closure to trawling in New River when it is opened between August 16 and November 30, and finally, SSNAs can only be opened to shrimp trawling from August 16 to May 14 each year.

Although shrimp effort varies annually with the abundance of shrimp, overall effort in the estuarine shrimp fishery has steadily declined in recent years in terms of number of trips and number of participants (Table 12.28 and Table 12.29). Otter trawl effort has decreased over time from a high of 15,482 trips in 1995 to a low of 3,004 trips in 2011. Skimmer trawl effort peaked in 2002 at 3,565 trips but has since decreased to 327 trips in 2011. Channel nets show a similar decrease from 2,589 trips in 1999 to a low of 531 trips in 2011. Participation in the estuarine shrimp fishery has decreased in the otter trawl, skimmer trawl and channel net fisheries. Cast net and shrimp pound net participation has been variable over time with no apparent trends. With the exception of one year, shrimp pound information is confidential (less than three dealers reporting). Otter trawl participation was the highest with 888 participants in the fishery in 1995. The fishery has experienced low participation for the last several years with a 66% decline to 301 participants in 2011. Skimmer trawl participation was the highest in 2000 with 180 participants but similar to otter trawls it has dropped 83% to 31 participants in 2011. Channel nets have also shown a steady decline in participation dropping 77% from 176 participants in 1995 to 40 participants in 2011.

Table 12.28 Number and average pounds, trips, and value in otter trawls, skimmer trawls, channel nets, cast nets and pound nets in state estuarine waters, 1994-2011.

Year	Otter Trawl			Skimmer Trawl			Channel Net			Cast Net			Shrimp Pound		
	Pounds	Trips	Value	Pounds	Trips	Value	Pounds	Trips	Value	Pounds	Trips	Value	Pounds	Trips	Value
1994	5,240,153	14,585	\$13,797,757	203,866	1,118	\$382,118	185,585	2,109	\$402,539	236	15	\$566	0	0	\$0
1995	5,729,152	15,482	\$13,759,068	424,181	1,563	\$760,945	272,892	2,279	\$568,260	1,266	36	\$2,645	1,680	13	\$4,226
1996	3,055,860	11,008	\$7,809,425	188,666	1,179	\$439,670	198,653	1,473	\$454,963	637	51	\$1,769	**	**	**
1997	4,911,799	12,702	\$12,958,128	339,056	2,203	\$763,231	191,188	2,088	\$459,963	70	36	\$380	**	**	**
1998	2,019,600	8,297	\$4,473,965	179,387	1,058	\$375,854	181,915	1,864	\$399,726	620	50	\$1,587	0	0	\$0
1999	5,275,158	10,817	\$12,928,539	599,465	2,080	\$899,582	284,257	2,589	\$571,077	4,936	63	\$5,600	0	0	\$0
2000	7,847,702	10,521	\$19,585,614	624,010	2,429	\$1,087,923	260,321	2,168	\$621,181	928	71	\$2,582	0	0	\$0
2001	3,493,218	7,734	\$8,506,491	314,994	1,765	\$497,427	185,277	1,623	\$394,717	289	140	\$2,316	0	0	\$0
2002	7,511,154	10,030	\$14,159,626	831,511	3,565	\$1,136,668	250,656	1,865	\$436,803	386	161	\$5,131	**	**	**
2003	3,179,629	6,682	\$6,011,535	475,582	2,535	\$714,348	255,892	1,697	\$420,083	271	105	\$7,822	0	0	\$0
2004	2,581,743	5,358	\$5,523,421	377,173	2,097	\$529,413	149,933	1,351	\$228,586	142	115	\$2,334	0	0	
2005	1,078,088	2,890	\$2,016,414	176,928	1,101	\$263,381	130,710	865	\$187,292	116	82	\$1,087	**	**	**
2006	2,891,435	3,255	\$5,059,891	686,475	1,344	\$590,720	181,102	897	\$227,972	41	20	\$635	637	9	\$907
2007	7,123,976	4,465	\$13,595,395	586,700	1,556	\$672,596	165,729	954	\$272,177	740	11	\$1,398	**	**	**
2008	6,764,108	4,206	\$13,516,404	365,331	935	\$432,017	253,530	1,101	\$336,822	531	16	\$1,184	**	**	**
2009	4,049,599	3,890	\$6,452,588	181,458	807	\$173,889	180,704	1,084	\$195,984	218	65	\$1,837	**	**	**
2010	4,280,703	3,946	\$7,649,074	284,972	1,095	\$384,020	129,865	1,063	\$182,808	126	37	\$1,321	**	**	**
2011	3,889,637	3,004	\$8,178,854	55,576	327	\$93,420	97,908	531	\$107,651	231	64	\$3,187	**	**	**
Average	4,495,706	7,715	\$9,776,788	383,074	1,598	\$566,512	197,562	1,533	\$359,367	655	63	\$2,410	257	2	\$642

**Confidential

Table 12.29 Number and pounds, trips, and participants in otter trawls, skimmer trawls, channel nets, cast nets in state estuarine waters, 1994-2011.

Year	Otter Trawl			Skimmer Trawl			Channel Net			Cast Net		
	Pounds	Trips	Participants	Pounds	Trips	Participants	Pounds	Trips	Participants	Pounds	Trips	Participants
1994	5,240,153	14,585	845	203,866	1,118	79	185,585	2,109	148	236	15	4
1995	5,729,152	15,482	888	424,181	1,563	128	272,892	2,279	176	1,266	36	14
1996	3,055,860	11,008	705	188,666	1,179	102	198,653	1,473	126	637	51	7
1997	4,911,799	12,702	722	339,056	2,203	143	191,188	2,088	136	70	36	6
1998	2,019,600	8,297	513	179,387	1,058	92	181,915	1,864	113	620	50	8
1999	5,275,158	10,817	667	599,465	2,080	155	284,257	2,589	120	4,936	63	8
2000	7,847,702	10,521	793	624,010	2,429	180	260,321	2,168	122	928	71	12
2001	3,493,218	7,734	553	314,994	1,765	135	185,277	1,623	97	289	140	11
2002	7,511,154	10,030	639	831,511	3,565	158	250,656	1,865	88	386	161	13
2003	3,179,629	6,682	439	475,582	2,535	130	255,892	1,697	86	271	105	12
2004	2,581,743	5,358	421	377,173	2,097	101	149,933	1,351	83	142	115	6
2005	1,078,088	2,890	272	176,928	1,101	72	130,710	865	57	116	82	5
2006	2,891,435	3,255	297	686,475	1,344	87	181,102	897	60	41	20	5
2007	7,123,976	4,465	338	586,700	1,556	84	165,729	954	67	740	11	9
2008	6,764,108	4,206	364	365,331	935	92	253,530	1,101	66	531	16	9
2009	4,049,599	3,890	340	181,458	807	60	180,704	1,084	60	218	65	7
2010	4,280,703	3,946	355	284,972	1,095	64	129,865	1,063	57	126	37	9
2011	3,889,637	3,004	301	55,576	327	31	97,908	531	40	231	64	10

High fuel prices and cheaper imports will most likely continue to keep new entrants out of the shrimp trawl fishery and it is not likely to grow significantly in the foreseeable future.

This same downward trend in effort is also seen in the ocean fishery, where the Exclusive Economic Zone (EEZ) off North Carolina is under the management of the South Atlantic Fisheries Management Council (SAFMC). In recent years, low shrimp prices, rising fuel costs, competition with imported products, and the recent hurricanes in the Gulf of Mexico have all impacted the shrimp fleets. Fishing effort has been reduced by as much as 50% for offshore waters of the Gulf of Mexico (GMFMC 2007) and by about 40% in the South Atlantic (NMFS 2012a). During AC discussion on the following options it will be necessary to distinguish if measures need to include ocean waters. For consistency with SAFMC measures the North Carolina coastal fishing waters less than 3 miles from shore have mirrored federal restrictions for the most part.

In considering ways to reduce bycatch, using time and seasonal restrictions for reducing effort in the shrimp trawl fishery is discussed herein. This issue paper will discuss seasonal closures like closing trawling an additional day of the week or from December or January through May, day/night closures, and restricting tow times for example.

IV. AUTHORITY

§ 113-134. Rules.

§ 113-182. Regulation of fishing and fisheries.

V. DISCUSSION

Seasonal Closure

One possible method of reducing bycatch of non-target species is to further restrict the amount of time that shrimp trawling is allowed. As previously stated, the two primary time restrictions in existence regarding the shrimp fishery are that shrimp trawls are not allowed from 9:00 p.m. on Fridays to 5:00 p.m. on Sundays and SSNAs can only be opened from August 16 to May 14 each year. Closure of additional time periods to shrimp trawling may reduce bycatch by reducing the amount of time that shrimping effort occurs.

An option to consider would be a shrimp trawl closure of several months to occur when shrimp landings were insignificant and juvenile fish were abundant. This would reduce the trawl effort when shrimp landings and value were not optimal.

Although all months could be considered for seasonal closure, as an example, the period of December through May could be closed to shrimp trawling to reduce the catches of juvenile finfish when the target shrimp were not as available. In some years, white shrimp may still be present in December and in warmer springs, pink shrimp may be available in May, but this would be a way to reduce effort. Little shrimp trawl activity occurs in internal coastal waters during December through April. In May, Pamlico, Core and Bogue sounds as well as Newport and North River fishermen begin trawling as pink shrimp that have overwintered make their way to the inlets and the ocean. Closures during June through November are not consistent with maximizing economic value of shrimp since these are the months when this annual crop is available and the most efficient way to harvest them is with otter and skimmer trawls.

Along with the regional differences in shrimp harvest times, there is seasonal variability in the availability of the three different shrimp species. Table 12.30 illustrates the variation in seasonal landings of the three shrimp species and unclassified shrimp by month from 2001 through 2011 (eleven year sum) in inside waters. Table 12.31 depicts average monthly trawl landings and effort in different waterbody groupings. Pink shrimp that have overwintered in sea grass beds may begin to migrate toward the ocean inlets in April and May as water temperatures rise. Extremely cold winters can reduce the pink shrimp crop significantly. The pink shrimp are mostly gone by July and brown shrimp are dominant from July to October. White shrimp appear in the southern part of the state in May or June as adults that have overwintered. They are present in the central waterbodies and Pamlico Sound in late summer through late fall, depending on cooling water temperatures. This seasonal availability means for example, that a closure in April and May would negatively impact the Pamlico, Core and Bogue sounds as well as the Newport and North River area, when pink shrimp are migrating to sea.

Table 12.30 Average pounds of shrimp landed in North Carolina's inside waters per month by species, 2001-2011.

Month	Brown Shrimp		Pink Shrimp		White Shrimp		Unclassified	
	Pounds	Percent	Pounds	Percent	Pounds	Percent	Pounds	Percent
January			16	4.02	345	84.44	47	11.54
February	10	27.94	25	70.89	0	1.16	-	-
March	30	1.51	1,385	70.07	557	28.18	5	0.23
April	146	0.86	13,070	76.64	3,809	22.33	29	0.17
May	4,018	7.91	45,386	89.33	921	1.81	482	0.95
June	168,585	66.01	84,211	32.97	530	0.21	2,083	0.82
July	1,522,408	94.45	39,584	2.46	5,288	0.33	44,664	2.77
August	1,112,668	88.10	1,556	0.12	121,105	9.59	27,650	2.19
September	255,364	40.29	844	0.13	372,954	58.85	4,617	0.73
October	67,734	11.79	2,050	0.36	500,650	87.18	3,854	0.67
November	11,348	4.94	1,723	0.75	215,762	94.00	711	0.31
December	262	1.81	40	0.28	14,098	97.22	101	0.70

Seasons in other states are based on the economic considerations of the availability of species of shrimp (mostly white shrimp). South Carolina allows trawling on their white shrimp crop from May through December in general trawl areas (nearshore ocean) and from September to December 15 downstream of channel net areas.

Georgia's white shrimp season opens as early as May 15 and can run as late as February. Florida, which manages by region, has a closed season from June through October and certain counties close in April and May. Alabama opens and closes its seasons by regulation. Mississippi opens in May or June and closes December 31 north of the WW and closes April 30 south of the IWW. Louisiana has a May through July inshore season and an August through December inshore season. Finally, Texas has a May through July season for early morning shrimping, an August through November season for daylight shrimping and a February through April season for nighttime only shrimping.

If a seasonal closure was implemented, conflicts could increase among trawlers who can switch gear and existing channel netters and shrimp pound operators since the productive sites for this

gear are already taken and additional fishermen moving in would be problematic. A statewide season opening day of June 1 could result in a “grand opening” event in certain waterbodies with many boats vying for limited space along the closure lines.

Additional Day of Week Closure

Adding another day of the week to the present closed trawling period is another time related bycatch reduction measure to consider. Shrimp trawling is closed from 9:00 p.m. on Fridays to 5:00 p.m. on Sundays in internal coastal waters. Friday could be considered as an additional day to be closed. The closure could be from 9:00 p.m. on Thursdays to 5:00 p.m. on Sundays.

The present Friday through Sunday evening closure evolved from a February 1984 petition from fishermen to close Core Sound from 8:00 a.m. Saturday to 6:00 a.m. Monday by proclamation. Although some fishermen and dealers complained that they needed shrimp for the Monday morning market and there was a fear of effort shifting to adjacent open areas, there was some support for a Sunday night closure. A proposal to close from Saturday morning through Monday morning by rule failed. Fishermen continued to request a weekend closure by proclamation and this was tried in July, 1984. Core Sound, North River, South River, Turnagain Bay, Rataan, Cedar, Long and West bays, Newport River and Adams Creek were closed from July 15 through December 31, 1984 and this was continued from that time on in some fashion. In 1993 the weekend closure was adjusted to begin one hour after sunset on Fridays and end one hour before sunset on Sundays. A 1993 effort by the Marine Fisheries Commission to extend the closure through Monday morning failed to go forward. Actual times (9 p.m. and 5:00 p.m.) were implemented in 2004 to avoid confusion with varying times found on sunrise/sunset tables.

Although an additional day added to the weekend closure, be it Friday or Monday, would reduce shrimp trawling effort, it is not possible to quantify the reduction in bycatch. A uniform number of shrimp are not caught each available trawling day so an additional closed day would not reduce bycatch proportionally. Regardless of the days of the week closed, it has been observed that the best catches of shrimp are on the night of the opening after that “rest period”. Johnson (2006) noted that twice as much shrimp were caught early in the five-day trawling week than later in the week in the coastal shrimp trawl fishery in NC, suggesting that time restrictions could further improve the efficiency of the shrimp fishery. An additional weekend closure day would be an option that would reduce effort, however reducing the number of days from five to four does not take into account days already lost to wind and weather, unfavorable tide and moon phases, etc. that may further impact shrimp catches. An additional day added to the weekend closure would reduce trawling effort by Recreational Commercial Gear License holders and part-time fishermen who shrimp trawl mainly around the weekends.

Day/Night Closures

Another way to reduce the amount of time shrimp trawling is allowed and perhaps reduce bycatch is to close areas during the daytime or nighttime hours. The habits of North Carolina’s three shrimp species determine when they are fished for now. In the central part of the state, brown and pink shrimp usually burrow into the substrate during the day and trawling for them usually occurs at night. Occasionally trawling for brown shrimp can occur during the daytime when waters are murky. These trips usually last one night or one day. White shrimp are found up in the water column and fishing for them occurs mainly at night. Larger trawlers in Pamlico Sound with the capacity to store ice usually stay out four or five days and tow day and night. Shrimp from the larger trawlers are usually landed on Thursdays and Fridays.

In 1997, many Sneads Ferry trawler fishermen requested opening the New River to daytime shrimp trawling only. This was not based on any biological information. Many of the local shrimpers preferred to go during the daytime and wanted to keep trawlers from neighboring areas out of there at night. Marine Fisheries Commission Rule 15A NCAC 03J .0208, effective in 1998, makes it unlawful to use trawl nets upstream of the Highway 172 bridge over New River from 9:00 p.m. through 5:00 a.m. when opened by proclamation from August 16 through November 30.

In North Carolina it is unlawful to trawl for shrimp in the Atlantic Ocean off Brunswick County from 9:00 P.M. to 5:00 A.M. each day [15A NCAC 03J .202 (8)]. This management measure was implemented in large part to reduce the bycatch of finfish in this gear. Ingraham (2003) examined this question by conducting a study of shrimp and finfish catch rates (day vs. night) in state waters from Topsail Inlet to Little River Inlet. Data from the study showed that finfish bycatch was higher at night than during the day. Of the nine commercially important finfish species caught, southern flounder, spot, Atlantic croaker, and southern kingfish catch rates were significantly higher at night. The catch of shrimp did not vary significantly between nighttime and daytime trawling, although catches were slightly higher during the day. South Carolina shrimp trawling has been closed at night since the 1970s, but that was enacted to keep North Carolina fishermen from catching brown shrimp at night because South Carolina fishermen wanted to work during the day, not for any biological reason (L. DeLancey. SC DNR, personal communication).

Table 12.31 Average landings and effort by month of all shrimp species by waterbody* from otter and skimmer trawls combined), 2002-2011.

Month	Pamlico Sound		Western Rivers		Central Sounds		Southern		Ocean <3 miles	
	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds	Trips	Pounds
January	1	341	-	-	-	-	-	-	28	35,844
February	0.2	11	-	-	-	-	0.2	19	18	32,622
March	0.4	19	-	-	-	-	14	1,305	10	11,349
April	5	1,579	1	132	1	132	102	10,944	29	5,096
May	19	14,614	5	841	5	841	253	33,904	103	53,249
June	84	116,739	89	34,216	89	34,216	386	93,122	170	138,253
July	611	1,393,375	194	85,330	194	85,330	494	130,019	350	247,899
August	615	1,056,351	122	33,984	122	33,984	529	144,646	278	131,156
September	264	430,894	52	14,612	52	14,612	409	125,427	324	217,701
October	251	496,953	14	6,659	14	6,659	208	45,164	463	342,944
November	88	205,238	2	2,035	2	2,035	51	13,061	294	241,485
December	9	12,334	0.1	3	0.1	3	2	168	67	79,160

*Waterbody Groupings:

Pamlico Sound

Western Rivers include Neuse, Bay, Pamlico and Pungo rivers

Central Sounds include Core and Bogue sounds, Newport, North and White Oak rivers

Southern includes all waters south of White Oak River

Ocean < 3 miles

Tow Times

One more way to consider reduction of effort in relation to time involves restricting tow times. A tow time limit of 45 minutes has been mentioned by the public. Although theoretically and commonsense-wise, it would appear that reducing tow times would reduce bycatch, in reality that does not necessarily occur. Reduced tow times would likely reduce bycatch mortality. Fish aggregations as well as shrimp aggregations are not uniformly distributed and each tow is different depending on depth, tide stage, moon phase bottom type and many other factors. Carothers and Chittendon (1985) found a significant linear relationship between catch and tow duration (i.e., the longer you tow, the more you catch). Their experiments tested tows of 5, 10, 15, 20, 25 and 30 minutes duration.

A tow time requirement would be very difficult to enforce without constant Marine Patrol oversight or costly Vessel Monitoring Systems. Tow times in the ocean were enforced from 1996 through 2005 under a now-expired Incidental Take Permit from National Marine Fisheries Service issued to trawlers from Browns Inlet to Rich Inlet due to the presence of grass (brown algae). This involved constant monitoring and observers and was very difficult to enforce. The timing of tows began when the otter trawl doors were lowered into the water and ended when they exited the water. Skimmer trawl tows could not be timed in that way since they are towed continuously and the tailbags are pulled in and emptied. Additional tows could be made to make up for the "lost effort" of limited tow times. Tow times have been suggested and may be effective in reducing bycatch mortality in individual tows. Recoupment of trawl times could be made up with additional tows.

VII. EVALUATION MATRICES

AC Evaluation of Effort Management for Bycatch Reduction in the North Carolina Shrimp Trawl Fishery

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	
1. <i>Status quo</i>	No change in existing about of bycatch. -	No additional loss or gain in revenue. N	No change in fishing practices. N	Commercial and recreational trawling will continue with no changes in season or conflicts. N	Same level of enforcement. Not evaluated	No proclamation or rule change needed. Not evaluated	
2. Implement seasonal closure (i.e. December or January through May) AC voted to not consider this option. Failed 4-4.	Bycatch from shrimp trawls eliminated during the months of closure. +	Loss of income due to reduced shrimp catch. -	May create effort shifts. May cause potential conflict between user groups. -	Trawlers who switch gears may conflict with present users of that gear. -	May increase enforcement efforts patrolling closed areas. Not evaluated	Implemented by proclamation authority or rule change if preferred. Not evaluated	
3. Add an additional day to the weekend closure in internal coastal waters	Some reduction in bycatch for an additional day per week. +	Trawlers may lose one additional day of income. May recoup, but loss of another day per week may be detrimental with lost trips due to weather, breakdowns, etc. Channel netters and offshore trawlers would favor this as it allows more shrimp to reach them. -	Increased effort to recoup lost day could create conflict between trawlers and other user groups. -	No impact with other fisheries. N	Same amount of enforcement on opening nights. Not evaluated	Rule change required. Not evaluated	

AC Evaluation of Effort Management for Bycatch Reduction in the North Carolina Shrimp Trawl Fishery

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
4. Close shrimp trawling at night in internal coastal waters.	Possible reduction in bycatch in certain areas. +	Potential loss of income due to reduced shrimp catch. Channel netters and offshore trawlers would favor this as it allows more shrimp to reach them. -	Loss of traditional fishery. Potential to increase conflict. -	Effort shifts to other fisheries may occur. -	Night patrol will need to be increased. Not evaluated	Implemented by proclamation authority or rule change if preferred. Not evaluated	Loss or gains could be species specific. Not evaluated
5. Implement a tow time limit in internal coastal waters. AC voted to not consider this option. Passed 6-2.	No impact on bycatch due to variable shrimp and fish distribution and the ability to recoup with additional tows. May reduce bycatch mortality.	No impact on harvest or income. Difficult to quantify due to variability.	Numerous complaints of violations are likely.	No Impact.	Difficult to enforce and time the tows.	Implemented by proclamation authority or rule change if preferred.	

DMF Evaluation of Effort Management for Bycatch Reduction in the North Carolina Shrimp Trawl Fishery

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	
1. <i>Status quo</i>	No change in existing bycatch. -	No additional loss or gain in revenue. N	No change in fishing practices. N	Commercial and recreational trawling will continue with no changes in season or conflicts. N	Same level of enforcement. N	No proclamation or rule change needed. +	.
2. Implement seasonal closure (i.e. December or January through May)	Bycatch from shrimp trawls eliminated during the time of closure. The magnitude of bycatch reduction depends on time selected. +	Loss of income due to reduced shrimp catch. -	May create effort shifts. May cause potential conflict between user groups. -	Trawlers who switch gears may conflict with present users of that gear. -	May increase enforcement efforts patrolling closed areas. -	Implemented by proclamation authority or rule change. -	
3. Add an additional day to the weekend closure in internal coastal waters	Reduces bycatch for an additional day per week. +	Trawlers may lose one additional day of income. May recoup, but loss of another day per week may be detrimental with lost trips due to weather, breakdowns, etc. Channel netters and offshore trawlers would favor this as it allows more shrimp to reach them. +/-	Increased effort to recoup lost day could create conflict between trawlers and other user groups. -	No impact with other fisheries. N	Same amount of enforcement on opening nights. N	Rule change required. -	

DMF Evaluation of Effort Management for Bycatch Reduction in the North Carolina Shrimp Trawl Fishery

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/Administrative	Other Impacts
4. Close shrimp trawling at night in internal coastal waters.	Possible reduction in bycatch in certain areas. Increase of bycatch mortality due to increase in air temperature +/-	Potential loss of income due to reduced shrimp catch. Channel netters and offshore trawlers would favor this as it allows more shrimp to reach them. +/-	Loss of traditional fishery. Potential to increase daytime shrimping conflict. -	Effort shifts to other fisheries may occur. -	Night patrol will need to be increased. -	Implemented by proclamation authority or rule change. -	Loss or gains could be shrimp and bycatch species specific. N
5. Implement a tow time limit in internal coastal waters	May reduce bycatch mortality. Minimal impact on bycatch due to variable shrimp and fish distribution and the ability to recoup with additional tows. +/-	Loss fishing time due to increase in number of haulbacks. This results in lost income. Difficult to quantify due to variability. -	Numerous complaints of violations are likely. Operator frustration due to decrease in efficiency. -	No Impact. N	Difficult to enforce and time the tows. Increase in number of responses to complaints on tow times -	Implemented by proclamation authority or rule change if preferred. -	

12.7 CHARACTERIZATION OF THE NORTH CAROLINA COMMERCIAL SHRIMP TRAWL FLEET

I. ISSUE

Characterize the otter and skimmer trawl fleet in the North Carolina shrimp fishery and determine the impacts of any potential restrictions on headrope length, vessel size, or number of nets towed.

II. ORIGINATION

Request by the public, Coastal Conservative Association North Carolina (CCA NC), Coastal Fisheries Reform Group (CFRG), Marine Fisheries Commission (MFC), Shrimp Advisory Committee (AC)

III. BACKGROUND

The North Carolina shrimp fleet consists of vessels of various sizes and configurations. Roughly 92% of North Carolina's shrimp is harvested using otter trawls (NCDMF 2012). Otter trawls derived their name from the two trawl doors (otter doors/boards) that attach to the bridle that are hydrodynamically designed to hold the wings of the net open (Jennings et al. 2001). As the net is pulled along the bottom, the otter boards plane in opposite directions holding the net open. The webbing or the "body" of the net is usually constructed of nylon or polyethylene mesh and is also held open by a series of buoys attached to a "headrope" and weighted "footrope". "Tickler chains" are attached between the otter doors in front of the footrope to agitate the bottom, spooking shrimp into the net. Larger nets may also be rigged with rollers on the footrope preventing it from digging into muddy bottoms (NMFS 2012). The footrope can also be fitted with "rockhoppers" made out of rubber bobbins that allow the trawl to bounce over obstructions (Jennings et al. 2001). While the configuration of otter trawls may vary from state-to-state, all otter trawls consist of a pair of otter doors, headrope, footrope, tickler chain, and mesh body with wings that funnel shrimp in to a "cod end" or "tail bag" (Figure 12.29).

There are a number of otter trawl designs used in the southeastern shrimp fishery which include, the flat trawl, two-seam trawl, four-seam trawl, and tongue or "mongoose" (Watson et al. 1984; NMFS 2012). Optimum fishing configuration of trawls may be changed to reduce the bycatch of jellyfish, sponges, bottom trash, and finfish (Watson 1984). Trawls may also be rigged for different substrates and target species. In North Carolina, conventional two-seam otter trawls are used for bottom-hugging pink and brown shrimp, while four seam and tongue trawls with adjustable headropes are used for white shrimp which have the ability to jump over two-seam trawls when disturbed (NCDMF 2012; NMFS 2012). While otter trawl design and construction may vary, headrope length is commonly used to define a single otter trawl's horizontal spread or size. However, footrope length may also be used to determine the horizontal spread of tongue nets that have adjustable headropes. In North Carolina, the size of a trawl is based on its headrope length. Headrope length is defined as the support structure for the mesh or webbing of a trawl that is nearest to the water surface when in use. Headrope length is measured from the outer most mesh knot at one end of the headrope following along the line to the outer most mesh knot at the opposite end of the headrope [15A NCAC 03I .0101(i)].

When otter trawls were first brought to North Carolina in the early 1920s, 15 to 20 ft skiffs powered by small gasoline engines were used to pull a single rig otter trawl (one net) with

headropes ranging in size from 25 to 50 ft. By the 1940s and 1950s, larger 40 to 60 ft diesel-powered vessels began pulling larger nets with headropes ranging in size of 60 to 65 ft (Maiolo et al. 1980; Maiolo 2004). As vessels became larger and more powerful, trawlers began using “double-barrel” and “four-barreled” rigs allowing a vessel to pull multiple smaller trawl nets (40-50 ft) with combined headrope lengths measuring up to 200 ft (NCDMF 2012; NMFS 2012). The use of smaller nets creates less drag, making vessels more fuel efficient. Watson (1984) notes that four-barrel trawls are able to sweep a larger total area per gallon of fuel than do double-barrel rigs. Additional benefits of double and four-barreled also include: (1) increased catch per unit of effort, (2) fewer handling problems with the smaller nets, (3) lower initial gear costs, (4) reduced costs associated with damage or loss of nets, and (5) greater crew safety (NMFS 2012).

The size of gear allowed in North Carolina’s shrimp fishery has been the subject of debate particularly with respect to trawls. Prior to the 2006 Shrimp FMP, there were size limits on channel nets and on recreational shrimp trawls (26 ft headrope length) used by recreational commercial gear license (RCGL) holders, but no restriction on the size of trawls used in the commercial shrimp fishery. At the time, many fishermen felt that there should be a maximum limit placed on the size of trawls particularly in some of the smaller water bodies. They cited it was unfair to allow larger vessels into these areas especially on opening days when many boats would crowd into an area. It was thought that the larger vessels took most of the shrimp, rendering areas unproductive for several days, and then left to fish in more open waters unworkable by the smaller vessels. In addition to fairness, management actions were put into place to reduce bycatch and decreased the affect of trawling on the habitat. Currently, it is unlawful to use shrimp trawls that have a combined headrope greater than 90 ft in the internal coastal waters of North Carolina, except in the Pamlico Sound and mouths of the Pamlico and Neuse rivers [15A NCAC 03L .0103(c)].

Otter Trawl Headrope/Footrope Regulations in Other States

Many states have enacted various regulations limiting maximum headrope length and trawl configuration (Table 12.32). Estuarine trawling is prohibited in much of South Carolina, however in designated areas fishermen may use shrimp trawls with a combined footrope length no greater than 220 ft and “try” nets or test nets cannot have a footrope length greater than 16 ft (SC 50-5770). In South Carolina, a majority of the fishermen use tongue nets with adjustable headrope to target white shrimp, thus the footrope is used to measure horizontal spread. In Georgia it is unlawful to fish for shrimp for human consumption with trawls having a total footrope length greater than 220 ft (OCGA 27-4-133). Georgia commercial and recreational bait shrimpers are further restricted to trawls with maximum footrope lengths of 20 ft and 10 ft, respectively. Georgia state law also prohibits trawlers targeting shrimp for human consumption from the 60 bait zones located in the middle and upper estuaries as well the sounds and its sounds are closed to trawling for shrimp taken for consumption. Florida’s net ban in 1994 limited the use of all nets over 500 square feet of mesh and reduced Florida’s shrimp fishery to a bait shrimp fishery; however, trawling for shrimp for human consumption still occurs on a small scale. In the nearshore and inshore waters of Florida where otter trawls are allowed, fishermen are limited to a single net with a headrope no greater than 10 ft. Two trawls may be used in certain nearshore and inshore regions of Florida, however combined headrope length cannot exceed 20 ft. Outside of these waters trawlers may use up to a single net with headrope no greater than 20 ft (FL 68B-31.009).

The Gulf States also have limited maximum headrope length and the number of nets fishermen are allowed to use. In Alabama, commercial vessels operating in Mobile Bay and its sounds are

limited to two trawls with a maximum combined headrope length of 50 ft. In the offshore waters of Alabama there is no restriction on trawl size or number of rigs. Alabama commercial and recreational bait trawlers are restricted to a single trawl with a maximum headrope length of 16 ft. In Mississippi, commercial shrimp trawlers operating in the internal waters are allowed to use a single trawl with a maximum headrope length of 50 ft or two 25 ft trawls. Recreational fishermen in Mississippi are limited to a 16 ft maximum headrope length. Much like Mississippi, commercial vessels fishing in the inshore waters of Louisiana are limited to a single net with a headrope of 50 ft or two nets with headropes not exceeding 25 ft each. However, in Breton and Chandeleur Sounds vessels may use two nets with a headrope of 65 ft (130 ft combined). Vessels fishing in Louisiana's territorial waters (from beach to 3 miles offshore) may use trawls with up to 130 ft of headrope and in the federal waters (EEZ) vessels may use up to four nets of any size. Recreational fishermen in Louisiana are limited to a single net with a maximum headrope length of 25 ft. Much like Florida, Texas implements its headrope and trawl configurations by region, season, and intended use (bait vs. consumption). However, the maximum allowable headrope length is also determined by door length in Texas; trawls with headrope lengths measuring 40 ft must have a door 3 to 4 ft in length and nets measuring 54 ft must have a door 10 ft or larger. In major bays of Texas commercial fishermen targeting penaeid shrimp may use a single net with a headrope measuring 40 to 54 ft during the Spring Open Season (May 15-July 15). During the Fall Open Season (Aug. 15-Nov. 30) they may use a single net with a headrope not exceeding 95 ft. During the Winter Open Season (Feb.1-April 15) Texas fishermen working south of the Colorado River are limited to a single net with a headrope measuring 40 to 54 ft. Commercial bait fishermen are also limited to a single net with a headrope measuring 40 to 54 ft; however, bait fishermen are allowed to use smaller mesh sizes and are required to meet additional requirements and trip limits. Commercial vessels operating in Texas inside 3 nautical miles may use two trawls with headrope lengths ranging from 71 to 89 ft based on door size. Vessels operating 3 to 9 nautical miles offshore are not limited by number of nets they can pull or headrope length.

Skimmer Trawl Headrope Regulations

While headrope length is most commonly associated with otter trawls, headrope length can also be used to describe the length of the support structure that the mesh or webbing attaches to that is nearest the surface of the water. Thus, the headrope length of most skimmer trawls is dictated by the length of the skimmer trawl frame. Most skimmer trawls consist of two frames mounted to each side of the vessel, net (mounted to each frame), sled, weighted shoes, tickler chain, and an "easy line" (Figure 12.30). When deployed, the nets are aligned perpendicularly to the vessel and held in place by two or more stays or cables that run to the bow (NMFS 2012). The weighted sleds or "bullets" also help to maintain the nets position in the water column, while the weighted skids or "shoes" allow the frame to ride along the bottom, rising and falling with the bottom contour. Shrimp are spooked into the net by the tickler chain as it agitates the bottom; the tickler chain is attached in front of the lead line or footrope. The easy or "lazy" line is attached to the skimmer trawl tailbag and allows the tailbag to be retrieved without stopping the vessel (Coale et al. 1994).

Very few states have specific regulations for skimmer trawl configuration in regards to net length and design. Mississippi's skimmer trawl regulations mirror their otter trawl regulations, limiting vessels to two nets with a 25 ft headrope on each diagonal arm (not to exceed a combined headrope length of 50 ft). In Florida, skimmers must be equipped with rollers and vessels are limited to two unconnected trawls with upper and lower horizontal beams that do not exceed 16 ft in length each. In most states where skimmer trawl net and frame lengths are not specified, headrope length is defined to include the length of supporting structure that is the nearest to the

surface of the water. In the internal coastal waters of North Carolina, skimmer trawls nets cannot exceed a combined headrope length of 90 ft, except in the Pamlico Sound and parts of the Pamlico and Neuse Rivers [15A NCAC 03L .0103(c)].

Characterization of North Carolina Trawler Configuration

Using data from the Commercial Fishing Vessel Registration (CFVR) license, North Carolina Trip Ticket Program (NCTTP), DMF characterization studies (Brown 2009, 2010a, 2010b) and phone surveys, headrope length data were compiled to characterize the North Carolina shrimp trawl fleet during 2010 and 2011. A CFVR is required if a vessel is going to be used in a commercial fishing operation. When fishermen apply or renew their CFVR to obtain their vessel ID or “P” number they are asked a series of survey questions pertaining to vessel length, type of gear, number of nets, headrope length, as well as other vessel specifications. Data from the NCTTP were analyzed to identify the vessels that operated in the commercial shrimp trawl fishery in 2010 and 2011. Once the vessels were identified, their respective survey data obtained from the CFVR was then extracted from the DMF License database. Some vessels entered the shrimp trawl fishery after they obtained their CFVR license. To obtain the gear characteristics for these vessels, NCTTP port agents conducted phone surveys of the vessel owners. The phone surveys and characterization studies were used to further verify the CFVR data. Using each vessel’s ID and landings data from the NCTTP, total shrimp landings (all three species combined) were calculated by gear, area, total headrope length (ft), number of nets, and vessel length (ft). In viewing the tables and figures keep in mind while this is the best available data, it still has several limitations: 1) gear data from the CFVR is for their predominant gear and variation in the use of different size or number of nets and rigs is not captured, and 2) in a similar manner only one predominant waterbody can be recorded on paper trip ticket forms. It should also be noted that estimated reductions calculated using vessel and headrope length frequency distributions can be influenced by bin size (10 ft increments); thus averages and modes listed in Table 12.33 calculated on a finer scale (1ft increments) may not correspond directly to the bin modes shown in the figures .

Pamlico Sound

The number of vessels using otter trawls in Pamlico Sound declined 4% from 220 in 2010 to 201 in 2011 (Table 12.33). Average vessel length ranged from 49 to 53 ft, while the most frequently occurring (mode) vessel length was 36 ft. Vessels 30 ft or less made up 13% of the fleet in 2010 and 21% in 2011, while boats larger than 90 ft or greater made up 6% in 2010 and 5% in 2011 (Figure 12.31). On average these vessels made 1,656 trips in 2010 and 1,502 trips in 2011. Average shrimp landings ranged from 2,317 lb/trip in 2010 to 2,419 lb/trip in 2011. The majority of the vessels operating in Pamlico Sound during 2010 and 2011 used double and four-barrel rigs. Vessels using double-barrel rigs made up 32% of the fleet in 2010 and 35% in 2011, those using four-barrel rigs made up 54% in 2010 and 46% in 2011. Boats using single rigs made up 14% in 2010 and 18% in 2011. The average total headrope length ranged from 117 ft in 2011 to 128 ft in 2010. The most frequently observed total headrope length was 180 ft in 2010 and 70 ft in 2011. However, the vast majority of the fleet used nets with headropes larger than 70 ft. Total headropes lengths measured 70 ft or greater made up 83% of the observations in 2010 and 77% in 2011 (Figure 12.32).

Very few fishermen used skimmer trawls in the Pamlico Sound during 2010 and 2011 (Table 12.33). Only two vessels were observed in 2010 and four in 2011. Average vessel length ranged from 24 to 34 ft. In 2011, three out of the four vessels observed were over 40 ft (Figure 12.33). Since only two vessels used skimmer rigs in the Pamlico Sound in 2010 their landings

data is confidential and cannot be shown. In 2011, four vessels made four trips landing an average of 175 lb/trip. All skimmers operating in the Pamlico Sound were double rigged. The average total headrope length ranged from 20 ft in 2010 to 46 ft in 2011. Total headrope lengths measuring 50 ft or greater were not observed in 2010, however they made up 50% of rigs in 2011 (Figure 12.34).

Neuse, Pamlico and Bay Rivers

In the Neuse, Pamlico, and Bay Rivers, 58 vessels made 377 trips, landing an average of 305 lb/trip of shrimp in 2010 (Table 12.33). In 2011, 49 vessels made 446 trips, landing an average of 235 lb/ trip. Average vessel length ranged from 30 ft to 31 ft. The most frequently observed vessel length observed in 2010 was 20 ft, dropping slightly in 2011 to 19 ft. Vessels 40 ft or less made up approximately 85% of the fleet during both years (Figure 12.35). Double-barrel rigs were predominately used in 2010 (57%) and 2011 (51%). Vessels using single rigs made up 38% of the fleet in 2010 and 43% in 2011. Less than 10% of the vessels used four-barreled rigs during 2010 and 2011. The average total headrope length ranged from 52 ft in 2011 to 55 ft in 2010. The most frequently observed total headrope length was 80 ft in 2010 and 30 ft in 2011. Vessels using total headrope lengths that were 90 ft or less made up 95% of observations in 2010 and 92% in 2011 (Figure 12.36).

Skimmer trawls landed on average 202 lb/trip in 2010 and 235 lb/trip in 2011. The number of vessels using skimmer trawls in the Neuse, Pamlico, and Bay Rivers ranged from 4 to 7 (Table 12.33). Average vessel length ranged from 22 to 28 ft. In 2010, the most frequently observed vessel length was 25 ft. In 2011, 71% of the vessels were approximately 30 ft in length and in 2010 all vessels were between 20 and 30 ft (Figure 12.37). All vessels were double rigged with total average headrope lengths and ranged from 21 to 27 ft. In 2010, the most commonly observed headrope length was 28 ft. In 2011 the number of vessels using total headropes length between 20 and 30 ft were equally distributed; however in 2010, 86% of the headropes were 30 ft or less (Figure 12.38).

Bogue and Core Sounds, Newport and North Rivers

The number of vessels using otter trawls in central internal waters of the state (Bogue Sound, Core Sound, Newport River, and North River) ranged from 43 to 67 (Table 12.33). In 2010, 553 trips landed an average of 199 lb/trip. While the number of trips fell almost 70% to 166 trips the landings increased slightly to 208 lb/trip in 2011. Average vessel length ranged from 28 to 29 ft. The most frequently occurring vessel length ranged from 21 to 22 ft. In 2011, there was a slight increase in the number of 30 ft vessels as well as 50 ft vessels (Figure 12.39). The number of vessels using single rigs increased from 45% in 2010 to 49% in 2011. Vessels using double-barrel rigs fell slightly in 2010 from 52% to 51% in 2011. Overall, very few boats used four-barrel rigs in this part of the state. Average total headrope lengths ranged from 46 to 47 ft. However, vessels using total headrope lengths that were 20 ft or less made up approximately 42% observations during both years (Figure 12.40). The most commonly observed total headrope length was 15 ft during both years.

In 2010, CFVR data indicated that there were 37 skimmers in central region internal water of the state; this number fell to 12 in 2011 (Table 12.33). Vessel length ranged from 28 to 29 ft. The most commonly reported vessel length was 25 ft in 2010 and 28 ft in 2011. No boats larger than 50 ft were observed (Figure 12.41). As with the other parts of the state, all skimmer rigs consisted of two rigs (Table 12.33). Total average headrope length was 29 ft in 2010 and increased slightly in 2011 to 32 ft. The most commonly reported head rope was 20 ft in 2010

and 24 ft in 2011. Total headrope lengths measuring 40 ft or greater made up on average 27% to 33% fleet gear in 2010 and 2011, respectfully (Figure 12.42). On average these vessels landed an average of 218 lb/trip of shrimp in 2010, average landings declined in 2011 to 154 lb/trip.

Southern Region

In 2010, 103 vessels landed an average of 155 lb/trip of shrimp using otter trawls in the Intracoastal Waterway (IWW) to the South Carolina state line and the New and Cape Fear Rivers (Table 12.33). Otter trawl landings fell roughly 32% in this part of the state in 2011. Average vessel length ranged from 22 to 23 ft during 2010 and 2011. The most commonly reported vessel length was 17 ft in 2010 and 19 ft in 2011. While vessels ranging in length of 40 to 70 ft made up only 8% of the fleet in 2010 and 14% in 2011 (Figure 12.43), it's important to note that the data used to characterize the fleet is based on its CFVR data and trip ticket landings and doesn't take into account area restrictions. Regulations limiting headropes and the inability to safely navigate the narrow waterway of areas such as the southern portion of the state restrict larger vessels. Vessels with total headrope lengths measuring 40 ft or less made up 83% of the fleet in 2010 and 77% of the fleet in 2011 (Figure 12.44).

In 2010, 26 vessels reported using skimmer rigs in the southern region; this number declined to 17 in 2011 (Table 12.33). Out of the 439 trips made in 2010, an average of 313 lb/trip of shrimp were landed. In 2011, 149 lb/trip of shrimp was landed. The average vessel length reported ranged from 30 to 33 ft. The most commonly observed vessel length reported ranged from 17 ft in 2010 to 38 ft in 2011. Skimmer vessels ranging in length of 40 to 50 ft made up 42% of the fleet in 2010 and 52% in 2011 (Figure 12.45). As with the other parts of the state all vessels used double rigs. Average total headrope lengths ranged in size from 40 to 42 ft, with a mode of 48 ft. Vessels with total headrope lengths measuring between 50 and 70 ft made up 39% of the fleet in 2010 and 53% in 2011 (Figure 12.46).

Atlantic Ocean

The number of vessels using otter trawls in the Atlantic Ocean declined 21% from 116 in 2010 to 92 in 2011 (Table 12.33). The average vessel length was 51 ft and the most commonly reported vessel size was 55 ft during 2010 and 2011. However, vessels between 60 and 90 ft made up roughly 50% of the fleet during both years and the number of 60 and 80 ft vessel both increased in 2011 (Figure 12.47). The majority of the vessels operating in the ocean used four-barrel rigs, with roughly 47% of the fleet using them. Double-barrel rigs were the second most commonly used configuration, with 33% using them in 2010 and 28% using them in 2011. An average of 120 ft of total headrope was used during 2010 and 2011. The most commonly observed total headrope value was 160 ft in 2010 and 200 ft in 2011. Vessels using total headropes less than 120 ft accounted for 44% of the fleet in 2010 and 46% in 2011 (Figure 12.48). Overall, vessels using otter trawls in the Atlantic Ocean landed an average of 772 lb/trip of shrimp in 2010 and 819 lb/trip in 2011. No landings were reported for skimmer trawls in the Atlantic Ocean during 2010 and 2011.

IV. AUTHORITY

- § 113-134. RULES
- § 113-173. RECREATIONAL COMMERCIAL GEAR LICENSE
- § 113-182. REGULATION OF FISHING AND FISHERIES
- § 143B-289.52. MARINE FISHERIES COMMISSION – POWERS AND DUTIES

V. DISCUSSION

In 2011, DMF sent out a press release and held several public meetings as part of the 5-year review of the Shrimp FMP. At those meetings and through public comment, several requests were made to limit inshore trawls to a headrope length of 50 ft. This request was later echoed by Coastal Conservation Association North Carolina (CCA NC) and its members following the DMF's initial plan to revise the FMP. Shortly after the formation of the Shrimp Advisory Committee (AC), a letter was sent from the Coastal Fisheries Reform Group (CFRG) requesting that all trawl nets in North Carolina's inshore coastal waters be limited to a single net with a maximum headrope size of 35 ft.

North Carolina's headrope regulations were put in place following the 2006 Shrimp FMP as a means to allocate the resource fairly amongst vessels of all sizes, reduce bycatch, and to limit the effects of trawling in prescribed areas. In other states, headrope limitations have also been used to reduce conflict and effort, specifically the fishing power of larger boats. While there has been no definitive data indicating there is more overall bycatch associated with larger trawlers, the general public perceives that larger vessels pulling double-barrel and four-barrel rigs are capable of removing more non-target species. Double-barrel and four-barrel rigs are capable of sweeping larger areas; however, the fishing power, efficiency and selectivity of the gear rely on more than just the length of its headrope. The measurement of a net's gape, measured by the horizontal spread and vertical distance between the headrope and footrope, affects not only the nets efficiency, but its ability to reduce unwanted bycatch (Watson et al. 1984). Just as there is no ideal trawl design or configuration for harvesting every species of shrimp in all substrates, there is no ideal design that excludes both demersal and pelagic species of fish (Watson et al. 1984; Harrington et al. 1985). While double and four-barrel rigs are capable of sweeping more area, making demersal species more vulnerable to the gear, the reduced vertical height of multiple smaller nets may reduce the bycatch of more pelagic species (S. Nichols, NMFS. pers. com. 1995). The use of smaller nets associated with double and four-barrel rigs may also allow larger fish to escape during haul back due to the shortened body length of the net; these reductions may be minimal for vessels pulling tongue trawls with adjustable headropes.

Using the distribution of vessels lengths reported in the CFVR and trip ticket data linked by a vessel's ID, reductions in effort were calculated for 2010 and 2011. While these data provide insight on the potential effects of regulations limiting vessel size, number of rigs and total headrope length, estimates of bycatch reduction cannot be calculated. Implementing a maximum vessel size would reduce the fleet size in North Carolina's internal waters. Overall, the average length of shrimp trawlers operating in the internal waters of North Carolina ranged from 22 to 53 ft and varied by water body and year (Table 12.33). Establishing a maximum vessel size would be difficult and the total reduction in fleet size may be minimal in certain areas. If vessels size was limited to 50 ft in internal coastal waters of North Carolina, 44% to 52% of the vessels operating in the Pamlico Sound would no longer be allowed to fish. In the southern portion of the state reductions in fleet size would range from 2% to 4% and 5% to 6% in the Neuse, Pamlico, and Bay Rivers (Figure 12.34 and Figure 12.41). Limiting the size of vessels using skimmer trawls to 50 ft or less would not reduce the fleet size based on the 2010 and 2011 data (Figures 4,8,10,12,14). Not only would it be difficult to determine the appropriate vessel size for each region of the state, but enforcement would be extremely difficult and would most likely cause shifts in effort. While current regulations limit the use of 90 ft headropes in the internal waters of North Carolina, there is not a direct limit on the size of vessels. However, larger vessels were typically found to have larger total headrope lengths (Figure 12.49). North Carolina General Statute 143B-289.52 (a)(1)a. provides the MFC with the authority to "authorize,

license, regulate, prohibit, prescribe, or restrict all forms of marine and estuarine resources in coastal fishing waters with respect to: time, place, character, or dimensions of any methods or equipment that may be employed in taking fish.” The Division is verifying whether a vessel would be included in the meaning of equipment in this statute.

Similar considerations apply when examining regulations limiting the total number of rigs or otter trawls a vessel may use. Eliminating the use of four-barrel rigs may reduce the fleet size by as much as 54% in the Pamlico Sound, while only reducing effort by as much as 6% in the Neuse, Pamlico, and Bay Rivers (assuming vessels do not re-rig) (Table 12.33). Restricting the use of double-barrel rigs in the Neuse, Pamlico, and Bay Rivers would have more impact, reducing the fleet size by as much as 57%. In the southern portion of the state, only 11% to 16% of the vessels reported using double and four-barrel rigs. In many parts of the state vessels using double and four-barrel rigs are limited to waters that allow the use of total headropes 90 ft or greater. Thus, potential reductions based on the number of rigs are confounded not only by regulations, but also by CFVR data that does not account for variation in fishing methods and gear quantities across different areas. Restricting the use of four-barrel rigs would only be effective at reducing the fleet size in the Pamlico Sound. Overall, restricting the total headrope length of otter trawls would essentially restrict the total number of rigs as well as vessel size in most parts of the state. Limiting skimmer trawls to single rigs, would not only eliminate the majority of North Carolina’s skimmer fleet, but would reduce the vessel’s ability to navigate.

The distribution of total headrope length (10 ft bins) was used to estimate the effect of reducing the maximum headrope length in the internal coastal waters of North Carolina. Establishing a 50 ft otter trawl total headrope limit would reduce the fleet gear by as much as 60% statewide during 2010 and 2011. However, this reduction would be even greater in the Pamlico Sound; reducing the fleet gear by 84% in 2010 and 79% in 2011 (Figure 12.33). In the Neuse, Pamlico, and Bay Rivers, estimated reductions would have been 48% in 2010 and 39% in 2011 (Figure 7). During both 2010 and 2011, the number of vessels with total headrope length greater than 50 ft would be reduced roughly 46% in the aggregate waters of Bogue and Core Sounds, Newport and North Rivers (Figure 11). The reductions in the southern portion of the state would be minimal for otter trawls at 9% in 2010 and 15% (Figure 12.43). However, a 50 ft headrope limit would reduce the fleet gear of skimmer vessels in the southern portion of the state by as much as 19% in 2010 and 24% in 2011 (Figure 12.45). In the Pamlico Sound, there were no skimmers observed with total headrope lengths over 50 ft in 2010 and in 2011 only two of four skimmer trawls observed had headropes larger than 50 ft (Figure 12.33). No skimmer trawls reported using total headropes larger than 50 ft in Neuse, Pamlico, and Bay River during both years of the survey (Figure 12.36).

Implementing a 35 ft maximum headrope length in the internal coastal waters of North Carolina would severely reduce the fleet size and the fishing power of the otter trawl fishery. As with a 50 ft maximum headrope length, the Pamlico Sound otter trawl fishery would see the greatest reductions at 92% in 2010 and 88% in 2011 (Figure 12.31). In the adjacent waters of the Neuse, Pamlico, and Bay Rivers estimated reductions for otter trawls would range from 61% in 2011 to 67% in 2010 (Figure 12.34). Roughly half of vessels using otter trawls in the central (Bogue and Core Sounds, Newport and North Rivers) and southern portions of the state would no longer be able to operate using their gear configurations as reported in 2010 and 2011 (Figure 12.39 and Figure 12.43). Vessels using skimmer rigs in the southern portion of the state would also see severe reductions ranging from 69% and 82% (Figure 12.45). Skimmers in the central portion of the state were estimated to have the second highest reductions at 27% in 2010 and 33% in 2011 (Figure 12.41). No vessels reported having had a total headrope greater than 35 ft in the Neuse, Pamlico and Bay Rivers in 2011 and only one vessel (14%) was observed in 2011, thus

estimated reductions would be minimal (Figure 12.37). There were also no skimmer vessels using headropes greater than 35 ft in the Pamlico Sound in 2010 and in 2011 there were only two vessels that would not be able to operate using their reported gear configuration (Figure 12.33). It is important to note that estimated reductions based on total headrope length and vessel size may be an overestimate as a result of bin size (10 ft increments). The estimated reductions presented in this paper do not represent reductions in fishing effort; they are merely a snap shot of the potential reductions in the fleet size and gear based on the vessel and gear configurations observed in 2010 and 2011. Hence, these reductions may not accurately reflect the current make up of the fishery or the years prior to 2010.

Regardless of vessel size, shrimp trawl design has evolved to improve the efficiency of the gear to capture shrimp and maximize area swept. Regulations limiting total headrope length will reduce the efficiency of both large and small vessels using nets with headropes larger than 35 ft. Thus, overall effort will likely be reduced due to a loss of fishing power and as fishermen leave the fishery because it is no longer economically feasible to continue. Not only will the current gear configuration used by many fishermen be obsolete, but operating costs will likely exceed the value of their catch. Shifts in effort may also occur putting more pressure on already over burdened fishing locations, leading to increased conflict and minimal reductions in bycatch. Fishermen attempting to compensate for lost catches as result of being forced to use less efficient gear may actually make more or longer tows, generating more bycatch per pound of shrimp landed. Reductions in bycatch may also be minimal if crews of larger vessels begin operating multiple smaller vessels, not only increasing effort (participants and trips) but the total headrope size of the fleet as a whole. There is also the potential for shifts in the species and size makeup of the bycatch. If larger vessels are forced out of the internal coastal waters into the ocean due to regulations that reduce total headrope length, more pressure may be put on the winter ocean spawners (spot, croaker, and flounder). While reducing headrope length has the potential to reduce bycatch associated with inshore trawling (Watson et al. 1984), the issue is extremely complex making it difficult to quantify its total impact on the fishery beyond reduction in effort; social, economic, and historical factors must also be examined. Unfortunately, all of the necessary data do not exist to adequately quantify the full extent that such regulations could have on levels of bycatch reduction and on the shrimp trawl fishery and its associated industries. Even after all the data are presented, there still will not be clear guidance concerning the issue. The essential decision will be the unquantified potential gain in some natural resources versus the losses to a major economically important fishery.

VII. EVALUATION MATRIX

AC Evaluation of Characterization of the North Carolina Commercial Shrimp Trawl Fleet							
Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/Administrative	Other Impacts
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. Continued reduction in effort may result in overall bycatch reduction in the fleet. -	Will not create shifts in effort to other fisheries. Maintains present market value of fishery. N	Allows flexibility of fishermen to continue to fish in their normal areas using their normal gears. N	Commercial and recreational fishing will continue with no changes in gear use or conflict. Potential recreational angling could remain stagnant if status quo continues. +/-	Same level of enforcement. Not Evaluated	No change in rule. Not Evaluated	Not Evaluated
2. Reduce maximum headrope length all internal coastal waters for commercial and recreational fisheries. AC added "for commercial and recreational fisheries".	May reduce bycatch from vessels using larger headropes. Reductions may be minimal as effort changes. Size and species makeup of bycatch may shift as effort changes. +/-	Overall loss in gear efficiency will likely result in a decrease in landings and income. Increases operating costs and reduces efficiencies for many shrimp fishing operations due to: 1) New gear purchase or reconfiguration 2) Increase in fuel cost 3) Increase travel time to non-restricted areas 4) Increase number of haul backs -	Prior efforts (time and money) expended to improve gear efficiencies lost. May reduce the availability of NC caught shrimp. May be favorably viewed by general public, while industry frustrated with additional restrictions. Remaining vessels may have less competition. +/-	May create conflicts with other commercial and recreational fishermen due to changes in effort. May improve recreational fishing. +/-	Increase in enforcement duties. Not Evaluated	Will require rule change. Not Evaluated	Not Evaluated

AC Evaluation of Characterization of the North Carolina Commercial Shrimp Trawl Fleet (August 15,2013)

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
<p>3. Implement a maximum number of rigs (i.e., double and four-barrel) in the internal coastal waters</p>	<p>May reduce bycatch from vessels using larger headropes. Reductions may be minimal as effort shifts. Size and species of bycatch makeup may shift as effort shifts.</p> <p align="center">+/-</p>	<p>Overall loss in gear efficiency will likely result in a decrease in landings and income. Increases operating costs and reduces efficiencies for many shrimp fishing operations due to:</p> <ol style="list-style-type: none"> 1) New gear purchase or reconfiguration 2) Increase in fuel cost 3) Increase travel time to non-restricted areas 4) Increase number of haul backs <p align="center">-</p>	<p>May reduce the availability of NC caught shrimp. May be favorably viewed by general public, while industry frustrated with additional restrictions. Remaining vessels may have less competition.</p> <p align="center">+/-</p>	<p>May create conflicts with other commercial and recreational fishermen due to changes in effort. May improve recreational fishing</p> <p align="center">+/-</p>	<p>Increase in enforcement duties.</p> <p align="center">Not Evaluated</p>	<p>Will require rule change.</p> <p align="center">Not Evaluated</p>	<p align="center">Not Evaluated</p>
<p>4. Implement a maximum vessel size in the internal coastal waters</p> <p>AC elected to not consider this option</p>	<p>Eliminates bycatch from larger vessels. Reductions may be minimal as effort changes. Size and species of bycatch makeup may shift as effort changes.</p>	<p>Increased operating cost as larger vessels are forced to travel further to fishing grounds. May create effort shifts into other fisheries. May result in reduced overall landings and income of fishermen and industry.</p>	<p>Reduces the availability of NC caught shrimp. Favorably viewed by general public, while industry frustrated with additional restrictions. Remaining vessels may have less competition.</p>	<p>May create conflicts with other commercial and recreational fishermen due to changes in effort.</p>	<p>Increase in enforcement duties.</p>	<p>Will require rule change. Authority to limit vessel size is unclear.</p>	

DMF Evaluation of Characterization of the North Carolina Commercial Shrimp Trawl Fleet (8/19/2013)

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
1. <i>Status quo</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. -	Will not create shifts in effort to other fisheries. Maintains present market value of fishery. N	Allows flexibility of fishermen to continue to fish in their normal areas using their normal gears. N	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. N	No change in rule. N	
2. Reduce maximum headrope length in all internal coastal waters for commercial and recreational fisheries.	May reduce bycatch from vessels using larger headropes. Reductions may be minimal as effort changes. Size and species makeup of bycatch may shift as effort changes. +/-	Overall loss in gear efficiency will likely result in a decrease in landings and income. Increases operating costs and reduces efficiencies for many shrimp fishing operations due to: 1) New gear purchase or reconfiguration 2) Increase in fuel cost 3) Increase travel time to non-restricted areas 4) Increase number of haul backs -	Prior efforts (time and money) expended to improve gear efficiencies lost. May reduce the availability of NC caught shrimp. May be favorably viewed by general public, while industry frustrated with additional restrictions. Remaining vessels may have less competition. +/-	May create conflicts with other commercial and recreational fishermen due to changes in effort. -	Increase in enforcement duties. -	Will require rule change. -	

DMF Evaluation of Characterization of the North Carolina Commercial Shrimp Trawl Fleet (8/19/2013)

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
3. Implement a maximum number of rigs (i.e., double and four-barrel) in the internal coastal waters	<p>May reduce bycatch from vessels using larger headropes. Reductions may be minimal as effort shifts. Size and species of bycatch makeup may shift as effort shifts.</p> <p align="center">+/-</p>	<p>Overall loss in gear efficiency will likely result in a decrease in landings and income. Increases operating costs and reduces efficiencies for many shrimp fishing operations due to:</p> <ol style="list-style-type: none"> 1) New gear purchase or reconfiguration 2) Increase in fuel cost 3) Increase travel time to non-restricted areas 4) Increase number of haul backs <p align="center">-</p>	<p>Prior efforts (time and money) expended to improve gear efficiencies lost. May reduce the availability of NC caught shrimp. May be favorably viewed by general public, while industry frustrated with additional restrictions. Remaining vessels may have less competition.</p> <p align="center">+/-</p>	<p>May create conflicts with other commercial and recreational fishermen due to changes in effort.</p> <p align="center">-</p>	<p>Increase in enforcement duties.</p> <p align="center">-</p>	<p>Will require rule change.</p> <p align="center">-</p>	
4. Implement a maximum vessel size in the internal coastal waters	<p>May reduce bycatch from larger vessels. Reductions may be minimal as effort changes. Size and species of bycatch makeup may shift as effort changes.</p> <p align="center">+/-</p>	<p>Increased operating cost as larger vessels are forced to travel further to fishing grounds. May create effort shifts into other fisheries. May result in reduced overall landings and income of fishermen and industry.</p> <p align="center">-</p>	<p>Reduces the availability of NC caught shrimp. Favorably viewed by general public, while industry frustrated with additional restrictions. Remaining vessels may have less competition.</p> <p align="center">+/-</p>	<p>May create conflicts with other commercial and recreational fishermen due to changes in effort.</p> <p align="center">-</p>	<p>Increase in enforcement duties.</p> <p align="center">-</p>	<p>Will require rule change. Authority to limit vessel size is unclear.</p> <p align="center">-</p>	

Table 12.32 Maximum commercial, bait, and recreational shrimp trawl configurations by state.

State	Commercial		Bait		Recreational		Additional Restrictions****
	Length (ft)	# of Nets	Length (ft)	# of Nets	Length (ft)	# of Nets	
North Carolina	90 [†]	unlimited	90 [†]	unlimited	26	1	Area [†] , Proclamation Authority SSNA
South Carolina	16* (try net), 220*						Season, area, time of day
Georgia	16* (try net), 220*		20*	1	10*	1	Closed Mar. 1-May 14, area closures
Florida (internal)	10	1 or 2 (10 ft nets)	10	1	16**	1	Body not to exceed 500 square feet
Florida (ocean)	20	1	20	1			Body not to exceed 500 square feet
Alabama (internal)	10 (try net), 50	2	16	1	16	1	Season, area (bait)
Alabama (offshore)	unlimited	unlimited					
Mississippi	50	1 or 2 (25 ft)			16	1	Season, area, door size
Louisiana (internal)	16 (try net), 50, 130 ^{††}	1 or 2 (25 ft), 2 (65 ft) ^{††}			25	1	Season, area, tow times (try nets)
Louisiana (3 mi offshore)	16 (try net), 130	unlimited			25	1	
Louisiana (EEZ)	unlimited	4			25	1	No night time by area
Texas (Bays - Spring)	21 (try net), 40-54 ^{***}	1	12 (try net), 40-54	1	20	1	Season (May 15 - July 15), area, door size
Texas (Bays - Fall)	21 (try net), 95	1	12 (try net), 40-54	1	20	1	Season (Aug. 15 - Nov. 30, area, door size
Texas (Bays - Winter)	21 (try net), 40-54 ^{***}	1	12 (try net), 40-54	1	20	1	Season (Feb.1 - April 15), area, door size
Texas (<3 mi offshore)	71-89 ^{***}	2					Season, time by area , door size
Texas (3-9 mi offshore)	unlimited	unlimited					Season, time by area (3-5 mi, no nights)

[†] It is unlawful to take shrimp with trawls which have a combined headrope of greater than 90 feet in internal coastal waters except:

(1) Pamlico Sound;

(2) Pamlico River downstream of a line from a point 35° 18.5882'N – 76° 28.9625'W at Pamlico Point; running northerly to a point 35° 22.3741'N - 6° 28.6905'W at Willow Point;

(3) Neuse River northeast of a line from a point 34° 58.2000'N – 76° 40.5167'W at Winthrop Point on the eastern shore of the entrance to Adam's Creek running northerly to a point 35° 01.0744' N – 76° 42.1550' W at Windmill Point at the entrance of Greens Creek at Oriental.

^{††} Breton and Chandeleur Sounds, Louisiana

* Footrope length

** Horizontal frame of skimmer (Florida further specifies use of roller in specific areas)

*** Maximum headrope length is specific to door length (ft); other trawls must have doors at least 3 ft long from the leading tip to trailing edge of door

**** In addition to no trawling in nursery areas, maximum mesh size restrictions (commercial and recreational), TED and BRD requirements (varies by state).

Table 12.33 North Carolina vessel and shrimp trawl configuration by area and year, 2010-2011.

Year	Trawl Type	Area Fished	Total Shrimp lb	Trips #	Average Shrimp (lb/trip)	Vessels #	Vessel Length		Total Headrope Length (all rigs)		Single Rig		Double-Barrel Rig		Four-Barrel Rig	
							Average ft	Mode ft	Average ft	Mode ft	#	%	#	%	#	%
2010	Otter	Pamlico Sound†	3,837,201	1,656	2,317	220	53	36	128	180	31	14%	71	32%	118	54%
2011	Otter	Pamlico Sound†	3,633,502	1,502	2,419	201	49	36	117	70	37	18%	71	35%	93	46%
2010	Otter	Neuse, Pamlico, Bay Rivers†	114,871	377	305	58	31	20	55	80	22	38%	33	57%	3	5%
2011	Otter	Neuse, Pamlico, Bay Rivers†	104,743	446	235	49	30	19	52	30	21	43%	25	51%	3	6%
2010	Otter	Bogue/Core/ Newport/North River	110,046	553	199	67	29	22	47	15	30	45%	35	52%	2	3%
2011	Otter	Bogue/Core/ Newport/North River	34,584	166	208	43	28	21	46	15	21	49%	22	51%	0	0%
2010	Otter	Southern	216,110	1,394	155	103	22	17	38	35	92	89%	7	7%	4	4%
2011	Otter	Southern	114,799	945	121	65	23	19	39	30	55	85%	9	14%	1	2%
2010	Otter	Ocean	1,253,754	1,623	772	116	51	55	120	160	23	20%	38	33%	55	47%
2011	Otter	Ocean	1,091,810	1,333	819	92	51	55	120	200	22	24%	26	28%	44	48%
2010	Skimmer	Pamlico Sound†	*	*	*	2	24	-	20	-	0	0%	2	100%	0	0.0%
2011	Skimmer	Pamlico Sound†	699	4	175	4	34	34	46	-	0	0%	4	100%	0	0.0%
2010	Skimmer	Neuse, Pamlico, Bay Rivers†	14,771	73	202	7	28	25	27	28	0	0%	7	100%	0	0.0%
2011	Skimmer	Neuse, Pamlico, Bay Rivers†	17,191	73	235	4	22	-	21	-	0	0%	4	100%	0	0.0%
2010	Skimmer	Bogue/Core/ Newport/North River	132,458	607	218	37	28	25	29	20	0	0%	37	100%	0	0.0%
2011	Skimmer	Bogue/Core/ Newport/North River	14,470	94	154	12	29	28	32	24	0	0%	12	100%	0	0.0%
2010	Skimmer	Southern	137,408	439	313	26	30	17	40	48	0	0%	26	100%	0	0.0%
2011	Skimmer	Southern	23,215	156	149	17	33	38	42	48	0	0%	17	100%	0	0.0%

* Confidential, 3 or less participants, vessels, or dealers

† It is unlawful to take shrimp with trawls which have a combined headrope of greater than 90 feet in internal coastal waters except:

(1) Pamlico Sound;

(2) Pamlico River downstream of a line from a point 35° 18.5882'N – 76° 28.9625'W at Pamlico Point; running northerly to a point 35° 22.3741'N - 6° 28.6905'W at Willow Point;

(3) Neuse River northeast of a line from a point 34° 58.2000'N – 76° 40.5167'W at Winthrop Point on the eastern shore of the entrance to Adam's Creek running northerly to a point 35° 01.0744' N – 76° 42.1550' W at Windmill Point at the entrance of Greens Creek at Oriental.




Figure 12.29 Schematic of an otter trawl and its components (from NMFS 2012).




Figure 12.30 Schematic of a skimmer trawl and its components (from NMFS 2012).




Figure 12.31 Length frequency distribution of vessels using otter trawls in the Pamlico Sound, 2010-11.




Figure 12.32 Length frequency distribution of total headrope length (ft) of otter trawls in the Pamlico Sound, 2010-11.




Figure 12.33 Length frequency distribution of vessels (ft) using skimmer trawls in the Pamlico Sound, 2010-11.




Figure 12.34 Length frequency distribution of total headrope length (ft) of skimmer trawls in the Pamlico Sound, 2010-11.




Figure 12.35 Length frequency distribution of vessels (ft) using otter trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.




Figure 12.36 Length frequency distribution of total headrope length (ft) of otter trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.




Figure 12.37 Length frequency distribution of vessels (ft) using skimmer trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.




Figure 12.38 Length frequency distribution of total headrope length (ft) of skimmer trawls in the Neuse, Pamlico and Bay Rivers, 2010-11.




Figure 12.39 Length frequency distribution of vessels (ft) using otter trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11.




Figure 12.40 Length frequency distribution of total headrope length (ft) of otter trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11.




Figure 12.41 Length frequency distribution of vessels (ft) using skimmer trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11.




Figure 12.42 Length frequency distribution of total headrope length (ft) of skimmer trawls in the Bogue and Core Sounds as well as the Newport and North Rivers, 2010-11.




Figure 12.43 Length frequency distribution of vessels (ft) using otter trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.




Figure 12.44 Length frequency distribution of total headrope length (ft) of otter trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.




Figure 12.45 Length frequency distribution of vessels (ft) using skimmer trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.




Figure 12.46 Length frequency distribution of total headrope length (ft) of skimmer trawls in the southern region of the state (New River, Cape Fear River, IWW to SC state line), 2010-11.




Figure 12.47 Length frequency distribution of vessels (ft) using otter trawls in the Atlantic Ocean, 2010-11.




Figure 12.48 Length frequency distribution of total headrope length (ft) of otter trawls in the Atlantic Ocean, 2010-11.




Figure 12.49 Scatter plot of vessel length (ft) and total headrope length (ft) of the North Carolina shrimp trawler fleet (all water bodies included), 2010-11.

12.8 AREA RESTRICTIONS TO REDUCE SHRIMP TRAWL BYCATCH IN NORTH CAROLINA'S INTERNAL COASTAL WATERS

I. ISSUE

The consideration of closing waterbodies to shrimp trawl gear in North Carolina

II. ORIGINATION

The Division of Marine Fisheries (DMF) Shrimp Plan Development Team (PDT) and the public

III. BACKGROUND

The estuarine system in North Carolina is the largest of any state along the Atlantic coast. Its coast is framed by a chain of low-lying barrier islands, from Virginia to Cape Fear River and makes up a diverse aquatic system of estuarine rivers, creeks, large sounds, and inlets totaling over 2.2 million acres of estuarine waters (Deaton et al. 2010; DMF unpublished data). The northern portion of these natural barrier islands are called the Outer Banks and separates the Albemarle-Pamlico sound complex from the coastal ocean. Along the southern coast, southwest of the Cape Fear River, the Intracoastal Waterway (IWW) creates an artificial extension of these barrier Islands. The topography of the three major capes (Cape Hatteras, Cape Lookout, and Cape Fear) has a major influence on adjacent ocean circulation. North Carolina is located at the convergence of two major ocean currents: the warm, north flowing Gulf Stream Current and the cool south-flowing Labrador Current that creates a mix of both northern and southern fish species in North Carolina waters. The convergence of currents and the diversity and abundance of habitat and species occurring in North Carolina's estuaries makes its coastal fisheries among the most productive in the United States.

Pamlico Sound is considered an extremely important area for both commercial and recreational fishing. It makes up approximately 56% of North Carolina's total coastal waters and contributes

23% of total commercial landings from state waters and 15% of recreational landings (DMF 2011; DMF unpublished data). Blue crab, brown and white shrimp, flounder and striped mullet are the top species commercially landed. Spotted seatrout, southern flounder, bluefish, spot and sheepshead were the most common recreational species caught and discarded in 2011 (DMF unpublished data). The extensive riverine and estuarine wetland communities, shallow nursery areas, diversity of habitats and salinity regimes provide for the disproportionately high productivity of the sound. Habitat features include extensive submerged aquatic vegetation (SAV) beds along the Outer Banks and along the rivers, intertidal and subtidal oysters and primary nursery areas. Pamlico Sound is a focal point for the shrimp, crab, and oyster fisheries as well as for other fisheries, both commercial and recreational, targeting southern flounder, spotted seatrout and red drum. All of these characteristics combined make this system important ecologically, economically and socially to the citizens of North Carolina.

The estuarine otter trawl fishery is the most efficient way to harvest shrimp in North Carolina and makes the shrimp fishery the second most valuable commercial fishery in the state behind the blue crab fishery (1994-2011 average shrimp dockside value \$9.8 million). The majority (89%) of the estuarine shrimp harvest in North Carolina comes from otter trawls. However, major concerns of otter trawls are the capture of non-target species and mortality associated with discarded fish. The amount of dead discards in the fishery may have an influence on the amount of resources available to recreational and commercial fishing. Economically valuable finfish species such as Atlantic croaker, southern flounder, spot, and weakfish are of special concern to all fishermen in North Carolina.

However, as noted in Section 6.3.5 of the draft Shrimp Revision, the resource conservation issues for these latter three species are for species that are covered in the North Carolina Interjurisdictional (IJ) FMP that selectively adopts management measures contained in approved federal council or ASMFC FMPs by reference as minimum standards. So, even with the stated goal of this shrimp amendment to 'minimize harvest of non-target species of finfish and crustaceans and protected, threatened, and endangered species, the extent and benefit of actions to be considered must be viewed in this broader ASMFC coast wide context.

Since 1978 almost one million acres of estuarine waters have been closed to trawling through fishery nursery area designations (primary and secondary nursery areas), military danger zones and restricted areas, and trawl net prohibited areas. This is approximately 45 percent of the estuarine waters. Another 65,000 acres of estuarine waters are closed some time during the year, either due to shrimp size management or areas classified as special secondary nursery areas (SSNA).

IV. AUTHORITY

§ 113-134. Rules.

§ 113-182. Regulation of fishing and fisheries

V. DISCUSSION

Prohibiting Shrimp Trawl Nets in Internal Coastal Waters

Area restrictions for trawling have been used to address allocation, resource protection, habitat protection and safety issues. The 2006 Shrimp Fishery Management Plan evaluated area restrictions as an option to reduce bycatch. During development of the 2006 FMP, area prohibitions were implemented and included closures in the IWW in the Wrightsville Beach area,

in the bays south of Fort Fisher and Bald Head Creeks, White Oak River above Hancock Point, the SSNA in Newport River, the banks side between Drum Inlet to Wainwright Island, Neuse River above Wilkinson Point, Pamlico River above Pamlico Point, and Pungo River above Wades Point. Approximately 92,000 acres of water were closed through implementation of the 2006 plan.

The value of shrimp trawl landings from estuarine waters have ranged from over \$14,000,000 in 2002 to around \$2,000,000 in 2005 and makes up between 75% and 80% of all shrimp landings in North Carolina (Table 12.34). Participation in the estuarine fishery has dropped approximately 66% since 1995 with effort in number of trips dropping approximately 81% since 1995 (Table 12.34).

Table 12.34 Number of pounds, trips, value and participation in the estuarine shrimp trawl fishery.

Estuarine Shrimp Trawl Fishery				
Year	Pounds	Participants	Trips	Value
1994	5,240,153	845	14,585	\$13,797,757
1995	5,729,152	888	15,482	\$13,759,068
1996	3,055,860	705	11,008	\$7,809,425
1997	4,911,799	722	12,702	\$12,958,128
1998	2,019,600	513	8,297	\$4,473,965
1999	5,275,158	667	10,817	\$12,928,539
2000	7,847,702	793	10,521	\$19,585,614
2001	3,493,218	553	7,734	\$8,506,491
2002	7,511,154	639	10,030	\$14,159,626
2003	3,179,629	439	6,682	\$6,011,535
2004	2,581,743	421	5,358	\$5,523,421
2005	1,078,088	272	2,890	\$2,016,414
2006	2,891,435	297	3,255	\$5,059,891
2007	7,123,976	338	4,465	\$13,595,395
2008	6,764,108	364	4,206	\$13,516,404
2009	4,049,599	340	3,890	\$6,452,588
2010	4,280,703	355	3,946	\$7,649,074
2011	3,889,637	301	3,004	\$8,178,854

While the declining value of shrimp, increasing market share of imported shrimp, regulatory changes and increased fuel prices have contributed to the decline in effort, prohibiting estuarine shrimp trawling would be detrimental to North Carolina's shrimp fishery. The closure of estuarine waters would result in the loss of the economic value to dealers, harvesters, and support industries through decreased revenue and income. In addition, less local North Carolina shrimp would be available to the public causing a higher dependence on shrimp landed from out of state and on those shrimp imported from other countries.

Prohibiting Shrimp Trawl Nets in Pamlico Sound and Adjacent Tributaries

Trawling in the Pamlico Sound has been controversial because of bycatch and discard of valuable juvenile and adult finfish. Pamlico Sound landings from shrimp trawls average 81% of internal coastal shrimp trawl landings since 1994. The amount of bycatch varies greatly from fisherman to fisherman, trip to trip and even tow to tow. Factors that influence bycatch include water temperature, water clarity, fishing location, amount of bycatch, tow time and gear configuration. Brown (2010) conducted a short term characterization study of the shrimp trawl fishery in the Pamlico Sound. During this six month study, conducted from July to December 2009, shrimp made up 23% of the total catch by weight. This study represents a 6-month snapshot in time of the Pamlico Sound shrimp fishery making this study temporally limited to that one summer and one fall season.

Atlantic croaker accounted for approximately 33% of the catch by weight, with spot and weakfish accounting for 13% and 6%, respectively. The majority of Atlantic croaker and spot were harvested in the summer months in the double seamed and four seamed trawls which are used to target brown shrimp. Other commercially and recreationally important species observed include southern, summer and gulf flounders species representing 1% of the catch by weight, as well as kingfishes and spotted sea trout representing 0.8% and 0.02%, respectively. Atlantic croaker had the largest amount of unmarketable discards by weight with all being discarded. Spot made up the second largest component of the unmarketable bycatch, 99% of the spot landed were discarded. All of the weakfish caught were undersized; length frequency distributions of discarded weakfish ranged from 70 mm to 150 mm. Weakfish represented the largest regulatory discards by weight for all net types.

The DMF has conducted a stratified-random trawl survey in the Pamlico Sound uninterrupted for twenty-five years. Sampling began in 1987 and was conducted over two weeks during the months of March, June, September, and December from 1987 to 1989. In 1990, sampling occurred over two weeks during the months of March, June, and September. From 1991 to present, the Pamlico Sound Survey has been conducted during the middle two weeks in June and September. From 1990-present, the sample area covers all of Pamlico Sound and its bays, Croatan Sound up to the Highway 64 Bridge, the Pamlico River up to Blounts Bay, the Pungo River up to Smith Creek, and the Neuse River up to Upper Broad Creek (DMF 2012).

The primary objective of this survey is to provide a multi-species long-term index of abundance for juvenile fish in Pamlico Sound and its coastal rivers. Data are used to calculate abundance indices for several recreationally and commercially significant species in Pamlico Sound, including: Atlantic croaker, spot, and weakfish and are produced annually. These juvenile abundance indices (JAI) estimates have been used in both state and federal stock assessments and management plans. This survey also provides data to evaluate other potential critical habitat areas in Pamlico Sound (DMF 2012).

Table 12.35 Weighted CPUE, standard error (SE), total number collected (n), mean size (mm), and size range for select species during 2011 in the Pamlico Sound Survey (DMF 2012).

Common Name	Month	CPUE	SE	n	Mean Size (mm)	SE	Min (mm)	Max (mm)
spot	Jun	552.0	66.3	30,396	106.4	0.6	58	188
	Sep	278.2	38.9	17,822	133.2	0.4	93	231
	All	415.1	40.4	48,218	116.3	0.4	58	231
Atlantic croaker	Jun	177.5	25.4	10,769	131.0	0.9	52	237
	Sep	118.8	15.6	5,581	169.1	0.3	145	200
	All	148.1	15.7	16,350	144.0	0.7	52	237
weakfish	Jun	37.9	11.4	1,908	197.7	1.3	52	288
	Sep	35.8	11.2	1,808	149.0	1.4	57	290
	All	36.9	7.9	3,716	174.0	1.2	52	290

In 2011, spot was the most abundant target species with an annual weighted CPUE of 415.1 individuals per sample (Table 12.35). Lengths ranged from 58-231 mm FL with a mean size of 116.3 mm FL. Atlantic croaker was the second most abundant target species with an annual weighted CPUE of 148.1 individuals per sample (Table 12.35). Lengths ranged from 52-237 mm TL with a mean size of 131.0 mm TL. Weakfish was the fourth most abundant target species behind brown shrimp with an annual weighted CPUE of 36.9 individuals per sample (Table 122). Lengths ranged from 52-290 mm TL with a mean length of 174.0 mm TL (DMF 2012.35).




Figure 12.50 Weighted CPUE of Atlantic croaker, spot, and weakfish from the NCDMF Pamlico Sound Survey and number of shrimp trawl trips (effort lagged 1 year) in Pamlico Sound and Neuse, Pamlico and Pungo rivers (DMF 2012).

Abundances of Atlantic croaker and spot are variable with all three species showing overall increases in CPUEs from 1991 to 2010 and dropping again in 2011 (Figure 12.50). Shrimping effort in Pamlico Sound has decreased by 65% since 1995 (Figure 12.50). Regression analysis indicates that there are no significant relationships of decreasing commercial trawling effort with CPUE of Atlantic croaker ($r^2=0.04$; $p=0.46$), spot ($r^2=0.02$; $p=0.63$) and weakfish ($r^2=0.03$; $p=0.50$), suggesting that other factors may influence juvenile abundances of these three finfish in Pamlico Sound. However, the number of trips used here is the number of trip tickets recorded by the Trip Ticket Program and does not take into account the number of tows, the number of trawls used, the amount of headrope, or the number of days fished per trip ticket.

The identification and designation of Strategic Habitat Areas (SHA) for marine and coastal fishery species is a critical component in the implementation of the North Carolina's approved Coastal Habitat Protection Plan (CHPP). SHAs are defined in the CHPP as specific locations of individual fish habitat or systems of habitats that have been identified to provide exceptional habitat functions or that are particularly at risk due to imminent threats, vulnerability, or rarity (Deaton et al. 2010). The nomination process is meant to identify a possible subset of high quality areas that will maintain the diversity of habitats, species and ecological functions found within the overall ecosystem (Deaton et al. 2010). The Pamlico Sound Region or Region 2 was examined for SHA identification and nomination through a scientifically based process using biological data and a consensus based approach of a regional expert panel beginning in early

2010 and was completed in 2011 (DMF 2011). During this process, fish abundance data from DMF's Pamlico Sound Survey, described above and covering the Pamlico Sound, and the Neuse, Pamlico and Pungo rivers were included as a data layer in the analysis. Based on a series of statistical analysis, two groups of species were used as the basis of creating a data layer of fish abundances. One group included spot, croaker, pinfish, pigfish, hogchoker, southern flounder, harvestfish, weakfish, blue crab, silver perch, and white shrimp, while the other group included fringed flounder, planehead filefish, mantis shrimp, spadefish, southern kingfish, striped anchovy, lesser blue crab, bay whiff, summer flounder, inshore lizardfish, pink shrimp and brown shrimp. The fish and habitat data were used as targets in a site selection Software program to select a subset of areas containing a diversity of high quality biological features (DMF 2011).




Figure 12.51 Strategic Habitat Area nominations and existing protected areas (DMF 2011)

Figure 12.51 illustrates those areas that have been nominated. The majority of the nominated areas occur along the edge of the sound including or adjacent to existing Primary and Secondary Nursery Areas, and the mouths of the rivers. These areas were selected because of their proximity to documented oyster habitat and/or their relatively high abundance of fish, based on the DMF data. Also with the inclusion of all designated PNA as a resource target, the model was influenced to provide connectivity with these PNA targets. The selected SHAs located in the center of Pamlico Sound consisted of soft bottom in waters greater than six feet in

depth that had relatively higher fish abundance, and trawling was the only documented alteration. Those SHAs located toward the mouths of the Pamlico and Neuse Rivers had subtidal oysters and SAV with bottom disturbing gear listed as the major alteration (DMF 2011).

The relatively greater amount of area selected as SHAs along the perimeter of the sound was due to the greater diversity of shallow productive habitats in those locations that support juvenile fish. These areas were also considered at greater risk from nearby activities that affect water quality such as development, marinas, and wetland ditching. In contrast, the center of Pamlico Sound had lower habitat diversity and fewer documented threats. These results indicate that the edges of Pamlico Sound, where benthic habitats and juvenile fish are more concentrated, may merit further protection from bottom disturbing fishing gear than the center of Pamlico Sound.

Prohibiting Shrimp Trawl Nets in Special Secondary Nursery Areas

Nursery Areas are fish habitat areas that for reasons such as food cover bottom type, salinity, temperature and other factors, young finfish and crustaceans spend the major portion of their initial growing season (15A NCAC 03I .0101(4)(f)). SNAs are those areas in the estuarine system where later juvenile development takes place. Populations are composed of developing sub-adults of similar size which have migrated from an upstream primary nursery area of the secondary nursery area located in the middle portion of the estuarine system. There are specific gear protections for designated PNAs such as the prohibition of the use of trawls, dredges, long haul, swipe seines, and mechanical methods for oysters and clams and the prohibition of trawls in SNAs.

Special Secondary Nursery Areas are SNAs where trawling may be allowed by the director through proclamation authority from August 16 through May 14 (Table 12.36). This enables fishermen to catch any shrimp late in the season that have not migrated out into the larger estuaries. There are approximately 37,400 acres of SSNAs located in Roanoke Sound, Pamlico and Pungo rivers, West Bay, Core Sound, North River, Newport River, New River, Chadwick Bay, IWW in Onslow/Pender County, Cape Fear River, Lockwood Folly River and Saucepan Creek in the Shallotte River. Of these areas, SSNAs in the Pamlico and Pungo rivers have not been open since 1990 (Table 12.36). Other areas in Pamlico and Pungo Rivers were reclassified as permanent secondary nursery areas because of having never been opened (Table 12.36). The North River SSNA was permanently closed in 1997 and Newport River SSNA was permanently closed through the 2006 Shrimp FMP. Both of these closures were due to the constant movement of lines. Permanent lines were established to eliminate this. Cape Fear, Lockwood Folly and Saucepan Creek SSNA also have not opened for many years.

In the 2006 FMP, it was recommended that Chadwick Bay be investigated to determine if it functioned as a secondary nursery area. Through DMF sampling, it was determined Chadwick Bay was a SSNA and closed by rule from May 15th through August 15th in April 2011.

Prohibiting shrimp trawls in SSNAs would eliminate bycatch in those areas and allow further protection of those juvenile finfish and shrimp using those areas before migration out into the sounds and ocean.

Table 12.36 Current and past designated special secondary nursery areas.

Current Rule ID 03R .0105	Description	Year Designated (reclassified)	Latest Year Opened	Proc Ref.	Comment
1 (a)	Outer Shallowbag Bay	2004	2013	SH-1-2013	Opened for peeler crab trawling. Will likely open August 18 for shrimp trawling
1 (b)	Kitty Hawk/Buzzard Bay	2004	2013	SH-1-2013	Opened for peeler crab trawling
2 (a)	Pungo Creek	1989	1990	SH-22-90	
2 (b)	Scranton Creek	1989	1990	SH-22-90	
2 (c)	Slade Creek	1989	1990	SH-22-90	
2 (d)	South Creek	1989	1990	SH-22-90	
2 (e)	Bond Creek	1989	1990	SH-22-90	
3 (a)	West Thorofare Bay	1986	2012	SH-15-2012	
3 (b)	Long Bay	1986	2012	SH-15-2012	
3 (c)	Turnagain Bay	1991	2012	SH-15-2012	
4 (a)	Cedar Island Bay	1986	2012	SH-15-2012	
4 (b)	Thorofare Bay	1986	2012	SH-15-2012	
4 (c)	Nelson Bay	1986	2012	SH-15-2012	
4 (d)	Brett Bay	1986	2012	SH-15-2012	
4 (e)	Jarrett Bay	1986	2012	SH-15-2012	
5 (a)	North River	1986	1997	SH-11-97	Closed through public negotiation
5 (b)	Ward Creek	1986	1997	SH-11-97	Closed through public negotiation
6	Newport River	1991	2006	SH-5-2006	Closed 2006 FMP
7	New River	1994	2012	SH-8-2012	
8	Chadwick Bay	2011	2012	SH-8-2012	Recommended in the 2006 FMP to investigate if functioned as a SSNA
9	IWW	1994	2012	SH-16-2012	
10	Cape Fear	1986	1987?	None?	
11	Lockwood Folly River	1986	1987?	None?	
12	Saucepan Creek	1986	1987?	None?	
03R .0104 3(c)	Upper Pamlico River	1989 (2004)	None	None	Reclassified to SNA
03R .0104 3(a)	Upper Pungo River	1989 (2004)	None	None	Reclassified to SNA
03R .0104 4(d)	Upper Broad Creek	1989 (2004)	None	None	Reclassified to SNA
03R .0104 4(e)	Goose Creek	1989 (2004)	None	None	Reclassified to SNA

Prohibiting Shrimp Trawl Nets in Brunswick County

The Brunswick County coastline stretches for approximately 33 miles and is bound by the Cape Fear River Inlet on the east end and by the Little River Inlet on the west end. Four barrier islands, all of which are densely developed are separated by five inlets along the coastline. The IWW channel from Sunset Beach Bridge to the South Carolina state line, including Eastern Channel, lower Calabash River and Shallotte River are frequently closed to trawling because of the abundance of small shrimp (Figure 12.52). The Shallotte River has been closed to shrimping since 1998, because shrimp rarely reach a large enough size to open and tend to remain at a 60-count (heads on) or greater size (Figure 12.52). The Division recommended that this area be permanently closed by rule during the development of the 2006 Shrimp FMP because of the abundance of small shrimp, but the Advisory Committee recommended *status quo* with the resulting management strategy being *status quo* (DMF 2006). Eastern Channel (Gause Creek), lower Calabash River, and Shallotte River have not been opened in recent years.

Logothetis and McCuiston (2006) described and quantified bycatch in the southern region of North Carolina, including the IWW in Brunswick County from April through November in 2004 and 2005. Bycatch generally were composed of juvenile to sub-adult fish with bycatch rates highest in April and May. Salinities were also found to affect bycatch rates with bycatch increasing with increasing salinity. The bycatch ratio reported was 0.68 in Brunswick County, meaning for every 1.0 lb of shrimp caught, there was 0.68 lb of bycatch caught. Although bycatch does not appear to be high, based on this ratio, the continuing occurrence of small shrimp in these areas warrants the consideration of making this area a permanent closure.




Figure 12.52 Brunswick County Shrimp Management Areas.

VII. EVALUATION MATRIX

AC Evaluation of Area Restrictions to Reduce Shrimp Trawl Bycatch in North Carolina's Internal Coastal Waters

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	
1. <i>Status quo.</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. Continued reduction in effort may result in overall bycatch reduction in the fleet. N	Will not create shifts in effort to other fisheries. Maintains present market value of fishery. N	Allows flexibility of fishermen to continue to fish in their normal areas using their normal gears. N	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. Not Evaluated	No change in rule. Not Evaluated	Not Evaluated
2. Prohibit all shrimp trawling in all internal coastal waters.	Eliminates bycatch from shrimp trawls and skimmer trawls in all Internal Coastal waters. +	Severe reduction of the second most commercially valuable fishery in the state. Will likely create effort shifts into other fisheries. Will likely create economic losses to coastal fishing communities as well as shrimp fishermen, dealers, and related industries. -	Loss of a historical fishery. May be perceived by some of the public as a step forward in improved bycatch reduction. Will likely be perceived by the commercial public as unjust management. Greatly reduces the availability of NC caught shrimp. -	Will likely create conflicts with other commercial and recreational fishermen due to shift in effort. May improve recreational fishing for croaker, spot, and weakfish. +/-	Initial increased level of enforcement. Not Evaluated	Will require a rule change. Not Evaluated	Not Evaluated
3. Prohibit all shrimp trawling in Pamlico Sound and tributaries.	Eliminates bycatch from shrimp trawls and skimmer trawls in Pamlico Sound. +	Loss of a valuable commercial fishery. Will likely create effort shifts. Will likely create economic losses to coastal fishing communities as well as shrimp fishermen, dealers, and related industries. -	Loss of a historical fishery. Perceived by some of the public as a step forward in improved bycatch reduction. Will likely be perceived by the commercial public as unjust management. Greatly reduces the availability of NC caught shrimp. -	Will likely create conflicts with other commercial and recreational fishermen due to shift in effort. May or may not improve recreational fishing for croaker, spot, and weakfish. +/-	Initial increased level of enforcement. Not Evaluated	Will require a rule change. Not Evaluated	Not Evaluated

AC Evaluation of Area Restrictions to Reduce Shrimp Trawl Bycatch in North Carolina's Internal Coastal Waters

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		Other Impacts
	Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	
4. Prohibit all shrimp trawling in portions of Pamlico Sound to allow a buffer from sensitive habitats.	Eliminates bycatch from shrimp trawls and skimmer trawls along edge of PNAs and SNA in Pamlico Sound.	Possible loss of some income to smaller boats. Shrimp crop will filter out possibly resulting in larger shrimp.	May gain public support. Industry view may be mixed.	May or may not improve recreational fishing for croaker, spot, and weakfish.	Closure lines may be difficult to enforce.	May be implemented through proclamation.	
	+	+/-	N	+/-	Not Evaluated	Not Evaluated	Not Evaluated
5. Prohibit all shrimp trawling in Special Secondary Nursery Areas.	Eliminates bycatch from shrimp trawls and skimmer trawls in Special Secondary Nursery Areas.	Loss of income in latter part of season. Shrimp crop will filter out resulting in larger shrimp.	May gain public support. Industry view may be mixed. Eliminates grand openings.	Will likely create conflicts with other commercial fishermen due to shift in effort. May or may not improve recreational fishing for croaker, spot, and weakfish.	Initial increased level of enforcement.	Will require a rule change.	
<i>Decreasing the duration that SSNAs are open may be another option. See Effort paper issue paper</i>	+	+/-	+/-	+/-	Not Evaluated	Not Evaluated	Not Evaluated
6. Prohibit shrimp trawling in the IWW channel from Sunset Beach Bridge to the South Carolina state line, including Eastern Channel, lower Calabash River and Shallotte River.	Minimal decrease in bycatch. No waste of small shrimp.	Some loss of income to fishermen.	Minimal impact.	May or may not improve recreational fishing for croaker, spot, and weakfish.	Increased level of enforcement.	Will require a rule change.	
	+	-	N	+	Not Evaluated	Not Evaluated	Not Evaluated

DMF Evaluation of Area Restrictions to Reduce Shrimp Trawl Bycatch in North Carolina's Internal Coastal Waters

Impacted Group	Impacted Group			Impacted Group	Impacted Group		Impacted Group
Management Option	Management Option	Management Option	Management Option	Management Option	Management Option	Management Option	Management Option
1. <i>Status quo.</i>	Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. -	Will not create shifts in effort to other fisheries. Maintains present market value of fishery. N	Allows flexibility of fishermen to continue to fish in their normal areas using their normal gears. N	Commercial and recreational fishing will continue with no changes in gear use or conflict. N	Same level of enforcement. N	No change in rule. N	
2. Prohibit all shrimp trawling in all internal coastal waters.	Eliminates bycatch from shrimp trawls and skimmer trawls in all Internal Coastal waters +	Severe reduction of the second most commercially valuable fishery in the state. Will likely create effort shifts into other fisheries. Will likely create economic losses to coastal fishing communities as well as shrimp fishermen, dealers, and related industries. -	Loss of a historical fishery. May be perceived by some of the public as a step forward in improved bycatch reduction. Will likely be perceived by the commercial public as unjust management. Greatly reduces the availability of NC caught shrimp. -	Will likely create conflicts with other commercial and recreational fishermen due to shift in effort. -	Initial increased level of enforcement. -	Will require a rule change. -	
3. Prohibit all shrimp trawling in Pamlico Sound and tributaries.	Eliminates bycatch from shrimp trawls and skimmer trawls in Pamlico Sound. +	Loss of a valuable commercial fishery. Will likely create effort shifts. Will likely create economic losses to coastal fishing communities as well as shrimp fishermen, dealers, and related industries. -	Loss of a historical fishery. Perceived by some of the public as a step forward in improved bycatch reduction. Will likely be perceived by the commercial public as unjust management. Greatly reduces the availability of NC caught shrimp -	Will likely create conflicts with other commercial and recreational fishermen due to shift in effort. -	Initial increased level of enforcement. -	Will require a rule change. -	

DMF Evaluation of Area Restrictions to Reduce Shrimp Trawl Bycatch in North Carolina's Internal Coastal Waters

Impacted Group	Impacted Group			Impacted Group	Impacted Group		Impacted Group
Management Option	Management Option	Management Option	Management Option	Management Option	Management Option	Management Option	Management Option
4. Prohibit all shrimp trawling in portions of Pamlico Sound to allow a buffer from sensitive habitats.	Eliminates bycatch from shrimp trawls and skimmer trawls along edge of PNAs and SNA in Pamlico Sound. +	Possible loss of some income. Shrimp crop will filter out possibly resulting in larger shrimp. -/+	May gain public support. Industry view may be mixed. +	May reduce conflict . +	Closure lines may be difficult to enforce. -	May be implemented through proclamation. +	
5. Prohibit all shrimp trawling in Special Secondary Nursery Areas. <i>Decreasing the duration that SSNAs are open may be another option. See Effort paper issue paper</i>	Eliminates bycatch from shrimp trawls and skimmer trawls in Special Secondary Nursery Areas. +	Loss of income in latter part of season. Shrimp crop will filter out resulting in longer season and larger shrimp. -/+	May gain public support. Industry view may be mixed. Eliminates grand openings. +	Will likely create conflicts with other commercial and recreational fishermen due to shift in effort. -	Initial increased level of enforcement. -	Will require a rule change. -	.
6. Prohibit shrimp trawling in the IWW channel from Sunset Beach Bridge to the South Carolina state line, including Eastern Channel, lower Calabash River and Shallotte River.	Minimal decrease in bycatch. No waste of small shrimp. +	Some loss of income to fishermen. -	Minimal impact. -	No impact. N	Increased level of enforcement. -	Will require a rule change. -	

12.9 REMOVAL OF THE SHRIMP TRAWL FROM THE RECREATIONAL COMMERCIAL GEAR LICENSE

I. ISSUE

The consideration of eliminating the shrimp trawl as an authorized gear from the Recreational Commercial Gear License

II. ORIGINATION

The Shrimp Fishery Management Plan Advisory Committee at its May 15, 2013 meeting.

III. BACKGROUND

On August 14, 1997, the Fisheries Reform Act (FRA) was signed into law. One aspect of this law was the creation of the Recreational Commercial Gear License (RCGL). According to the Fisheries Moratorium Steering Committee (MSC), a group that provided the recommendations for the FRA, the purpose of creating this license was to: (1) allow individuals and families who have traditionally accessed the State's public trust fishery with commercial gear to supply themselves with fresh seafood; (2) limit the effort that may be expended by this class of fishermen both individually and as a group; and (3) implement the principle that all persons who harvest state public trust resources pay for that privilege by investing in coastal fisheries conservation and management (Moratorium Steering Committee, 1996). A statutory sunset clause in the FRA was put in place to test the RCGL concept, as well as other license types and would have expired if unsuccessful. DMF began selling this license July 1, 1999.

RCGL allows recreational fishermen to use limited amounts of commercial gear to harvest seafood for their personal consumption including a shrimp trawl with a maximum headrope length of 26 feet. Seafood harvested under this license cannot be sold. RCGL holders are limited to the same bag and size limits as Coastal Recreational Fishing License (CRFL) holders. The 2006 Shrimp Fishery Management Plan (FMP) added two new allowable RCGL gears, one shrimp pound and a 26 foot skimmer trawl. The FMP also limited all recreational harvesters, including RCGL holders to 48 quarts of head-on (32 quarts of head-off) shrimp per day, greatly reducing the harvest in some areas. If there are two valid license holders on board a vessel, then the shrimp possession limit may be doubled. The MFC also passed a rule allowing mechanical retrieval of the gear as long as a Turtle Excluder Device was properly installed in the trawl; prior to the FMP, shrimp trawls could only be retrieved by hand.

Many of the species taken by recreational users of commercial gear are included in fisheries management plans. Until 2002, the influence that RCGL holders may have on these species was unknown. Two surveys were used to collect information from RCGL holders; a socioeconomic survey, conducted in 2001, 2004, and 2007, and catch and effort surveys conducted monthly from 2002 through 2008. Both of these surveys were terminated in 2008 due to budget constraints.

IV. AUTHORITY

§ 113-134. Rules.

§ 113-173. Recreational Commercial Gear License.

§ 113-182. Regulation of fishing and fisheries.

IV. DISCUSSION

With the exception of 2002, the number of RCGLs sold on a fiscal basis has declined each year from 2000 through 2011 (Table 12.37); with a 29% decline overall. The largest single year decline occurred in 2011 (12%) followed by 2001 (8%). In 2009 and 2010 there was an average of 3.35% increase in sales. Twenty-five counties consistently comprise approximately 85% of the total number of RCGLs purchased each year.

Table 12.37 Number of license sales of Recreational Commercial Gear Licenses, 2002 through 2011. (fiscal year, July 1 through June 30).

Fiscal Year	Number of RCGLs Sold	Percent Change from Previous Sales Year
2000	6,740	
2001	6,202	-8.0
2002	6,300	1.6
2003	6,157	-2.3
2004	5,868	-4.7
2005	5,653	-3.7
2006	5,368	-5.0
2007	5,134	-4.4
2008	5,113	-0.4
2009	5,280	3.3
2010	5,458	3.4
2011	4,802	-12.0

Typical RCGL holders were married Caucasian males with an average age of 56. Findings from license sales statistics and the three socioeconomic surveys conducted in 2001, 2004, and 2007 indicated that coastal counties, in particular, southern coastal counties, substantially contributed to the overall number of RCGL holders.

The top three gears used by RCGL holders fishing in all regions of the coast were crab pot, small mesh gill net, and large mesh gill net. Shrimp trawls were the fourth most common gear used in the Pamlico, Southern, and Central Regions while fish pots were the fourth most common gear used in the Northern Region. On average the highest number of trips using shrimp trawls from 2002 to 2008 occurred in the Pamlico region, followed by the southern region, the central region, and the northern region (Table 12.38). In the Pamlico region, the number of trips ranged from 1,127 (2005) to 2,384 (2002), averaging 1,642 per year from 2002 to 2008. In the southern region, the number of trips ranged from 355 (2007) to 1,123 (2002), averaging 586 trips per year. An average of 413 trips a year were made in the central region, ranging from 132 (2008) to 1,070 (2002). In the Northern region, the number of trips ranged from 50 (2006) to 911 (2004). Overall, the highest number of trips made by RCGL holders using shrimp trawls was observed in 2002; the lowest was observed in 2007.

RCGL holders harvested an average of 52,352 pound of shrimp a year from 2002 to 2008 (Table 12.39). The highest landings occurred in 2002 (101,766 lb), followed by 2008 (54,359 lb) and 2003 (50,961 lb). RCGL holders harvested an average of 16.8 pounds of shrimp per trip from 2002 to 2008 (Table 3). The highest number of pounds of shrimp per trip was observed in 2009 (22.3 lb/trip), followed by 2006 (20.3 lb/trip) and 2002 (19.1 lb/trip).

Table 12.38 Number of trips by shrimp trawl by region, 2002-2008.

Year	Region				Total
	Southern	Central	Pamlico	Northern	
2002	1,123	1,070	2,384	742	5,319
2003	711	246	1,448	348	2,753
2004	392	318	2,122	911	3,743
2005	553	365	1,127	387	2,432
2006	471	464	1,441	50	2,426
2007	355	295	1,510	69	2,229
2008	500	132	1,464	337	2,433
Mean	586	413	1,642	406	3,048

Table 12.39 Harvest (lb) and pounds per trip of shrimp by RCGL gear from 2002-2008.

Year	Pounds	Pounds/trip
2002	101,766	19.1
2003	50,961	18.5
2004	43,698	9.3
2005	32,542	13.4
2006	49,362	20.3
2007	33,778	15.2
2008	54,359	22.3
Mean	52,352	16.8

When compared to North Carolina's commercial shrimp harvest statistics from the NC Trip Ticket Program the average yearly RCGL harvest contributes a little over 1.0% to the overall harvest of shrimp, but contribute 37% of the shrimp trawl effort in number of trips (Table 12.40).

Table 12.40 Commercial estuarine shrimp trawl harvest and Recreational Commercial shrimp trawl harvest, 2002-2008.

Year	Estuarine Commercial		RCGL	
	Pounds	Trips	Pounds	Trips
2002	7,511,154	10,030	101,766	5,319
2003	3,179,629	6,682	50,961	2,753
2004	2,581,743	5,358	43,698	3,743
2005	1,078,088	2,890	32,542	2,432
2006	2,891,435	3,255	49,362	2,426
2007	7,123,976	4,465	33,778	2,229
2008	6,764,108	4,206	54,359	2,433
Mean	4,447,162	5,269	52,352	3,048

However, it should be noted that the vast majority of the RCGL trips are single day trips, while commercial trip duration is quite variable depending on the location fished. Overall commercial trip duration has consistently averaged slightly more than 2 days across all areas.

Limited discard information is available for blue crab, Atlantic croaker, spot, flounder and shrimp which was consistently reported by those RCGL surveyed (Table 12.41). Several other species including Atlantic menhaden, pigfish, white perch, American shad, speckled trout, Spanish mackerel, sharks and rays reported too sporadically for an estimate.

Table 12.41 Recreational Commercial trawl discard numbers and trips, 2002-2008.

Year	Blue Crab		Atlantic Croaker		Spot		Flounder		Shrimp	
	Trips	Discarded	Trips	Discarded	Trips	Discarded	Trips	Discarded	Trips	Discarded
2002	3,301	96,404	560	26,197	322	9,949	2,011	9,949	5,050	1,397
2003	1,723	34,819	530	17,100	571	4,868	1,000	4,868	2,737	6,273
2004	2,583	39,480	781	21,438	611	12,896	996	12,896	3,655	4,313
2005	1,685	36,602	398	11,959	235	1,631	396	1,631	2,437	1,102
2006	1,448	11,875	582	34,605	476	18,339	605	18,339	2,352	13,028
2007	917	8,394	296	7,362	289	3,149	372	3,149	2,048	4,939
2008	1,172	29,301	256	8,161	270	8,013	540	8,013	2,252	6,165

Resource or conflict issues related to the RCGL since its implementation have been minimal. There have been instances, as with all gear, where the user was not acting responsibly. Reports to the DMF have ranged from shrimpers harvesting over the legal limit, improperly marked gear, and the illegal sale of RCGL harvested shrimp.

VII. EVALUATION MATRIX

DRAFT

AC Evaluation of Removal of the shrimp trawl from the recreational commercial gear license							
Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
<p>1. <i>Status quo</i></p> <p>(As evaluated in skimmer trawls and other gear paper on May 15, 2013)</p>	<p>Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. Effort reduction has resulted in reduced bycatch and will continue. Gear is more effective, even if effort is reduced.</p> <p style="text-align: center;">+/-</p>	<p>Will not create shifts in effort to other fisheries. Maintains present market value of fishery.</p> <p style="text-align: center;">+/-</p>	<p>Allows flexibility of use of gears in the fishery.</p> <p style="text-align: center;">+</p>	<p>Commercial and recreational fishing will continue with no changes in gear use or conflict.</p> <p style="text-align: center;">Not evaluated</p>	<p>Same level of enforcement.</p> <p style="text-align: center;">Not evaluated</p>	<p>Continued proclamation authority. No rule change needed.</p> <p style="text-align: center;">Not evaluated</p>	<p>Allows for further characterization and bycatch reduction studies to fill data gaps prior to new regulations.</p> <p style="text-align: center;">Not evaluated</p>
<p>2. Eliminate otter trawls as an authorized gear of the Recreational Commercial Gear License</p> <p>AC Voted to not evaluate after presentation</p>	<p>Minimal impact to the existing amount of bycatch because of smaller percentage of fishery.</p>	<p>Loss of a food source thereby possibly increasing food expense. Value and function of gear purchased for RCGL shrimp fishery will be lost.</p>	<p>Removes ability of recreational fishermen to efficiently harvest larger quantities of shrimp for personal consumption.</p>	<p>May cause shift into other RCGL gear users.</p>	<p>Same level of enforcement.</p>	<p>Rule change required. Loss of license fees but possible increase in Standard Commercial Fishing License fees.</p>	

DMF Evaluation of Removal of the shrimp trawl from the recreational commercial gear license

Impacted Group	Shrimp Fishery			Other Fisheries	Agency		
Management Option	Bycatch Reduction Impact	Economic Impact	Social Impact	Inter-fishery Impact	Enforceability	Authority/ Administrative	Other Impacts
<p>1. <i>Status quo</i></p> <p>(As evaluated in skimmer trawls and other gear paper on May 15, 2013)</p>	<p>Continues the existing amount of bycatch and bycatch mortality in the shrimp fishery. Effort reduction has resulted in reduced bycatch and will continue. Gear is more effective, even if effort is reduced.</p> <p align="center">+/-</p>	<p>Will not create shifts in effort to other fisheries. Maintains present market value of fishery.</p> <p align="center">+/-</p>	<p>Allows flexibility of use of gears in the fishery.</p> <p align="center">+</p>	<p>Commercial and recreational fishing will continue with no changes in gear use or conflict.</p> <p align="center">Not evaluated</p>	<p>Same level of enforcement.</p> <p align="center">Not evaluated</p>	<p>Continued proclamation authority. No rule change needed.</p> <p align="center">Not evaluated</p>	<p>Allows for further characterization and bycatch reduction studies to fill data gaps prior to new regulations.</p> <p align="center">Not evaluated</p>
<p>2. Eliminate otter trawls as an authorized gear of the Recreational Commercial Gear License</p>	<p>Minimal impact to the existing amount of bycatch because of smaller percentage of fishery.</p> <p align="center">+</p>	<p>Loss of a food source thereby possibly increasing food expense. Value and function of gear purchased for RCGL shrimp fishery will be lost.</p> <p align="center">-</p>	<p>Removes ability of recreational fishermen to efficiently harvest larger quantities of shrimp for personal consumption.</p> <p align="center">-</p>	<p>May cause shift into other RCGL gear users.</p> <p align="center">N</p>	<p>Same level of enforcement.</p> <p align="center">N</p>	<p>Rule change required. Loss of license fees but possible increase in Standard Commercial Fishing License fees</p> <p align="center">-</p>	

DRAFT

12.10 BYCATCH MANAGMENT RECOMMENDATIONS

12.10.1 Trawling in the New River above the Highway 172 Bridge

Advisory Committee Recommendation

Allow skimmer and otter shrimp trawling in the New River special secondary nursery area (above the Highway 172 Bridge).

Division Recommendation

Status quo (Continue to prohibit otter trawls in the New River special secondary nursery area above the Highway 172 Bridge)

Marine Fisheries Commission Preferred Management Strategy

Status quo (Continue to prohibit otter trawls in the New River special secondary nursery area above the Highway 172 Bridge)

12.10.2 Evaluation of the skimmer trawl and other gears used for shrimping in North Carolina

Advisory Committee and Division Recommendation

Allow hand cast netting of shrimp in all closed areas and increase the limit to four quarts, heads on per person. **Division added "heads on"**.

Advisory Committee Recommendation

Require a fishing license from DMF to fish a cast net.

Division Recommendation

Status quo on a license requirement to fish a cast net for shrimp

Marine Fisheries Commission Preferred Management Strategies

Allow hand cast netting of shrimp in all closed areas and increase the limit to four quarts, heads on per person.

Status quo on a license requirement to fish a cast net for shrimp

12.10.3 The use of TEDs in commercial skimmer trawl operations

Advisory Committee Recommendation

Status quo

Division Recommendation

Upon federal adoption of TEDs in skimmer trawls, the division will support the federal requirement (Rule 15A NCAC 03L .0103 (g) allows for state enforcement).

Marine Fisheries Commission Preferred Management Strategy

Upon federal adoption of TEDs in skimmer trawls, the division will support the federal requirement.

12.10.4 Consideration of a commercial live bait shrimp fishery in North Carolina

Advisory Committee Recommendation

Establish a permitted live shrimp bait fishery and for DMF to craft the guidelines and permit fees after reviewing permitted operations in other states.

Division Recommendation

Status quo (continue to manage the live shrimp bait fishery the same as food shrimp fishery).

Marine Fisheries Commission Preferred Management Strategy

Establish a permitted live shrimp bait fishery and for DMF to craft the guidelines and permit fees after reviewing permitted operations in other states.

12.10.5 Gear Modifications in North Carolina shrimp trawls to reduce finfish bycatch

Advisory Committee Recommendations

Allow any federally certified BRD in all NC internal and offshore waters.

Update and certify bycatch reduction devices through the state bycatch reduction program.

Convene an ongoing stakeholder workgroup charged with suggesting new trawl gear or trawl gear modification.

Initiate industry testing of new or modified bycatch reduction devices and gear modifications under the supervision of the DMF. After testing and collection of scientific data, regulations should be implemented to require or allow such devices or modifications to be used in NC internal and offshore waters.

Test a three-inch bar-spaced turtle excluder device to see if it can be certified as a bycatch reduction device.

Allow the shrimp industry a two year period to test bycatch reduction devices.

Division Recommendations

Allow any federally certified BRD in all NC internal and offshore waters.

Update the scientific testing protocol for the state BRD certification program.

Convene a stakeholder group to initiate industry testing of minimum tail bag mesh size, T-90 panels, skylight panels, and reduced bar spacing in TEDs to reduce bycatch to the extent practicable.

- Upon securing funding, testing in the ocean and internal waters will consist of three years of data using test nets compared to a control net with a Florida Fish Eye, a federally approved TED, and a one and a half inch tailbag.
- Results should minimize shrimp loss and maximize reduction of bycatch of finfish. Promising configurations will be brought back to the MFC for consideration for mandatory use.
- This stakeholder group may be partnered with DMF and Sea Grant.
- Members could consist of fishermen, net/gear manufacturers and scientist/gear specialists.

Require either a T-90 panel/ square mesh tailbag or other applications of square mesh panel (e.g., skylight panel), reduced bar spacing in a TED, or another federal or state certified BRD in addition to existing TED and BRD requirements in all skimmer and otter trawls.

Marine Fisheries Commission Recommendations **requested review by public, regional and standing committee*

*Convene a stakeholder group to initiate a three year study to test minimum tail bag mesh size, T-90 (square mesh) panels, skylight panels, reduced bar spacing in TEDs and any other new methods of reducing unwanted finfish bycatch to achieve a minimum of a 40 percent reduction of finfish by weight.

- Compare these to a control net with a Florida fish eye, a federally approved TED, and a one and half inch mesh tail bag.
- The stakeholder group should partner with DMF and Sea Grant to help secure funding for the study.
- If the 40 percent target reduction by weight in finfish is not achieved, further restrictions will be placed on the shrimp trawl industry to achieve the 40 percent reduction.
- Additional restrictions on the shrimp trawl industry will be reviewed and discussed at that time.

Marine Fisheries Commission Preferred Management Strategies

Allow any federally certified BRD in all NC internal and offshore waters.

Update the scientific testing protocol for the state BRD certification program.

Convene a stakeholder group to initiate industry testing of minimum tail bag mesh size, T-90 panels, skylight panels, and reduced bar spacing in TEDs to reduce bycatch to the extent practicable with a target of 40% reduction.

- Upon securing funding, testing in the ocean and internal waters will consist of three years of data using test nets compared to a control net with a Florida Fish Eye, a federally approved TED, and a one and a half inch tailbag.

- Results should minimize shrimp loss and maximize reduction of bycatch of finfish. Promising configurations will be brought back to the MFC for consideration for mandatory use.
- This stakeholder group may be partnered with DMF and Sea Grant.
- Members could consist of fishermen, net/gear manufacturers and scientist/gear specialists.

Require either a T-90 panel/ square mesh tailbag or other applications of square mesh panel (e.g., skylight panel), reduced bar spacing in a TED, or another federal or state certified BRD in addition to existing TED and BRD requirements in all skimmer and otter trawls.

12.10.6 Effort Management for bycatch reduction in the North Carolina shrimp trawl fishery

Advisory Committee Recommendation

Status quo (no changes in season, weekend or nighttime fishing)

Division Recommendation

Status quo (no changes in season, weekend or nighttime fishing)

Marine Fisheries Commission Preferred Management Strategy

Status quo (no changes in season, weekend or nighttime fishing)

12.10.7 Characterization of the North Carolina commercial shrimp trawl fleet

Advisory Committee Recommendation

Status quo (no additional maximum combined headrope requirements)

Division Recommendation

In order to put a cap on fleet capacity as a management tool, establish a maximum combined headrope length of 220 feet in all internal coastal waters where there are no existing maximum combined headrope requirements (i.e., 90 foot requirement).

Marine Fisheries Commission Preferred Management Strategy

In order to put a cap on fleet capacity as a management tool, establish a maximum combined headrope length of 220 feet in all internal coastal waters where there are no existing maximum combined headrope requirements (i.e., 90 foot requirement). Allow a phase out period of two years.

12.10.8 Area restrictions to reduce shrimp trawl bycatch in North Carolina's internal coastal waters

Advisory Committee and Division Recommendation

Prohibit shrimp trawling in the IWW channel from the Sunset Beach Bridge to the SC line, including Eastern Channel, lower Calabash River and Shallotte River.

Division Recommendation

Recommend the MFC Habitat and Water Quality Committee considers changing the designation of special secondary nursery areas that have not been opened to trawling since 1991 to permanent secondary nursery areas. Based on the outcome of AC input, rule changes may follow under the authority of the Shrimp FMP.

Marine Fisheries Commission Preferred Management Strategies

Recommend the MFC prohibit shrimp trawling in the IWW channel from the Sunset Beach Bridge to the SC line, including Eastern Channel, lower Calabash River and Shallotte River.

Recommend the MFC Habitat and Water Quality Committee considers changing the designation of special secondary nursery areas that have not been opened to trawling since 1991 to permanent secondary nursery areas.

12.11 PROPOSED RULES

12.11.1 Trawling in the New River above the Highway 172 Bridge

This rule change reflects the current management (status quo) of prohibiting otter trawls above the Highway 172 Bridge and removes it from proclamation.

15A NCAC 03J .0208 NEW RIVER

~~It is unlawful to use trawl nets upstream of the Highway 172 Bridge over New River from 9:00 P.M. through 5:00 A.M. when opened by proclamation from August 16 through November 30.~~

- (a) It is unlawful to use trawl nets except skimmer trawls upstream of the Highway 172 Bridge over New River.
- (b) It is unlawful to use skimmer trawls upstream of the Highway 172 Bridge over New River from 9:00 P.M. through 5:00 A.M. when opened by proclamation from August 16 through November 30.

*History Note: Authority G.S. 113-134; 113-182; 143B-289.52;
Eff. August 1, 1998;
Amended Eff. April 1, 2015, August 1, 2004.*

12.11.2 Evaluation of the Skimmer Trawl and other gears used for shrimping in North Carolina

Allow hand cast netting of shrimp in all closed areas and increase the limit to four quarts heads on per person.

15A NCAC 03L .0105 RECREATIONAL SHRIMP LIMITS

It is unlawful to:

- (1) Possess from open harvest areas more than 48 quarts, heads on or 30 quarts, heads off, of shrimp per person per day or if a vessel is used, per vessel per day for recreational purposes except as provided in 15A NCAC 03O .0303(e) and (f).
- (2) ~~Take or possess shrimp from areas closed to the taking of shrimp except two quarts of shrimp per person per day may be taken while fishing in a closed area with a cast net.~~
- (3) Take or possess more than four quarts, heads on or 2.5 quarts, heads off, of shrimp per person per day from areas closed to the taking of shrimp pursuant to 15A NCAC 03L .0101.

History Note: Authority G.S. 113-134; 113-182; 143B-289.52;
 Eff. April 1, 2009;
 Amended Eff. April 1, 2015; June 1, 2013.

12.11.3 Gear Modifications in the North Carolina Shrimp Trawl to Reduce Finfish Bycatch

Require either a T-90 panel/ square mesh tailbag or other applications of square mesh panel (e.g., skylight panel), reduced bar spacing in a TED, or another federal or state certified BRD in addition to existing TED and BRD requirements in all skimmer and otter trawls. May be implemented through 15A NCAC 03J .0104 (d)

15A NCAC 03I .0101 DEFINITIONS

All definitions set out in G.S. 113, Subchapter IV and the following additional terms apply to this Chapter:

- (1) Enforcement and management terms:
 - (a) Commercial Quota. Total quantity of fish allocated for harvest by commercial fishing operations.
 - (b) Educational Institution. A college, university or community college accredited by an accrediting agency recognized by the U.S. Department of Education.
 - (c) Internal Coastal Waters or Internal Waters. All coastal fishing waters except the Atlantic Ocean.
 - (d) Length of finfish.
 - (i) Curved fork length. A length determined by measuring along a line, tracing the contour of the body from the tip of the upper jaw to the middle of the fork in the caudal (tail) fin.
 - (ii) Fork length. A length determined by measuring along a straight line the distance from the tip of the snout with the mouth closed to the middle of the fork in the caudal (tail) fin, except that fork length for billfish is measured from the tip of the lower jaw to the middle of the fork of the caudal (tail) fin.
 - (iii) Pectoral fin curved fork length. A length of a beheaded fish from the dorsal insertion of the pectoral fin to the fork of the tail measured along the contour of the body in a line that runs along the top of the pectoral fin and the top of the caudal keel.
 - (iv) Total length. A length determined by measuring along a straight line the distance from the tip of the snout with the mouth closed to the tip of the compressed caudal (tail) fin.
 - (e) Recreational Possession Limit. Restrictions on size, quantity, season, time period, area, means, and methods where take or possession is for a recreational purpose.
 - (f) Recreational Quota. Total quantity of fish allocated for harvest for a recreational purpose.
 - (g) Regular Closed Oyster Season. March 31 through October 15, unless amended by the Fisheries Director through proclamation authority.
 - (h) Seed Oyster Management Area. An open harvest area that, by reason of poor growth characteristics, predation rates, overcrowding or other factors, experiences poor utilization of oyster populations for direct harvest and sale to licensed dealers and is designated by the Marine Fisheries Commission as a source of seed for public and private oyster culture.

(2) Fishing Activities:

- (a) Aquaculture operation. An operation that produces artificially propagated stocks of marine or estuarine resources or obtains such stocks from permitted sources for the purpose of rearing in a controlled environment. A controlled environment provides and maintains throughout the rearing process one or more of the following:
 - (i) food,
 - (ii) predator protection,
 - (iii) salinity,
 - (iv) temperature controls, or
 - (v) water circulation,utilizing technology not found in the natural environment.
- (b) Attended. Being in a vessel, in the water or on the shore and immediately available to work the gear and within 100 yards of any gear in use by that person at all times. Attended does not include being in a building or structure.
- (c) Blue Crab Shedding. The process whereby a blue crab emerges soft from its former hard exoskeleton. A shedding operation is any operation that holds peeler crabs in a controlled environment. A controlled environment provides and maintains throughout the shedding process one or more of the following:
 - (i) food,
 - (ii) predator protection,
 - (iii) salinity,
 - (iv) temperature controls, or
 - (v) water circulation,utilizing technology not found in the natural environment. A shedding operation does not include transporting pink or red-line peeler crabs to a permitted shedding operation.
- (d) Depuration. Purification or the removal of adulteration from live oysters, clams, and mussels by any natural or artificially controlled means.
- (e) Long Haul Operations. Fishing a seine towed between two boats.
- (f) Peeler Crab. A blue crab that has a soft shell developing under a hard shell and having a white, pink, or red-line or rim on the outer edge of the back fin or flipper.
- (g) Possess. Any actual or constructive holding whether under claim of ownership or not.
- (h) Recreational Purpose. A fishing activity that is not a commercial fishing operation as defined in G.S. 113-168.
- (i) Shellfish marketing from leases and franchises. The harvest of oysters, clams, scallops, mussels, from privately held shellfish bottoms and lawful sale of those shellfish to the public at large or to a licensed shellfish dealer.
- (j) Shellfish planting effort on leases and franchises. The process of obtaining authorized cultch materials, seed shellfish, and polluted shellfish stocks and the placement of those materials on privately held shellfish bottoms for increased shellfish production.
- (k) Shellfish production on leases and franchises:
 - (i) The culture of oysters, clams, scallops, and mussels, on shellfish leases and franchises from a sublegal harvest size to a marketable size.
 - (ii) The transplanting (relay) of oysters, clams, scallops and mussels from areas closed due to pollution to shellfish leases and franchises in open waters and the natural cleansing of those shellfish.
- (l) Swipe Net Operations. Fishing a seine towed by one boat.
- (m) Transport. Ship, carry, or cause to be carried or moved by public or private carrier by land, sea, or air.
- (n) Use. Employ, set, operate, or permit to be operated or employed.

(3) Gear:

- (a) Bunt Net. The last encircling net of a long haul or swipe net operation constructed of small mesh webbing. The bunt net is used to form a pen or pound from which the catch is dipped or bailed.

- (b) Channel Net. A net used to take shrimp which is anchored or attached to the bottom at both ends or with one end anchored or attached to the bottom and the other end attached to a boat.
- (c) Commercial Fishing Equipment or Gear. All fishing equipment used in coastal fishing waters except:
 - (i) Cast nets;
 - (ii) Collapsible crab traps, a trap used for taking crabs with the largest open dimension no larger than 18 inches and that by design is collapsed at all times when in the water, except when it is being retrieved from or lowered to the bottom;
 - (iii) Dip nets or scoops having a handle not more than eight feet in length and a hoop or frame to which the net is attached not exceeding 60 inches along the perimeter;
 - (iv) Gigs or other pointed implements which are propelled by hand, whether or not the implement remains in the hand;
 - (v) Hand operated rakes no more than 12 inches wide and weighing no more than six pounds and hand operated tongs;
 - (vi) Hook-and-line and bait-and-line equipment other than multiple-hook or multiple-bait trotline;
 - (vii) Landing nets used to assist in taking fish when the initial and primary method of taking is by the use of hook and line;
 - (viii) Minnow traps when no more than two are in use;
 - (ix) Seines less than 30 feet in length;
 - (x) Spears, Hawaiian slings or similar devices, which propel pointed implements by mechanical means, including elastic tubing or bands, pressurized gas or similar means.
- (d) Corkline. The support structure a net is attached to that is nearest to the water surface when in use. Corkline length is measured from the outer most mesh knot at one end of the corkline following along the line to the outer most mesh knot at the opposite end of the corkline.
- (e) Dredge. A device towed by engine power consisting of a frame, tooth bar or smooth bar, and catchbag used in the harvest of oysters, clams, crabs, scallops, or conchs.
- (f) Fixed or stationary net. A net anchored or staked to the bottom, or some structure attached to the bottom, at both ends of the net.
- (g) Fyke Net. An entrapment net supported by a series of internal or external hoops or frames, with one or more lead or leaders that guide fish to the net mouth. The net has one or more internal funnel-shaped openings with tapered ends directed inward from the mouth, through which fish enter the enclosure. The portion of the net designed to hold or trap fish is completely enclosed in mesh or webbing, except for the openings for fish passage into or out of the net (funnel area).
- (h) Gill Net. A net set vertically in the water to capture fish by entanglement by the gills in its mesh as a result of net design, construction, mesh size, webbing diameter or method in which it is used.
- (i) Headrope. The support structure for the mesh or webbing of a trawl that is nearest to the water surface when in use. Headrope length is measured from the outer most mesh knot at one end of the headrope following along the line to the outer most mesh knot at the opposite end of the headrope.
- (j) Hoop Net. An entrapment net supported by a series of internal or external hoops or frames. The net has one or more internal funnel-shaped openings with tapered ends directed inward from the mouth, through which fish enter the enclosure. The portion of the net designed to hold or trap the fish is completely enclosed in mesh or webbing, except for the openings for fish passage into or out of the net (funnel area).
- (k) Lead. A mesh or webbing structure consisting of nylon, monofilament, plastic, wire or similar material set vertically in the water, held in place by stakes or anchors to guide fish into an enclosure. Lead length is measured from the outer most end of the lead along the top or bottom line, whichever is longer, to the opposite end of the lead.

- (l) Mechanical methods for clamming. Dredges, hydraulic clam dredges, stick rakes and other rakes when towed by engine power, patent tongs, kicking with propellers or deflector plates with or without trawls, and any other method that utilizes mechanical means to harvest clams.
 - (m) Mechanical methods for oystering. Dredges, patent tongs, stick rakes and other rakes when towed by engine power and any other method that utilizes mechanical means to harvest oysters.
 - (n) Mesh Length. The ~~diagonal~~ distance from the inside of one knot to the outside of the ~~other~~ opposite knot, when the net is stretched hand-tight in a manner that closes the mesh opening.
 - (o) Pound Net Set. A fish trap consisting of a holding pen, one or more enclosures, lead or leaders, and stakes or anchors used to support the trap. The lead(s), enclosures, and holding pen are not conical, nor are they supported by hoops or frames.
 - (p) Purse Gill Nets. Any gill net used to encircle fish when the net is closed by the use of a purse line through rings located along the top or bottom line or elsewhere on such net.
 - (q) Seine. A net set vertically in the water and pulled by hand or power to capture fish by encirclement and confining fish within itself or against another net, the shore or bank as a result of net design, construction, mesh size, webbing diameter, or method in which it is used.
- (4) Fish habitat areas. The estuarine and marine areas that support juvenile and adult populations of fish species, as well as forage species utilized in the food chain. Fish habitats as used in this definition, are vital for portions of the entire life cycle, including the early growth and development of fish species. Fish habitats in all coastal fishing waters, as determined through marine and estuarine survey sampling, include:
- (a) Anadromous fish nursery areas. Those areas in the riverine and estuarine systems utilized by post-larval and later juvenile anadromous fish.
 - (b) Anadromous fish spawning areas. Those areas where evidence of spawning of anadromous fish has been documented in Division sampling records through direct observation of spawning, capture of running ripe females, or capture of eggs or early larvae.
 - (c) Coral:
 - (i) Fire corals and hydrocorals (Class Hydrozoa);
 - (ii) Stony corals and black corals (Class Anthozoa, Subclass Scleractinia); or
 - (iii) Octocorals; Gorgonian corals (Class Anthozoa, Subclass Octocorallia), which include sea fans (*Gorgonia* sp.), sea whips (*Leptogorgia* sp. and *Lophogorgia* sp.), and sea pansies (*Renilla* sp.).
 - (d) Intertidal Oyster Bed. A formation, regardless of size or shape, formed of shell and live oysters of varying density.
 - (e) Live rock. Living marine organisms or an assemblage thereof attached to a hard substrate, excluding mollusk shells, but including dead coral or rock. Living marine organisms associated with hard bottoms, banks, reefs, and live rock include:
 - (i) Coralline algae (Division Rhodophyta);
 - (ii) *Acetabularia* sp., mermaid's fan and cups (*Udotea* sp.), watercress (*Halimeda* sp.), green feather, green grape algae (*Caulerpa* sp.) (Division Chlorophyta);
 - (iii) *Sargassum* sp., *Dictyopteris* sp., *Zonaria* sp. (Division Phaeophyta);
 - (iv) Sponges (Phylum Porifera);
 - (v) Hard and soft corals, sea anemones (Phylum Cnidaria), including fire corals (Class Hydrozoa), and Gorgonians, whip corals, sea pansies, anemones, *Solenastrea* (Class Anthozoa);
 - (vi) Bryozoans (Phylum Bryozoa);
 - (vii) Tube worms (Phylum Annelida), fan worms (*Sabellidae*); feather duster and Christmas treeworms (*Serpulidae*), and sand castle worms (*Sabellaridae*);
 - (viii) Mussel banks (Phylum Mollusca: Gastropoda); and
 - (ix) Acorn barnacles (Arthropoda: Crustacea: *Semibalanus* sp.).

- (f) Nursery areas. Those areas in which for reasons such as food, cover, bottom type, salinity, temperature and other factors, young finfish and crustaceans spend the major portion of their initial growing season. Primary nursery areas are those areas in the estuarine system where initial post-larval development takes place. These are areas where populations are uniformly early juveniles. Secondary nursery areas are those areas in the estuarine system where later juvenile development takes place. Populations are composed of developing sub-adults of similar size which have migrated from an upstream primary nursery area to the secondary nursery area located in the middle portion of the estuarine system.
- (g) Shellfish producing habitats. Those areas in which shellfish, such as clams, oysters, scallops, mussels, and whelks, whether historically or currently, reproduce and survive because of such favorable conditions as bottom type, salinity, currents, cover, and cultch. Included are those shellfish producing areas closed to shellfish harvest due to pollution.
- (h) Strategic Habitat Areas. Locations of individual fish habitats or systems of habitats that provide exceptional habitat functions or that are particularly at risk due to imminent threats, vulnerability, or rarity.
- (i) Submerged aquatic vegetation (SAV) habitat. Submerged lands that:
 - (i) are vegetated with one or more species of submerged aquatic vegetation including bushy pondweed or southern naiad (*Najas guadalupensis*), coontail (*Ceratophyllum demersum*), eelgrass (*Zostera marina*), horned pondweed (*Zannichellia palustris*), naiads (*Najas* spp.), redhead grass (*Potamogeton perfoliatus*), sago pondweed (*Stuckenia pectinata*, formerly *Potamogeton pectinatus*), shoalgrass (*Halodule wrightii*), slender pondweed (*Potamogeton pusillus*), water stargrass (*Heteranthera dubia*), water starwort (*Callitriche heterophylla*), waterweeds (*Elodea* spp.), widgeongrass (*Ruppia maritima*) and wild celery (*Vallisneria americana*). These areas may be identified by the presence of above-ground leaves, below-ground rhizomes, or reproductive structures associated with one or more SAV species and include the sediment within these areas; or
 - (ii) have been vegetated by one or more of the species identified in Sub-item (4)(i)(i) of this Rule within the past 10 annual growing seasons and that meet the average physical requirements of water depth (six feet or less), average light availability (secchi depth of one foot or more), and limited wave exposure that characterize the environment suitable for growth of SAV. The past presence of SAV may be demonstrated by aerial photography, SAV survey, map, or other documentation. An extension of the past 10 annual growing seasons criteria may be considered when average environmental conditions are altered by drought, rainfall, or storm force winds.

This habitat occurs in both subtidal and intertidal zones and may occur in isolated patches or cover extensive areas. In defining SAV habitat, the Marine Fisheries Commission recognizes the Aquatic Weed Control Act of 1991 (G.S. 113A-220 et. seq.) and does not intend the submerged aquatic vegetation definition, or rules 15A NCAC 03K .0304, .0404 and 03I .0101, to apply to or conflict with the non-development control activities authorized by that Act.

- (5) Licenses, permits, leases and franchises, and record keeping:
 - (a) Assignment. Temporary transferal to another person of privileges under a license for which assignment is permitted. The person assigning the license delegates the privileges permitted under the license to be exercised by the assignee, but retains the power to revoke the assignment at any time, is still the responsible party for the license.
 - (b) Designee. Any person who is under the direct control of the permittee or who is employed by or under contract to the permittee for the purposes authorized by the permit.
 - (c) For Hire Vessel. As defined by G.S. 113-174 when the vessel is fishing in state waters or when the vessel originates from or returns to a North Carolina port.

- (d) Holder. A person who has been lawfully issued in their name a license, permit, franchise, lease, or assignment.
- (e) Land:
 - (i) For commercial fishing operations, when fish reach the shore or a structure connected to the shore.
 - (ii) For purposes of trip tickets, when fish reach a licensed seafood dealer, or where the fisherman is the dealer, when the fish reaches the shore or a structure connected to the shore.
 - (iii) For recreational fishing operations, when fish are retained in possession by the fisherman.
- (f) Licensee. Any person holding a valid license from the Department to take or deal in marine fisheries resources.
- (g) Master. Captain of a vessel or one who commands and has control, authority, or power over a vessel.
- (h) New fish dealer. Any fish dealer making application for a fish dealer license who did not possess a valid dealer license for the previous license year in that name or ocean pier license in that name on June 30, 1999. For purposes of license issuance, adding new categories to an existing fish dealers license does not constitute a new dealer.
- (i) North Carolina Trip Ticket. Paper forms provided by the Division, and electronic data files generated from software provided by the Division, for the reporting of fisheries statistics, which include quantity, method and location of harvest.
- (j) Office of the Division. Physical locations of the Division conducting license and permit transactions in Wilmington, Washington, Morehead City, Columbia, Roanoke Island and Elizabeth City, North Carolina. Other businesses or entities designated by the Secretary to issue Recreational Commercial Gear Licenses or Coastal Recreational Fishing Licenses are not considered Offices of the Division.
- (k) Responsible party. Person who coordinates, supervises or otherwise directs operations of a business entity, such as a corporate officer or executive level supervisor of business operations and the person responsible for use of the issued license in compliance with applicable statutes and rules.
- (l) Tournament Organizer. The person who coordinates, supervises or otherwise directs a recreational fishing tournament and is the holder of the Recreational Fishing Tournament License.
- (m) Transaction. Act of doing business such that fish are sold, offered for sale, exchanged, bartered, distributed or landed.
- (n) Transfer. Permanent transferal to another person of privileges under a license for which transfer is permitted. The person transferring the license retains no rights or interest under the license transferred.

History Note: Authority G.S. 113-134; 113-174; 143B-289.52; Eff. January 1, 1991; Amended Eff. March 1, 1995; March 1, 1994; October 1, 1993; July 1, 1993; Recodified from 15A NCAC 03I .0001 Eff. December 17, 1996; Amended Eff. April 1, 1999; August 1, 1998; April 1, 1997; Temporary Amendment Eff. May 1, 2000; August 1, 1999; July 1, 1999; Amended Eff. August 1, 2000; Temporary Amendment Eff. August 1, 2000; Amended Eff. April 1, 2015; April 1, 2011; April 1, 2009; October 1, 2008; December 1, 2007; December 1, 2006; September 1, 2005; April 1, 2003; April 1, 2001.

12.11.4 Characterization of the North Carolina Commercial Shrimp Trawl Fleet

In order to put a cap on fleet capacity as a management tool, establish a maximum combined headrope length of 220 feet in all internal coastal waters where there are no existing maximum combined headrope requirements with a two year phase out period.

15A NCAC 03L .0103 PROHIBITED NETS, MESH SIZES AND AREAS

(a) It is unlawful to take shrimp with nets with mesh lengths less than the following:

- (1) Trawl net - one and one-half inches;
- (2) Fixed nets, channel nets, float nets, butterfly nets, and hand seines - one and one-fourth inches; and
- (3) Cast net - no restriction.

(b) It is unlawful to take shrimp with a net constructed in such a manner as to contain an inner or outer liner of any mesh size. Net material used as chafing gear shall be no less than four inches mesh length except that chafing gear with smaller mesh may be used only on the bottom one-half of the tailbag. Such chafing gear shall not be tied in a manner that forms an additional tailbag.

(c) It is unlawful to take shrimp with trawls which have a combined headrope of greater than 90 feet in internal coastal ~~fishing waters~~ except in the following areas:

- (1) Pamlico Sound; north of a line beginning at a point 35° 48.5015' N - 75° 44.1228' W on Roanoke Marshes Point, running southeasterly to a point 35° 44 .1710'N - 75°31 .0520'W on the north point of Eagle Nest Bay;
- (2) Core Sound and all adjacent coastal fishing waters south of a line beginning at a point 34° 59.7942' N - 76° 14.6514' W on Camp Point; running easterly to a point 34° 58.7853' N - 76° 09.8922' W on Core Banks; to the South Carolina State line.
- ~~(2)~~(3) Pamlico River ~~downstream-upstream~~ of a line from a point 35° 18.5882'N – 76° 28.9625'W at Pamlico Point; running northerly to a point 35° 22.3741'N – 76° 28.6905'W at Willow Point;
- ~~(3)~~(4) Neuse River ~~northeast southwest~~ of a line from a point 34° 58.2000'N – 76° 40.5167'W at Winthrop Point on the eastern shore of the entrance to Adam's Creek running northerly to a point 35° 01.0744' N – 76° 42.1550' W at Windmill Point at the entrance of Greens Creek at Oriental.

(d) Beginning January 1, 2017 it is unlawful to take shrimp with trawls which have a combined headrope of greater than 220 feet in internal coastal fishing waters in the following areas:

- (1) Pamlico Sound; south of a line beginning at a point 35° 48.5015' N - 75° 44.1228' W on Roanoke Marshes Point, running southeasterly to a point 35° 44 .1710'N - 75°31 .0520'W on the north point of Eagle Nest Bay and north of a line beginning at a point 34° 59.7942' N - 76° 14.6514' W on Camp Point; running easterly to a point 34° 58.7853' N - 76° 09.8922' W on Core Banks.
- (2) Pamlico River downstream of a line from a point 35° 18.5882'N – 76° 28.9625'W at Pamlico Point; running northerly to a point 35° 22.3741'N – 76° 28.6905'W at Willow Point;
- (3) Neuse River northeast of a line from a point 34° 58.2000'N – 76° 40.5167'W at Winthrop Point on the eastern shore of the entrance to Adam's Creek running northerly to a point 35° 01.0744' N – 76° 42.1550' W at Windmill Point at the entrance of Greens Creek at Oriental.

~~(d)~~(e) It is unlawful to use a shrimp trawl in the areas described in 15A NCAC 03R .0114.

~~(e)~~(f) It is unlawful to use channel nets except as provided in 15A NCAC 03J .0106.

~~(f)~~(g) It is unlawful to use shrimp pots except as provided in 15A NCAC 03J .0301.

~~(g)~~(h) It is unlawful to use a shrimp trawl that does not conform with the federal rule requirements for Turtle Excluder Devices (TED) as specified in 50 CFR Part 222.102 Definitions, 50 CFR Part 223.205 (a) and Part 223.206 (d) Gear Requirements for Trawlers, and 50 CFR Part 223.207 Approved TEDs. Copies of these rules are available via the Code of Federal Regulations posted on the Internet at <http://www.gpoaccess.gov/cfr/index.html> and at the Division of Marine Fisheries, P.O. Box 769, Morehead City, North Carolina 28557 at no cost.

*History Note: Authority G.S. 113-134; 113-182; 143B-289.52;
Eff. January 1, 1991;
Amended Eff. April 1, 2015; April 1, 2009; July 1, 2006.*

12.11.5 Area Restrictions to Reduce Shrimp Trawl Bycatch in North Carolina's Internal Coastal Waters

Prohibit shrimp trawling in the IWW channel from the Sunset Beach Bridge to the SC line, including Eastern Channel, lower Calabash River and Shallotte River.

15A NCAC 03R .0114 SHRIMP TRAWL PROHIBITED AREAS

The shrimp trawl prohibited areas referenced in 15A NCAC 03L .0103(d) are delineated in the following coastal water areas:

- (1) Pungo River- all waters upstream of a line from a point 35° 23.3166'N – 76° 34.4833'W at Wades Point; running westerly to a point 35° 23.6463'N – 76° 31.0003'W on the north shore of the entrance to Abels Bay.
- (2) Pamlico River- all waters upstream of a line from a point 35° 20.5108'N – 76° 37.7218'W on the western shore of the entrance to Goose Creek; running northeasterly to a point 35° 23.3166'N – 76° 34.4833'W at Wades Point.
- (3) Neuse River- all waters upstream of a line from a point 34° 56.3658'N – 76° 48.7110'W at Cherry Point; running northerly to a point 34° 57.9116'N – 76° 48.2240'W at Wilkerson Point.
- (4) Lower Shallotte River- all waters upstream of a line beginning at a point 33° 54.832' N– 78° 22.363' W on the west side of Shallotte River; running southeasterly to a point 33° 54.621' N - 78° 21.796' W on the east side of the river.
- (5) Eastern Channel – all waters of Eastern Channel east and north of a line beginning at a point 33° 52.676' N - 78° 28.735' W at Jinks Creek); running southerly to a point 33° 52.585' N - 78° 28.671' W at Tubbs Inlet; and south and west of a line beginning at a point 33° 53.627' N - 78° 26.633' W; running easterly to a point 33° 53.648' N - 78° 26.554' W.
- (6) Sunset Beach- all waters of the IWW west of a line beginning at a point 33° 52.921' N – 78° 30.709' W on the north end of state road No. 1172 bridge; running southerly to a point 33° 52.830' N – 78° 30.650' W at the south end of the bridge.
- (7) Calabash River- all waters west of a line beginning at a point 33° 53.444' N - 78° 32.968' W on the north end of state road No. 1164 bridge; running southerly to a point 33° 53.350' N – 78° 32.972' W at the south end of the bridge .

History Note: Authority G.S. 113-134; 113-182; 143B-289.52;
 Eff. July 1, 2006.
Amended Eff. April 1, 2015

12.11.6 Additional Rule Changes to Address Clarity and Consistency

Proposed rule changes listed below reflect current management from the 2006 Shrimp FMP and do not change the intent of current rules. These rule changes are proposed to address rule clarity and consistency in the MFC rulebook.

This rule change reflects the current management of proclamation authority to manage shrimping and makes this consistent with other proclamation authority rules.

15A NCAC 03L .0101 SEASON SHRIMP HARVEST RESTRICTIONS

~~It is unlawful to take shrimp with nets until the Fisheries Director, by proclamation, opens the season in various waters. Proclamations may specify any hours of day or night or both and any other conditions appropriate to management of the fishery. If sampling indicates primarily undersized shrimp or juveniles of any other species of major economic importance, the Fisheries Director may close such waters to shrimping and prohibit the use of nets for any purpose except cast nets as provided in 15A NCAC 3L .0102. Prominent landmarks or other permanent type markers shall be considered when establishing closure lines even if such lines extend beyond the area of concern.~~

The Fisheries Director may, by proclamation, impose any or all of the following restrictions on the taking of shrimp:

- (1) Specify time;
- (2) Specify area;
- (3) Specify means and methods;
- (4) Specify season;
- (5) Specify size; and
- (6) Specify quantity.

History Note: Authority G.S. 113-134; 113-182; 113-221; 143B-289.52;
 Eff. January 1, 1991.

This rule change reflects the current management of keeping otter trawls and skimmer trawls as shrimp trawls as authorized commercial fishing gear by providing consistency within the MFC Rule Book of skimmer trawls being managed as shrimp trawls.

15A NCAC 030 .0302 AUTHORIZED GEAR

(a) The following are the only commercial fishing gear authorized (including restrictions) for use under a valid Recreational Commercial Gear License:

- (1) One seine 30 feet or over in length but not greater than 100 feet with a mesh length less than 2 1/2 inches when deployed or retrieved without the use of a vessel or any other mechanical methods. A vessel may be used only to transport the seine;
- (2) One shrimp trawl with a headrope not exceeding 26 feet in length per vessel.
- (3) With or without a vessel, five eel, fish, shrimp, or crab pots in any combination, except only two pots of the five may be eel pots. Peeler pots are not authorized for recreational purposes;
- (4) One multiple hook or multiple bait trotline up to 100 feet in length;
- (5) Gill Nets:
 - (A) Not more than 100 yards of gill nets with a mesh length equal to or greater than 2 1/2 inches except as provided in (C) of this Subparagraph. Attendance is required at all times;
 - (B) Not more than 100 yards of gill nets with a mesh length equal to or greater than 5 1/2 inches except as provided in (C) of this Subparagraph. Attendance is required when used from one hour after sunrise through one hour before sunset in internal coastal fishing waters east and north of the Highway 58 Bridge at Emerald Isle and in the Atlantic Ocean east and north of 77° 04.0000' W. Attendance is required at all times in internal coastal fishing waters west and south of the Highway 58 Bridge at Emerald Isle and in the Atlantic Ocean west and south of 77° 04.0000' W; and
 - (C) Not more than 100 yards of gill net may be used at any one time, except that when two or more Recreational Commercial Gear License holders are on board, a maximum of 200 yards may be used from a vessel;
 - (D) It is unlawful to possess aboard a vessel more than 100 yards of gill nets with a mesh length less than 5 1/2 inches and more than 100 yards of gill nets with a mesh length equal to or greater than 5 1/2 inches identified as recreational commercial fishing equipment when only one Recreational Commercial Gear License holder is on board. It is unlawful to possess aboard a vessel more than 200 yards of gill nets with a mesh length less than 5 1/2 inches and more than 200 yards of gill nets with a mesh length equal to or greater than 5 1/2 inches identified as recreational commercial fishing equipment when two or more Recreational Commercial Gear License holders are on board;
- (6) A hand-operated device generating pulsating electrical current for the taking of catfish in the area described in 15A NCAC 03J .0304;
- ~~(7) Skimmer trawls not exceeding 26 feet in total combined width.~~
- ~~(8)~~(7) One pound net used to take shrimp with each lead 10 feet or less in length and with a minimum lead net mesh of 1 1/2 inches, and enclosures constructed of net mesh of 1 1/4 inches or greater and with all dimensions being 36 inches or less. Attendance is required at all times and all gear must be removed from the water when not being fished. Gear is to be marked and set as specified in 15A NCAC 03J .0501.

(b) It is unlawful to use more than the quantity of authorized gear specified in Subparagraphs (a)(1) through (a)(8) of this Rule, regardless of the number of individuals aboard a vessel possessing a valid Recreational Commercial Gear License.

(c) It is unlawful for a person to violate the restrictions of or use gear other than that authorized by Paragraph (a) of this Rule.

(d) Unless otherwise provided, this Rule does not exempt Recreational Commercial Gear License holders from the provisions of other applicable rules of the Marine Fisheries Commission or provisions of proclamations issued by the Fisheries Director as authorized by the Marine Fisheries Commission.

History Note: Authority G.S. 113-134; 113-173;

Temporary Adoption Eff. August 9, 1994, for a period of 180 days or until the permanent rule becomes effective, whichever is sooner;
Eff. February 1, 1995;
Temporary Amendment Eff. August 1, 1999; July 1, 1999;
Amended Eff. August 1, 2000;
Temporary Amendment Eff. August 1, 2000;
Amended Eff. April 1, 2015; April 1, 2009; July 1, 2006; November 1, 2005; August 1, 2002.

13.0 LITERATURE CITED

- Adkins, B. E., R.M. Harbo, and N. Bourne. 1983. An evaluation and management considerations of the use of a hydraulic clam harvester on intertidal clam populations in British Columbia. Canadian Manuscript Reports Fisheries Aquatic Science 1716: 38.
- Alverson, D. L. and S. E. Hughes, 1996. Bycatch: from emotion to effective natural resource management. Reviews in Fish Biology and Fisheries 6: 443–462.
- Alverson, D.L., M.H. Freeberg, S.A. Murawski and J.G. Pope. 1994. A global assessment of fisheries bycatch and discards. FAO Fisheries Technical Paper. No. 339, Rome, FAO. 233p.
- ASMFC (Atlantic States Marine Fisheries Commission). 1994. Acronyms, abbreviations, and technical terms used in ASMFC fishery management programs. Special Report No. 33 of the Atlantic States Marine Fisheries Commission, Washington D.C. 22p.
- ASMFC (Atlantic States Marine Fisheries Commission) 1996a. Fishery management plan and addendum 1 for scup. ASMFC FMR No. 26. Atlantic States Marine Fisheries Commission, Washington D.C.
- ASMFC (Atlantic States Marine Fisheries Commission) 1996b. Fishery management plan for black sea bass. ASMFC FMR No.28. Atlantic States Marine Fisheries Commission, Washington D.C.
- ASMFC (Atlantic States Marine Fisheries Commission) 1996c. Amendment 3 to the interstate fishery management plan for weakfish. ASMFC FMR No. 27. Atlantic States Marine Fisheries Commission, Washington D.C.
- ASMFC (Atlantic States Marine Fisheries Commission). 2000. Evaluating fishing gear impacts to submerged aquatic vegetation and determining mitigation strategies. ASFMC Habitat Management Series 5. 38p.
- ASMFC (Atlantic States Marine Fisheries Commission). 2003. Atlantic Croaker 2003 Stock Assessment Report. Atlantic States Marine Fisheries Commission. Washington, D.C. 151p.
- ASMFC (Atlantic States Marine Fisheries Commission). 2004. Atlantic Croaker 2004 Stock Assessment Supplement. Atlantic States Marine Fisheries Commission. Washington, D.C. 188p.
- ASMFC (Atlantic States Marine Fisheries Commission). 2009. Weakfish 2009 Stock Assessment Report. Atlantic States Marine Fisheries Commission. Washington, D.C. 396p.
- ASMFC (Atlantic States Marine Fisheries Commission) 2011a. 2010 Review of the Atlantic States Marine Fisheries Commission Fishery Management Plan for Spot (*Leiostomus xanthurus*) 2009 Fishing Year. Atlantic States Marine Fisheries Commission. Washington DC. 15p

- ASMFC (Atlantic States Marine Fisheries Commission). 2011b. Omnibus Amendment to the Interstate Fishery Management Plans For Spanish Mackerel, Spot, and Spotted Seatrout. Atlantic States Marine Fisheries Commission. Washington, D.C. 161p.
- Bales, J. D. and D.J. Newcomb. 1996. North Carolina wetland resources. p. 297-302 *in* R.M. Hirsch (dir). National Water Summary on Wetland Resources. U.S. Geological Survey, Atlanta, GA, USGS Water-Supply Paper 2425.
- Bellido, J. M., Santos, M.B., Pennino, M.G. Valeiras X., and G. J. Pierce. 2011. Fishery discards and bycatch: solutions for an ecosystem approach to fisheries management? *Hydrobiologia* 670: 317-333.
- Bielsa, L.M., W.H. Murdich, and R.F. Labisky. 1983. Species profiles: life histories and environment requirements of coastal fish and invertebrates (south Florida) – pink shrimp. U.S. Fish and Wildlife Service FWS/OBS-82/11.17. U.S. Army Corps of Engineers, TR EL-82-4. 21 pp.
- Brinson, M. M. 1977. Decomposition and nutrient exchange in litter in an alluvial swamp forest. *Ecology* 58: 601-609.
- Broome, J.D., Anderson, J.W., and D. W. Anderson. 2011. By-catch volume reduction through turtle excluder device (TED) reduced grid spacing. North Carolina Sea Grant Project # 10-FEG-03. 37p.
- Brown, K.B. 2006. Evaluation of Experimental Shrimp Pots from Carteret County to Brunswick County, North Carolina Completion report for NOAA award no. NA 05 NMF 4741003 North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, 19p.
- Brown, K.B. 2009. Characterization of the near-shore commercial shrimp trawl fishery from Carteret County to Brunswick County, North Carolina Completion report for NOAA award no. NA05NMF4741003 North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, 29p.
- Brown, K.B. 2010a. Compare catch rates of shrimp and bycatch of other species in standard (control) and modified (experimental) otter trawls in the Neuse River and Pamlico Sound, North Carolina Completion report for NOAA award no. NA08NMF474076 North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, 22p.
- Brown, K.B. 2010b. Characterization of the inshore commercial shrimp trawl fishery in Pamlico Sound and its tributaries, North Carolina Completion report for NOAA award no. NA05NMF4741003 North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, 28p.
- Brown, S. S., G.R. Gaston, C.F. Rakocinski, and R.W. Heard. 2000. Effects of sediment contaminants and environmental gradients on macrobenthic community trophic structure in Gulf of Mexico estuaries. *Estuaries* 23(3): 411-424.
- Browder, J.A., Zein-Eldin, Z., Criales, M.M., Robblee, M.B. Wong, S., Jackson, T.L. Johnson, D., 2002. Dynamics of pink shrimp (*Farfantepenaeus durarum*) recruitment potential in relation to salinity and temperature in Florida Bay. *Estuaries* 25: 1355-1371.

- Burkholder, J. M., K.M. Mason, and H.B. Glasgow Jr. 1992b. Water-column nitrate enrichment promotes decline of eelgrass *Zostera marina*: Evidence from seasonal mesocosm experiments. *Marine Ecology Progress Series* 81(2): 163-178.
- Buzzelli, C. P., R.A. Luetlich Jr., S.P. Powers, C.H. Peterson, J.E. McNinch, J.L. Pinckney, and H.W. Pearl. 2002. Estimating the spatial extent of bottom water hypoxia and habitat degradation in a shallow estuary. *Marine Ecology Progress Series* 230: 103-112.
- Byrne, D. M. 1995. The effect of bulkheads on estuarine fauna: a comparison of littoral fish and macroinvertebrate assemblages at bulkheaded and non-bulkheaded shorelines in a Barnegat Bay Lagoon. *Second Annual Marine Estuarine Shallow Water Science and Management Conference*: 53-56.
- Caillouet, C.W. Jr., R.A. Hart, J.M. Nance. 2008. Growth overfishing in the brown shrimp fishery of Texas, Louisiana, and adjoining Gulf of Mexico EEZ. *Fisheries Research* 92: 289-302.
- Carothers, P.E. and Chittendon, M.E., Jr. 1985. Relationships between trawl catch and tow duration for penaid shrimp. *Transactions of the American Fisheries Society* 114: 851-856.
- Carr, W.E. and C.A. Adams. 1973. Food Habits of juvenile marine fishes occupying seagrass beds in the estuarine zone near Crystal River, Florida. *Transactions of the American Fisheries Society* 102: 511-540.
- Chevront, B. 2002. A Social and Economic Analysis of Commercial Fisheries of Core Sound, North Carolina. North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, Morehead City, NC. (NOAA Award No. NA16FG1220-1).
- Chevront, B. 2003. A Social and Economic Analysis of Commercial Fisheries in North Carolina: Beaufort Inlet to the South Carolina State Line. North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, Morehead City, NC. (NC Technical Assistance to the South Atlantic Fisheries Management Council, Task 5: NEPA Related Activities, Contract No. SA-03-03-NC).
- Chopin, F.S. and T. Arimoto. 1995. The condition of fish escaping from fishing gears-a review. *Fisheries Research* 21:315-327.
- Chmura, G. L. and N.W. Ross. 1978. Environmental impacts of marinas and their boats. Rhode Island Sea Grant, Narragansett, RI, P675; RIU-T-78-005.
- Coale, J. S., R. A. Rulifson, J. D. Murray, and R. Hines. 1994. Comparisons of shrimp catch and bycatch between a skimmer trawl and an otter trawl in the North Carolina inshore shrimp fishery. *North American Journal of Fisheries Management* 14:751-768.
- Coen, L. D. and M.W. Luckenbach. 1998. Developing success criteria and goals for evaluating shellfish habitat restoration: ecological function or resource exploitation? *Goal Setting and Success Criteria for Habitat Restoration Conference*, January 13-15. Abstract only.
- Coen, L. E., M.W. Luckenbach, and D.L. Breitburg. 1999. The role of oyster reefs as essential fish habitat: a review of current knowledge and some new perspectives. p. 438-454 *in* L.R. Benaka (ed.). *Fish habitat: Essential fish habitat and rehabilitation*. American Fisheries Society, Bethesda, MD, Symposium 22, 459p.

- Coen, L. E., M.W. Luckenbach, and D.L. Breitburg. 1999. The role of oyster reefs as essential fish habitat: a review of current knowledge and some new perspectives. p. 438-454 in L.R. Benaka (editor). Fish habitat: Essential fish habitat and rehabilitation. American Fisheries Society, Bethesda, MD, Symposium 22, 459p.
- Collie, J. S., S.J. Hall, M.J. Kaiser, and I.R. Poiners. 2000. A quantitative analysis of fishing impacts on shelf-sea benthos. *Journal of Animal Ecology* 69: 785-798.
- Connell, B. and T. Murphey. 2004. A preliminary evaluation on the effects of dock shading on density and coverage of shoal grass (*Halodule wrightii*). DMF, Unpublished report.
- Cooper, S. R. and G.S. Brush. 1991. A 2500 year history of anoxia and eutrophication in the Chesapeake Bay. *Science* 254: 992-1001.
- Courtney A., J Haddy, M. Campbell, D. Roy, M. Tonks, S. Gaddes, K. Chilcott, M. O'Neil, I. Brown, M. McLennan, J. Jebreen, C. van der Geest, C. Rose, S. Kistle, C. Turnbull, P. Kyne, M. Bennet, and J. Taylor. 2007. Bycatch weight, composition and preliminary estimates of the impact of bycatch reduction devices in Queensland's trawl fishery. Queensland Department of Primary Industries and Fisheries. Department of Anatomy and Developmental Biology, University of Queensland. Australia Maritime College. Project no. 2000/170.
- Crosson, Scott .2007a. A Social and Economic Analysis of Commercial Fisheries in North Carolina: Albemarle and Pamlico Sounds. NC Department of Environment and Natural Resources, Division of Marine Fisheries, Morehead City, NC. (NOAA Award No. NA16FW1543), 25p.
- Crosson, Scott 2007b. A Social and Economic Analysis of Commercial Fisheries in North Carolina: Core Sound. NC Department of Environment and Natural Resources, Division of Marine Fisheries, Morehead City, NC. (NOAA Award No. NA16FW1543), 32p.
- Crosson, S. 2009. A Social and Economic Analysis of Commercial Fisheries in North Carolina. NC Department of Environment and Natural Resources, Division of Marine Fisheries, Morehead City, NC. (NOAA Award No. NA05NMF4741003), 26p.
- Crowder, L. B. and S. A. Murawski. 1998. Fisheries bycatch: Implication for management. *Fisheries* 24(6): 8-17.
- Currin, C. A., S.Y. Newell, and H.W. Paerl. 1995. The role of standing dead *Spartina alterniflora* and benthic microalgae in salt marsh food webs: considerations based on multiple stable isotope analysis. *Marine Ecology Progress Series* 121: 99-116.
- Dahl, T. E. 1990. Wetlands - losses in the United States, 1780's to 1980's. U.S. Fish and Wildlife Service, Washington, D.C., Report to Congress, 13p.
- Dauer, D. M., J.A. Ranasinghe, and S.B. Weisberg. 2000. Relationships between benthic community condition, water quality, sediment quality, nutrient loads, and land use patterns in Chesapeake Bay. *Estuaries* 23(1): 80-96.
- Davies, R. W. D., S. J. Cripps, A. Nickson & G. Porter, 2009. Defining and estimating global marine fisheries bycatch. *Marine Policy* 33: 661-672.

- DeAlteris, J., L. Skrobe, and C. Lipsky. 1999. The significance of seabed disturbance by mobile fishing gear relative to natural processes: A case study in Narragansett Bay, Rhode Island. *American Fisheries Symposium* 22: 14.
- Deaton, A.S., W.S. Chappell, K.Hart, J. O'Neal, B. Boutin. 2010. North Carolina Coastal Habitat Protection Plan. North Carolina Department of Environment and Natural Resources. Division of Marine Fisheries, NC. 639 pp.
- Deehr, R.A. 2012. Measuring the ecosystem impacts of commercial shrimp trawling and other fishing gear in Core Sound, North Carolina using ecological network analysis. Dissertation. East Carolina University, Coastal Resources Management Program. 410pp.
- DEHNR (NC Department of Environment Health and Natural Resources). 1990. North Carolina coastal marinas: water quality assessment. Department of Environment Health and Natural Resources, Raleigh, NC, 90-01, 69p.
- DeLancey, L.B., Jenkins, J.E., Maddox M. B., Whitaker J. D., Wenner E.L. 2005. Field observations on white shrimp, *Litopenaeus setiferus*, during spring spawning season in South Carolina, U.S.A., 1980-2003. *Journal of Crustacean Biology* 25: 212-218.
- Dennison, W. C., R.J. Orth, K.A. Moore, J.C. Stevenson, V. Carter, S. Kollar, P.W. Bergstrom, and R. Batiuk. 1993. Assessing water quality with submerged aquatic vegetation. *Bioscience* 43: 86-94.
- Diamond S.L. 2003. Estimation of bycatch in shrimp trawl fisheries: a comparison of estimation methods using field data and simulated data. *Fishery Bulletin* 101(Supplement 3), 484–500.
- Diamond-Tissue, S. L. 1999. Characterization and estimation of shrimp trawl bycatch in North Carolina waters. Doctorate dissertation, North Carolina State University, Department of Zoology, Raleigh, NC 27695. 54 p.
- Diamond-Tissue, S.L. 1999b. The population effects of shrimp trawl bycatch on Atlantic croaker. Phd. Dissertation(draft). North Carolina State University, Depart.of Zoology, Raleigh, NC 27695. 58p.
- Durako, M. J. 1994. Seagrass die-off in Florida Bay (USA): changes in shoot demographic characteristics and population dynamics in *Thalassia testudinum*. *Marine Ecology Progress Series* 110: 59-66.
- DWQ (North Carolina Division of Water Quality). 2000. A citizen's guide to water quality management in North Carolina. DENR, Div. Water Quality, Planning Branch, Raleigh, NC, 156p.
- DWQ (North Carolina Division of Water Quality). 2011. Annual report of fish kill events. Department of Natural Resources, Raleigh, NC, 19p.
- Dyer, K. R. and R.J. Orth. 1994. Changes in fluxes in estuaries: implications from science to management. Olsen and Olsen, Fredenburg, Denmark.

- Eby and Crowder. 2002. Hypoxia-based habitat compression in the Neuse River Estuary: context-dependent shifts in behavioral avoidance thresholds. *Canadian Journal of Fisheries and Aquatic Sciences* 59(6): 952-965.
- Ellifrit, N. J., M.S. Uoshinaka, and D.W. Coon. 1972. Some observations of clam distribution at four sites on the Hook Canal, Washington. *Proceedings of the National Shellfish Association* 63: 7.
- Ellis, W. 1995. The intertidal distribution of the daggerblade grass shrimp: effects of vegetation assessed using bottomless liftnet samples. Twenty-third Benthic Ecology Meeting.
- EPA (U.S. Environmental Protection Agency). 2001. Hydromodification chapter factsheet. United States Environmental Protection Agency. Available: <http://www.epa.gov/OWOW/NPS/MMGI/hydro.html> (December 2001).
- Epperly, S. P. and S. W. Ross. 1986. Characterization of the North Carolina Pamlico-Albemarle estuarine complex. National Marine Fisheries Service - Southeast Fisheries Center, Beaufort, NC, NMFS-SEFC-175, 55p.
- Evans, D. W., R.D. Kathman, and W.W. Walker. 2000. Trophic accumulation and depuration of mercury by blue crabs (*Callinectes sapidus*) and pink shrimp (*Penaeus duorarum*). *Marine Environmental Research* 49(5): 419-434.
- Feierabend, S. J. and J.M. Zelazny. 1987. Status report on our nation's wetlands. National Wildlife Federation, Washington, D.C., 50p.
- Ferguson, R. L. and L.L. Wood. 1994. Rooted vascular aquatic beds in the Albemarle-Pamlico estuarine system. NMFS, NOAA, Beaufort, NC, Project No. 94-02, 103 p.
- Fonseca, M. S. 1996. The role of seagrasses in nearshore sedimentary processes: a review. p. 261-286 *in* C. Roman and K. Nordstrom (editors). *Estuarine Shores: Hydrological, Geomorphological and Ecological Interactions*. Blackwell, Boston, MA.
- Fonseca, M. S., W. J. Kenworthy, and G. W. Thayer. 1998. Guidelines for the conservation and restoration of seagrasses in the United States and adjacent waters. NOAA Coastal Ocean Office, Silver Springs, Md., NOAA Coastal Ocean Program Decision Analysis Series No. 12, 222p.
- Funderburk, S. L., J.A. Mihursky, S.J. Jordan, and D. Riley. 1991. Habitat requirements for Chesapeake Bay living resources. Habitat Objectives Workgroup, Living Resources Subcommittee and Chesapeake Research Consortium with assistance from Maryland Department of Natural Resources, Solomons, MD.
- Garbisch, E. W., P.B. Woller, W.J. Bostian, and R.J. McCallum. 1973. Biotic techniques for shore stabilization. p. 405-407 *in* L.E. Cronin (editor). *Estuarine Research*. Academic Press Inc., New York, NY, II.
- Garcia, A. 1989. White shrimp (*Penaeus setiferus*) recruitment overfishing. *Marine and Freshwater research*. 47:59-65.
- Georgia. 2012. Rules and Regulations of the State of Georgia.

- Gibson, M.R. 1994. Alternative estimates of weakfish bycatch in shrimp trawl fisheries in the south Atlantic using shrimp effort and relative weakfish abundance data. Report to the Atlantic States Marine Fisheries Commission.
- Gilmore, G. and Trent X. 1974. Abundance of benthic macroinvertebrates in natural and altered estuarine areas. NMFS, NOAA Technical Report NMFS SSRF - 677.
- GMFMC (Gulf of Mexico Fishery Management Council). 2007. Amendment 27 To The Reef Fish Fishery Management Plan and Amendment 14 to the Shrimp Fishery Management Plan with a Supplemental Environmental Impact Statement, Regulatory Impact Review, and Regulatory Flexibility Act Analysis. Gulf of Mexico Fishery Management Council. June 2007.
- Godcharles, M. F. 1971. A study of the effects of a commercial hydraulic clam dredge on benthic communities in estuarine areas. Florida Department of Natural Resources, St. Petersburg, FL, 51p.
- Goldsborough, W. J. and W.M. Kemp. 1988. Light responses of submersed macrophytes: implication for survival in turbid waters. *Ecology* 69: 1775-1786.
- Grabowski, J. H., D. Pettipas, M.A. Dolan, A.R. Hughes, and D.L. Kimbro. 2000. The economic and biological value of restored oyster reef habitat to the nursery function of the estuary. NC Sea Grant, Morehead City, NC, FRG # 97-EP-6, 29p.
- Graff, L. and J. Middleton. 2003. Wetlands and fish: catch the link. NOAA, National Marine Fisheries Service, Silver Springs, MD, 48p.
- Gray, J. S., R. S. Wu, and Y.Y. Or. 2002. Effects of hypoxia and organic enrichment on the coastal marine environment. *Marine Ecology Progress Series* 238: 249-279.
- Guthrie, J. F. and C. W. Lewis. 1982. The clam-kicking fishery of North Carolina. *Marine Fisheries Review* 44(1): 16-21.
- Hackney, C. T., J. Grimley, M. Posey, T. Alphin, and J. Hyland. 1998. Sediment contamination in North Carolina's estuaries. Center for Marine Science Research, UNC-W, Wilmington, NC, Publication #198, 59p.
- Hackney, C. T., M.H. Posey, S.W. Ross, and A.R. Norris. 1996. A review and synthesis of data on surf zone fishes and invertebrates in the South Atlantic Bight and the potential impacts from beach renourishment. Prepared for Wilmington District, US Army Corps of Engineers. UNC-Wilmington, Wilmington, NC, 111 p.
- Hall, M.A., D.L. Alverson, and K.I. Metuzals. 2000. By-catch: problems and solutions. *Marine Pollution Bulletin* 41(1-6):204-219.
- Harrington, J. M., R. A. Myers, and A. A. Rosenberg. 2005. Wasted fishery resources: Discarded by-catch in the USA. *Fish Fish.* 6:350-361.
- Harrington, D., J. Watson, L. Parker, J. Rivers, and C. Taylor. 1988. Shrimp Trawl Design and Performance. Marine Extension Bulletin Number 12. Athens, GA: University of Georgia, Georgia Seagrass College Program.

- Hataway, D. 2010. Data Summary for 2" Bar Spacing TEDs compared to 4" Bar Spacing TEDs. Bycatch Reduction Engineering Project.
- Hines, K. L., R. A. Rulifson, and J. D. Murray. 1999. Performance of low-profile skimmer trawls in the inshore shrimp fishery of North Carolina. *North American Journal of Fisheries Management*. 19:569–580.
- Hawkins, J. H. 1980. Investigations of anadromous fishes of the Neuse River, North Carolina. DMF, Morehead City, NC, Special Science Report No. 34, 111p.
- Hettler, W. F. J. and D.L. Barker. 1993. Distribution and abundance of larval fish at two North Carolina inlets. *Estuarine, Coastal, and Shelf Sciences* 37: 161-179.
- Holland, B. F., Jr. 1988. Evaluation of Certified Trawl Efficiency Devices (TEDS) in North Carolina's Nearshore Ocean. N. C. Department of Natural Resources and Community Development, Division of Marine Fisheries, Completion Report, Project 2-439-R, 38 pp.
- Ianuzzi, T. J., M.P. Weinstein, K.G. Sellner, and J.C. Barrett. 1996. Habitat disturbance and marina development: An assessment of ecological effects. I. Changes in primary production due to dredging and marina construction. *Estuaries* 19(2A): 257-271.
- IMPLAN PRO version 2.0 (2000). Stillwater, MN: Minnesota IMPLAN Group.
- Ingraham, B. 2003. Night Vs. Day Bycatch Comparison for Shrimp Trawling in the Southern District of North Carolina. North Carolina Fisheries Resource Grant. FRG-98-FEG-46.
- Jackson, J. B. C., M. Kirby, W.H. Berger, K.A. Bjorndal, L.W. Botsford, B.J. Bourque, R.H. Bradbury, R. Cooke, J. Erlandson, J.A. Estes, T.P. Hughes, S. Kidwell, C.B. Lange, H.S. Lenihan, J.M. Pandolfi, C.H. Peterson, R.S. Steneck, M.J. Tegner, and R.R. Warner. 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293: 629-638.
- Jennings S., M.J. Kaiser and J.D. Reynolds. 2001. *Marine Fisheries Ecology*.
- Johnson, G. A. 2003. The role of trawl discards in sustaining blue crab populations. North Carolina Fisheries Resource Grant. FRG-99-EP-07.
- Johnson, S.K. 1978. Handbook of Shrimp Diseases. Sea Grant College Program. TAMU-SG-75-603. Texas A&M University, College Station. 23p.
- Jones, R. A. and T.M. Sholar. 1981. The effects of freshwater discharge on estuarine nursery areas of Pamlico Sound. NC Division of Marine Fisheries, Morehead City, NC, Project CEIP 79-11, 60p.
- Kennedy, F.S. and D.G. Barber. 1981. Spawning and recruitment of pink shrimp, *Penaeus duorarum*, off Eastern Florida. *Journal of Crustacean Biology*. 1:474-485.
- Kelleher, K., 2005. Discards in the world's marine fisheries. An update. FAO Fisheries Technical Paper No. 470. FAO, Rome: 131 p.
- Kenworthy, W. J. and D.E. Haurert. 1991. The light requirements of seagrasses: proceedings of a workshop to examine the capability of water quality criteria, standards and

- monitoring progress to protect seagrasses. National Oceanic and Atmospheric Administration, Beaufort, NC, Tech. Memo. NMFS-SEFC-287, 181p.
- Kirby, J., W. Maher, and F. Krikowa. 2001. Selenium, cadmium, copper, and zinc concentrations in sediments and mullet (*Mugil cephalus*) from the southern basin of Lake Macquarie, NSW Australia. Archives of environmental contamination and toxicology 40(2): 246-256.
- Knott, D.M., P. L. Fuller, A. J. Benson, and M.E. Neilson. 2012. *Penaeus monodon*. USGS Nonindigenous Aquatic Species Database, Gainesville, FL. U.S. Geological Survey. Available: <http://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=1209> (May 2011).
- Knuckey, I., R. Hudson, M. Koopman, S. Skoljarev, and J. Moore. 2008. Trials of T-90 mesh configuration in the Great Australian Bight Trawl Fishery. Australian Government. Fisheries Research and Development Corporation. FRDC Project 2007/063).
- Knutson, P. L. 1977. Planting guidelines for marsh development and bank stabilization. US Army Corps of Engineering Research Center, Fort Belvoir, Va.
- Kwon, Y. and C. Lee. 2001. Ecological risk assessment of sediment in wastewater discharging area by means of metal speciation. Microchemical Journal 70: 255-264.
- Larson, S.C, M.J. Van Den Avyle, E.L. Boseman, Jr. 1989. Species profiles: life histories and environment requirements of coastal fish and invertebrates (South Atlantic) – brown shrimp. U.S. Fish and Wildlife Service FWS/OBS-82/11.90. U.S. Army Corps of Engineers, TR EL-82-4. 14p.
- Lathman, F. F. 1951. Evidence of fish loss due to shrimping in Pamlico Sound. Committee Report to Atlantic States Marine Fisheries Commission, November 1, 1951. Appendix B, 4p.
- Lenihan, H. S. and C.H. Peterson. 1998. How habitat degradation through fishery disturbance enhances impacts of hypoxia on oyster reefs. Ecological Applications 8(1): 128-140.
- Lenihan, H. S., C.H. Peterson, J.E. Byers, J.H. Grabowski, and G.W. Thayer. 2001. Cascading of habitat degradation: oyster reefs invaded by refugee fishes escaping stress. Ecological Applications 11(3): 764-782.
- Lenihan, H. S. and F. Micheli. 2000. Biological effects of shellfish harvesting on oyster reefs: resolving a fishery conflict by ecological experimentation. Fishery Bulletin 98: 86-95.
- Lenihan, H. S., F. Micheli, S.W. Shelton, and C.H. Peterson. 1999. The influence of multiple environmental stressors on susceptibility to parasites: an experimental determination with oysters. Limnology and Oceanography 44: 910-924.
- Livingstone, D. R. 2001. Contaminant-stimulated reactive oxygen species production and oxidative damage in aquatic organisms. Marine Pollution Bulletin 42(8): 656-666.
- Loflin, R. K. 1995. The effects of docks on seagrass beds in the Charlotte Harbor estuary. Florida Scientist 58(2): 198-205.

- Logothetis, E. and D McCuiston. 2006. An assessment of the bycatch generated in the inside commercial shrimp fishery in southeastern North Carolina, 2004 & 2005. North Carolina Sea Grant Fisheries Resource Grant Program, Project #05-EP-04. 87 pp.
- Lowery, J. and K.T. Paynter. 2002. The importance of molluscan shell substrate. National Marine Fisheries Service, Unpublished report. 17p.
- Luetlich, R. A., J.E. McNinch, J.L. Pinckney, M.J. Alperin, C.S. Martens, H.W. Paerl, C.H. Peterson, and J.T. Wells. 1999. Neuse River estuary modeling and monitoring project, final report: Monitoring phase. Water Resources Research Institute, Raleigh, NC, 190p.
- Main, J., and G. Sangster. 1988. Scale damage and survival of young gadoid fish escaping from the codend of a demersal trawl. In: Proceedings of a workshop on "Trawl Net Selectivity and the Survival of Fish escaping from codends." University of Rhode Island Graduate School of Oceanography, May 16-17, 1988. Narragansett, Rhode Island. National Sea Grant Depository Publication #RIUW-88-002.
- Maiolo, J.R. 1981. "Historical Development of the Shrimp Fishery. In The Sociocultural Context and Occupational and Marketing Structures of the North Carolina Shrimp Fishery. Second Year Report to the NUC Sea Grant, Vol. 1.
- Maiolo, J.R. 2004. Hard Times and a Nickel a Bucket: Struggle and survival in North Carolina's shrimp industry. Chapel Hill, NC: Chapel Hill Press.
- Maiolo, J.R., J. Bort, W. Still, M. Fisch, R. Faisan. 1980. Historical development of the shrimp fishery (The sociocultural context and occupational and marketing structure of the North Carolina shrimp fishery. Greenville, NC: East Carolina University.
- Mangel, M. 1993. Effects of high seas driftnet fisheries on the Northern right whale dolphin *Lissodelphis borealis*. *Ecological Applications*. 3(2), 221-229.
- Manisseri, M. K. and N.R. Menon. 1995. Copper-induced damage to the hepatopancreas of the penaeid shrimp *Metapenaeus dobsoni* - an ultrastructural study. *Diseases of Aquatic Organisms* 22(1): 51-57.
- Manisseri, M. K. and N.R. Menon. 2001. Effect of sublethal copper on growth efficiency of the shrimp, *Metapenaeus dobsoni*. *Journal of the Marine Biological Association of India* 43(1-2): 81-90.
- Marburger, J. E., W.E. Johnson, T.S. Gross, D.R. Douglas, and J. Di. 2002. Residual organochlorine pesticides in soils and fish from wetland restoration areas in central Florida. *Wetlands* 22(4): 705-711.
- Marcus, J. M. and T.P. Stokes. 1985. Polynuclear aromatic hydrocarbons in oyster tissue around three coastal marinas. *Bulletin of Environmental Contamination and Toxicology* 35: 835-844.
- Matheson, R. E. Jr., D.K. Camp, S.M. Sogard, and K.A. Bjorgo. 1999. Changes in seagrass-associated fish and crustacean communities on Florida Bay mud banks: the effects of recent ecosystem changes? *Estuaries* 22(2B): 534-551.

- Matoura, R. F. C. and E.M.C. Woodward. 1983. Conservative behavior of riverine dissolved organic carbon in the Severn estuary: chemical and geochemical implications. *Geochimica Cosmochimica Acta* 47: 1293-1309.
- McCoy, E.G. 1968. Movement, growth and mortality of brown shrimp (*Penaeus aztecus*) marked and released in Swanquarter Bay, Pamlico Sound North Carolina. North Carolina Department of Conservation and Development, Division of Commercial and Sports Fisheries, Special Scientific Report No. 15, 26p.
- McCoy, E.G. 1972. Dynamics of North Carolina Commercial Shrimp Populations. North Carolina Department of Natural and Economic Resources, Division of Commercial and Sports Fisheries, Special Scientific Report No. 21, 53p.
- McCoy, E.G. and J.T. Brown. 1967. Migration and Growth of Commercial Penaeid Shrimps in North Carolina. Ann. Rep., Spec. Sci. Rep. 11, North Carolina Department of Conservation and Development, Division of Commercial and Sports Fisheries, 29 p.
- McKenna, S. and A.H. Clark. 1993. An examination of alternative fishing devices for the estuarine shrimp and crab trawl fisheries. Final Report to the Albemarle/Pamlico Estuarine Study, Project No. 93-11, 34p.
- McKenna, S. and J.T. Camp. 1992. An examination of the blue crab fishery in the Pamlico River estuary. Final Report to the Albemarle/Pamlico Estuarine Study, Project No. 92-08, 101p.
- McKenna, S. A. and J. P. Monaghan, Jr. 1993. Gear development to reduce bycatch in the North Carolina trawl fisheries. Completion Report for Cooperative Agreement No. NA90AA-SK052 to Gulf and South Atlantic Fisheries Development Foundation, Contract No. 43-01, North Carolina Department of Environment, Health, and Natural Resources, Division of Marine Fisheries.
- McKenna, S. A., G. Judy, C. P. Lewis, and J. Schoolfield. 1996. Evaluation of trawl efficiency device/bycatch reduction device in estuarine and nearshore waters of North Carolina. Completion Report NOAA, No. NA 47FF0016, North Carolina Department of Environment, Health, and Natural Resources, Division of Marine Fisheries, 37p.
- Meyer, D. L., E.C. Townsend, and P.L. Murphey. 1996. Final report for the project evaluation of restored wetlands and enhancement methods for existing restorations. National Oceanic and Atmospheric Administration, Office of Habitat Conservation Restoration Center, Silver Springs, MD.
- Micheli, F. M. and C. H. Peterson. 1999. Estuarine vegetated habitats as corridors for predator movement. *Conservation Biology* 13(4): 869-881.
- Minello, T. J. 1999. Nekton densities in shallow estuarine habitats of Texas and Louisiana and the identification of Essential Fish Habitat. p. 43-75 *in* Benaka, L. R. ed. *Fish Habitat: Essential Fish Habitat and Rehabilitation*. American Fisheries Society, Bethesda, Maryland, 459p.
- Minello, T.J. and R.J. Zimmerman. 1983. Fish predation on juvenile brown shrimp, *Penaeus aztecus* Ives: The effect of simulated *Spartina* structure on predation rates. *Journal of Experimental Marine Biology and Ecology* 72: 211-231.

- Minello, T.J., R.J. Zimmerman, E.X. Martinez .1989. Mortality of young brown shrimp, *Penaeus aztecus*, in estuarine nurseries. Transactions of American Fisheries Society 118: 693–708.
- Mitsch, W. J. and J.G. Gosselink. 1993. Wetlands, Second Edition. Van Nostrand Reinhold, New York, NY, Second Edition, 772p.
- Mock, C. R. 1966. Natural and altered estuarine habitats of penaeid shrimp. Proceedings Gulf Caribbean Fish Institute 19th Annual Session: 86-98.
- MSC (Moratorium Steering Committee). 1996. Final report of the Moratorium Steering Committee to the Joint Legislative Commission on Seafood and Aquaculture of the North Carolina General Assembly. N.C. Sea Grant College Program, Raleigh, NC, NC-SG-96-11, 155p.
- Muncy, R.J. 1984. Species profiles: life histories and environment requirements of coastal fish and invertebrates (south Atlantic) – white shrimp. U.S. Fish and Wildlife Service FWS/OBS-82/11.27. U.S. Army Corps of Engineers, TR EL-82-4, 19p.
- Murawski, S.A. 1995. Biological implications of bycatch. In Proceedings of the East Coast Bycatch Conference, April 7-8, 1995, p. 31-39.
- Murphey, P. L. and M. S. Fonseca. 1995. Role of high and low energy seagrass beds as nursery areas for *Penaeus duorarum* in North Carolina. Marine Ecology Progress Series 121: 91-98.
- Murray, J. D., J. J. Bahen, and R. A. Rulifson. 1991. Management Considerations for By-Catch in the North Carolina and Southeast Shrimp Fishery. Fisheries 17(1): 21-26.
- Murray, J. D., J. L. Gearhart, R. A. Rulifson, and C. W. Wescott. 1995. Introduction of large mesh webbing in the belly and wings of traditional shrimp trawls to reduce bycatch in inshore waters. Saltonstall-Kennedy Final Report, Project NA37FD008801, February 1995, 75p.
- Nance, J. M. (Editor). 1998. Report to Congress. Southeastern United States Shrimp Trawl Bycatch Program, 154p.
- Nance, J.M., C.W. Cailouet Jr., R.A. Hart. Size-composition of annual landings in the white shrimp, *Litopenaeus setiferus*, fishery of the Northern Gulf of Mexico, 1960-2006: Its trend and relationships with other fishery-dependent variables. Marine Fisheries Review. 72: 1-13.
- Newell, R. I. E. 1988. Ecological changes in the Chesapeake Bay: are they the result of overharvesting the American oyster? p. 536-546 in M.P. Lynch and E.C. Krome (eds.). Understanding the estuary: advances in Chesapeake Bay research. Chesapeake Bay Research Consortium, Baltimore, MD, Publication 129.
- NCDMF. North Carolina Division of Marine Fisheries Bycatch Reduction Device Certification Procedures.

- NCDMF (North Carolina Division of Marine Fisheries). 1991. Shrimp and crab trawling in North Carolina's estuarine waters. North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, 154p.
- NCDMF (North Carolina Division of Marine Fisheries). 1999. Shrimp and crab trawling in North Carolina's estuarine waters. DENR, Morehead City, NC, Report to NC Marine Fisheries Commission, 121p.
- NCDMF (North Carolina Division of Marine Fisheries). 2001. North Carolina oyster fishery management plan. N.C. Department of Environment and Natural Resources, Division of Marine Fisheries, 225p.
- NCDMF (North Carolina Division of Marine Fisheries). 2005. North Carolina southern flounder fishery management plan. North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, 335p.
- NCDMF (North Carolina Division of Marine Fisheries). 2006. North Carolina Fishery Management Plan for Shrimp. North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries.
- NCDMF (North Carolina Division of Marine Fisheries). 2011. Strategic Habitat Area Nominations for Pamlico Sound System, North Carolina (Region 2) Final. North Carolina Department of Environment and Natural Resources. Division of Marine Fisheries, NC. 128 pp
- NCDMF (North Carolina Division of Marine Fisheries). 2012. North Carolina Fishery Management Plan for Shrimp Revision. North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries.
- NCDMF (North Carolina Division of Marine Fisheries). 2012. Pamlico Sound Independent assessment. North Carolina Department of Environment and Natural Resources. Division of Marine Fisheries, NC. Grant NC-F-70128 pp.
- NCFA (North Carolina Fisheries Association). 2004. Press Release. Commerce finds four more countries dumped shrimp in U.S. Market. Available: <http://www.ncfish.org/article.asp?id=123> (July 2004).
- NMFS (National Marine Fisheries Service). 1995. Cooperative research program addressing finfish bycatch in the Gulf of Mexico and South Atlantic shrimp fisheries. NMFS Southeast Regional Office, St. Petersburg, Florida, 67p.
- NMFS and USFWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service). 1991a. Recovery Plan for U.S. Population of Atlantic Green Turtle. National Marine Fisheries Service, Washington D.C., 52p.
- NMFS and USFWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service). 1991b. Recovery Plan for U.S. Population of Loggerhead Turtle. National Marine Fisheries Service, Washington D.C., 64p.
- NMFS and USFWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service). 1992. Recovery Plan for Leatherback Turtles in the U.S. Caribbean, Atlantic, and Gulf of Mexico. National Marine Fisheries Service, Washington, D.C., 65p.

- NMFS and USFWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service). 1993a. Recovery Plan for the Kemp's Ridley Sea Turtle. National Marine Fisheries Service, St. Petersburg, Florida, 40p.
- NMFS and USFWS (National Marine Fisheries Service and U.S. Fish and Wildlife Service). 1993b. Recovery Plan for Hawksbill Turtles in the U.S. Caribbean Sea, Atlantic Ocean, and Gulf of Mexico. National Marine Fisheries Service, St. Petersburg, Florida, 52p.
- NOAA. 2004. Status of Bycatch Reduction Device (BRD) Performance and Research in North-Central and Western Gulf of Mexico. United States Department of Commerce. National Oceanic and Atmospheric Administration. National Marine Fisheries Service. Southeast Fisheries Science Center. Mississippi Laboratories. Pascagoula, Mississippi.
- National Oceanographic and Atmospheric Administration (NOAA). 2006. Bycatch in the South Atlantic, Gulf of Mexico, and Caribbean. Southeast Bycatch Workshop Proceedings May 16-18, 2006.
- NOAA. 2008. Bycatch Reduction Device Testing Manual. National Marine Fisheries Service. Southeast Regional Office. St. Petersburg, Florida
- NOAA (National Oceanic and Atmospheric Administration). 2010. List of Fisheries for 2011. Federal Register 75:215 (8 November 2010): 68468-68504.
- NOAA (National Oceanic and Atmospheric Administration). 2011. Fisheries Products. NOAA Fisheries: Office of Science and Technology. Available: http://www.st.nmfs.noaa.gov/st1/fus/fus10/07_supply2010.pdf (May 2012).
- National Marine Fisheries Service (NMFS). 2012a. Reinitiation of Endangered Species Act Section 7 Consultation on the Continued Implementation of the Sea Turtle Conservation Regulations, as Proposed to Be Amended, and the Continued Authorization of the Southeast U.S. Shrimp Fisheries in Federal Waters under the Magnuson-Stevens Act. Biological Opinion. May 8, 2012.
- NMFS (National Marine Fisheries Service). 2012. ESA Section 7 consultation on the continued implementation of the sea turtle conservation regulations, as proposed to be amended, and the continued authorization of the southeast U.S. shrimp fisheries in federal waters under the Magnuson-Stevens Act. Biological Opinion.
- Noble, E. B. and R.J. Monroe. 1991. Classification of Pamlico Sound Nursery Areas: Recommendations for Critical Habitat Criteria. North Carolina Department of Environment, Health, and Natural Resources, Morehead City, NC, A/P Project No. 89-09, 70p.
- North Carolina Sea Grant. 1997. Coastal water quality. NC State University, Raleigh, NC, UNC-SG-97-04, 72p.
- North Carolina State Demographics (2012). North Carolina Office of State Budget and Management. Available: <http://demog.state.nc.us>
- Orbach, M. K. and J. C. Johnson 1988. Transformation of Fishing Communities: A public policy perspective. In Marine Resource Utilization: A conference on social science issues. Proceedings of the Mississippi-Alabama Sea Grant Consortium, Mobile, Alabama.

- O'Rear, C. W. 1983. A study of river herring spawning and water quality in Chowan River. NC Department of Natural Resources and Community Development, DMF, Raleigh, NC, Complete Report, Project AFC-17, 31p.
- Orth, R. J., J. Simons, J. Capelli, V. Carter, L. Hindman, S. Hodges, K. Moore, and N. Rybicki. 1986. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries - 1985. US EPA, Washington, DC, Final report.
- Orth, R. J., K.A. Moore, and J.F. Nowak. 1990. Monitoring seagrass distribution and abundance patterns: A case study from the Chesapeake Bay. *in* US Fish and Wildlife Service. Federal coastal wetland mapping. U.S. Fish and Wildlife Service, Biological Report 90(18): 111-123.
- Paerl, H. W., J. Pinckney, J. Fear, and B. Peierls. 1998. Ecosystem response to internal watershed organic matter loading: Consequences for hypoxia in the eutrophying Neuse River Estuary, North Carolina. *Marine Ecological Progress Series* 166: 17-25.
- Paerl, H. W., M.M. Mallin, C.A. Donahue, M. Go, and B.L. Peierls. 1995. Nitrogen loading sources and eutrophication of the Neuse River, North Carolina: direct and indirect roles of atmospheric deposition. UNC - Chapel Hill, Water Resources Research Institute, Chapel Hill, NC, Publication 291.
- Pate, P. P. Jr. and R. Jones. 1981. Effects of upland drainage on estuarine nursery areas of Pamlico Sound, North Carolina. UNC Sea Grant , Raleigh, NC, Pub. No. UNC-SG-WP-10, 24p.
- Pattilo, M. E., D. M. N. T.E. Czapla, and M.E. Monaco. 1997. Distribution and abundance of fishes and invertebrates in Gulf of Mexico estuaries. Volume II: Species life history summaries. NOAA/NOS Strategic Environmental Assessment Division, Silver Springs, MD, ELMR Rep. No. 11. 377p.
- Pearce, K.B., D.W. Moyer, and S.K. Strasser. 1988. Evaluation of trawl excluder devices in the Pamlico Sound shrimp fishery. Albemarle-Pamlico Estuarine Study Rep. No.88-07. 46p.
- Peterson, B. J. and R.W. Howarth. 1987. Sulfur, carbon, and nitrogen isotopes used to trace organic matter flow in the salt-marsh estuaries of Sapelo Island, Georgia. *Limnology and Oceanography* 32(6): 1195-1213.
- Peterson, C. H., H.C. Summerson, and S.R. Fegley. 1987. Ecological consequences of mechanical harvesting of clams. *Fisheries Bulletin* 85(2): 281-298.
- Peterson, C. H., J.H. Grabowski, and S.P. Powers. 2003. Quantitative enhancement of fish production by oyster reef habitat: restoration valuation. *Marine Ecology Progress Series* 264: 249-264.
- Peterson, C. H. and N.M. Peterson. 1979. The ecology of intertidal flats of North Carolina: A community profile. U.S. Fish and Wildlife Service, OBS-79/39, 73 p.
- Peterson, M. S., B.H. Comyns, J.R. Hendon, P.J. Bond, and G.A. Duff. 2000. Habitat use by early life-stages of fishes and crustaceans along a changing estuarine landscape:

- difference between natural and altered shoreline sites. *Wetland, Ecology, and Management* 8(2-3): 209-219.
- Peuser, R. 1996. Estimates of finfish bycatch in the south Atlantic shrimp fishery. Final report Southeastern Area Monitoring and Assessment Program SEAMAP-south Atlantic Committee, Shrimp Bycatch Work Group, 62p.
- Price, A.B. and J.L. Gearhart. 2011. Evaluations of turtle excluder device (TED) performance in the U.S. southeast Atlantic and Gulf of Mexico skimmer trawl fisheries. NOAA technical Memorandum NMFS-SEFSC-615,15p.
- Pulver, J. R., E. Scott-Denton and J.A. Williams. 2012. Characterization of the U.S. Gulf of Mexico skimmer trawl fishery based on observer coverage. NOAA Technical Memorandum NMFS-SEFSC-636, 27 p.
- Purvis, C.E., and E.G. McCoy. 1972. Overwintering Pink Shrimp (*Penaeus duorarum*) in Core and Pamlico Sounds, N.C. North Carolina Department of Natural and Economic Resources, Division of Commercial and Sports Fisheries, Special Scientific Report No. 21, 53p.
- Riggs, S. R., E.R. Powers, J. T. Bray, P.M. Stout, C. Hamilton, D. Ames, R. Moore, J. Watson, S. Lucas, and M. Williamson. 1989. Heavy metal pollutants in organic rich muds of the Pamlico River estuarine system: their concentration, distribution, and effects upon benthic environments and water quality: Albemarle-Pamlico Estuarine Study. Project No. 89-06. US EPA and NC DNRCD, Raleigh, NC, 108p.
- Riggs, S. R., J.T. Bray, E.R. Powers, C. Hamilton, D. Ames, D. Yeates, K. Owens, S. Lucas, J. Watson, and M. Williamson. 1991. Heavy metal pollutants in organic-rich muds of the Neuse River Estuary: their concentration and distribution. Albemarle-Pamlico Estuarine Study Report. Project no. 90-07. DENR, Raleigh, 168p.
- Roblee, M. B. and W.J. DiDomenico. 1992. Seagrass die-off in Florida Bay, Everglades National Park. *Park Science* 11: 21-23.
- Roelofs, E. W. 1950. Observations of the capture of small fish by the shrimp trawls. Annual Report, Institute of Fisheries Research UNC, Morehead City, NC: 111-115.
- Ross, S. W. and S.P. Epperly. 1985. Chapter 10: Utilization of shallow estuarine nursery areas by fishes in Pamlico Sound and adjacent tributaries, North Carolina. p. 207-232 *in* A. Yanez-Arancibia (ed.). *Fish Community Ecology in Estuaries and Coastal Lagoons: Towards and Ecosystem Integration*. DR (R) UNAM Press, Mexico, 654p.
- Rozaz, L.P., and T.J. Minello. 2011. Variation in penaeid shrimp growth rates along an estuarine salinity gradient: implications for managing river diversions. *Journal of Experimental Marine Biology and Ecology* 397: 196-207.
- Rozas, L. P. and R.J. Zimmerman. 2000. Small-scale patterns of nekton use among marsh and adjacent shallow nonvegetated areas of the Galveston bay estuary, Texas (USA). *Marine Ecology Progress Series*. 193: 217-239.

- Rozas, L. P. and W.E. Odum. 1987. The role of submerged aquatic vegetation in influencing the abundance of nekton on contiguous tidal freshwater marshes. *Journal of Experimental Marine Biology and Ecology* 114(2-3): 289-300.
- Rudershausen, P. J., and J. F. Weeks. 1999. Mechanical efficiency of experimental skimmer trawl frames. North Carolina Sea, Project#. 99-FEG-31, 24 p.
- SAFMC (South Atlantic Fishery Management Council). 1993. Fishery management plan for the shrimp fishery of the South Atlantic region including a final environmental impact statement and regulatory impact review. SAFMC, Charleston, SC, 184 pp. + appendices.
- SAFMC (South Atlantic Fishery Management Council). 1998. Final habitat plan for the South Atlantic region: Essential Fish Habitat requirements for fishery management plans of the South Atlantic Fishery Management Council. SAFMC, Charleston, SC.
- Sanger, D. M. and A.F. Holland. 2002. Evaluation of the impacts of dock structures on South Carolina estuarine environments. SC Department of Natural Resources, Marine Resources Research Institute, Charleston, SC, Tech. Rep. No. 99, 82p.
- Saoud, P., and D. A. Davis. 2003. Salinity tolerance of brown shrimp *Farfantepenaeus azetecus* as it relates to postlarval and juvenile survival, distribution, and growth in estuaries. *Estuaries* 26: 970-974.
- Schoof, R. 1980. Environmental impact of channel modification. *Water Resources Bulletin* 16(4): 697-701.
- Schueler, T. R. 1997. Impact of suspended and deposited sediment. *Watershed Protection Techniques* 2(3): 443-444.
- Scott-Denton, E., P. Cryer, J. Gockett, M. Harrelson, K. Jones, J. Nance, J. Pulver, R. Smith, and J.A. Williams. 2006. Skimmer Trawl Fishery Catch Evaluations in Coastal Louisiana, 2004 and 2005. *Marine Fisheries Review* 2006, Vol 68 (1-4): 30 - 35.
- Scientific Council on Amphibians and Reptiles. 2011. Reevaluation of Status Listings for Jeopardized Amphibians and Reptiles in North Carolina. Submitted to the Nongame Wildlife Advisory Committee of the North Carolina Wildlife Resources Commission.
- Seldberg, C. D., L.A. Eby, and L.B. Crowder. 2001. Hypoxia in the Neuse River Estuary: Responses of blue crabs and crabbers. *North American Journal of Fisheries Management* 21(2): 358-366.
- Sessions, K. and T. Thorpe. 2006. Catch potential and condition of shrimp and by-catch associated with a new commercial shrimp set design and new RCGL shrimp set design. North Carolina Sea Grant Project # 05-FEG-12. 48p.
- Shafer, D. J. 1999. The effects of dock shading on the seagrass *Halodule wrightii* in Perdido Bay, Alabama. *Estuaries* 22(4): 936-943.

- Shervette, V.R. and F. Gelwick. 2008. Relative nursery function of oyster, vegetate marsh edge, and nonvegetated bottom habitats for juvenile white shrimp (*Litopenaeus setiferus*). *Wetlands Ecology Mangement* 16: 405-419.
- Simpson, D.G. 1990. A study of Marine Recreational Fisheries in Connecticut. Federal Aid in Sport Fish Restoration Project F-54-R, Job 8 Final Report. Connecticut Department of Energy and Environmental Protection, Bureau of Fisheries and Wildlife, Division of Marine Fisheries, 3p.
- Smith, E.M. and P.T. Howell. 1987. The effects of bottom trawling on American lobster, *Homarus americanus*, in Long Island Sound. *Fisheries Bulletin* 85(4): 737-744.
- Smolowitz, R. 1983. Mesh size and the New England groundfishery-applications and implications. NOAA Tech Report. NMFS SSRF-771.
- STAC (Sea Turtle Advisory Committee). 2006. Sea turtle interactions with North Carolina fisheries review and recommendations to the North Carolina Marine Fisheries Commission. North Carolina Department of Natural Resources, Division of Marine Fisheries, 72p.
- St. Amant, L.S., and J.G. Broom, and T.B. Ford. 1965. Studies of the brown shrimp, *Penaeus aztecus*, in Barataria Bay, Louisiana. 1965. *Bulletin of Marine Science of the Gulf and Caribbean*. 18: 1-16.
- Steele, P. 2002. Stock assessment profile for the penaeid shrimp fisheries of the southeastern United States and Gulf of Mexico. Report to the Florida Marine Fisheries Commission. 227p.
- Stevenson, J. C., L.W. Staver, and K.W. Staver. 1993. Water quality associated with survival of submerged aquatic vegetation along an estuarine gradient. *Estuaries* 16(346-361).
- Takade-Heumacher, H. and C. Batsavage. 2009. Stock status of North Carolina southern flounder (*Paralichthys lethostigma*). North Carolina Department of Environment and Natural Resources, Division of Marine Fisheries, Morehead City, NC. 93 p.
- Tarplee, W. H. Jr., D.E. Louder, and A.J. Weber. 1971. Evaluation of the effects of channelization on fish populations in North Carolina's coastal plain streams. North Carolina Wildlife Resources Commission, Raleigh, NC.
- Teal, J. 1962. Energy flow in salt marsh macrophyte production: a review. *Ecology* 43: 614-624.
- Tenore, K. R. 1972. Macrobenthos of the Pamlico River estuary, North Carolina. *Ecological Monographs* 42: 51-69.
- Thayer, G. W., W.J. Kenworthy, and M.S. Fonseca. 1984. The ecology of eelgrass meadows of the Atlantic coast: a community profile. U.S. Fish and Wildlife Service, FWS/OBS-84/02, 147p.
- Theiling, D. 1988. Assessment of participation and resource impact of shrimp baiting in coastal South Carolina during 1987. S.C. Marine Research Center Technical Report No. 69. 41p.

- Turner, R. E. 1977. Intertidal vegetation and commercial yields of Penaeid shrimp. Transactions of the American Fisheries Society 106(5): 411-416.
- Twilley, R. R., W.M. Kemp, K.W. Staver, J.C. Stevenson, and W.R. Boynton. 1985. Nutrient enrichment of estuarine submersed vascular plant communities. 1. Algal growth and effects on production of plants and associated communities. Marine Ecology Progress Series 23: 179-191.
- Uncles, R. J., J.A. Stephens, and T.Y. Woodrow. 1988. Seasonal cycling of estuarine sediment and contaminant transport. Estuaries 11: 108-116.
- U.S. Department of Commerce. 2003. Fisheries of the United States, 2002. Current Fishery Statistics No. 2002. National Oceanic and Atmospheric Administration, National Marine Fisheries Service.
- Vaughan, D.S. 1994. Incorporation of shrimp trawl bycatch in the assessment of Atlantic weakfish stock, 1982-1992. Report to ASMFC weakfish Technical Committee, National Marine Fisheries Service, Beaufort, N.C. Lab.
- Vaughan, D.S., and J.M. Nance. 1998. Estimates of bycatch of mackerel and cobia in U.S. south Atlantic shrimp trawls. Report to Gulf of Mexico and South Atlantic Fishery Management Councils, 25 pp.
- Vaughan, D.S., R.J. Seagraves and K. West. 1991. An assessment of the status of the Atlantic weakfish stock, 1982-1988. Special Report No. 21. Atlantic States Marine Fisheries Commission.
- Voudrias, E. A. and C.L. Smith. 1986. Hydrocarbon pollution from marinas in estuarine sediments. Estuarine Coastal Shelf Sciences 22: 271-284.
- Walker T., J. Newman, and I. Knuckley. 2010. Promoting industry uptake of gear modifications to reduce bycatch in the South East trawl fishery. Australian Government: Fisheries Research and Development Corporations. Project no. 2001/006.
- Wannamaker, C. M. and J.A. Rice. 2000. Effects of hypoxia on movements and behavior of selected estuarine organisms from the southeastern United States. Journal of Experimental Marine Biology and Ecology 249: 145-163.
- Wassenberg, T.J. and B.J. Hill. 1989. The effect of trawling and subsequent handling on the survival rates of the by-catch of Prawn trawlers in Moreton Bay, Australia. Fisheries Research 7: 99-110.
- Waters, C. T. and C.D. Thomas. 2001. Shoreline hardening effects on associated fish assemblages in five North Carolina coastal rivers. North Carolina Wildlife Resources Commission, Raleigh, NC, 20p.
- Watson, J. W. 1984. *Configurations and relative efficiencies of shrimp trawls employed in southeastern United States waters*. US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.

- Weis, J. S. and P. Weis. 1989. Effects of environmental pollutants on early fish development. *Aquatic Sciences* 1(1): 45-55.
- Weis, J. S., P. Weis, and T. Proctor. 1998. The extent of benthic impacts of CCA-treated wood structures in Atlantic coast estuaries. *Archives of Environmental Contamination and Toxicology* 34(4): 313-322.
- Wendt, P. H., R.F. Van Dolah, M.Y. Bobo, and J.J. Manzi. 1990. Effects of marina proximity on certain aspects of the biology of oysters and other benthic macrofauna in a South Carolina estuary. South Carolina Wildlife and Marine Resources Department, Charleston, SC, South Carolina Marine Resources Center Tech. Rep. No. 74, 49p.
- Whitaker, D. J., J.E. Jenkins and L.B. DeLancey. (1991). Catch rates and size of white shrimp caught by cast net with different mesh sizes. South Carolina Marine Research Center Technical Report No. 77. 18 p.
- Wilbur, A. R. and M.W. Pentony. 1999. Human-induced nonfishing threats to essential fish habitat in the New England region. p. 299-321 *in* L.R. Benaka (ed.). *Fish Habitat: Essential Fish Habitat and Rehabilitation*. American Fishery Society, Silver Springs, MD, Symposium 22, 459p.
- Williams, A.B. 1965. Marine decapods crustaceans of the Carolinas. *U.S. Fish and Wildlife Service Fisheries Bulletin* 65:1-298.
- Williams, B. 1990. A survey of shrimp cast netters in Georgia during 1989. Georgia Department of Natural Resources. Coastal Resources Division. Project NA90AA-H-SK042.
- Wolff, M. 1972. A study of the North Carolina scrap fishery. North Carolina Department of Natural and Environmental Resources, Division of Commercial and Sport Fisheries, Special Scientific Report No. 20, 27p.
- Zimmerman, R. T., T. Minello, and M. Gastiglione. 1989. Oyster reef as habitat for estuarine macrofauna. National Oceanic and Atmospheric Administration, Springfield, VA, NOAA Technical Memorandum NMFS-SEFC-249.

14.0 APPENDICES

14.1 APPENDIX1- AMENDMENT 1- PUBLIC COMMENT

Public comment and recommendations by five different advisory committees are required as part of the FMP process to ensure adequate input from interested citizens. Both Shrimp AC and DMF recommendations regarding the management of bycatch in the shrimp fishery were vetted through the MFC's Southern AC, Northern AC, Habitat and Water Quality AC, Finfish AC and the Shellfish/Crustacean AC during January 2014.

Trawling in the New River above the Highway 172 Bridge

Shrimp FMP Advisory Committee Recommendation

Recommend the MFC allow skimmer and otter shrimp trawling in the New River special secondary nursery area.

Division Recommendation

Status quo

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as DMF	Same as DMF	Same as DMF	Motion to support DMF, failed 3-3-1	Same as DMF

Evaluation of the skimmer trawl and other gears used for shrimping in North Carolina

Shrimp FMP Advisory Committee Recommendations

Allow hand cast netting of shrimp in all closed areas and increase the limit to four quarts per person.

Recommend the MFC require a fishing license from DMF to fish a cast net.

Division Recommendation

Status quo

MFC Advisory Committee Recommendations

Four quart limit

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as AC	Same as AC	Abstain	Same as AC	Same as AC

Cast net license

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Motion to support AC, failed 3-3-2	<i>Status quo</i>	Motion to support suggestion of a license, failed 1-2-4	Motion to support license, failed 2-5	No Action

The use of TEDs in commercial skimmer trawl operations

Shrimp FMP Advisory Committee Recommendation

No Recommendation

Division Recommendation

Upon federal adoption of TEDs in skimmer trawls, the division will support the federal requirement. Rule 15A NCAC 03L .0103 (g) allows for state enforcement.

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as DMF	Same as DMF	Abstain	Same as DMF	Same as DMF

Consideration of a commercial live bait shrimp fishery in North Carolina

Shrimp FMP Advisory Committee Recommendation

Recommend NC establish a permitted live shrimp bait fishery and for DMF to craft the guidelines and permit fees after reviewing permitted operations in other states.

Division Recommendation

Status quo

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as AC	<i>Status quo</i>	Same as AC	<i>Status quo</i>	<i>Status quo</i>

Gear Modifications in North Carolina shrimp trawls to reduce finfish bycatch

Shrimp FMP Advisory Committee Recommendation

Recommend the Marine Fisheries Commission allow any federally certified BRD in all NC internal and offshore waters.

Recommend the MFC update and certify bycatch reduction devices through the state bycatch reduction program.

Recommend the MFC convene an ongoing stakeholder workgroup charged with suggesting new trawl gear or trawl gear modification.

Initiate industry testing of new or modified bycatch reduction devices and gear modifications under the supervision of the NC DMF. After testing and collection of scientific data, regulations should be implemented to require or allow such devices or modifications to be used in N.C. internal and offshore waters.

Recommend the MFC test a three-inch bar-spaced turtle excluder device to see if it can be certified as a bycatch reduction device.

Recommend the MFC allow the shrimp industry a two year period to test bycatch reduction devices.

Division Recommendations

Allow any federally certified BRD in all NC internal and offshore waters.

Update the scientific testing protocol for the state BRD certification program.

Convene a stakeholder group to initiate industry testing of minimum tail bag mesh size, T-90 panels, skylight panels, and reduced bar spacing in TEDs.

- Upon securing funding, testing in the ocean and internal waters will consist of three years of data using test nets compared to a control net with a Florida Fish Eye, and a federally approved TED.
- Results should minimize shrimp loss and maximize reduction of bycatch of finfish. Promising configurations will be brought back to the MFC for consideration for mandatory use.
- This stakeholder group may be partnered with DMF and Sea Grant.
- Members could consist of fishermen, net/gear manufacturers and scientist/gear specialists.

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as DMF	Same as DMF	Same as DMF	Same as DMF	Same as DMF

Shrimp FMP Advisory Committee Recommendation

No recommendation

Division Recommendation

Require either a T-90 panel/ square mesh tailbag or other applications of square mesh panel (e.g., skylight panel), reduced bar spacing in a TED, or another federal or state certified BRD in addition to existing TED and BRD requirements in all skimmer and otter trawls.

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as DMF	<i>Status quo</i>	Abstain	Same as DMF	Same as DMF but to require one year of testing

Effort Management for bycatch reduction in the North Carolina shrimp trawl fishery

Shrimp FMP Advisory Committee Recommendation

No Recommendation

Division Recommendation

Status quo

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
No Discussion	No Discussion	Abstain	No Discussion	No Discussion

Characterization of the North Carolina commercial shrimp trawl fleet

Shrimp FMP Advisory Committee Recommendation

No Recommendation

Division Recommendation

In order to put a cap on fleet capacity as a management tool, establish a maximum combined headrope length of 220 feet in all internal coastal waters where there are no existing maximum combined headrope requirements (i.e., 90 foot requirement).

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Do not support DMF recommendation	<i>Status quo</i>	Abstain	Same as DMF but include a three year phase in period	Same as DMF

Area restrictions to reduce shrimp trawl bycatch in North Carolina’s internal coastal waters

Shrimp FMP Advisory Committee and DMF Recommendation

Recommend to the MFC to prohibit shrimp trawling in the IWW channel from the Sunset Beach Bridge to the SC line, including Eastern Channel, lower Calabash River and Shallotte River.

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as DMF	Same as DMF	Same as DMF	Same as DMF	Same as DMF

Shrimp FMP Advisory Committee Recommendation

No recommendation

Division Recommendation

Recommend the MFC Habitat and Water Quality Committee considers changing the designation of special secondary nursery areas that have not been opened to trawling since 1991 to permanent secondary nursery areas.

MFC Advisory Committee Recommendations

Southern AC	Northern AC	Habitat AC	Finfish AC	Shell/Crust AC
Same as DMF	Same as DMF	Same as DMF	Motion to support DMF, failed 3-3-1	Same as DMF

The Marine Fisheries Commission at its November meeting requested that an additional recommendation also be included during the public comment period.

Marine Fisheries Commission Recommendation

Convene a stakeholder group to initiate a three year study to test minimum tail bag mesh size, T-90 (square mesh) panels, skylight panels, reduced bar spacing in TEDs and any other new methods of reducing unwanted finfish bycatch to achieve a minimum of a 40 percent reduction of finfish by weight.

- Compare these to a control net with a Florida fish eye, a federally approved TED, and a one and half inch mesh tail bag.
- The stakeholder group should partner with DMF and Sea Grant to help secure funding for the study.
- If the 40 percent target reduction by weight in finfish is not achieved, further restrictions will be placed on the shrimp trawl industry to achieve the 40 percent reduction.
- Additional restrictions on the shrimp trawl industry will be reviewed and discussed at that time.

Southern AC

Do not support the MFC recommendation because of no support for the 40% reduction due to no scientific basis and also to not support the remaining portions of the MFC's motion because they are redundant to earlier recommendations.

Northern AC

Strongly reject the MFC recommendation for the 40% reduction due to no scientific basis and also to not support the remaining portions of the MFC's motion because they are redundant to earlier recommendations.

Habitat and Water Quality AC

Convene a stakeholder group to initiate a three year study to test minimum tail bag mesh size, T-90 (square mesh) panels, skylight panels, reduced bar spacing in TEDs and any other new methods of reducing unwanted finfish bycatch to achieve a reduction in bycatch.

- Compare these to a control net with a Florida fish eye, a federally approved TED, and a one and half inch mesh tail bag.
- The stakeholder group should partner with DMF and Sea Grant to help secure funding for the study.

Finfish AC

Strongly reject the MFC recommendation for the 40% reduction due to no scientific basis and also to not support the remaining portions of the MFC's motion because they are redundant to earlier recommendations.

Shellfish and Crustacean AC

Do not support the MFC recommendation because recommendations are already addressed in the DMF recommendation.

Public Comment

Southern Advisory Committee (January 8)

Buzzy Frederick- He participated in skimmer trawl testing of TEDs. Tests showed a 25% to 45% decrease in bycatch. In 2011 bycatch decreased between 23% and 43% in skimmer rigs and in 2011 there was a 27% to 45% decrease in bycatch. When combining 25 boats in both the Gulf and in NC the overall reduction was between 25% and 43% reduction.

Birdie Potter-Please do not support the MFC's motion to reduce bycatch by 40%. If we do not reach it, the MFC will do it biologically.

Chris McCaffity- Fisheries are at a cross roads and will be lost. Fishermen are beat down with so many regulations (gamefish, petition). It is harassment. Stop pushing regulations; give everyone a chance to work together.

Bradley Styron- Forty percent reductions in bycatch is unreasonable and what baseline are we working from? We need to know what is in the system. How many fish will live to maturity? There is no basis for the 40%. The industry is always proactive. Go back 70 years. There has been a 90% decrease through attrition, gear modifications and other things. Fishermen are always looking to get rid of bycatch. It is inefficient to fish with bycatch in your net. Bycatch is area specific and can even be hard to find sometimes to compare gear. There is no way to gage bycatch and 40% is arbitrary and unattainable.

Steve Weeks- *read off a prepared statement handed out to AC. See below.*

Brent Fulcher- He agrees with Bradley and supports the stakeholder group. He disagrees with anything more than a 30% reduction. The 40% is not achievable and is unreasonable.

Bill Hooper- He agrees with the DMF recommendations. The 40% is unachievable. He proposes a more reasonable reduction and to get funding. There is no evidence of bycatch impacting species. The MFC is overstepping its power in light of biological evidence.

Ken Sieglar- The southern area of Brunswick County is like the White Oak River when the shrimp get a certain size and leave. When they leave they do not get caught. Size does not matter down there.

Steve Parrish- We still need to improve the gear to reduce bycatch without losing shrimp bycatch. We tried the 1" bar in the TED and it excluded bycatch but lost a lot of shrimp. The MFC should reconsider its recommendation and give us a reasonable goal. Mr. Parrish is in favor of the stake holder group and volunteers to participate.

Northern Advisory Committee (January 9)

Terry Pratt- There is no baseline to reduce 40% from. The fishery has reduced bycatch. It used to be 10:1, now it is 4:1. Every BRD was developed by fishermen. Fishermen attempt to reduce bycatch.

Steve Weeks- *read off a prepared statement handed out to AC. See below.*

Brent Fulcher- *read off handout prepared by Jerry Schill. See below.*

Brent Fulcher-Bycatch is a lot of things. Some is sellable, it is not all dead. The T-90, the reduced bar TEDs, all of the BRDs were developed by the industry. He agrees with the DMF's recommendation to form a stakeholder group to test new BRDs and TEDs. He disagrees with anything over 30%. Support reducing bycatch with minimum shrimp loss. Forty percent is not achievable.

Greg Judy- Was involved with BRD development I NC while he worked with the DMF. When you reduced weakfish by 50%, you reduced the spot and croaker by 70%. The 40% reduction that the MFC is asking for is unachievable; you will lose too many shrimp. Amendment 6 of the SA shrimp plan stated there is no evidence of bycatch having an adverse effect on weakfish, spot, or croaker. Thus the extra 40% reduction on top of the 30% reduction we have already achieved is too much and unfair.

Bill Hooper- Agrees the DMF needs to work with the fishermen and the industry will find a way to reduce bycatch. We can build better BRDs, but we should not be held to a number that is unachievable. TEDs have been shown to reduce bycatch as much as 40%; however, an additional 40% on top of the 30% that is in place now is unachievable without shrimp loss. Biological measures to impose if the reductions are not met are no more than a threat without a scientific basis to support such a number. There will have to be too many assumptions to establish a benchmark.

Glenn Skinner- I m disappointed with the current MFC, there is no data to support the 40% number and they are setting the industry up for failure.

Habitat and Water Quality Committee (January 13)

Jerry Schill- *read off a prepared statement handed out to AC. See below.*

Greg Judy-BRDs decrease gray trout by 50%. He tested lots of BRDs. A 50% reduction in gray trout equals about a 60% reduction in other species. An additional 40% reduction is unattainable. If you get a 40% reduction, you will lose too many shrimp. SAMFC shrimp plan also states that there is no evidence that trawling is having an impact on bycatch of weakfish, spot and croaker. In 1975-1989 average landings of shrimp were 4.9 million. There were no BRD or TED requirements. During the same time frame the average landings of croaker were 6.7 million, spot 5.4 million, and weakfish 1.8 million. It makes you wonder if shrimp trawling was the culprit, there is something else limiting these fisheries. I ask you to not support the additional 40% reduction as suggested by the MFC.

Finfish Committee (January 14)

Clarence Fredrick- Decisions that are being developed will have impacts. You need to address the potential impacts of a live bait fishery. They should be treated like other commercial shrimpers and they should not be allowed to trawl in closed areas. A Sunday evening, 5:00 tow is the best tow. The industry manages itself. I am also against the closure in the south; it will hurt recreational shrimpers too. If the shrimp are small, the commercial fishermen will not go because the shrimp will not sell. Studies conducted by NMFS, testing TEDs in skimmer reduced bycatch 28% to 45%. We are reducing bycatch now. The shrimpers are also studying the 3" bar in the TED; as a whole the industry is working to reduce bycatch.

Shellfish/Crustacean (January 16)

Doug Cross-The 40% motion from Joe Shute is not achievable. Originally, there were discussions with commissioners about what were good reductions. Somehow it evolved into a target before there was data or logic to support it. He is totally against the 40%. We should not set a target without the science to back it up. There should be no consequences if there is no scientific data to support reductions. He supports bycatch reduction but we should not have a target first. That is putting the cart before horse. He did not see how any committee could support this recommendation. Let us eliminate bycatch as much as humanly possible. Do not set an arbitrary number that is not achievable. We do not need a dooms day effect in the 40% is not met.

Presented to Northern, Southern, Habitat, and Crustacean ACs

North Carolina Fisheries Association, Inc. (Jerry Schill, Brent Fulcher)

There is one thing that is striking to me since getting back involved in fisheries issues after an absence of personal involvement for almost 9 years. I was with the North Carolina Fisheries Association for 18 years, from 1987 till 2005, and during that time served on numerous boards, committees and panels including 6 years on the South Atlantic Fishery Management Council and several years on the Joint Legislative Commission on Seafood & Aquaculture.

The one striking issue that I referred to is the loss of memory of what has transpired over the years in fishery management, in this case, in the shrimp bycatch history. Last summer in Raleigh it was crystal clear that many who were proposing gamefish or were involved with the banning of inside shrimping, had little to no experience in fisheries issues. There were others who knew better but had selective memories. I say that because I heard directly or indirectly that fishermen needed to compromise, and that opinion seemed to be based upon an assumption that commercial fishermen have not conceded anything when sitting at the table to discuss a myriad of issues, including bycatch. So let's talk about that.

When I started with NCFCA in 1987 I had no preconceived notions about anything regarding fishing issues. I had no bias from a trawling, gillnetting, bottom fishing, crabbing or even from a hook and line recreational fishing perspective. I didn't do any of it. My only connection to fishing was as a consumer. I was, however, attuned to the political winds and the public perception of a number of issues including the protection of turtles and bycatch issues. And so were many fishermen who made their living on the water. In the late 80s, the debate wasn't really about whether or not to protect turtles for example, but how to do it. We learned quickly that the federal government shoving a particular method down the fishermen's throat wasn't going to do it. It was only after fishermen got involved with the planning and testing was there success in coming up turtle excluder devices that worked.

And the same is true with bycatch reduction. Whether you prefer to call them BRDs, bycatch reduction devices, or FEDs, fish excluder devices, they came to be because they were developed and tested by fishermen. (We prefer BRDs because the word FED is so negative!)

In the case of North Carolina, many shrimpers used them voluntarily. Why? Because they worked! It's also important to note that North Carolina was the first state to mandate their use in shrimp trawls and it was OK with the shrimpers because most were using them anyway!

So why is this stuff important? I'll paraphrase something I heard the late Congressman Walter Jones, Sr., say at a Merchant Marine and Fisheries Committee meeting a long time ago: **“NO fisheries regulation will work without the acceptance, albeit grudgingly, of the fishermen being regulated.”**

In the mid 80s, if any of the self-described conservationists who were complaining about bycatch in the shrimp trawl fishery could have envisioned North Carolina shrimpers helping to develop and actually use reduction devices that actually worked such as what we're using now, they would have thought they died and went to heaven!

Now, does that mean we should just forget about it and do nothing to reduce it even further? Absolutely not! But I would strongly caution this committee, when you make your recommendations to the Marine Fisheries Commission, not to push for any particular percentage reduction that is unattainable or worse yet, any threat of what you'll do to the fishermen if they don't reach that unattainable goal, because you will be doomed to fail from the start! (UNLESS the ultimate goal is the elimination of the shrimpers in the first place.)

Such a mandate back in the 80s would have doomed the process. Cooperation is the key, and yes, I know full well about those who claim otherwise. Just remember that they weren't sitting at the table back then and really have no idea what went on. The process I'm speaking about is not easy, but if our goal is to truly reduce bycatch even more, rather than sabre rattling, then the cooperative approach is the only way to be effective.

The North Carolina Fisheries Association appeals to you to go the route that was taken for many years and that is the fostering of cooperation on the studies and testing of ANY gear in our efforts to conserve our resources while allowing a proud and noble fishing tradition to continue.

Fishing families and the seafood consumers are counting on you!

Presented to Northern and Southern ACs

North Carolina Fisheries Association, Inc. (Steve Weeks)

Commercial fisherman want to reduce bycatch for the viability of all finfish species, not just by reducing bycatch in the shrimp fishery, but in all fisheries, recreationally and commercially.

The North Carolina Fisheries Association is opposed to the Commission's recommendation requiring a minimum 40% reduction of finfish by weight within a 3 year period.

The Association supports the Division's recommendations to:

1. Convene a stakeholder group to initiate industry testing of bycatch reduction devices;
2. To allow any Federally certified bycatch reduction devices in all internal and offshore waters;
3. Upon Federal adoption of turtle excluder devices in skimmer trawls, state enforcement of the Federal requirement; and
4. Updating the scientific testing protocol for the state bycatch reduction device certification program.

The shrimp industry since the introduction of the bycatch reduction device has reduced bycatch by approximately 1/3. Before a bycatch reduction device can receive Federal certification it must reduce bycatch by a minimum of 30%. In addition to a bycatch reduction device, all commercial shrimp trawls require a turtle excluder device. Recent testing by the National Marine Fisheries in conjunction with a North Carolina skimmer trawl fisherman has indicated that turtle excluder devices also reduce bycatch.

In addition, there has been a substantial reduction in effort in the shrimp industry. In 1994 there were 14,585 shrimp trips with otter trawls, in 2011 there were 3,004 trips, a 485% reduction. In 1994 there were 1,118 shrimp trawl trips, in 2011 there were 327 trips, a 340% reduction. In 1994 there were 2,109 channel net trips, in 2011 there were 531 trips, a reduction of 397%.

The recommended 40% reduction over a 3 year time period is not achievable without destroying the North Carolina shrimp industry, is without scientific basis and is arbitrary and capricious.

Scientifically the Division has not established an accurate finfish to shrimp, bycatch ratio. Director Daniel, at the hearing in Raleigh in August 2013, testified that there was no accurate basis in the bycatch ratio the State of North Carolina was using. This ratio is considerably higher than the ratio of all South Atlantic States according to data from National Marine Fisheries, Southeast Region.

Steven Parrish, a net maker and designer with significant experience and expertise in bycatch reduction devices spoke before the Southern Advisory Committee on January 8, 2014. Mr. Parrish advised that through his experience and testing a 40% additional reduction in bycatch was not achievable without a significant loss of the shrimp catch.

The reduction of bycatch in the shrimp fishery should be achieved in a reasonable manner and over a reasonable time period with cooperation between the State and the

fisherman. It took years to develop, test and implement the fish excluder devices currently in use, which have to be Federally approved. There are currently Federally approved finfish excluder devices that the State of North Carolina will not allow the fisherman in State waters to use.

Kevin Brown with the Division, who is in charge of the bycatch reduction program, advised the Southern Advisory Committee on January 8, 2014 that he questions whether he has adequate personnel and resources to verify the data required to substantiate whether or not any reduction mandated by the Commission has been met.

In 2013 NOAA Fisheries awarded 16 grants totaling nearly \$2.4 million as a part of its Bycatch Reduction Engineering Program. Bycatch of various species – whether fish, marine mammals, or turtles – can have significant, biological, economic, and social impacts. Preventing and reducing bycatch is a shared goal of fisheries managers, the fishing industry, and the environmental community.

NOAA Fisheries' Bycatch Reduction Engineering Program provides funds critical to key partners for the research and development of innovative approaches and strategies for reducing bycatch.

Mr. Brown advised the Southern Advisory Committee on January 8, 2014 that he has applied for grants to assist in the research and testing of bycatch reduction devices.

This advisory committee should recommend the above recommendations of the Division and should not recommend the Commission's recommendation of a 40% bycatch reduction over a three year period as there is no scientific basis for said reduction and said reduction is not achievable without significant monetary loss to the North Carolina shrimp industry, which has traditionally been one of the more lucrative fisheries in this State.

Additional Public input (emails)

United National Fisherman's Association (received January 8)

Do not need a number or percentage for the amount of by catch.

Reduction in shrimp Trawls. DIVISION & National Marine Fisheries Service needs to document the by Catch reduction from Turtle Excluder Devices. No credit is given for large fish excluded by Turtle Excluder Devices. {Loss of income to fishermen}

NOW! The committee should address the by catch from recreational fish that can be reduced by changing the regulations on all regulated recreational fish.

A possession limit for each species in total length in inches to be retained, all fish must be retained, thus allowing low income fishing from the bank access to fish for food.

In brought over the rail. Require barb less hooks on all recreational vessels, only allow barbed hooks on piers bridges or shore. Consider barbs on hook side away from shank.

Trout as an example would have a 60 inch total length limit, thus allowing 5 12 inch fish

Or 6 10 inch fish NO BY CATCH NO CATCH & RELEASE!

ELIMINATE BY CATCH OR HOOK & RELEASE IN RECREATIONAL FISHING.

Coastal Fishery Reform Group (received January 16)

The Coastal Fisheries Reform Group (CFRG) is a coalition of recreational coastal fishermen, who support sound management of our marine fisheries based upon the best available science. We represent many thousands of fishermen from across the state who fish in our coastal waters. We have had over 127,000 hits on our blog site (<http://cfrgnc.blogspot.com/>) where we have discussed coastal fisheries issues since 2009. In the role as a voice for the average salt water fisherman, we submit the following comments on the proposed Shrimp FMP amendment

to reduce finfish bycatch that the Marine Fisheries Commission will consider at their February 2014 meeting.

We strongly believe that the draft amendment to the shrimp plan, which includes only proposals for industry testing of bycatch reduction devices, updating testing protocols for the state bycatch reduction device certification program, and requiring additional bycatch reduction devices in all shrimp trawl nets, falls woefully short of an acceptable proposal to amend the Shrimp FMP to reduce bycatch.

The Shrimp FMP Advisory Committee met several times over the course almost a year and many additional, significant measures were discussed and considered. The proposals emerging from the study are almost meaningless and will do little if anything to reduce finfish bycatch in shrimp trawling operations. The recommended amendment contains no options for gear restrictions, no time closures, no areas closures, and no target reduction in bycatch. The Shrimp FMP should be amended to include goals, timetables, and management measures to accomplish significant by-catch reduction and an aggressive data collection and analysis program to monitor the success of management actions taken over the next five year period.

We quote here from the draft amendment (page 65 Section **6.3 Shrimp Trawl Bycatch**): *“As perhaps the prime example of the new policy positions, the re-authorized Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA) contains a National Standard (#9) requiring bycatch minimization (USDOC 1996). National Standard 9 states: “Conservation and management measures shall, to the extent practicable, (A) minimize bycatch and (B) to the extent bycatch cannot be avoided, minimize the mortality of such bycatch.” Additionally, in 1991 the MFC adopted a policy directing the DMF to establish the goal of reducing bycatch losses to the absolute minimum and to consciously incorporate that goal into all of its, management considerations (Murray et al. 1991).”*

The CFRG urges the MFC to amend the Shrimp FMP to include the following provisions:

1. Limit all trawl nets in inshore coastal waters (especially Pamlico Sound) to a maximum headrope size of 110 feet and only allow two nets per boat. This would remove the large nets and their excessive bycatch but would allow the small trawlers that have shrimped in our sounds for generations to continue working uninterrupted. These smaller shrimp boats are mostly local boats, with local crews that sell their catch at local fish houses in North Carolina. Such a rule change would greatly benefit the vast majority of North Carolina shrimpers while truly helping our coastal economies and our marine resources by significantly reducing bycatch.
2. Limit tow times to 60 minutes. This would allow for some bycatch to be released alive and also increase the chance of sparing any endangered turtles which are entrapped in the net.
3. Delay shrimp season until the shrimp size has reached the level of having 36 to 41 (or lower) shrimp per pound. This would postpone the harvest of shrimp and allow juvenile finfish to grow larger and have more of a chance of escaping shrimp trawls. These fish would also have more time to move out of their nursery areas where the trawlers are now working. In addition, this change would cause the shrimp to be larger when they are harvested and market value would be greater, thereby benefiting shrimp fishermen.
4. Establish exclusion zones around both sides of our inlets where trawlers with headropes exceeding 110 feet would not be allowed. This would allow juvenile finfish that are transitioning to a life in the open ocean to escape our sounds without being killed by a shrimp trawler. These fish become concentrated when they are near the inlets and are especially vulnerable to trawlers until they can disperse into the ocean.

Now is the time to get serious about the finfish decimation caused by the current activities of shrimp trawlers in the inshore waters of NC. Destruction of fisheries resources of this magnitude cannot be tolerated any longer. If the proposed amendment to the Shrimp FMP is adopted as presented, the schedule for meaningful action will be delayed for years while we look for the magic solution that is right before us now. Establish some realistic goals, implement some meaningful management measures, set a timetable for implementation, evaluate improvements in terms of bycatch reduction, and make subsequent changes as dictated by results.

Joe Albea
On behalf of Coastal Fisheries Reform Group

North Carolina Wildlife Federation Camo Coalition (received January 16)

Dr. Daniel:

I have closely followed the course of the current attention given bycatch in the Shrimp FMP from the first proposal to simply revise the FMP to the decision to amend the FMP and all of the work of the Advisory Committee that was appointed and has worked for about a year to review the bycatch issue and make proposals to amend the FMP to address bycatch.

If I correctly interpret the draft amendment being considered now, the recommendations are limited to an industry study of bycatch reduction devices, updating testing protocols for the state bycatch reduction device certification program, and requiring additional bycatch reduction devices in all shrimp trawl nets. These limited proposals are not at all significant if we are trying to reduce bycatch from shrimp trawling in a timely and effective way. Many reasonable and effective means to reduce bycatch are available now. If we limit the amendment to these points, then bycatch will not come up again for five years. We need to at least adopt a list of alternative management measures such as gear restrictions, maximum tow times, area closures around inlets and a schedule for implementation and a plan for evaluation in terms of reduction of bycatch.

Essential to any deliberate, serious plan of action to address a natural resource issue as crucial as the unacceptable bycatch of immature finfish in shrimp trawls is a plan of action with a slate of management choices, a timetable of implementation and evaluation, and a target level of success. The draft amendment has none of these elements. The Advisory Committee discussed many of the management measures that could have given promise toward achieving a meaningful reduction in bycatch, but none of these actions are before the Commission for consideration. The omission of potential actions that could work flaws the whole process.

Now is the time to begin an approach toward reduction of shrimp trawling bycatch. We do not need more studies that will continue to show that finfish mortality is significant and the effect on their stocks is uncertain. We do not need to evaluate bycatch reduction devices as the main thrust of bycatch reduction. A risk of doing nothing substantial at this good opportunity is the possibility of a far more precipitous and radical change similar to what happened with the sea turtle and the gill nets.

Dick Hamilton

Dr. Daniel:

Reference: Proposed amendment Shrimp FMP to Reduce Bycatch

The North Carolina Wildlife Federation is a statewide, non-profit, conservation organization dating back to 1945 dedicated to the professional management of our fish and wildlife resources based upon scientific principles. We represent many thousands of fishermen from across the state who fish in our coastal waters. It is within this purview that we submit the following comments on the proposed Shrimp FMP amendment to reduce finfish bycatch that the Marine Fisheries Commission will consider at their February 2014 meeting.

We strongly believe that the draft amendment to the shrimp plan falls far short of addressing the serious problem of bycatch in the shrimp trawling industry. It seem as if the draft amendment includes only proposals for industry testing of by-catch reduction devices, updating testing protocols for bycatch reduction device certification program, and requiring additional by-catch reduction devices in all shrimp trawl nets. These limited proposals will do little if anything to reduce by-catch.

The Shrimp FMP Advisory Committee met several times over the course almost a year and many additional, significant measures were discussed and considered. The recommended amendment contains no options for gear restrictions, no time closures, no areas closures, and no target reduction in bycatch. We believe definite, measurable reductions must be implemented by and for the commercial trawlers. The Shrimp FMP should be amended to include definitive goals, timetables, and management measures to accomplish significant by-catch reduction and an aggressive data collection and analysis program to monitor the success of management actions taken over the next five year period.

NCWF urges the MFC to amend the Shrimp FMP to include the following provisions:

1. Limit all trawl nets in inshore coastal waters (especially Pamlico Sound) to a maximum headrope size of 110 feet and only allow one net per boat. This would remove the large nets and their excessive bycatch but would allow the small trawlers that have shrimped in our sounds for generations to continue working uninterrupted. These smaller shrimp boats are mostly local boats, with local crews that sell their catch at local fish houses in North Carolina. Such a rule change would greatly benefit the vast majority of North Carolina shrimpers while truly helping our coastal economies and our marine resources by significantly reducing bycatch.
2. Limit tow times to 45 minutes. This would allow for some bycatch to be released alive and also increase the chance of sparing any endangered turtles which are entrapped in the net.
3. Delay shrimp season until the shrimp size has reached the level of having 36 to 41 (or lower) shrimp per pound. This would postpone the harvest of shrimp and allow juvenile finfish to grow larger and have more of a chance of escaping shrimp trawls. These fish would also have more time to move out of their nursery areas where the trawlers are now working. In addition, this change would cause the shrimp to be larger when they are harvested and market value would be greater, thereby benefiting shrimp fishermen.
4. Establish exclusion zones around both sides of our inlets where trawlers with headropes exceeding 90 feet would not be allowed. This would allow juvenile finfish that are transitioning to a life in the open ocean to escape our sounds without being killed by a shrimp trawler. These fish become concentrated when they are near the inlets and are especially vulnerable to trawlers until they can disperse into the ocean.

Now is the time to get serious about the finfish decimation caused by the current activities of shrimp trawling in the inshore waters of NC. Destruction of fisheries resources of this magnitude cannot and should not be tolerated any longer as it is in nobody's interests. If the proposed amendment to the Shrimp FMP is adopted as presented, the schedule for meaningful action will be delayed for years while we look for solutions, which in our opinion, are right before

us now. We urge you to establish realistic goals, implement meaningful, deliberate, and measurable reductions by the industry, management measures, set a timetable for implementation, evaluate improvements in terms of by-catch reduction, and make subsequent changes as dictated by results.

Thank you for considering our requests,

Tim Gestwicki
CEO
North Carolina Wildlife Federation

Email received January 16

Please accept this communication as unconditional support of the Coastal Fisheries Reform Group's proposal, dated this day, regarding the referenced matter. My personal experience in a Marine Fisheries advisory capacity fosters my concern that, by the time any effective action is taken, it will be too late for the resource. Let's get on with a common sense approach, and do it now! Sincerely, Frank Liggett

Email received January 16

To whom this may concern: I fully support the CFRG's position on the proposed amendment to reduce shrimp trawling bycatch in NC. I urge you to do everything in your power to look at the science, and the reality, of the unconscionable damage that shrimp trawling is daily perpetuating on our environment, and act accordingly. Thank you, Lee Dunn, Beaufort

Phone call received January 23

Joe Buck called and suggested that shrimping should not be allowed at night, because you catch a lot more bycatch at night compared to day time shrimping.

Email received January 31

- * Limit all trawl nets in inshore coastal waters (especially Pamlico Sound) to a maximum headrope size of 110 feet and only allow two nets per boat.
- * Limit tow times to 60 minutes to allow some by-catch to be released alive and increase the chance of sparing endangered turtles trapped in the nets.
- * Delay shrimp season until the shrimp size has reached the level of having 36 to 41 (or less) shrimp per pound. This would postpone the harvest of shrimp and allow juvenile finfish to grow larger and have more of a chance of escaping shrimp trawls. These fish would also have more time to move out of their nursery areas where trawlers now work.
- * Establish exclusion zones around both sides of inlets where trawlers with headropes exceeding 110 feet wouldn't be allowed.

"Now is the time to get serious about the finfish decimation caused by the current activities of shrimp trawlers in the inshore waters of North Carolina," CFRG wrote. "Destruction of fisheries resources of this magnitude cannot be tolerated any longer."

CFRG also asked the Commission to establish "realistic goals, implement some meaningful management measures, set a timetable for implementation, evaluate improvements in terms of by-catch reduction, and make subsequent changes as dictated by results." it's time to get serious about the damage being done to our nurserys in nc thank you paul brown 600 n. rocky river rd. sanford n.c.

Email received January 31

Mr. Laughridge, It's time to end inshore shrimp trawling. Not reduce, not alter limits, no other half-steps, END. The old song of jobs lost doesn't cut it anymore. The damage trawling does FAR outweighs the benefit and this can and has been proven in many ways. If you would like I can easily support this position with data. The snowball is growing if you haven't noticed. Long overdue change is coming to NC fisheries management. I hope to be able to count on you to finally take the big picture in to account and join every other east coast state and BAN inshore trawling. The reasons they banned it are 100% applicable here too. Thanks, Bruce

Email received January 31

Mr. Rose, It's time to end inshore shrimp trawling. Not reduce, not alter limits, no other half-steps, END. The old song of jobs lost doesn't cut it anymore. The damage trawling does FAR outweighs the benefit and this can and has been proven in many ways. If you would like I can easily support this position with data. The snowball is growing if you haven't noticed. Long overdue change is coming to NC fisheries management. I hope to be able to count on you to finally take the big picture in to account and join every other east coast state and BAN inshore trawling. The reasons they banned it are 100% applicable here too. Thanks, Bruce

Email received January 31

The CFRG recommendations on shrimp trawling would be a great step in the right direction. Better still, STOP ALL TRAWLING INSIDE! Farm raised shrimp is the way to go. Recreational fishing has become so bad now that it is barely worth trying in NC coastal waters. Friends and I have been fishing at the coast for more than 50 years and have found it difficult to catch enough fish for one dinner in the last several years. Our long trips to Florida for great fishing would end if NC would copy Florida's approach to marine management. rcaproger@aol.com

Email received January 31

I have fished the North Carolina coast for over 30 years and have seen how the shrimp trawlers have destroyed the NC fishing. I would pay triple for shrimp or fish than to continue seeing the shrimp trawlers continue to destroy the fishing for not only myself but for all future fishermen. First of all the Commission or personnel involved in making changes to the fishing laws that have interest or own commercial fishing license should be banned from voting on changes to the fishing regulations. I really believe that if the N.C. Marine Fisheries Commission continue allowing nets and shrimp trawlers to continue in North Carolina, fishing for everyone will be destroyed. Looks at Virginia and how they banned nets and shrimp trawlers – five years after the ban, fishing is plentiful. Thanks for your time, Charles Brown, 109 Holly Creek Rd, Morrisville, NC 27560

Email received January 31

This, if all reports are even half way accurate, is a travesty...there is absolutely no way that killing all those small spots, croakers, and weakfish can have anything but a disastrous effect on those species....it does not take a fisheries biologist to recognize the terrible practice needs to be stopped or at least drastically changed. Shrimp trawling by catch is the 600 lb. gorilla in the room and he isn't going away. I ask the MFC to address this issue and do what is right for the resource, for a change. I, and a great number of others, am fed up with the lack of foresight and

continued lack of courage that has allowed this issue to fester like a boil on the reputation of fisheries management in our state. Thank you, Hubert Parrott

Email received January 31

The people of NC should know about the by catch situation, if they did maby their would be better managment by the marine fishries. This has been going on for many years and has had a big impact on the poor fish stock situation. Billy Reavis [bjreavis@gmail.com

Email received February 1

Ms. Fish: It is obvious that the SMP put forth by the Marine Fisheries Comm. is but a stopgap measure and will do nothing to actually limit the killing of juvenile finfish by trawlers in the sounds of NC. The coastal Fisheries Reform Group has listed a number of operating procedures and rules that IF ENACTED will actually give a more realistic chance for the survival of some of the affected finfish. I urge you to hear these proposals as a sincere effort to address the massive bycatch problem that retards any future growth of NC's fish stocks. Sincerely, Neil M. Smith, 486 Tom Absher RD. Scottville, NC 28672, neilmlynn@skybest.com

Email received February 13

To: Fishwatch@noaa.gov

Good Morning,

Please take the following into consideration when you rate the impact of otter trawling caught NC shrimp for your FishWatch Bulletin.

The Atlantic States Marine Fisheries Commission is charged with managing weakfish (gray trout). The current weakfish stock status is severely depleted with the adult spawning population at levels so low, that further declines are expected. In fact, in 2003, the ASMFC projected a greater than 90% chance that weakfish biomass could fall to zero by 2015. In 2008, weakfish biomass was less than 4.5-million pounds, a 96% drop in thirty years. In a 2009 report, the ASMFC Weakfish Technical Committee states "Unless there has been a steady rise in weakfish juvenile discards since 1999, the emergence of a demographic bottleneck is consistent with enhanced predation (e.g. spiny dogfish and striped bass) on smaller weakfish."

What is a weakfish's #1 predator in NC? The Spiny Dogfish? The Striped Bass? The Pamlico Sound shrimp boat?

NCDMF Director Daniel has publicly stated that 4.5 to 1 is a clear and well established bycatch ratio in NC's shrimp industry. In 2008, NC landed 9.4-million pounds of shrimp. Approximately 68% of NC shrimp landings come from the Pamlico Sound system. According to NCDMF studies, juvenile weakfish represent approximately 7% of trawling bycatch in the Pamlico Sound. In those studies, weakfish bycatch averaged 27.5-fish per pound. In 2008, NC shrimp trawlers killed 55-million Pamlico Sound weakfish as bycatch, 2-million pounds. The total east coast weakfish spawning stock biomass was only 4.5-million pounds in 2008.

Bycatch is not only affecting weakfish, but also spot and croaker stocks- once both important seafood staples. Those two stocks had historical low catches in 2012. NC spot landings have dropped from 7.1-million pounds to less than 500,000-pounds, a 93% decline. NC croaker landings have dropped from 21.1-million pounds to 3.1-million pounds, an 85% decline.

On February 19-21, the NCMFC will meet in Morehead City and the topic of shrimp bycatch will be discussed. While there is always a possibility that meaningful change will come from that meeting, history will show such wishful thinking is doubtful. It appears the NCMFC is going to vote to "study" the issue for three more years, a vote for maintaining status quo. During the

next three years, NC shrimpers will continue to trawl in critical habitat nursery areas important to weakfish, Atlantic croaker, spot, southern flounder and blue crab. Important finfish and crab stocks will continue to decline as trawling bycatch kills 1.0 to 1.5 billion juvenile species in those three years.

I understand that it is difficult for outside agencies to control what happens in the territorial waters of NC, but those agencies can at minimum- acknowledge the problem, document it and suggest improvements to NC's unsustainable fishery practices, practices that are not only detrimental to NC fish stocks, but interstate fish stocks.

Thank you for your time and consideration. Sincerely, Rick Sasser, Goldsboro, NC,
rick.sasser@hotmail.com

Email received February 17

Can all of bycatch be converted to Economic Value & nothing was returned to water, would By Catch remains a problem in shrimp fishery? Product would be worth 10 to 12.5 cent per pound. Pass question on to whom ever. Thank you, James Fletcher, United National Fisherman's Association, 123 Apple Rd. Manns Harbor NC 27953, Phone: 252-473-3287 Cell: 757-435-8475, Fax: 252-473-4969

14.2 APPENDIX 2 – REVISION REVIEW PROCESS

MEMORANDUM

TO: Management Review Team

FROM: Shrimp Plan Development Team

DATE: May 3, 2012

SUBJECT: Amendment versus Revision of the Shrimp FMP

The Shrimp Plan Development Team (PDT) has met and discussed the question of whether or not the Division Shrimp Fishery Management Plan (FMP) should be amended or revised. The PDT has met on three occasions to discuss management issues that have developed since the implementation of the 2006 Shrimp FMP. These issues include:

1. Restricted Trawl Area Offshore of Bogue Banks
2. Permanent Shrimp Line in Newport River
3. Trawling in New River above the Highway 172 Bridge
4. Volumetric Measurement of Shrimp

Each issue was addressed by the PDT in issue papers (see attached), including management options. After thorough discussion, no management changes were recommended for three of the four issues listed above. The fourth issue; Volumetric Measurement of Shrimp was discussed and two options were suggested for consideration by the Management Review Team (MRT). Both options require a rule change and it is unclear to the PDT whether these suggested rule changes constitute a change in management of the shrimp fishery.

To insure the public was aware that the Shrimp FMP was under review, a press release was sent out on November 7th, 2011 requesting public comment as part of our review process to determine whether to proceed with an amendment or revision of the FMP. The last day for comment by the public was December 2nd, 2011. The PDT received five comments. Each comment was reviewed and addressed by the PDT (see attached). A sixth comment was received on December 9th, after the deadline and after the PDT's review of the other comments. This comment was forwarded to the PDT and is also included within the public review document but was not thoroughly reviewed by the PDT outside of email.

After careful consideration of the issues listed above and of the public comments, the Shrimp PDT recommends to the MRT that the Shrimp FMP should proceed as a revision.

/plm
Enclosures

PERMANENT SHRIMP LINE IN NEWPORT RIVER

November 15, 2011

I. ISSUE

In the 2006 Shrimp Fishery Management Plan (FMP) a permanent closure line was established at the Penn Point-Hardesty Farm line. Fishermen who fish in Newport River would like the Division to repeal this rule in Newport River.

II. ORIGINATION

A request was made by the Newport River fishermen.

III. BACKGROUND

The Newport River is a relatively small estuary of about 63 square miles located north of Morehead City in Carteret County. Average depth is less than three feet with a maximum depth in natural channels of six feet and 40 feet in the dredged channels near the State Port. The western portion of the Newport River has bottoms composed of silts, clays and oyster rocks and the eastern part is composed of a firm sand bottom. There is a Primary Nursery Area (PNA) and a Special Secondary Nursery Area (SSNA) located in the western portion as well.

Before the 2006 FMP, the Newport River had a long history of disagreements concerning the best location of a shrimping closure line. Lines used in the past were the Hardesty Farm line, the White Rock line (SSNA line) and the Turtle Rock line (PNA line). During this long period of conflict that peaked in the mid-1980s, the line would move several times during a season in response to requests by fishermen and the variation in shrimp size. By October of each year the river would open to the PNA line with the opening of the SSNA by proclamation. Based on input from the public, the Shrimp Advisory Committee, the Division of Marine Fisheries and the Marine Fisheries Commission (MFC), a permanent trawl nets prohibited line was established from Penn Point to Hardesty Farms (15A NCAC 03R .0106) (Figure 1).

Shrimp harvest generally begins in June with the presences of brown shrimp and can continue into November and sometimes as late as December if white shrimp are abundant. The primary conflict has historically occurred in the fall, between fishermen, who generally wanted the Hardesty Farm line established because shrimp that have migrated down are a more marketable size. This line also allows for more maneuverability for large shrimping vessels, while other fishermen with smaller vessels, preferred the White Rock line (SSNA) in order to access the shrimp and harvest the majority of them, before the shrimp moved down to the Hardesty Farm line. The White Rock line is located in shallow water, where the larger boats are unable to work because only a small portion of the White Rock line is deep enough for trawling.

The western half of the Newport River above the Hardesty Farm permanent closure line contains sites where significant shellfish management efforts have occurred over the past 35 years. Natural oyster rocks extend from the Cross Rock in the western part of the river through White Rock located at the mouth of Harlowe Creek. Oyster rocks also exist along the shores of Newport River Marshes and the entrance to Core Creek. The Division has planted approximately 201,514 bushels of cultch material in the western portion of Newport River above the Hardesty Farm line since 1981 and 22,990 from 2006 through 2011. These plantings have

expanded the natural rocks (Flat Rock, White Rock, Turtle Rock and the Bullseye Rock). There are also 15 active leases above the shrimp line totaling 103 acres.




Figure 1. Newport River Shrimp Lines

III AUTHORITY

G.S. 113-182.1 FISHERY MANAGEMENT PLANS

North Carolina Marine Fisheries Commission Rules for Coastal Fishing Waters (15A NCAC) 15A NCAC 03R .0106 TRAWL NETS PROHIBITED

IV. DISCUSSION

Since 2006, the implementation of the Hardesty Farms line has been successful because it protects small shrimp that move out of Harlowe Creek in the early summer and provides a buffer when the abundance of juvenile shrimp, heavy rainfall or strong northerly winds pushes the shrimp downstream of their normal location. The permanent line has also eliminated the costs and time spent by division staff sampling this area in response to requests to move the shrimp closure lines. Total landings in the Newport River do not appear to have been impacted by the establishment of the permanent line (Figure 2). Total fall landings in the Newport River for the years when the lines moved (2001-2005) ranged from 4% to 33%. After implementation of the permanent line in 2006, total fall landings ranged from 15% to 23%. Number of trips during the same time frame has averaged 200 per year from 2001 to 2005, while number of trips averaged 109 from 2006 through 2010. Overall, fall landings have remained stable, averaging approximately 20% of the total landings in Newport River before and after the implementation of the permanent line (Figure X).




Figure 2. Percent total landings (lbs.) from October through December in Newport River

Juvenile spot, croaker, brown shrimp, blue crab and southern flounder utilize the PNA and SSNA habitats in Newport River. Trawling is prohibited in PNAs; however, the Fisheries Director may open SSNAs by proclamation from August 16 through May 14. With the implementation of the Hardesty Farm line in rule, the Fisheries Director no longer has the authority to open the Newport River SSNA. This protects leases and other oyster resources

from being trawled over or covered in sediment. This was a frequent concern for both lease holders and resource enhancement staff of the division when the SSNA was opened to trawling.

Small mesh gill net attendance rules have been impacted by the implementation of the Hardesty Farm line causing confusion to fishermen because of a contradiction in gill net attendance requirements. Rule 15A NCAC 03R .0112 (b) (1) states that in areas that are prohibited to trawl nets, permanent secondary nursery areas and in PNAs, small mesh gill nets must be attended from May 1st through November 30th. However according to 15A NCAC 03R .0112 (b) (5) gill net attendance is from May 1st through September 1st within 50 yards of the shore line. The Rules Review Team will be addressing this contradiction.

Proponents for trawling above the permanent line and in the SSNA cite the lack of growth of remaining shrimp due to falling water temperatures and the need to stir up sedimentation by trawling to remove silt from the upper river and that stirring the bottom removes silt (at least at ebbing tides), keeps it oxygenated (or alive), and exposes old oyster rocks and plantings to new spat set the following spring. Fishermen have noted an increase in siltation since the area has been closed to trawling.

V. PROPOSED RULE(S)

VII. MANAGEMENT OPTIONS/IMPACTS

(+ Potential positive impact of action)

(- Potential negative impact of action)

1. *Status quo* (permanent line at Hardesty Farm)
 - + Eliminates grand openings
 - + Protection of shellfish plantings, natural oyster rocks and leases
 - + Longer and deeper line for less congestion when trawling
 - +/- Trawling in SSNA is prohibited
 - No flexibility of management by proclamation
 - Continued confusion of small mesh gill net attendance

2. Remove permanent line at Penn Point-Hardesty Farm
 - + Provides flexibility in managing around variable conditions (excessive rainfall, early migration)
 - + Access to resource by a variety of users
 - + Able to open the SSNA by proclamation
 - No confusion on small mesh gill net attendance
 - Does not minimize harvest of small shrimp and bycatch
 - Does not prevent damage to shellfish plantings, natural rocks and leases
 - Labor intensive and expensive to sample
 - Necessitates "grand openings"

VIII. RECOMMENDATION

PDT: *Status quo* (permanent line at Hardesty Farm)

Prepared by: Trish Murphey
Trish.Murphey@ncdenr.gov
(252)-726-7021
November 15, 2011

Revised:

NOTICE OF TEXT ATTACHMENT

#6 – Explain Reason for Proposed Action:

MFC Rulebook Index Worksheet

Rule	Rulebook Page #	Subject	Index Entry	Add/Delete

TRAWLING IN NEW RIVER ABOVE THE HIGHWAY 172 BRIDGE

November 15, 2011

I. ISSUE

Request to reexamine the provision in the 2006 Shrimp Fishery Management Plan (FMP) which prohibits the use of otter trawls upstream of the Highway 172 Bridge over the New River.

II. ORIGINATION

Request by the New River shrimp and crab trawlers

III. BACKGROUND

The use of otter trawls upstream of the Highway 172 Bridge was phased out in 2010 following the adoption of the 2006 Shrimp FMP. Those who wished to continue to harvest shrimp in the waters above the 172 Bridge were allowed a four year grace period to convert to skimmers. Subsequently, crab trawls were also phased out of this area as part of the 2006 Shrimp FMP. Prior to the 2006 Shrimp FMP, crab trawlers would often fish above the Highway 172 Bridge to target flounder more so than crab; however, stricter minimum size limits for flounder made it economically unfeasible for crab trawlers to harvest only crabs in this area. Currently, the waters upstream of the Highway 172 Bridge are only open to boats equipped with skimmer rigs.

The waters upstream of the Highway 172 Bridge (Figure 1) were designated by rule as a Special Secondary Nursery Area (SSNA) in 1996. The areas of the SSNA that are impacted by the trawling opening include the river above the bridge up to the marked closure line running from Grey's Point to the opposite side of the river. Trawling in any of the tributary creeks is prohibited. The river consists mostly of shallow bays with the exception of the marked navigation channel. Bottom types range from sand and sand/mud to live shell bottom. The Division of Marine Fisheries (DMF) actively manages seven Shellfish Management Areas (SMAs) in this portion of New River.

Data from the DMF Trip Ticket Program were used to describe the commercial shrimp fishery in New River from 1994 to 2010 (Tables 1-5). Landed bycatch by gear was calculated and ratios (in pounds) of marketable bycatch relative to shrimp catch were also calculated for the four main gears: channel nets, otter trawls, skimmers, and the various miscellaneous gears (cast nets, gill nets, etc). Marketable bycatch from skimmers was consistently lower than with the other gears. Marketable bycatch landings in channel nets were also low, with the exception of 2000-2002 when significant amounts of blue crabs were landed in this fishery. In 2005, trip limits were put in place to restrict harvest of crabs in channel nets in the first crab fmp.pg.18 O3J.0106 (h). During this three-year period, ratios of pounds of shrimp per pound of marketable bycatch in the channel nets were 4:1, 2:1, and 3:1 respectively. These bycatch ratios apply only to the portion of bycatch retained and sold.

Discarded bycatch is much more difficult to quantify because of the lack of data in most areas. However, during 2003-2009, DMF staff sampled the study area for shrimp management purposes using a 25-foot, 4-seam otter trawl. This gear was not equipped with a turtle excluder or a finfish excluder. Catches were separated into four categories: commercial finfish, non-

commercial finfish, invertebrates, and shrimp. Each component was weighed and bycatch percentages were derived for each year (Table 6). Tow times ranged from one to 10 minutes. The primary objective of the sampling was to determine if the shrimp were large enough to warrant an opening or a closing but the weights of all the biomass components were recorded.

Overall, finfish accounted for 39.6% of the total biomass, with shrimp representing 51.2% of the weight, and invertebrates making up the remaining 9.1% of the weight from 2004 to 2009. Total bycatch ranged from 42.0% (2006) to 97.1% (2008).

The number of trips by the major shrimp gears indicates a decrease in effort for all gears from 1994 to 2010 (Figure 2). Following the ban of trawls above the Highway 172 Bridge, the numbers of otter trawl trips and participants dropped significantly in the New River. Prior to the ban, only 10 trips were made by seven participants in 2009 and only 13 trips were made by seven participants in 2010. The use of channel nets and skimmer rigs increased slightly in 2010. Prior the 2006 Shrimp FMP, channel nets were fished in the waters above and below the Highway 172 Bridge while skimmer effort was focused more in the SSNA located above the bridge. Currently, channel nets are only allowed to be set above the 172 Bridge Channel when the river opens to trawling by proclamation. Channel nets show the most consistency in the mean number of pounds harvested per trip while skimmers and otter trawls show similar year-to-year fluctuations; skimmers generally harvest more shrimp per trip (Figure 3). Landings from skimmers have shown a marked increase since 1994 reflecting the increased popularity of this gear, especially in the capture of white shrimp during the late summer and early fall (Figure 4). However, the variability of catches between all the gears is expected and is a result of year class strength.

The number of trips made by crab trawls also indicates a decrease in effort from 1994 to 2010 (Table 5). Following the adoption of the 2006 Shrimp FMP, there were no reported trips from 2007 to 2009 (Figure 5). In 2010, 32 trips were made by nine participants below the 172 bridge, landing 23,383 pounds of crab. Prior to the trawl ban above the Highway 172 Bridge, mean catch per trip ranged from 64 to 725 pounds from 1994 to 2006. In 2010, an average of 731 pounds of crab per trip was landed below the Highway 172 Bridge, well above the 262 pound per trip average observed from 1994 to 2006 when trawls were allowed above the bridge (Figure 6).

IV. AUTHORITY

G.S. 113-182.1 FISHERY MANAGEMENT PLANS

15A NCAC 03J .0104 TRAWL NETS

15A NCAC 03J .0208 NEW RIVER

15A NCAC 03L .0101 SEASON

15A NCAC 03N .0105 PROHIBITED GEAR, SECONDARY NURSERY AREAS

V. DISCUSSION

As part of the 2006 Shrimp FMP, otter trawls were prohibited by proclamation upstream of the Highway 172 Bridge in the New River beginning in 2010. Subsequently, this also prohibited the use of crab trawls, eliminating a traditional Sneads Ferry fishery, prompting the remaining members of that fishery to question the prohibition. The area above the Highway 172 Bridge is still designated as a SSNA, but the use of otter trawls is prohibited to minimize waste/bycatch and disturbance to the bottom. Additionally, trip ticket harvest data indicates that following the prohibition of otter trawls in New River SSNA, otter trawl bycatch has been reduced significantly

while the mean catch per trip (lbs) for shrimp has remained fairly high for the rest of the river (Table 2; Figure 3). The trip ticket data also indicates that skimmers are more effective at catching the target species than conventional otter trawls (Table 3). A skimmer trawl study conducted by Sea Grant found that skimmers were much more effective on white shrimp than otter trawls in water less than 12 feet (most of the water above the bridge in the New River) and in some cases out-fishing otter trawls as much as five to one (Coale, et al. 1994). The majority of the shrimp openings in the New River SSNA are for white shrimp, since by late summer most of the brown shrimp have already migrated.

Continuing to prohibit the use of all trawls, including crab trawls, above the Highway Bridge 172 protects the New River SSNA from bottom disturbing activities. Given the inherent design of most crab trawls (heavy-framed gears designed to dig into the substrate) the effect they have on the benthos is no different than that of otter trawls used to take shrimp. In some cases their effect on the benthos is worse due to their added weight. In addition, trip ticket data indicates that the highest mean catch per trip (lbs) occurred in 2010 when trawlers were not allowed access to the New River SSNA; higher than that of all the years prior to the trawl ban (Table 2). The implementation of the rule that prohibits otter trawls above the Highway 172 Bridge has been successful; opening the waters to crab and shrimp trawls would only reverse the progress made in the 2006 Shrimp FMP.

VI. PROPOSED RULE(S)

VII. MANAGEMENT OPTIONS/IMPACTS

(+ potential positive impact of action)
(- potential negative impact of action)

1. *Status quo* (prohibit trawls as an allowable gear in New River SSNA)
 - + Benefit to existing Shellfish management areas
 - + Encourage the use of a more efficient gear for harvesting white shrimp
 - + Reduction in waste/fish kills, especially on opening day
 - + Added protection for sub-legal flounder in New River SSNA
 - Eliminates part of a traditional Sneads Ferry fishery in this SSNA
 - Difficult to catch shrimp in a few deep-water spots
 - Financial hardship on trawlers who would likely convert to skimmers
2. Allow all trawlers in New River SSNA
 - + Allows prosecution of traditional fishery in SSNA
 - + Possible decreased financial hardship
 - No immediate remedy for waste/fish kills on opening day
 - No benefits to SMAs
 - Increased harvest on opening day, reduced price at market
3. Prohibit all trawlers and skimmers in New River SSNA
 - + Bycatch issue completely eliminated
 - + Potential for healthier shellfish/finfish stocks
 - Eliminates potential lucrative opening days for fishermen
 - Eliminates traditional Sneads Ferry fishery in this SSNA

VIII. RECOMMENDATION

PDT recommendation: *Status quo*

Prepared by: Chris Stewart
Chris.Stewart@ncdenr.gov
(910)-796-7370
November 15, 2011

Revised: November 15, 2011
December 9, 2011

Literature Cited

Coale, J.S., R. A. Rulifson, J. D. Murray, and R. Hines. Comparisons of shrimp catch and bycatch between a skimmer trawl and an otter trawl in the North Carolina inshore shrimp fishery. *North American Journal of Fisheries Management* 14: 751-768.

MFC Rulebook Index Worksheet

Rule	Rulebook Page #	Subject	Index Entry	Add/Delete

Table 1. Catch and effort data on shrimp and landed bycatch for channel nets in New River, 1994-2010 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lbs)	Sold bycatch (lbs)	Mean catch per trip (lbs)	Ratio shrimp of sold to bycatch
1994	37	544	47,556	747	87	64
1995	39	850	87,536	1,435	103	61
1996	36	585	62,590	1,894	107	33
1997	44	1,122	86,610	3,065	77	28
1998	29	856	80,714	428	94	189
1999	40	1,453	124,727	4,444	86	28
2000	45	1,380	163,109	38,998	118	4
2001	41	1,112	137,595	79,793	124	2
2002	38	1,257	163,831	61,907	130	3
2003	33	835	100,667	1,685	121	60
2004	32	570	59,799	4,370	105	14
2005	19	126	15,379	886	122	17
2006	18	206	57,011	240	277	238
2007	15	255	36,742	1,043	144	35
2008	14	168	40,892	750	243	55
2009	10	118	16,558	259	140	64
2010	19	322	39,297	1,279	122	31

Table 2. Catch and effort data on shrimp and landed bycatch for otter trawls in New River, 1994-2010 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lbs)	Sold bycatch (lbs)	Mean catch per trip (lbs)	Ratio shrimp of sold to bycatch
1994	120	807	53,787	7,115	67	8
1995	152	1,186	152,285	12,142	128	13
1996	96	508	42,113	3,941	83	11
1997	109	828	79,788	3,721	96	21
1998	109	569	109,034	4,875	192	22
1999	141	755	77,956	4,537	103	17
2000	157	614	163,640	7,479	267	22
2001	70	186	14,926	4,389	80	3
2002	76	445	91,652	4,710	206	19
2003	67	247	39,264	5,612	159	7
2004	62	174	32,618	4,085	187	8
2005	26	58	11,820	1,528	204	8
2006	21	88	26,029	666	296	39
2007	36	71	21,117	1,735	297	12
2008	19	36	11,499	1,127	319	10
2009	7	10	1,016	30	102	34
2010	10	13	3,450	5	265	690

Table 3. Catch and effort data on shrimp and landed bycatch for skimmer in New River, 1994-2010 (courtesy of DMF trip ticket program).

Year	Participants	Trips	Shrimp (lbs)	Sold bycatch (lbs)	Mean catch per trip (lbs)	Ratio shrimp of sold to bycatch
1994	5	12	1,468	7	122	226
1995	25	85	21,554	0	254	0
1996	34	224	42,677	267	191	160
1997	41	341	75,029	188	220	400
1998	43	302	69,396	13	230	5,338
1999	49	449	68,813	222	153	310
2000	77	615	155,949	2,508	254	62
2001	44	306	36,043	1,879	118	19
2002	51	832	173,091	1,701	208	102
2003	55	564	89,780	1,356	159	66
2004	37	432	82,384	385	191	214
2005	24	155	21,714	307	140	71
2006	15	169	76,501	121	453	632
2007	27	265	93,094	152	351	611
2008	20	148	48,834	12	330	4,246
2009	9	42	4,973	3	118	1,658
2010	16	297	102,032	330	344	309

Table 4. Catch and effort data on shrimp and landed bycatch for other gear (cast nets, gill nets, etc.) in New River, 1994-2010 (courtesy of DMF trip ticket program). *Data confidential due to less than three participants reporting landings.

Year	Participants	Trips	Shrimp (lbs)	Sold bycatch (lbs)	Mean catch per trip (lbs)	Ratio shrimp of sold to bycatch
1994	2	2	*	*	*	1
1995	24	162	12,837	11,043	79	1
1996	12	20	884	1,528	44	1
1997	11	53	2,934	4,394	55	1
1998	3	6	130	442	22	0
1999	5	10	387	553	39	1
2000	11	18	1,041	827	58	1
2001	7	9	519	819	58	1
2002	5	5	209	184	42	1
2003	5	16	670	27	42	25
2004	6	5	100	710	20	0
2005	4	4	594	1,039	149	1
2006	10	64	4,870	349	76	14
2007	6	16	790	2,100	49	0
2008	3	6	329	631	55	1

Table 4 (cont).

Year	Participants	Trips	Shrimp (lbs)	Sold bycatch (lbs)	Mean catch per trip (lbs)	Ratio shrimp of sold to bycatch
2009	1	1	*	*	*	0
2010	3	3	140	104	47	1

Table 5. Catch and effort data on crab and landed bycatch for crab trawls in New River, 1994-2010 (courtesy of DMF trip ticket program). *Data confidential due to less than three participants reporting landings.

Year	Participants	Trips	Crab (lbs)	Sold bycatch (lbs)	Mean catch per trip (lbs)	Ratio crab of sold to bycatch
1994	7	35	10,848	492	310	22
1995	15	94	33,616	3,512	358	10
1996	14	47	8,284	519	176	16
1997	14	187	33,196	2,777	178	12
1998	10	62	3,988	373	64	11
1999	12	32	23,214	489	725	48
2000	11	42	17,643	555	420	32
2001	16	103	17,476	446	170	39
2002	13	77	12,190	183	158	67
2003	15	101	18,732	459	185	41
2004	23	159	41,192	863	259	48
2005	14	125	28,060	113	224	248
2006	2	5	*	*	*	17
2010	9	32	23,383	61	731	386

Table 6. Percent of weight in pounds of trawl biomass caught in the DMF 25-foot, 4-seam otter trawl in New River 2003-2009.

Year	Shrimp	Invertebrate	Commercial Finfish	Non-commercial Finfish	Combined Finfish	All Bycatch
2003	29.9	18.0	36.3	15.7	52.0	70.1
2004	40.2	14.2	24.7	20.9	45.6	59.8
2005	40.8	13.9	30.9	14.3	45.2	59.2
2006	58.0	6.2	27.7	8.1	35.8	42.0
2007	38.7	9.0	40.6	11.7	52.3	61.3
2008	2.9	7.8	23.2	66.1	89.3	97.1
2009	1.4	8.2	43.5	38.6	82.1	90.3
All	51.2	9.1	23.3	16.3	39.6	48.8




Figure 1. Map of the New River showing the areas for the PNAs and SSNA as well as the other trawl closure line.




Figure 2. Shrimp trips by gear in New River, 1994-2010.




Figure 3. Mean catch of shrimp in New River, 1994-2010.




Figure 4. Total catch of shrimp in pounds by gear in the New River, 1994-2010.




Figure 5. Number of trips and participants in the New River crab trawl fishery, 1994-2010.




Figure 6. Pounds of crab landed and number of pounds of crab landed per trip in the New River crab trawl fishery, 1994-2010. Dotted line represents the average pounds per trip landed from 1994-2010.

VOLUMETRIC MEASUREMENT OF SHRIMP

December 9, 2011

I. ISSUE

Under Rule 15A NCAC 03L .0105 (2) recreational fishermen using cast nets in closed areas to harvest shrimp are limited to 100 shrimp per person per day, it was requested that a volumetric measurement of shrimp be used in place of counts to check individuals.

II. ORIGINATION

The North Carolina Marine Patrol

III. BACKGROUND

The cast net fishery was originally developed for live bait fishermen who wanted to capture shrimp for bait. Overtime the fishery has evolved into a means of capturing shrimp for personal consumption and for sale. This rule has been in place since 1985, with very few changes made since its inception; however, the number of participants in the cast net fishery for shrimp in these closed areas has drastically increased. As a result of this increase in recreational consumption harvest for shrimp harvested with cast net gears, it has become increasingly difficult and dangerous for Marine Patrol officers to enforce the 100 count rule. This is mainly due to the nature of this fishery, where a number of fishermen will work together and combine their harvest in a single large container. This will often time lead to a Marine Patrol officer having to count shrimp while surrounded by numerous fishermen. Establishing a volumetric measurement would help to alleviate this problem.

IV. AUTHORITY

G.S. 113-134 Rules

G.S. 113-182 Regulation of Fishing and Fisheries

G.S. 143B-289.52 Marine Fisheries Commission—Powers and Duties

15A NCAC 03L.0101 Season

15A NCAC 03L.0105 Recreational Shrimp Limits

V. DISCUSSION

The Marine Patrol would like to use a volumetric measurement of shrimp instead of an individual count of 100 heads on shrimp per person. This method would be a more efficient and effective way to check individuals who harvest shrimp with a cast net in closed areas, allowing officers to check more individuals. This method would also make it safer for each officer working alone, who may need to determine the quantity of shrimp harvested for a number of recreational cast netters. The count of 100 shrimp takes a lot of time, especially when there are extra ordinary circumstances involved, i.e., weather, numerous fishermen, language barriers, time of day, live shrimp, size, and location. By using a volumetric measurement of shrimp, an officer would not have to bend down and expose their backs, weapon and other body parts in a manner that could leave them vulnerable to attack. The Marine Patrol would like to use a half-gallon bucket, approximately 80-120 shrimp depending on size as the standard volumetric measurement of shrimp taken by cast nets in a closed area. This proposed measure would still allow fishermen to retain shrimp for bait purposes and promote officer safety.

VI. PROPOSED RULE

15A NCAC 03L .0105 RECREATIONAL SHRIMP LIMITS

It is unlawful to:

(1) Possess more than 48 quarts, heads on or 30 quarts, heads off, of shrimp per person per day or if a Vessel is used, per vessel per day for recreational purposes except as provided in 15A NCAC 03O .0303 (e) and (f).

Option 1

(2) Take or possess shrimp taken from any area closed to the taking of shrimp except ~~100 shrimp per one half gallon~~ of heads on shrimp person per day may be taken while fishing in a closed area with a cast net. Individual limits shall be kept in separate one half-gallon containers.

Option 2

(2) Take or possess shrimp taken from any area closed to the taking of shrimp except 100 shrimp per person per day may be taken while fishing in a closed area with a cast net. Individual limits shall be kept in separate containers.

*History Note: Authority G.S. 113-134; 113-182; 143B-289.52;
Eff. April 1, 2009.*

VII. PROPOSED MANAGEMENT OPTIONS

- (+ Potential positive impact of actions)
- (- Potential negative impact of actions)

1. *Status quo* (continue to limit fishermen to 100 shrimp in closed areas)
 - + Still allows shrimp to be taken for bait purposes only
 - + No equipment to be purchased
 - + No grey areas, either 100 shrimp or not
 - Very time consuming when dealing with large groups
 - Potential safety issues when working alone
2. Use a volumetric measurement of shrimp (limit fishermen to one half gallon heads on shrimp)
 - + Officers less vulnerable to attacks
 - + More effective use of time, allowing more people to be checked
 - + Standardization of measurement throughout the State
 - + No numbers to keep track of
 - Shrimp may exceed 100 or less depending on size/count
 - Larger containers may promote the use of cast nets to take shrimp in closed areas for consumption and not for bait purposes
 - Extra gear to carry
3. Limit fishermen to 100 shrimp in closed areas but require individuals to keep their catch in individual containers
 - + Still allows shrimp to be taken for bait purposes only
 - + No equipment to be purchased
 - + No grey areas, either 100 shrimp or not
 - Very time consuming when dealing with large groups
4. Eliminate the ability to take or possess 100 shrimp per person per day in areas closed to shrimping.

- + Officers less vulnerable to attack
- + Less time consuming for officers to enforce
- + No equipment to purchase
- Eliminates a bait fishery

VIII. RECOMMENDATIONS

PDT: Use a volumetric measurement of shrimp (limit fishermen to one half gallon heads on shrimp) OR;
 Limit fishermen to 100 shrimp in closed areas but require individuals to keep their catch in individual containers

Prepared by: Lieutenant Michael S. Ervin
 Michael.ervin@ncdenr.gov
 (910)-796-7286

Revised: December 9, 2011

NOTICE OF TEXT ATTACHMENT

MFC Rulebook Index Worksheet

Rule	Rulebook Page #	Subject	Index Entry (Bold major headings)	Add/Delete
03L.0105(2)	40	Recreational Shrimp Limits		

RESTRICTED TRAWL AREA OFFSHORE BOGUE BANKS

November 15, 2011

I. ISSUE

Request to investigate the history of existing Marine Fisheries Rule 15A NCAC 03J .0202 (3), which prohibits trawling within one-half mile of shore in the Atlantic Ocean between Beaufort Inlet and Salter Path with the intention of changing that distance to one-fourth of a mile.

II. ORIGINATION

Marine Fisheries Commissioner in January of 2009

III. BACKGROUND

The Marine Fisheries Commission (MFC) has made use of its authority to regulate fishing practices in areas of existing or potential conflict between user groups. This prohibition of trawling within one-half mile of shore in the Atlantic Ocean between Beaufort Inlet and Salter Path originated in 1967 in an effort to separate conflicting user groups and residents along the shore from shrimp trawlers. Until a few years ago, there were four ocean piers (TripleS, Oceanana, Sportsman's and Sheraton), several public swimming areas (Fort Macon State Park, Oceanana, Atlantic Beach Circle, Sheraton), Fort Macon State Park, and several beachfront developments (Tar Landing, Sea Spray, Place at the Beach, etc.). The prohibition of trawling within one-half mile of the beach separated trawlers from physical proximity to the ocean fishing piers and helped reduce the amount of bycatch discarded from the trawlers that washed up on the beach during prevailing southwest winds in the summer, disturbing Park officials, beach and pier anglers, beachgoers and residents. Due to the removal of two of the four fishing piers, it has been requested that the one-half mile distance from shore in rule be reduced to one-fourth of a mile to allow for white shrimp harvest during years when they are present along the beach.

IV. AUTHORITY

G. S. 113-134 RULES

G.S. 113-182 REGULATION OF FISHING AND FISHERIES

G.S. 143B-289.52 MARINE FISHERIES COMMISSION –POWERS AND DUTIES

North Carolina Marine Fisheries Commission Rules for Coastal Fishing Waters (15A NCAC)

03J .0104 TRAWL NETS

03J .0202 ATLANTIC OCEAN NET RULES

V. DISCUSSION

The request to reduce the distance that Atlantic Ocean shrimp trawlers (by virtue of the mesh sizes in the rule) must stay offshore of the eastern portion of Bogue Banks was made due to a reduction in the number of ocean fishing piers present from four to two. Two piers still exist (Oceanana and Sheraton) in that area as does the state park, swimming beaches and more numerous condominium developments. Public sentiment against trawling has become increasingly negative in the past few years and bycatch washed up on the beach is less well-tolerated (based on complaints) by beachgoers and residents. Therefore, even though the

number of piers has decreased by two, the remaining two piers, increased houses and condominiums, the state park and public and private swimming areas that are present still cause the same concerns that led to the 1967 closure. With a legislative study ongoing to look at banning trawling, relaxing restrictions that have been effective is not advised.

It must be noted that while examining this issue, the Shrimp FMP PDT found a contradiction to other existing rules regarding mesh size. This contradiction will be examined by the Rules Advisory Team and addressed in a separate issue paper.

VI. PROPOSED MANAGEMENT OPTIONS

(+ Potential positive impact of action)

(- Potential negative impact of action)

1. *Status quo* (prohibit shrimp trawling within one half mile of shore in the Atlantic Ocean between Beaufort Inlet and Salter Path)
 - + No change in distance from shore requirement
 - + No potential increase in complaints from pier patrons, swimmers and beachgoers
 - Continued restriction of area that shrimp trawlers have access to
2. Allow shrimp trawling within one quarter mile of shore in the Atlantic Ocean between Beaufort Inlet and Salter Path
 - + Additional area opened to shrimp trawling in that area
 - Potential increase in numbers of complaints received due to increased bycatch and physical interaction with pier patrons and beachgoers.

VII. RECOMMENDATION

PDT: Due to the remaining two fishing piers along that stretch of shoreline, increased condominium development and the continued presence of the state park and swimming beaches (Oceanana, Fort Macon State Park, Dunes Club, the Circle, etc.), it is recommended that the rule remain *status quo*.

Prepared by: David L. Taylor

David.L.Taylor@ncdenr.gov

252-808-8074

October 7, 2011

NOTICE OF TEXT ATTACHMENT

#6 – Explain Reason for Proposed Action:

**15A NCAC 03J .0202 ATLANTIC OCEAN
MFC Rulebook Index Worksheet**

Rule	Rulebook Page #	Subject	Index Entry (Bold major headings)	Add/Delete
03J .0202 (3)	20	Atlantic Ocean Trawling		




Figure 1. Bogue Banks Trawl Restricted Area 15A NCAC 03J .0202 (3)

Public Input for Shrimp FMP review 2011

Press Release sent November 7, 2011

MOREHEAD CITY – The N.C. Division of Marine Fisheries is asking the public to submit comments on issues they would like to see addressed in an upcoming Shrimp Fishery Management Plan.

The division is beginning a mandated five-year review of the N.C. Shrimp Fishery Management Plan that was adopted by the N.C. Marine Fisheries Commission in 2006. The agency is soliciting public comment as part of an internal process to determine what procedural method to take in reviewing the plan.

If changes in management strategies or rules are needed, the division will pursue a plan amendment, where division staff and an advisory committee develop positions on specific issues that need to be addressed. If changes in management strategies are not required, the division will proceed with a revision, which is a more abbreviated process that involves updating data and fishery information contained in the plan.

Written comments will be accepted until Dec. 2 and should be addressed to Trish Murphey, N.C. Division of Marine Fisheries, P.O. Box 769, Morehead City, N.C. 28557 or sent by email to Trish.Murphey@ncdenr.gov or to Chris Stewart, N.C. Division of Marine Fisheries, 127 Cardinal Dr., Wilmington, N.C. 28405 or sent by email to Chris.Stewart@ncdenr.gov.

State law requires the division to prepare a fishery management plan for adoption by the Marine Fisheries Commission for all commercially and recreationally significant species or fisheries that comprise state coastal waters. These plans provide management strategies designed to ensure long-term viability of the fishery. State law also requires the division to review each fishery management plan every five years.

###

nr-53-11

From the Public

1. You want to finally start managing the shrimp-- EASY! Ban all trawling in the waterways and nurseries, for years I have watched the tiny shrimp get murdered in the waterways, for the most part they are so small these guys cull thru them for just the big ones and shovel the smaller ones overboard, along with all the juvenile dead fish, what a waste. I also have observed the boats on the outside shoveling and culling thru the smaller shrimp when the market is flooded, I remember years back when marine fisheries had the inside and outside closed and kept sampling till the shrimp reached a predetermined size on the outside, these were white shrimp, that was a win win, everybody made the best money they had in years because the shrimp were a decent count and brought a fair price. In summary, ban all trawling on the inside, it will help the fishery and the shrimp season, allow both commercial and recreational to bait and cast net for their shrimp, eliminates the bycatch

and people will not be killing the juvenile (popcorn) shrimp, which in turn will make it profitable for the outside boats, anyone wanting to commercial trawl can afford a real shrimp boat now., Also control the outside trawling, keep it closed until the white or brown shrimp reach a predetermined size.

PDT Response

Currently, it is unlawful to use trawl nets for any purpose in primary nursery areas (15A NCAC03N .0104) and permanent secondary nursery areas [15A NCAC03N .0105(a)]. In areas considered special secondary nursery areas (SSNA), it is unlawful to use trawl nets except by proclamation from the Fisheries Director from August 16 through May 14 [15A NCAC03N .0105(b)]. Management rationale for this rule included minimizing bycatch by delaying the trawl opening date to reduce the finfish bycatch and to reduce user conflicts. Shrimp abundance, count size, growth, as well the abundance of other economically important species (crabs and finfish) are sampled prior to August 16 depending on waterbody. Once it has been established that the shrimp are of a fair marketable size and bycatch is minimal, a proclamation is issued opening that area. Generally the ocean, sound, and major rivers remain open to shrimping year round except when extreme environmental conditions lead to significant percentages of small shrimp in that area. Allocating the resource solely to commercial ocean trawlers does not consider the needs of all user groups and does not provide sufficient opportunities for recreational and commercial shrimpers to optimize the use of the resource, thus not meeting the goals of the Shrimp FMP. In addition, a legislative study committee has been appointed to look at trawling in North Carolina. This includes all trawling, not just shrimp trawling and therefore will not be further addressed in this FMP.

NC Shrimp Fishery Management Plan 2006

http://portal.ncdenr.org/c/document_library/get_file?uuid=7dc55c67-c6df-4a39-9ffc-32471c055c23&groupId=38337

See Shrimp Trawl Bycatch p 90

See Shrimp Management by Size in North Carolina estuaries p 92

See Shrimp Management in the Atlantic Ocean p 110

2. Address the ongoing problem of shrimp size (or count): as re-opening and closing of the season.

I guess what I am looking for is a solution for a waste of our resource. Each year it seems to be in a different water body so my suggestion is to have a state-wide count rule. What I mean by waste is the low price that small shrimp bring. It seems that all other species have a size limit. I think one could fit the shrimp fishery. My suggestion of size would be 36/40 heads on minimum.

PDT Response

In North Carolina, shrimp develop at different growth rates depending on water temperature and salinity. Post-larvae shrimp are carried by wind driven currents from the ocean to the upper reaches of the estuaries, where in several weeks they develop into juveniles. As growth increases, shrimp migrate to the deeper, saltier waters of the sound and eventually to the ocean. As shrimp migrate to the ocean, they enter areas that are open or may be opened by the DMF to the harvest of shrimp. These areas include bays, creeks, rivers, sounds and the ocean. Sampling is conducted by the DMF staff to determine if an area should be opened or closed, based primarily on size and count. Over time, target sizes for opening different waterbodies have evolved and allows for better flexibility of management for both recreational and commercial shrimpers than what a single statewide count size will allow.

NC Shrimp Fishery Management Plan 2006

http://portal.ncdenr.org/c/document_library/get_file?uuid=7dc55c67-c6df-4a39-9ffc-32471c055c23&groupId=38337

See Shrimp Management by Size in North Carolina Estuaries p 92

3. Changes were adopted in 2006 for the boundary lines for trawling in the Neuse River. They were extended down river about a mile to Wilkerson's Point (around the ferry landing). I would like to see the boundary go back to where they were prior to 2006. This allowed us to harvest shrimp around the mouth of Baird and Slocum creeks. I am still allowed to pull a crab trawl but not a shrimp trawl in this area. Impact in these areas is minimal because of shallow waters. Only small boats harvest shrimp in this area. Please present this in the upcoming Shrimp Management plan. I pulled a shrimp trawl around the mouth of Bairds Creek for thirty years until they closed it. Thanks for your time and voice.

PDT Response

The 2005 Southern Flounder FMP recommended that shrimp trawling be examined with the goal of reducing the impact of incidental shrimp trawl bycatch on juvenile southern flounder. The restrictions that resulted from the 2006 Shrimp FMP established the shrimp trawl line at Wilkerson Point and Cherry Point. Prior to this, no line had been established. Crab trawling was not impacted due mainly to the difference in mesh sizes between shrimp and crab trawls. Crab trawls must have a minimum of four inch (stretch mesh) webbing while shrimp trawls have a minimum mesh length of one and one half inches, allowing fewer juvenile flounder to escape. The 2009 DMF stock assessment indicated that the southern flounder stock is overfished and overfishing is occurring and since the stock is heavily influenced by recruitment, it is felt that this measure should remain in place. The Marine Fisheries Commission has had a standing policy since 1991 to "establish the goal of reducing bycatch losses to the absolute minimum and to consciously incorporate that goal into all its management considerations".

NC Shrimp Fishery Management Plan 2006

http://portal.ncdenr.org/c/document_library/get_file?uuid=7dc55c67-c6df-4a39-9ffc-32471c055c23&groupId=38337

See Southern Flounder Bycatch in the Inshore Shrimp Trawl Fishery p 91

See Shrimp Management in Neuse River p 106

4. Phone in with input about the (1) amount of puppy drum in the southern rivers and their predatory impacts on shrimp in the rivers. There are "acres of schools" of puppy drum in the rivers. One fisherman this person mentioned, was fishing for mullet and it took 4.5 hours to remove the drum from the net. (2) New River has become a cleaner river and that we should

consider opening New River further up (halfway to Jacksonville) to skimmer trawls. This person is a supporter of skimmer trawls.

PDT Response

Red drum prey upon numerous species of shrimp, crab, and fish at various stages of their life history. Other species of fish also prey upon penaeid shrimp making it hard to quantify exactly how much is consumed by each species and what effect it has on the year-to-year fluctuations in shrimp abundance. There also has been a strong body of scientific evidence indicating that as red drum grow, their diets shift from shrimp and crab to primarily fish. In many cases age 1-2 red drum consumed fish more frequently, in greater numbers, and in greater volume than either shrimp or crabs alone. Given that the 2009 Atlantic States Marine Fisheries Commission stock assessment for red drum indicates that the current regulations have been effective at preventing overfishing, it is unlikely that these regulations will be changed to protect species outside of red drum.

Division of Water Quality staff from the Wilmington Regional Office agree that the New River is cleaner in some aspects, noting that there have been fewer algal blooms resulting from nutrient loading associated with municipal discharge. However, the shrimp line established at Grey Point was created to protect smaller shrimp until they grow to a harvestable size and not due to water quality. Based on the 2006-2010 DMF juvenile shrimp management sampling, shrimp biomass was typically higher for stations below the current line. In other regions of the state fishermen have also expressed their displeasure in the constant movement of shrimp lines, the line established at Grey Point is well marked (telephone poles and orange barrels), easily enforceable by Marine Patrol, and has been widely accepted by fishermen since the 1980s. NC Red Drum Fishery Management Plan Amendment I 2008

http://portal.ncdenr.org/c/document_library/get_file?uuid=cd9ce130-c426-40d7-b9d1-ecf446dec77e&groupId=38337

See Stock Status of Northern Red Drum Stock (attached at the end)

NC Shrimp Fishery Management Plan 2006

http://portal.ncdenr.org/c/document_library/get_file?uuid=7dc55c67-c6df-4a39-9ffc-32471c055c23&groupId=38337

See Shrimp Management in New River Above the Highway 172 Bridge p 96

5. Over the years, one of NCDMF's answers to shrimp management has been to close certain areas of NC's coastal waters. While that may be an effective management effort for some areas over a given period of time; I am convinced that closing an area and keeping it closed for years and years is not a real good shrimp management practice. Maybe it was for awhile, but by keeping these areas closed on a permanent basis is having a negative impact on the resource, the areas closed, and the fishermen.

Maybe it's time to try some other management practices such as reopening some of these areas on certain days of the week and restricting the size and type of trawls used in these areas. Also, in some cases we may be better served by allowing the resource to manage itself.

PDT Response

Many of the lines in existence now have resulted from years of practical experience, sampling and public input. The North River lines, for example were the result of a public meeting in Harkers Island in June of 1997. At that meeting, fisherman expressed displeasure over the constant movement of the lines at that time and negotiated acceptable lines to be made "permanent" with DMF staff. While extreme rainfall events occasionally force small shrimp

downstream of the lines, overall the lines serve the purpose of protecting smaller shrimp until they grow to a harvestable size and migrate out of the creeks, into the river and out of the inlet. The question of "cultivating" or "turning over" the bottom in these long-closed areas is one which has its proponents and detractors. A large body of research exists (although very little of it local) documenting negative effects of damage caused by the physical trawl and resulting sedimentation to seagrass beds, oyster rocks, coral reefs and larval fish and shellfish

NC Shrimp Fishery Management Plan 2006
http://portal.ncdenr.org/c/document_library/get_file?uuid=7dc55c67-c6df-4a39-9ffc-32471c055c23&groupId=38337

See Environmental Factors p 59

See Management of Trawling for Habitat Protection p 90

See Shrimp Management in Bogue Sound and North River p 104

6. *Unfortunately I just got my Beacon Newspaper in the mail, and thus missed the deadline for input on shrimp issue [Dec 2]. However, if there is interest in hearing some thoughts at this late date, I offer the following.*

In southern Brunswick County, Marine Fisheries identifies the side of the intercoastal waterway as "primary nursery", no trawling allowed, in recognition that the side of the waterway is nursery area. However MF allows trawling in the middle of the intercoastal. The obvious question is ..."what happens as the tide goes out", and the obvious answer is all the marine life that MF is attempting to protect in the primary nursery flows with the outgoing tide to the middle of the waterway, and thus is subject to trawling. Clearly the rules in this instance work in conflict with each other. Thus, a SOLUTION. What if trawling wasn't allowed 3 hours before low tide and 3 hours after low tide? This would prevent juvenile marine life from direct exposure to trawling in the middle of the waterway during low water.

The same issue impacts juvenile marine life leaving creeks out to the waterway on a outgoing tide. SOLUTION: Don't allow trawling within 1000 yards of a creek leading into the intercoastal. Sharing my thoughts; sorry I missed the deadline, but hope you can pass on to the MF folks.

PDT Response

This management strategy would be difficult to enforce and would further restrict shrimpers in the Brunswick county area, basically closing shrimping in the IWW. Large portions of the IWW around the Yaupon Beach Bridge (Yellow Banks) would be closed to shrimping as well a majority of the waterway below the Shallotte Inlet to the South Carolina state line. Most fishermen in Brunswick County tend to work around the tides. In North Carolina, the tides follow a semi-diurnal pattern; with two high tides and two low tides each day at different heights. Only allowing fishermen to fish three hours before and after each low tide would subsequently limit fishing to 12 hours. Further compounding the problem is the fact that the tide cycle shifts about an hour forward each day, forcing fishermen to work at night if they are to catch the right tide. More fishermen on the water at night could lead to potential safety issues and increased user conflict among boaters and shrimpers alike.

NC Shrimp Fishery Management Plan 2006

http://portal.ncdenr.org/c/document_library/get_file?uuid=7dc55c67-c6df-4a39-9ffc-32471c055c23&groupId=38337

See Shrimp Management in Brunswick County p 101

2012 Shrimp FMP Revision Committee Review Table

At the MFC meeting in August of 2012, the MFC voted to send the 2012 Shrimp FMP Revision to regional and standing committee review as well as public review. The revision was presented to the Southern Regional AC, the Northern Regional AC, the Habitat and Water Quality AC and the Crustacean/Shellfish AC during September and October of 2012. Table 13.1 summarizes recommendations of the different ACs. The DMF changed its recommendation from revise the plan to amend the plan after following the AC meetings and public comment (Table 13.1).

Table 13.1 Advisory Committee recommendations on whether to revise or amend the Shrimp FMP

ISSUE	DMF	SOUTHERN Sept 19	NORTHERN Sept 27	HABITAT AND WATER QUALITY Oct 2	CRUSTACEAN/SHELL FISH Oct 2
Recommend the revision or the amendment of the 2006 Shrimp FMP. The 2012 revision of the plan had data up through 2010, latest fishery trends, a new protected resources section and the continuation of management strategies put in place through the 2006 Shrimp FMP.	<p>May 2012 Recommend to revise the Shrimp FMP</p> <p>Oct 2012 Recommend to amend the Shrimp FMP</p>	<p>Recommend an amendment to the Shrimp FMP. Recommend to investigate the use of otter trawls upstream of the 172 Bridge over New River and adding a special license for bait shrimp fishery.</p> <p>150 public attended</p>	<p>Recommend to revise the Shrimp FMP</p> <p>6 public attended</p>	<p>Recommend to revise the Shrimp FMP</p> <p>4 public attended</p>	<p>Recommend to revise the Shrimp FMP.</p> <p>55 public attended</p>

Public Comment

Bycatch

Bycatch is no different from what it was years ago.

There are less shrimpers today, thus their impact is less. Among those who are left, effort is down. We can't afford to go unless we know there is a market.

This is a good plan. We catch 25 lbs of shrimp and catch 25 lbs of bycatch, 85% of the bycatch lives. I have caught 5-6 speckled trout per season and have never caught a drum. I catch mostly croakers and pinfish. When you catch a lot of shrimp, there is very little bycatch.

Ninety percent of bycatch is tongue fish and pinfish.

A 2004 bycatch study showed Cape Fear had a bycatch ratio of 0.38:1, Williams Landing had a ratio of 0.7:1, and Brunswick County had a ratio of 1.55:1.

I do all I can to return bycatch to the water.

Otter trawls, skimmer trawls, and channel nets all have bycatch. There is bycatch in the ocean too.

We have BRDs in our nets to let out the bycatch.

I throw out flounders and crabs first, croakers live a while.

I worked with Logothetis on bycatch study. There was less than 1 lb of total bycatch to 1 lb of shrimp and 0.5 lbs of finfish bycatch to 1 lb of shrimp.

There are four species (spot, croaker, weakfish, and flounder) that are concerned" or "depleted." To meet the objectives of the plan, the Division needs to amend the plan and study the best course of action.

A revision means status quo. I want an amendment. We are wasting spot, weakfish and croaker. You have to go to VA to catch croaker. Limit inshore trawls to 50 ft headrope length.

Yes there is bycatch. Efforts have been made to reduce bycatch. There were nine active packing houses, now there are three. There is less bycatch now because effort is down. Trawlers are bigger and come from out of state. Look at headrope length.

There are 480 million to 500 million juvenile finfish killed annually. We need to minimize bycatch. Look at 50 ft headrope limit.. Amend FMP to address the issue of bycatch of shrimp trawlers operating in our inshore waters.

Bycatch has been addressed. PNAs were recommended by the NCFR. There are TEDs and BRDs that were developed through cooperation of the industry. Can't say it is good enough, but the industry has done things to address bycatch.

Need to amend the plan. There are other things we can do to reduce bycatch while sustaining and perpetuating the industry. Reduce internal trawling and study its effects.

Need to amend the plan. There are multiple gear types to replace trawling. Limiting the size of the gear may also work. The plan lacks adequate options.

Spot fishing is down 90%, Croaker are plentiful in VA, weakfish landings are down; all the states south of us are reporting higher catches than NC.

Using numbers from the DMF website, finfish landings appear to be down 90% based on best years (70s & 80s) and worst years (2006-2009). Water quality is an issue, but bycatch is problem. Amend.

Looking at the landings data for spot, croaker, and weakfish as indicators of the stock's health is misleading; there are other reasons beside bycatch driving why landings down. The loss of the flynet fishery south of Hatteras, less effort, and new regulations are just a few of the things that have caused the landings to go down. It's not that the fish aren't there, they are not being caught.

If we push trawlers out 3 miles, bycatch will still be a problem and we will then have to deal with super draggers.

The industry is trying to reduce bycatch, we would be willing to give up head rope lengths before be pushed out 3 miles.

We need to implement a maximum head rope length of 50 feet in certain areas to reduce the amount of juvenile finfish being killed.

We need to amend, to address both commercial and recreational bycatch. Cast nets catch bycatch too.

Regulations

More rules will kill us.

Do not need any more changes. Do not like what has been done.

The shrimp fishery needs zero management.

We do not need any more regulations. Our rights are being violated. We are overregulated now. You are discriminating against otter trawls. They should be allowed to work above the 172 bridge just like skimmer trawls.

There are too many rules. We do not need any more closures.

I disagree with allowing skimmer trawls above the 172 bridge and not otter trawls.

If skimmer trawls can go up New River, then otter trawls should be allowed too.

Shrimping regulates itself.

The goal says to consider all user groups. I am a live bait shrimp dealer. You need to consider a special license for bait shrimp fishermen. I pull a 15 ft trawl and pull 10 minute tows. There is very little bycatch. You have bait shrimping licenses in other states. Managing by size does not allow for a bait shrimp fishery to exist. Can we look into what other states have done and how they do their licensing?

I am concerned about the turtles. I see a lot of turtles that have been hit by boats.

You can't manage by size because it changes from year to year and a set size or count may not work every year. At beginning of the year smaller shrimp are worth more and fishermen should be allowed to catch and sell these.

Socio/Economic

We are losing our heritage. Our rights are being destroyed.

There are far less of us than in the past. Our impact can't be that great. Pamlico Sound is not like it used to be. There are fewer boats everywhere.

You need a historical perspective of bycatch, the industry has greatly diminished.

You worry about the mortality of finfish, what about the mortality of fishermen?

If you cut out shrimping, then we will be sitting on the streets collecting food stamps.

It's the large fish houses and bigger boats that are making all the money. More regulations will only hurt the smaller operations. If you change anything, you will put the small boats out of business.

There are very few of the younger generations fishing; there is no use for them to get into the fishery because of all the regulations. In the 1980s, 95% of the fishermen relied on all their income to come from fishing. Now it's more like 51%. Everyone else has to rely on other sources of income.

If shrimp are not available in a particular area, then we don't go, our boats burn too much fuel to go look for them.

Small shrimp are worth more, early in the season, when they are the first on the market. Once those shrimp size up, the larger shrimp from Pamlico sound flood the district and our shrimp aren't worth anything when their shrimp are selling for \$2 a pound. When the shrimp get bigger, they fall out to the ocean, where we can't catch them. we should have access to them.

Effort is down, probably due to economics (rising fuel / operations costs, imports) more so than management measures.

I have seen a lot of impacts from the shrimp fishery. Finfish numbers are down. I make a living off the recreational fishery. Boat manufacturers, other tackle store owners are being impacted. You need to amend the shrimp plan. My tackle shop is losing money as a result of the poor fishing in North Carolina, bycatch and its effects need to be addressed.

Imports are killing us, fuel prices are killing use, we don't need any new regulations; North Carolina consumers want wild caught shrimp.

There are forces at work trying to end shrimping in NC, fishermen need to strike together and join up with the North Carolina Shrimpers Association

Habitat and Water Quality

We need to clean out the creeks. Increased development and population growth are degrading the habitat.

The New River bottom needs to be stirred up. There are no brown shrimp anymore.

The sewage plant from Holly Ridge is filling New River.

Sewer plant outlets and pump-out stations need to be examined; their water is overflowing into our PNAs. Pollution also enters our nursery areas from Marines washing boats down.

Sludge from the Marine Corp Base and Jacksonville is in the New River. The river needs turning up to get the stuff off the bottom.

If you don't turn the bottom over, New River won't produce.

If you close Pamlico Sound, then you need to just give up sportfishing and crabbing. You need to drag the bottom. Look at Rose Bay. It has not been dragged for years and now there is very little in it.

The habitat and water quality is bad in Neuse River.

Pamlico Sound is the second largest nursery area. Need to do something, so amend the plan.



CCA North Carolina
Board of Directors
Jim Hardin
Chairman
Greg Hurt
President
Bud Abbott
Vice President
Ty Coyll
Vice President
Dick Hamilton
Vice President
Seth Vernon
Vice President
Bud Abbott
Treasurer
Bill Austin
Dr. Nick Blackerby
Andrew Boyd
Craig Boyd
Mel Broughton
Bill Brown
Arthur Brownell
Billy Byrd
Frank Colonna
Jay Dall
Jorry Eaton
Dennis Edwards
Ken Eller
Dave Hawkins
Harris Huddle
Mike Kalot
John Knight
Dr. Bill Lambeth
Chuck Laughridge
Bob Lorenz
Bill Mandulak
Larry Mize
Edwin Newell
Bo Nowell
Ken Oppenheim
Bert Owens
Dale Petty
Alex Posey
Jeff Prince
John Rea
Chris Richey
Rob Spelr
Matt Tynes
Warren Ward
Chris West
Rex Willis III
Donald Willis, Jr

Stephen Ammons
Executive Director

Cindy Davis
Office Administrator

The Coastal Conservation Association North Carolina has reviewed the proposed revision to the Shrimp Fisheries Management Plan. CCA NC is very concerned that the proposed Shrimp FMP does not meaningfully address high levels of bycatch in the otter trawl fishery in our sounds and estuaries. The discarded bycatch of juvenile finfish ranges from two to four times as much as the harvest of shrimp by weight. This translates into an unacceptable waste of immature croaker, spot, weakfish and southern flounder that are discarded dead. These species represent both commercially and recreationally important species to North Carolina fishermen. Two of these species are listed as depleted and two are listed as species of concern.

In addition to bycatch issues in the shrimp otter trawl fishery, this bottom disturbing gear has the potential to limit productivity of certain areas of our sounds and estuaries by increasing turbidity and reducing bottom structure. This could again limit the productivity of areas of our sounds.

CCA NC urges the NC DMF to pursue a full amendment process for the Shrimp FMP. This amendment should consider multiple gear types that might have less bycatch such as skimmer trawls, channel trawls or shrimp pots where appropriate. Limitation in the size of otter trawls could limit bycatch and reduce destruction of bottom habitat and structure. Other options could be to examine time and area closures to protect important juvenile species as well as potential improvements in Bycatch Reduction Devices (BRDs).

Finally, while the Shrimp FMP recognizes the problems of discarded bycatch, there are no objective, measurable reduction targets for otter trawl bycatch. If you do not have some idea of what success should look like, and you do not quantify it, you are unlikely to achieve it

CCA NC believes a full amendment to the Shrimp FMP is the only possible way to fully examine all these issues.

Dedicated to the Conservation of North Carolina's Marine Resources
4809 Hargrove Road, Suite 123 Raleigh, NC 27616
(919) 781-3474 (FISH) Fax (919) 781-3475 contact@ccanc.org
www.ccanc.org



September 26, 2012

NC Division of Marine Fisheries
Attn: Nancy Fish
P.O. Box 769
Morehead City, NC 28557

Dear Ms. Fish,

The Coastal Fisheries Reform Group (CFRG) is a coalition of recreational coastal fishermen, who support sound management of our marine fisheries based upon the best available science. We represent many thousands of fishermen from across the state who fish in our coastal waters. We have had 122,533 visits to our blog site (<http://cfrgnc.blogspot.com/>) where we have discussed coastal fisheries issues since 2009. In the role as a voice for the average salt water fisherman, we submit the following comments on the proposed Shrimp FMP revision that the Marine Fisheries Commission will consider at their November meeting.

First and foremost, the by-catch of juvenile finfish taken in otter trawls used to harvest shrimp in North Carolina inshore waters is excessive and potentially harmful to the life cycles of some of the important finfish that are being killed and discarded. Recent studies by DMF personnel revealed alarming numbers of juvenile finfish in the by-catch of otter trawls with estimates in the range of a combined 300,000,000 young spot, croaker, and weakfish taken each year. In our opinion, this by-catch is excessive given the relative small annual harvest of shrimp (6 million pounds valued at about \$11,000,000.) Economic studies have shown the value of recreational fisheries to be many times greater than the commercial harvest. In the shrimp trawl fishery, you have the harvest of one commercial species (shrimp) cutting into the productivity of three species of fin fish (spot, croaker, and weakfish) whose recreational value is much greater than the product taken. This equation cannot be balanced in any way that you try to solve it.

While many options exist, some of which are described in the current Draft Shrimp FMP revision, to reduce the trawling by-catch without compromising the annual harvest of shrimp in North Carolina, the proposal recommends that none of them be implemented as management measures in the next five years. Instead, the recommendation is to adopt a status quo position with regard to by-catch. This "No Action" is unacceptable given the severity of the problem and the overwhelming evidence that otter trawling for shrimp is inherently destructive to the habitat and to a major portion of the finfish resource. We think the preferred management action should be to move trawls out of inshore waters altogether and as soon as possible due to the damage they do to the bottom, the sedimentation they cause, and the destruction of way too many juvenile finfish. Moving trawlers three miles offshore has been the solution in most Atlantic and Gulf States.

(continued)

The Shrimp FMP should be amended (not revised) to include goals, timetables, and management measures to accomplish significant by-catch reduction within the next five year period. We quote here from the proposed revision:


"As perhaps the prime example of the new policy positions, the re-authorized Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA) contains a National Standard (#9) requiring bycatch minimization (USDOC 1996). National Standard 9 states: "Conservation and management measures shall, to the extent practicable, (A) minimize bycatch and (B) to the extent bycatch cannot be avoided, minimize the mortality of such bycatch." Additionally, in 1991 the MFC adopted a policy directing the DMF to establish the goal of reducing bycatch losses to the absolute minimum and to consciously incorporate that goal into all of its, management considerations (Murray et al. 1991)."

It is time to follow the national and state policies pointing to the importance of reducing by-catch. We can start with an amendment to the Shrimp FMP that does this in an effective way.

One additional factor that needs to be incorporated into the Shrimp FMP for the next five years is the continuation of the requirement that Turtle Excluding Devices be used in shrimp trawls. This requirement should be expanded to all types of trawls operating in waters where endangered and threatened sea turtles are found. In areas where, and times when, sea turtles are especially abundant, tow times for shrimp trawling should be reduced to a period that will prevent mortality of any sea turtle that may be engaged.

Thank you for the opportunity to comment on the Shrimp FMP. Please consider and share our points as you decide how to proceed in the important process of reviewing the Plan.

Sincerely,



Joe Alpea
Coastal Fisheries Reform Group



*Bycatch photograph
taken week of
September 17, 2012
just outside New
River Inlet, NC.*



North Carolina
Wildlife Federation
Affiliated with the National Wildlife
Federation

2155 McClintock Rd.
St.
Charlotte, NC 28205

1024 Washington
St.
Raleigh, NC 27605

September 26, 2012

Louis Daniels, PhD
Director Division of Marine Fisheries
P.O. Box 769
Morehead City, NC 28557

Dear Dr. Daniel:

The North Carolina Wildlife Federation is a statewide nonprofit conservation organization created in 1945 and dedicated to the professional management of our fish and wildlife resources based upon the best available science. We represent many thousands of fishermen from across the state who fish in our coastal waters. In that role, we submit the following comments on the proposed Shrimp FMP revision that the Marine Fisheries Commission will consider at their November meeting.

First and foremost, the by-catch of juvenile finfish taken in otter trawls used to harvest shrimp in about one half of North Carolina inshore waters is excessive and potentially harmful to the life cycles of the important finfish taken and discarded as by-catch. Recent studies by DMF personnel reveal alarming numbers of juvenile finfish in the by-catch of otter trawls with estimates in the range of 300,000,000 young spot, croaker, and weakfish taken each year. In our opinion, this by-catch is excessive given the relative small harvest of shrimp (6 million pounds annually valued at about \$11,000,000.)

Many options exist to reduce the trawling by-catch without compromising the annual harvest of shrimp in North Carolina. Such management measures as time and area closures and restrictions, water depth restrictions for otter trawls, alternate gear and gear modification, limits on the size of trawls, and other measures, some of which are included in the proposal. We think the ultimate goal should be to move trawls out of inshore waters altogether due to the damage they do to the bottom, the sedimentation they cause, and the by-catch and discard of way too many juvenile finfish. Moving trawlers three miles offshore has been the solution in most Atlantic and Gulf States.

The Shrimp FMP should be amended (not revised) to include goals, timetables, and management measures to accomplish significant by-catch reduction within the next five year period. We quote here from the proposed revision:

"As perhaps the prime example of the new policy positions, the re-authorized Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA) contains a National Standard (#9) requiring bycatch minimization (USDOC 1996). National Standard 9 states: "Conservation and management measures shall, to the extent practicable, (A) minimize bycatch and (B) to the extent bycatch cannot be avoided, minimize the mortality of such bycatch." Additionally, in 1991 the MFC adopted a policy directing the DMF to establish the goal of reducing bycatch losses to the absolute minimum and to consciously incorporate that goal into all of its, management considerations (Murray et al. 1991)."

It is time to follow the national and state policies pointing to the importance of reducing by-catch. We can start with an amendment to the Shrimp FMP that does this in an effective way.

We have no criticism of the remainder of the proposal developed by the DMF shrimp management team and presented for public comment. All three species of shrimp are being managed well based upon surveys and science and are in the "viable" category. Environmental and habitat protections that establish nursery areas and maintain water quality are paramount to the well being of shrimp.

One additional factor needs to be incorporated into the Shrimp FMP for the next five years, which is continuation of the requirement that Turtle Excluding Devices be used in shrimp trawls. This requirement should be expanded to all types of trawls operating at times and in waters where endangered and threatened sea turtles are found. In areas where and at times when sea turtles are especially abundant, tow times for shrimp trawling should be reduced to a period that will prevent mortality of sea turtle that may be engaged.

Thank you for the opportunity to comment on the Shrimp FMP. Please consider and share our points as you decide how to proceed in the important process of reviewing the Plan.

Sincerely,



Tim Gestwicki
Chief Executive Officer
North Carolina Wildlife Federation

From: Byrd, Johnny [<mailto:JByrd@precisionwalls.com>]
Sent: Monday, August 27, 2012 8:59 AM
To: bizzyb@coastalnet.com; Fish, Nancy
Subject: upcoming NCMFC discussions about shrimp harvest

Gentlemen and ladies,

It is my understanding that MFC is currently reviewing inshore (sounds and ICW) shrimp trawling.

Due to work and family constraints, I will not be able to attend the meetings but I deserve for my thoughts to be part of the overall picture.

Some of my earliest memories are of North Carolina saltwater fishing with my family on surf, pier and boat. I have personally experienced the wonderful saltwater fishing that we have had in the past and the heartbreak it has become in the past couple of decades. I am not a scientist but I have read reports from scientists about our waters and waters in other places and they ALL have extremely similar results, TREMENDOUS bycatch for otter trawls. I actually had the opportunity to go out on an otter trawler on three different occasions back in the mid 70's and my personal experience all three times were heartbreaking to say the least. Thousands, probably hundreds of thousands of baby spot, croaker and quarter sized flounder dumped dead back over the side for on those trips about 6 or 7 baskets of shrimp. This was in the New River on the north end of Topsail.

Years of this activity has GOT to be taking a tremendous toll on the health of these and other fish in our waters. We MUST take steps to curtail or even prohibit inshore otter trawling before we pass the point of no return for species much like river herring and weakfish may already be.

PLEASE PLEASE, I and others like me are begging for resource based management so that my 4 granddaughters and my future generations can at least partially experience the same pleasures I did as a child.

Thank you and may the good Lord guide you on this matter.

Johnny R. Byrd
Native North Carolinian

From: Eddie Eatmon [<mailto:beeatmon@gmail.com>]

Sent: Monday, August 27, 2012 3:52 PM

To: bizzyb@coastalnet.com; Fish, Nancy

Subject: Shrimp Trawl Meeting Comments

I am a commercial fisherman and a recreational fisherman. I am in full support of banning wasteful fishing practices and shrimp trawling is the most wasteful of them all. I have been on shrimp trawl boats and it seems like we are killing 10-20lbs of small fish and crabs for every pound of shrimp we get. We had one tow that I bet we threw back 500-1000 dead flounder from 3in-8in long. It's unbelievable that NC still allows this. I'm almost positive that we are one of the last states to allow this fishing practice inshore. What's crazy is that someone can get a \$150 fine for a 14.75in flounder that they plan to eat while a commercial shrimper might kill 10,000 flounder a night that will be pushed overboard dead. There are far more environmentally friendly ways to catch shrimp without these trawls. I've seen shrimp channel nets work in topsail beach that catch 99.9% shrimp with the few fish that it does catch can be thrown back alive. Or why don't we push shrimp farming to so that our natural shrimp can help feed the depeted fish populations in NC? Dragging chains and nets along the bottom and destorying our delicate estuaries in the process is far to wasteful and destructful to be legal.

-Bryce Eatmon

From: Tom Canady [<mailto:tomcanady@bellsouth.net>]

Sent: Tuesday, August 28, 2012 7:25 AM

To: Fish, Nancy; bizzyb@coastalnet.com

Cc: beeatmon@gmail.com; 'RCANADY'

Subject: North Carolina Fishery Resources

I am a concerned citizen and tax payer in Pender,Hyde,Bladen and New Hanover counties. I have spent my entire life on the coast in these communitites. I have never seen the Jolly Roger

Pier never produce a King Mackerel by the end of eight months of a calendar year! What is up with that? Beach renurishment? Commercial netting. I bought a season pass; I have only fished once due to the lack of fish!!! Trawling is destroying the fishery we have in the Pamlico Sound. I have seen it go to pot in the last several years. We have got to do something to back this off to give our fisheries a chance to survive. If we miss this opportunity, we want have to worry about the commercial fisherman losing their income. We will have to worry about how to sustain our general population that the esturine resource sustains in some form or fashion. I am sure my comments from this email will be to no avail. But I feel I have to make the effort. I am also sure the people I am sending it to are more aware of the issues than I and have the facts to back it up. But until politics and political greed do not make the decisions for our best interests, it will probably not change.

Thank you,

Tom Canady
Barker & Canady Custom Homes, Inc.
President
910-612-4442 (mobile)
910-509-2014 (office)
www.barkercanady.com
[Follow us on FACEBOOK!](#)



From: Brown, Charles (HNP) [<mailto:CharlesL.Brown@pgnmail.com>]
Sent: Tuesday, September 11, 2012 10:00 AM
To: Fish, Nancy
Subject: Shrimp Trawling Public Meeting

I regret that I will not be able to attend the meeting concerning the hearing in the Shrimp Trawling issue. This allowance of shrimp trawling has been destroying the fishing in North Carolina for years and I think it is time that the fishermen have their rights to eliminate this terrible practice. Not only should shrimp trawling be disallowed but all type of nets for fish should also be disallowed in the state of North Carolina. I would recommend that the NCDMF look at the Virginia fishing areas and see how the numbers of fish have recovered since they have eliminated shrimp trawling and fish nets. As many of us have had to look for a different job or profession, it is time for the shrimp trawlers to do so likewise. Please case my vote to completely eliminate all shrimp trawling and types of fish netting in the state of North Carolina.

Thanks, Charles Brown, NCW lifetime license #16292

Charles L. Brown
(w) 919-362-2184, (c) 919-812-5310
email: charles.brown2@duke-energy.com



From: Humphries, Scott M (GE Power & Water) [<mailto:scott.m.humphries@ge.com>]
Sent: Tuesday, September 11, 2012 10:10 AM
To: Fish, Nancy
Subject: Shrimp Trawling Mtg

You have my vote to discontinue using shrimp trawling or any netting in the inland waters. I've been a recreational fisherman for over 40 years and have witnessed the continuing decline of inshore fishing. Thank you

Scott

Scott Humphries
Manufacturing Project Manager
GE Hitachi Nuclear Energy

T 910-819-1011
M 910-616-7073
F 910-341-2618
D *292-1011

3901 Castle Hayne Rd
PO Box 780, M/C B-11
Wilmington, N.C. 28401

From: w kerner [<mailto:kernerw@suddenlink.net>]
Sent: Tuesday, September 11, 2012 4:18 PM
To: Fish, Nancy
Subject: comments on internal waters trawling

Good Afternoon,
Regarding trawling in internal waters we have a brief comment. We go out several times a year on Pamlico Sound and almost every time we have gone in the last couple of years (recreational fishing) we have caught **nothing!** We have very experienced fishermen in our little group of friends and we even used chum, etc... However, we regularly see the trawlers working in the area. Now, anyone with any common sense can figure out that if there are no fish in such an area- where the environment should be holding tons of fish; then it can mean only one thing. The commercial fisherman/ trawlers are catching everything in their path. We don't want to deny those people a living but we all have to adapt to changing industries and environments. Certainly the scientists can figure out a way that recreational fishermen (a.k.a. "taxpayers" too) can coexist and be happy. But right now it seems the trawlers are destroying recreational fishing in the Pamlico Sound area.

Thank you.
Walt Kerner

From: Bert Owens [<mailto:ocean@embarqmail.com>]
Sent: Tuesday, September 11, 2012 4:03 PM
To: Fish, Nancy
Subject: inshore trawling

Nancy: Like many, I have reviewed the finfish by catch associated with Shrimp trawling in our inshore waters. The numbers are alarming and to assume they are of no effect to these species would be a mistake and poor stewardship of our resources. Fortunately, all other states to our South all the way to Mexico also have trawling. It would be prudent to look at their actions, including allowed gear, and learn what we could do here. A table of allowed gear, amount of gear and seasons etc. by state prepared by the DMF and distributed at the public meetings would be helpful. Whether I can make any of the meetings or not please add my voice to those calling for an **amendment** to the Shrimp FMP. Thank you.

Bert Owens
Crystal Coast **CCA**

From: Bob Fuller [<mailto:rhf.antares@gmail.com>]
Sent: Tuesday, September 11, 2012 7:27 PM
To: Fish, Nancy
Cc: Jim White
Subject: Shrimp trawling hearings

Ms. Fish:

With regard to the shrimp trawling meetings scheduled in the next few weeks, can you advise me what alternatives or options your department is proposing to solve this problem. Thank you very much for your courtesy and consideration.

Capain Bob Fuller

From: Mark Cable [<mailto:macable@uncg.edu>]
Sent: Tuesday, September 11, 2012 1:58 PM
To: Fish, Nancy
Subject: Ban Shrimp Trawling in Internal Waters in North Carolina !!!

Recent studies by the North Carolina Division of Marine Fisheries (NCDMF) showed that shrimp trawlers kill an estimated 500 million (half a billion!) small fish every year. Most of this "bycatch" consists of weakfish, croaker, and spot, and these finger-sized fish are destroyed before they reach breeding age, guaranteeing that the fish populations in our sounds cannot recover. I would ask that North Carolina follow the example of every other state on the east coast and "**Ban Shrimp Trawling in Internal Waters**" and help to keep our fish stocks at sustainable levels.

--

Mark Cable
Facilities Operations
University of North Carolina at Greensboro

From: HreStore1@aol.com [<mailto:HreStore1@aol.com>]
Sent: Tuesday, September 11, 2012 1:21 PM
To: Fish, Nancy
Subject: Shrimp

Dear Ms. Fish:

I agree with the following and therefore request an end to this process:

North Carolina is the last state on the Atlantic Coast to allow commercial shrimp trawling in internal waters. As other states banned this terribly destructive practice, their trawlers came to our sounds.

Recent studies by the North Carolina Division of Marine Fisheries (NCDMF) showed that shrimp trawlers kill an estimated 500 million (half a billion!) small fish every year. Most of this "by catch" consists of weakfish, croaker, and spot, and these finger-sized fish are destroyed before they reach breeding age, guaranteeing that the fish populations in our sounds cannot recover.

Jay Shoffner

From: Scott Gould [<mailto:sgould@capefearperio.com>]
Sent: Tuesday, September 11, 2012 12:11 PM
To: Fish, Nancy
Subject: shrimp trawling

To whom it may concern,

I want to start by saying that I'm not against commercial fishing. But, I've hoped for a long time that this destructive practice might come to an end in north Carolina inland waters. Before I moved to the coast and actually saw the effects of inshore trawling, this issue had little effect on me. Then, I started spending a lot of time on the water in July and August in the Pamlico sound. I would repeatedly see 25 – 50 boats scavenging the sound floor, running back and forth across the sound. We would ride for miles and see thousands of dead fish (spot, gray and speckled trout, redfish, and croakers) floating on the surface that were discarded as by-catch, not to mention the seabed destruction which is supposed to naturally clean our waters (oysters, grass, etc). I thought to myself, there has to be a better way. I'm not against commercial fishing at all, but this type of practice is destructive and results in wanton waste of resources that are supposed to be shared by all North Carolinians. I would like to voice my opposition to allowing continued inshore trawling of any kind in North Carolina, but I am not against trawling in the ocean waters.

Can we really ignore that all of the other Atlantic States have already agreed to ban shrimp trawling inshore? Sometimes, it's embarrassing to be from NC with the way that we treat our coastal resources. Please help make this long overdue change effective. I appreciate your consideration. Sincerely, Scott Gould

M. Scott Gould
7010 Market Street
Wilmington, NC 28411

From: ncparrott@gmail.com [<mailto:ncparrott@gmail.com>]

Sent: Tuesday, September 11, 2012 10:53 AM

To: Fish, Nancy

Cc: Bert

Subject: Shrimp trawling

Nancy----i believe that inshore shrimp trawling is having a negative impact on several finfish species. I would suggest that this practice be eliminated if possible ---bycatch figures from your own scientists have shown how. Destructive this method of shrimp trawling really is..... Best regards, hubert parrott Sent from my U.S. Cellular BlackBerry® smartphone

From: Neil Smith [<mailto:neilmlynn@skybest.com>]
Sent: Wednesday, September 12, 2012 8:05 AM
To: Fish, Nancy
Subject: inshore trawling

Dear Ms. Fish

It is not hard to see that inshore trawling should be banned in N.C.'s waters. We are the only state on the eastern seaboard who still allows this destructive practice, and scientific studies show that this practice and gill-netting contribute to a fish mortality rate in young fish that prevents our fish stocks from ever recovering to a sustainable point. Recreational fishing certainly brings more revenue to NC than all the commercial fishing put together, considering that many of the commercial fishermen are not even native North Carolinians, and are sending their money out of state. This is an issue that is a hot topic on any NC fishing pier, and it has been recognized for years that something needs to be done. Please give this matter your full attention. BAN GILL NETTING IN NC.
THANKYOU

Neil M. Smith
Scottville, NC

From: Ann Wisz [<mailto:kawisz@bellsouth.net>]
Sent: Wednesday, September 12, 2012 1:58 PM
To: Fish, Nancy
Subject: Commercial Shrimp Trawling

Please consider our comments at the upcoming public hearings regarding commercial shrimp trawling, as we are unable to attend. We must not lose this opportunity to stop commercial shrimp trawling. It is destroying many small fish every year, including spot, croaker and others, before they can reproduce and increase their numbers. These fish are vital to the anglers and tourism numbers in NC. Let us join the ranks of states on the Atlantic Coast which have stopped this practice which is so harmful to our fisheries.

Thank you very much.

Ken & Ann Wisz
1645 Kings Landing Road
Hampstead, NC 28443

From: Harry Archer III [<mailto:harcher3@gmail.com>]
Sent: Wednesday, September 12, 2012 2:13 PM
To: Fish, Nancy
Subject: Shrimp Trawling

Ms. Fish,

I cannot attend the public hearing on 9/19 in Wilmington. I would like to state that I am opposed to any and all trawling for any species in all internal waters, including the Pamlico and Albemarle Sounds. As someone who used to work in the shrimping business in Morehead City, I have seen the destruction trawlers reap on juvenile fish.

I support the banning of all netting, including trawls and gill nets, in our internal waters.

When you travel to other states like Fla, Ga, Texas, LA and SC (to spend money to fish), their sounds and rivers are cleaner and the fishing is light years better than we have in NC. It is very sad for NC natives who love our coastal sound waters.

Thanks, Harry Archer

Harry Archer III
910-791-7880 Office
910-470-7974 Cell



From: Gardner Reynolds [<mailto:info@bartoncreekcreative.com>]
Sent: Wednesday, September 12, 2012 1:54 PM
To: Fish, Nancy
Subject: shrimp and southern flounder management plans

Nancy,

I highly oppose shrimpers killing the fish that they do and would vote to have major restrictions on shrimping and get the guys into fish guiding like other states like Texas and Louisiana. They make more money and it makes sense for our fishery.

I can't even catch a nice flounder anymore.

BORN AND RAISED in NC.

Gardner Reynolds
Owner & Marketing Director
Barton Creek Creative, LLC

919.844-9492 Raleigh Office
919.749.3177 Mobile

info@bartoncreekcreative.com
www.bartoncreekcreative.com

From: Howie Hink [<mailto:howiehink@gmail.com>]

Sent: Wednesday, September 12, 2012 9:24 PM

To: Fish, Nancy

Subject: Shrimp Trawling

Thank you for taking time to look at my opinion. I own a trawl and one year, I used it. Got very disgusted with the by catch and the grass that was torn up from the bottom. Got very few shrimp. Please move the trawls out into the ocean where they will catch bigger shrimp and not baby flounders, trout, croakers etc. Thank you.

Howard Hink

PS Threw the doors away and kept the net for decoration. Never again.

From: Sheryl and Walton Joyner [<mailto:wjoyner262@aol.com>]

Sent: Wednesday, September 12, 2012 6:05 PM

To: Fish, Nancy

Subject: Inshore shrimp trawling

I will not be able to attend the upcoming hearing, but I wanted to at least have a say in the debate. I have lived and fished in NC all my life. I have seen the devastation that inshore "otter" trawls create first hand as a fisherman and a past recreational shrimper. It appalled me 40 yrs. ago and it stuns me that in this day and age that it continues. Bottom trawling is destructive and efficiently destroys nearly everything in its path. It rips up grassbeds, dislodges shellfish and crabs, and kills juvenile finfish by the 1000's at a time. Inshore botom trawling is no different than clear cutting a forest over and over again....except that it is nearly invisible to those not around or "in the know". I don't want the shrimping industry in NC to be shut down by any means, but I think that the time is long overdue for elimination of that part the industry that has been destroying statewide resources for the benefit of only a few.

Walton Joyner
Raleigh

From: Richard J Sessoms [<mailto:sessomsr@centurylink.net>]

Sent: Wednesday, September 12, 2012 5:42 PM

To: Fish, Nancy

Subject:

I rode in a shrimp boat once.after about about three pulls I asked to be taken back to the dock.The "bycatch" of immature fish was more than I could Watch. There we many species of fish that were just raked overboard. None of these fish were living. We need to take the shrimp trawls out of the inland waterway and other nursery areas and let the fish population have a chance to replenish itself. Everyone who goes to the coast fishing is aware of this situation and would like to see it stopped. we have too many politicians who depend on the commercial fishermans vote making the decisions about what is happenoing in our inland waters.Our neighbors to the north and south have have made the necessary changes to their trawling laws. It's time we did likewise.

Richard J. Sessoms

Magnolia NC 28458

Cell # 910 284 1900

From: lwf0831@suddenlink.net [<mailto:lwf0831@suddenlink.net>]

Sent: Wednesday, September 12, 2012 4:57 PM

To: Fish, Nancy

Subject: ShrimpTrawls

I grew up in Hyde county and have seen first hand what shrimp trawls do to our fisheries. We should not sacrifice flounder ,trout, and other species of fin fish for the sake of shrimp harvest in our inland waters. I recommend that we abolish shrimp trawling in inland waters
Sinderely, Lonnie Foreman

From: Woodard [<mailto:woodard@esn.net>]
Sent: Thursday, September 13, 2012 2:11 PM
To: Fish, Nancy
Subject: trawls

I am very much against fish trawls in inland waters (rivers & sounds} I saw this first hand today in white oak river. Shrimp trawls were trawling river channel 24 hrs straight mudding up water and wiping out every thing as they went. This waste has got to stop Woodard Jackson 919 658 5565 woodard@esn.net former advisory member

From: Sandlin Gaudette [<mailto:sandlingaudette@yahoo.com>]

Sent: Thursday, September 13, 2012 11:02 AM

To: Fish, Nancy

Subject: NC Shrimp Trawling

Ms. Fish,

Please help stop the shrimp trawling in North Carolina. I love shrimp but not at the expense of our environment. We have to change our "backwards" ways in this state and this is one way to do just that.

My family and I have lived in North Carolina for many many generations and we want to leave our children with the rich resources this state has to offer for many more generations to come.

Kindest Regards,

Sandlin A. Hunter

4516 Touchstone Forest Rd.

Raleigh, NC 27612

919.427.0191

From: E T Weaver [<mailto:deereman@skybest.com>]

Sent: Thursday, September 13, 2012 9:31 AM

To: Fish, Nancy

Subject: Shrimp Trawling Comment

Ms. Fish,

N.C. Marine Fisheries needs to put a stop to shrimp trawling in internal waters until (if ever) a way is found to reduce destruction of other species (bycatch). The economic value of our marine resources for recreational use must be acknowledged. Inside waters are the key to North Carolina's reputation for "the place to go". We need to show everyone that NC intends to protect and make best use of our resources.

Thank you, E.T. Weaver Jefferson, NC

From: groberson@ec.rr.com [<mailto:groberson@ec.rr.com>]

Sent: Thursday, September 13, 2012 7:53 AM

To: Fish, Nancy

Subject: shrimp trawling

Please use the power of your office to stop this trawling in internal waters; even the NCDMF says this is harmful. I am a 67yr old rec fisherman around Morehead City and know some of the small time shrimpers that will be hurt in the pocketbook if not put out of business but we need to protect the environment. Thanks in advance for any assistance, Gary Roberson.

From: Vernon Hunter [<mailto:vhunter@copycei.com>]

Sent: Thursday, September 13, 2012 4:13 PM

To: Fish, Nancy

Subject: Shrimp Trawling

Dear Ms Fish,

I own property in Beaufort and Hyde County and grew up spending summers as a child at my parent's cottage at Emerald Isle. I love shrimp. But the price we are paying for inshore trawling is just too much. I have commercial fished for a living in Alaska and NC have seen clean fisheries. This is not one of them.

This fishery is destroying our inshore fishery in NC. The millions of pounds of bycatch including immature croaker, grey trout, speckled trout, red drum, turtles, black drum, flounder and untold baitfish (menhaden, pinfish, spot, etc) is not worth any price for shrimp.

This fishery also destroys the filter feeders that clean the water - oysters, hard bottom areas, crabs, and the ecosystem on the sound bottom. It is a matter of time before this fishery is banished inshore. If the public really had any idea of the wanton waste of this fishery the outcry

would be deafening. I would really like my daughter to see a glimmer of what the fishery in the Pamlico Sound used to be, can be. Please help this happen.

Sincerely,
-R Vernon Hunter
1925 Sunset Drive
Raleigh, NC 27608
AND
5510 Sidney Road
Belhaven, NC 27810

From: dbuck16@embarqmail.com [<mailto:dbuck16@embarqmail.com>]

Sent: Saturday, September 15, 2012 7:55 AM

To: Fish, Nancy

Subject: Shrimping laws

I am 64 and have fished our coastal waters extensively. I have been on shallow water shrimp trawlers and the by catch is sickening! This is an obvious destrucktive practice. I can't belive it has been allowed to continue. Thank you.

Sent from my U.S. Cellular BlackBerry® smartphone

From: Wille Zee [<mailto:ddcarver123@yahoo.com>]

Sent: Sunday, September 16, 2012 7:05 AM

To: Fish, Nancy

Subject: trawlers

we the people of nc would love to see the trawlers move out and leave the beaches clean.We need to educate them they need to learn fish farming.If we dont do this now we will end up like other places that have dead zoons (no Fish).I,ve seen them dump out dead fish for miles just to have a few shrimp.It makes me sick.There is only a few people still doing it also trawlers come from other places just because we are so relaxed about laws concerning them.WE NEED TO STOP ESP> THE STOP NETS AND LIMIT OTHER NETS ALSO!!!!!!protect and perserve our fish and coast.

From: Billy Reavis [<mailto:bjreavis@gmail.com>]

Sent: Sunday, September 16, 2012 9:27 PM

To: Fish, Nancy

Subject: shrimp trawling

I've been fishing in the area of core sound for to many years to tell. But you can be assured that I have seen the effects of shrimp trawling. This practice should have been stopped many years ago it will take to recover'. there is years so much by kill. Also if nothing is done about the practice of long haul, which is killing many juvenile fish. The sound should be considered a nursery area.

From: bensdaddy@suddenlink.net [<mailto:bensdaddy@suddenlink.net>]

Sent: Monday, September 17, 2012 8:00 PM

To: Fish, Nancy

Subject: Public Comment Sought on Shrimp Trawling and Bycatch

Dear Ms. Fish,

I am pleased to see that North Carolina Marine Fisheries Commission is looking into the damage that shrimp trawling does to fish species when done in internal waters. I am opposed to this practice by commercial fishermen and hope that the NCMFC can come up with a plan to eliminate it.

I grew up in Southwest Florida and experienced first hand what damage unrestrained netting in internal waters can do to internal waters. I can remember the days when it was considered a waste of time to fish in Tampa Bay or Sarasota Bay back in the 70's. Now that I live in North Carolina, many people now tell me that it is a waste of my time to fish in Bogue Sound and that I would be much better off planning fishing trips to Virginia.

When Florida eliminated the commercial netting in its internal waters, the recreational fishing is outstanding. I hope that the NCMFC will see that shrimp trawling in these areas is devastating the fish populations and will restrict or eliminate this practice within a mile of shore.

Again, I thank the NCMFC for looking into this and hope they have the courage and the sense of duty to our state to make the right choice.

Sincerely,

Brack Craven

Greenville, NC

From: BC [<mailto:bcraver4@triad.rr.com>]
Sent: Monday, September 17, 2012 11:28 AM
To: Fish, Nancy
Cc: bC
Subject: ?

Nancy,

I wish that trawlers stripping our coast line of fish would be forced to further out off our coast line. I do not live there but have visited our coast a few times a year for years to fish . I find fewer fish every year . Would your please try to find more balance between tourism and commercial fishers so we could enjoy a few self caught fish for ourselves. Thanks Bud C.

From: Bruce [<mailto:bdrmjm@charter.net>]
Sent: Tuesday, September 18, 2012 10:48 AM
To: Fish, Nancy
Subject: Shrimp Trawling

Dear Ms. Fish:

I am writing to you to make a comment on the current Shrimp Fishery in North Carolina. While I believe that shrimping is a significant part of our North Carolina Commercial Fishery, the current method of the use of Otter Trawls is one of the most devastating to our environment and to the current and future health of our overall fisheries. Each year hundreds of millions of fingerling sized fish are wiped out and the trawls continue to destroy the bottom structure. Submerged aquatic vegetation never has a chance to recover as the bottom is continuously scraped. There are other methods of shrimping that allow netting without scraping the bottom where many of the fingerling fish reside and our aquatic vegetation grows. I would recommend we eliminate Otter Trawls completely and set up areas of our sound where shrimping is not allowed until the vegetation is healthy again. Please take the necessary action to protect the future of the precious fisheries resource in North Carolina.

Sincerely
Bruce D. Matthews
Manteo, NC

Sent: Tuesday, September 18, 2012 5:36 PM
To: Fish, Nancy
Subject: Against shrimp net trawling

Dear Ms Fish,

I am a resident of North Carolina (Troy, N.C.) and a recreational saltwater fisherman. Please vote to stop the destructive practice of shrimp net trawling in our NC waters. The juvenile fish by-catch that is wasted by the shrimp trawlers costs the State far more economic dollars than the few pounds of shrimp that are caught and sold. Please help our State and vote "No" to shrimp net trawling.

Sincerely,

Richard Capel (Troy, NC)

From: Mark Cable [<mailto:macable@uncg.edu>]
Sent: Wednesday, September 19, 2012 11:04 AM
To: Fish, Nancy
Subject: Reopen and Amend the "Shrimp Management Plan" !!!

In the latest studies by the DMF, the by catch ratio was typical (approximately 4:1) and indicates that an estimated 24 million pounds of finfish by catch has occurred annually for the last five years. The study, which is considered valid and solid by DMF scientists, also determined that the finfish killed in the trawlers were small, averaging around 20 fish per pound. Multiplying 20 fish per pound by 24 million pounds allows us to get a rough estimate of 480 million juvenile finfish that are killed annually by trawlers operating in North Carolina waters.

I am asking the MFC to reopen and **amend** the Shrimp FMP to better address the issue of by catch by shrimp trawlers operating in our in shore waters.

--

Mark Cable
Facilities Operations
University of North Carolina at Greensboro

From: Eb Pesci [<mailto:ebpesci@gmail.com>]
Sent: Wednesday, September 19, 2012 9:23 PM
To: Fish, Nancy
Subject: Amendment of the shrimp FMP

Dear Ms. Fish, Please forward this message to the members of the Marine Fisheries Commission. Thank you.

Dear Marine Fisheries Commission Members,

I am writing this letter to ask the MFC to reopen and amend the shrimp FMP. Studies by the NCDMF have clearly shown that shrimp trawl bycatch has averaged approximately 25 million pounds per year for the last five years. This equates to hundreds of millions of small finfish that are wasted before ever reaching maturity. The NCDMF studies have also shown that most of this bycatch consists of croaker and spot. In addition, there is also a large number of weakfish, a depleted species, included in this waste. It is time for this waste to stop. I am asking for an amendment to the shrimp FMP and I'm also asking for shrimp trawling inside of the ocean to be ceased immediately. Most other states and many countries have stopped or severely restricted inshore trawling but we continue to allow our fish nurseries to be raided. Others have reaped a huge bounty by removing trawlers (an 8 fold increase in biomass was seen in an Italian trawl ban area) yet we continue to have only the memories of sounds full of plentiful fish. Those days are gone, and there is no doubt that inshore shrimp trawling holds part of the blame.

It's time to start working for the resource and provide our marine environment with an opportunity to rebuild what was once a world class fishery. Please end inshore trawling immediately.

Thank you,

Eb Pesci
Greenville, NC

From: Richard Fulton [<mailto:rfulton1955@embarqmail.com>]
Sent: Wednesday, September 19, 2012 8:41 PM
To: Fish, Nancy
Cc: Tim Hergenrader
Subject: Shrimp FMP

Please reopen and **amend** the Shrimp FMP. The waste here is deplorable. It is time we changed our ways so that future generations can enjoy OUR resources.

Thank you,

Richard C. Fulton
101 Birch Rd
Hubert, NC

From: Russell Long [<mailto:fishingruss@gmail.com>]
Sent: Wednesday, September 19, 2012 3:15 PM
To: Fish, Nancy
Subject: shrimp trawling in our state waters

This wasteful practice has to stop. I went shrimping once and I was appalled by the by catch. This is not a tolerable means of gathering seafood anymore. Things have to change in NC. I would rather buy imported shrimp than local because I know how destructive our practice is.

Russell Long PO Box 1954 Kill Devil Hills, NC 27948

From: Will Brown [mailto:w_h_brown@yahoo.com]
Sent: Wednesday, September 19, 2012 2:43 PM
To: Fish, Nancy
Subject: shrimp trawling and bycatch

Dear Ms. Fish,

I am writing you about the current discussions of bycatch in the shrimp trawling industry. I know that NC is the only state on the east coast to allow trawling in internal waters. I believe that all internal (non-ocean) waters should be closed to trawling because this method of fishing is too devastating to the environment. It destroys countless numbers of juvenile trout, croaker, and spot, and forever changes the contours of the bottom, removing vegetation and flattening ridges. In the 1970's and very early 1980's my family used to catch quality gray trout in the Pamlico sound, but for the last 30 years, the fish have been difficult to find and small. Current recreational regulations are severely restrictive, but even these cannot allow the species to rebound with the current shrimp trawling industry wreaking havoc on this species' population. Shrimp trawling in internal waters should be banned! the shrimping industry will still have productive waters to fish, the ocean! Our sounds and estuaries are too important to all of NC's fish species. Thank you,

Will Brown
8813 New Forest Dr
Wilmington NC 28411
910-540-0332

From: Betts Tackle LTD [<mailto:bettstackle@bettstackle.net>]
Sent: Wednesday, September 19, 2012 2:02 PM

To: Fish, Nancy
Subject: Inland Shrimping

Marine Fisheries Commission

- I know you have correct information
- I know you have received this information again and again by hundreds of concerned people.
- You know N.C. is the only state that allows shrimping in the internal waters

I ask you to rise above politics and personal interest. I believe if you really care about the Marine Resources of North Carolina, you will stop this inland practice of shrimping.

Don Betts
Betts Tackle Ltd.
1701 West Academy St.
Fuquay Varina, NC 27526
Phone 919-552-2226
Fax 919-552-3423
bettstackle@bettstackle.net

From: Joe Lail [<mailto:lumberped@charter.net>]
Sent: Wednesday, September 19, 2012 1:22 PM
To: Fish, Nancy
Subject: Inshore trawling.

As a recreational fisherman of 30 years , it has become abundantly clear , that year over year our fish stocks have been depleted. Just a brief glimpse at the figures posted by the NCDENR in it's Marine Recreational Finfish Harvest report shows astronomical declines between the years of 1989 to 2010.

One particular example would be the reported catch of "Spot". The reported catch of this fish in 1999 was down 220% from the total reported in 1989. To go further , one just has to look at the catch reported in 2010 which was down 159% from the total reported in 1999. Total decline of the reported catch over this 20 year period amount to 2,089,739 fish.

While 2 million may seem like a small number , this is just one species that represents a huge cross section of the finfish decline, and these are only recreationally reported numbers. I chose the spot as a good example of a finfish that migrates to inland waters to mature. This is important, as the Spot is a staple in the diet of many other larger inshore fish that will in turn range into pelagic waters and become part of the diet for many pelagic specie. Thus , the decline of the "Spot" can immediately have an impact on the mortality rate of much larger and economically important specie. The "Spot " is only one example of the drastic decline of finfishes. Another example and perhaps ecologically, and economically more significant would be the "Atlantic Menhaden". Menhaden stocks are at a tipping point of un-sustainability up and down the eastern seaboard. This has occurred largely due to commercial overfishing of adult populations. However , the problem can only be compounded exponentially by reducing their numbers when taking them from their nursery grounds as juveniles. This happens predominantly as by-catch , but the "Menhaden" is a very delicate specie that has a very high morbidity rate when handled in any manner. There are countless other specie , even outside of finfish that are impacted by inshore trawling that have an astounding impact on the long term health of our marine eco-system.

By curtailing or altogether banning the trawling of inshore waters we begin to re-stabilize the nursery environment for countless numbers of species , thus ensuring a brighter economic future for all fisheries. While the immediate economic impact for this may be sharp for many , by adapting , we can ensure a more long term and sustainable economy for all of us that earn a living or just plain enjoy fishing our coastal waters. Understanding the plight of our commercial shrimpers in this situation is paramount. Many of our shrimpers are family businesses that have existed for decades off of the sea , and we cannot arbitrarily just throw them to the wind. But, with that being said ; all business environments change , and business models must change to adapt or cease to exist. A business is no different than a specie in the wild , when tampered with or manipulated in any way it will either adapt to it's environment , or decline and perish. The advantage that our shrimpers have , that wild creatures do not , is the ability to think long term and plan for a change that is coming , and that is exactly what will get them thru this potential change.

Economic planning is essential when considering changing business models. By looking at traditionally accepted supply and demand principles, we can help North Carolina shrimpers prepare for the future.

Currently , shrimpers are facing extreme pressure from imported, low priced and inferior shrimp. To make this transition , shrimpers may have to look several business cycles into the future. By removing a portion of North Carolina's shrimp from the market , the supply side will be somewhat shortened , and perhaps raise the prices proportionally.

This could have a two pronged effect:

Foreign suppliers may increase their shipments to take advantage of this , but this could also make the "offshore" taken shrimp more profitable (higher margins) for the local fishermen. This could help give the local fishermen an influx of cash to help them thru the transitional period. A longer term effect could be that foreign shrimp may eventually flood the market when trying to take advantage of the higher margins. When the market becomes flooded , prices will plummet and the margins will decrease

making the market less attractive to importers. Eventually forcing importers out entirely , making them opt to sell their goods in their local market with lower over head costs. As this happens it would give our local fishermen a chance to take advantage of this and fill the gap in supply with profitable shrimp taken from our local "offshore" waters. Optimally , by this time the local fishermen will have figured out alternative means to bridge the gap in seasonal fluctuation of prices by supplying different products to market , without flooding any one item in particular.

This would allow for a profitable business model , and yet preserve our fisheries for the decades to come.

Any way we look at it , our current path is not sustainable , a change must be made and made sooner rather than later. For each season that passes the problem is only compounded and will eventually result in a situation where the shrimper may have no product to catch , making their demise inevitable. In any of life's endeavors there is a price to pay. The choice is clear here ; continue to "play" now and pay the balloon payment at the end , or pay a small price immediately for the ability to continue to "play" indefinitely.

Thank You For Your Time

Joe Lail
Concerned Angler

From: James Coleman [<mailto:colemanj56@hotmail.com>]
Sent: Thursday, September 20, 2012 9:51 AM
To: Fish, Nancy; bill.cook@ncleg.net; marian.mclawhorn@ncleg.net; edith.warren@ncleg.net; The Honorable Clark Jenkins; louis.pate@ncleg.net
Subject:

Date: September 20, 2012
To: Ms. Nancy Fish, NCDMF/NCDENR
From: James Coleman, Greenville, NC colemanj56@hotmail.com
Re: Shrimp Trawling

I am writing to request that policies be enacted as soon as possible to prohibit inshore shrimp trawling activities in the North Carolina inshore waters. The irrefutable facts are that trawling activities in our waters directly lead to the killing massive numbers of vital species of feed- and game-fish at the fingerling stage. Decimation of these populations not only affects fish populations in our waters, but populations along the entire Eastern seaboard, because of the major role the vast estuary systems of North Carolina play as breeding and nursing areas for Atlantic Ocean fish species. Executive and legislative boards in other states controlling similar bodies of water such as Delaware River and Bay and Chesapeake Bay and its tributaries have taken seriously their responsibilities as stewards of our natural resources and have wisely restricted certain practices in these bodies of water. It is time now for North Carolina policy-makers to study the FACTS and ignore the anecdotal "old fisherman" myths and make a forward-thinking decision which can do nothing but help the future of the fishing and shell fishing industries – both commercial and recreational – in North Carolina.

PLEASE – amend the Shrimp FMP now!

Thank you.

From: Maxwell, Galen S Civ USAF ACC 4 FSS/FSMM [<mailto:galen.maxwell@seymourjohnson.af.mil>]
Sent: Thursday, September 20, 2012 1:44 PM
To: Fish, Nancy
Subject: Public Comment Sought on Shrimp Trawling and Bycatch

Ms Fish;

Since I cannot travel to one of the public meetings I would like to offer my input via email:

North Carolina is the last state on the Atlantic Coast to allow extensive commercial shrimp trawling in internal waters. Recent studies by the North Carolina Division of Marine Fisheries (NCDMF) indicate that shrimp trawlers destroy small fish every year and by-catch needs to be reduced. A significant portion of this "by-catch" consists of weakfish, croaker, and spot (recreational pan fish). These finger-sized fish are destroyed before they reach breeding age, and as a recreational salt water fishing advocate, this remains my concern. I would like to see the N.C. Marine Fisheries Commission switch from a revision to an amendment process, so that

the proposal to ban inshore commercial trawling altogether in NC waters can at least be debated.

Thank you.

V/r,
Galen S. Maxwell
Goldsboro NC
Member, CCANC

From: Greg Hefner [<mailto:gahefner@hotmail.com>]
Sent: Thursday, September 20, 2012 2:37 PM
To: Fish, Nancy; Tim Hergenrader
Subject: Excessive by catch of juvenile finfish in otter trawls used to harvest shrimp

Nancy,

I am emailing you to ensure my comments will be added to the public record concerning the by catch of juvenile finfish in otter trawls used to harvest shrimp.

I strongly urge the MFC to **reopen and amend** the Shrimp FMP! I understand that the DMF is only recommending a revision of the current Shrimp FMP. When is this state going to wake up and realize that we are wiping out our natural resources by allowing this to continue? Why has all the other South Eastern states amended or completely stopped in shore netting of all kinds? Why does North Carolina think they are so different?

It is very easy for any citizen who follows these matters to see and understand that our policies are driven by money and greed, not by science and evidence! It makes me shameful to call myself a North Carolinian when these practices continue to contribute to the few instead of the greater good for all, and most importantly our declining natural resources!

I am an avid recreational fisherman that has fished these waters for over 25 years. When I go out for a day of rod and reel fishing and struggle to catch a single fish in our inshore waters, there is a major problem occurring! I am by no means an expert, just your average everyday guy that enjoys time out on the water fishing with friends and family. But, it gets very frustrating when you have more no catch days than days of catching. Especially when a day on the water usually costs me well over 200 dollars when you add it all up, gas, tackle, bait, and food. Fifteen, even ten years ago this was unheard of. To go out in these waters, this huge **NURSERY** area of ours, and not catch fish means there are huge problems brewing. I could go on and on, but I think I have made my point and I'll leave it at that!

Once again, I strongly urge the MFC to reopen and amend the Shrimp FMP!

Thank you for your time,

Greg Hefner
Retired Marine
Sneads Ferry, NC

Courage is endurance for one moment more...

From: tthompson19 [<mailto:tthompson19@ctc.net>]

Sent: Thursday, September 20, 2012 4:10 PM

To: Fish, Nancy

Subject: Trawlers

Its been said over and over....but ill say it again....The amount of money the recreational fishermen put into, a Wide array of the economy, BY FAR, outweighs what the commercial fishermen do for our economy! Not to mention the commercial fishermen depleting a WIDE array of fish species!. There is NOT anywhere near "good" versus bad that the commercial fishermen are doing! Get the politics out of the equation! Let common sense be the deciding factor! Its for all good reasons all other atlantic coast states have stopped inshore trawlers. Lets get er done before its too late! Please! Skip the gay marriage issues and get to something important to us all! Our Economy! Our future! Our fish! TimmyT

From: Ted Mayer [<mailto:tedmdds@nc.rr.com>]

Sent: Thursday, September 20, 2012 8:29 PM

To: Fish, Nancy

Subject: Public comments against shrimp trawling in internal waters

Please include my comments on the topic of shrimp trawling. North Carolina is the last state on the Atlantic Coast to allow commercial shrimp trawling in internal waters.

Recent studies by the North Carolina Division of Marine Fisheries (NCDMF) showed that shrimp trawlers kill an estimated 500 million (half a billion!) small fish every year. Most of this "by catch" consists of weakfish, croaker, and spot, and these finger-sized fish are destroyed before they reach breeding age, guaranteeing that the fish populations in our sounds cannot recover.

Concerning the issue of excessive by catch of juvenile finfish in otter trawls used to harvest shrimp at the upcoming meeting I request the MFC to reopen and **amend** the Shrimp FMP. The Division of Marine Fisheries (DMF) is recommending that the Shrimp FMP only be revised and this revision is not adequate. If the Shrimp FMP is merely revised and not amended, then we will be stuck with no meaningful change to the use of otter trawls for five more years.

As a recreational fishermen, a citizen of Craven County, a

healthcare professional and a member of the NC Commission for Public Health I have a concern for our marine environment who would like to see our fisheries improve.

Regards,

Theodore F. Mayer
4008 Marina Townes
New Bern, NC 28560

tedmdds@nc.rr.com

From: Bill Gibson [<mailto:wgibsonlg@nc.rr.com>]
Sent: Saturday, September 22, 2012 1:23 PM
To: Fish, Nancy
Subject: Shrimp MFP

To not open the shrimp MFP for discussion/debate/review is irresponsible in light of the abosultely high numbers of small finfish destroyed as bycatch. No matter the fish, it is part of the food chain. Decimate the small end of the food chain and there is less food for larger fish. The larger fish either do not flourish or they move elsewhere. Decimate the small finfish, some of which grow to larger gamefish, and there are fewer gamefish. It is very simple. One does not need to have a Phd to see and understand. MFC is responsible for the health of our fisheries and to watch the decline and do nothing to one significant factor perpetuating the decline is reprehensible.

WH Gibson
NC Native

From: John Rakoci [<mailto:john@rakoci.com>]
Sent: Sunday, September 23, 2012 8:40 AM
To: Fish, Nancy
Subject: Comment

Hard to believe the only state that still permits this is asking for comments. First, I only buy local caught sea food. I used to be a charter boat captain on Lake Erie. The commercial people do toss the bycatch back into the ocean and ICW. No reason as nearly all of it is dead or will die. If commercial fisherman are professionals they should be offshore a minimum of 3 miles. This is important not only for the long term survival of the entire fishery but for safety reasons too. A strong commercial fishery is important to NC and those that make their living fishing and in related industry. What is mostly seen inshore are those with minimal investment in boats and equipment. Last Thursday there were three shrimpers between the Ocean Isle Beach bridge and Sunset Beach bridge.

That area of the ICW is narrow and there are quite a few boaters on the water still.

Commercial gigging- that is a sad practice!

Within 2 miles on that same day, there were 3 gill netters. Another very sad practice for all but the gulls and pelicans that feast on that dead bycatch. I live on the coast. The strength of the commercial lobby is well known. The influence of the coastal legislators is also strong. Expecting change is not very promising. There must be a rec/comm compromise or both will lose. If the animal rights activists and anti-fishing/hunting groups get involved everyone including the fishery loses.

John Rakoci

From: Paul Brown [<mailto:paulbbrown@windstream.net>]

Sent: Tuesday, September 25, 2012 9:02 PM

To: Fish, Nancy

Subject: shrimp amendment

the bycatch is unacceptable we can't keep killing these fish just to catch shrimp. this needs to be fixed now and not put on the back burner any more. the fish can't wait any more for someone to take action.

From: Ron Bennett

Sent: Tuesday, September 25, 2012 2:51 PM

To: nancy.fish@ncdenr.org

Subject: Shrimp FMP

Dear Ms Fish,

I strongly believe we need to amend the current regulations to ban shrimp trawling in our sounds. Killing over 500 Million fish as "by catch" of shrimping is unacceptable and one reason our fisheries are in such poor shape relative to most coastal states that have already banned this type of activity.

Thank you for entering my comments into the Public Record.

"Recent studies by the North Carolina Division of Marine Fisheries have shown that approximately 78 percent (by weight) of what is caught in shrimp trawls is untargeted "by catch" that will die before being shoveled overboard. More shockingly, the annual by catch from NC shrimp trawlers consists of over 500 million individual small fish."

Ron Bennett
117 Pudding Pan Lane
Kitty Hawk, NC 27949
252-261-4863

From: Stephen Martin [<mailto:stephenmartin@suddenlink.net>]

Sent: Tuesday, September 25, 2012 1:15 PM

To: Fish, Nancy
Subject: Shrimp Trawling Bycatch

I am unable to attend any of the public meetings scheduled to address this issue but wish to register my opinion. At the very least the subject should be reopened so that further research and evidence can be brought into play before the issue is buried for several more years. North Carolina has been my home for only 6 years and my time here has confirmed often the decision to move to the state. One thing which consistently disappoints is the blind eye turned toward the steady and unnecessary decimation of fish stocks in our state. Whether it is the wanton waste produced by in-shore gill netting, the high-grading by striped bass netters or the even more dramatic destruction of fish stocks by shrimp trawling in-shore, North Carolina seems always to disappoint when it comes to reasonable protections for marine resources. And it seems that the proud tradition of coastal fishing families in our state is the banner waved to justify the blind eye turned time after time. I moved here from Louisiana where there is an equally strong and proud tradition of commercial fishing stretching back many generations in the same families. The difference is that Louisiana (and nearly every other coastal state from here to Texas) recognizes the obvious: if destructive practices continue the commercial fishermen will eventually catch the last fish and in the meantime will have driven the far more economically significant recreational fishery away. So, rather than quibble about how many pounds of juvenile finfish it is okay to sacrifice for each pound of shrimp taken, why not talk about which of the already available techniques for limiting bycatch are to be mandated for general use. If not, I can guarantee that there are enough angry, frustrated recreational fishermen out there that no politician supporting the status quo will ever get elected again. Thanks for listening.

Steve Martin
1003 Coopers Ct.
Trent Woods, NC 28562

From: John Canup [<mailto:jcanup@suddenlink.net>]
Sent: Tuesday, September 25, 2012 10:01 AM
To: Fish, Nancy
Subject: Trawling in internal waters

I totally support banning trawling in internal waters in North Carolina. I have personally witnessed the mass destruction of small fish (bycatch) generated by shrimp trawling in Core and Pamlico Sounds. Also, yesterday (September 24, 2012) while I was trout fishing near Chapel Creek in Bay River I observed a net boat working a set net. He removed three turtles in the first sixty yards of the net. I had been fishing there for approximately 45 minutes before he arrived and had seen no turtle activity—therefore I assume they were dead. Our sounds and rivers are fish nurseries and we must protect them to sustain our fisheries resource.

From: Mike Hadley [<mailto:mikehadley@geoguys.net>]
Sent: Tuesday, September 25, 2012 8:20 AM
To: Fish, Nancy
Subject: Trawling Nets

Nancy,
I am not able to attend my area meeting in New Bern, NC and would like to voice my experience/opinion about shrimp trawling nets inside coastal waters. Years ago with my ex-father in law, I helped him on his small boat pulling a shrimp net in the intercoastal waterway between Sneads Ferry & Surf City. I witness the destruction of juvenile

fish that in numbers and weight accounted for the majority of the catch. At the time I thought nothing of it since I was young and we were part of what I thought was a very small number of boats pulling shrimp nets. Then I witnessed opening day one year in the New River and I saw at least 100 boats pulling shrimp nets in the river. It then hit me! If we are destroying small fish by the hundreds, then how many juvenile fish are going to be killed in the next few hours? 3,000, 4000, or 5000 lbs.? That was when I quit helping on the shrimp boat and started asking my then father in law to quit. He told me that the river and waterways would never run out of fish. That was the mindset then and I do believe that some people still think that way. Those are the people that need to be educated about the declining fish populations that we are experiencing now. For some reason, I have seen more menhaden in the waterways this year than I have seen in the past 5 years. Is this just coincidental or is it because we have new rules covering catching menhaden within 3 miles of the NC coast? I vote to ban all trawl nets in any interior coastal waterway. RIGHT NOW!!!!

Mike Hadley
128 Magens Way
Cedar Point, NC 28584
252-393-6382

From: John Trueblood [<mailto:jrdntrue@gmail.com>]

Sent: Wednesday, September 26, 2012 1:24 PM

To: Fish, Nancy

Subject: shrimp FMP

For the record:

I urge MFC to reopen the shrimp FMP and amend the plan regarding the "by catch" issue.

Thank You.

John R. Trueblood,

Recreational fisherman

Sent: Wednesday, September 26, 2012 10:08 AM

To: Fish, Nancy

Subject: shrimp management plan

i am asking the marine fisheries commission to please amend the shrimp fisheries management plan so me and my children and grandchildren will be able to enjoy and catch fish in pamlico sound like we once did 40 years ago .thank you

From: Lee Dunn [<mailto:lhdunn@comcast.net>]

Sent: Wednesday, September 26, 2012 9:22 AM

To: Fish, Nancy

Subject: Shrimp FMP comments: please meaningfully regulate bycatch

I cannot be present at any of the public hearings on the Shrimp FMP, but I want to enter my sentiments about the plan, and about the great state of North Carolina's fisheries management in general, into the public record:

I am in 100% agreement with the position of the Coastal Fisheries Reform Group, who's memo and position on this issue you are no doubt familiar. The present Shrimp FMP does not seriously address the bycatch issue, and to allow the status quo to continue without critical review is disdainful to NC citizens and blind to the resource. The statistics on bycatch clearly speak for themselves, though this issue alone is far from the only problem with the destructive otter trawl shrimping effort going on in NC's inland waters.

Could it be possible that there is a new day dawning in regard to marine fisheries in NC? Could NC actually move out of the dark ages and towards managing it's marine resources with respect to science, logic, and the fact that it belongs to all of NC's citizens, not just to fish house owners? Is it possible for North Carolina rise above it's current status as the laughing stock of coastal fisheries management in the U.S.?

I sure hope so, and it seems that the tide of awareness and concern is rising. To that end, as an initial step, please do everything in your power to support implement regulations to curtail this destructive fishery.

Thank you,

Lee H. Dunn

Beaufort

From: William Mayo [<mailto:william.mayo@rockymountnc.gov>]

Sent: Wednesday, September 26, 2012 7:03 AM

To: Fish, Nancy

Subject: Restrict/ban coastal trawling

Nancy,

I'm emailing in support of RESTRICTING or BANNING commercial trawling and other netting in NC's coastal waters. While I understand the detrimental effect my position poses to those who fish commercially, I also acknowledge that if the State doesn't make some changes, commercial nor recreational fishermen will have any resources to pursue. It is time to take a bold step to protect our declining natural resources. The many recreational fishermen in the State have been dictated to by the few commercial fishermen and their political allies for too long.

Thank you,

--

Trey Mayo, Fire Chief
City of Rocky Mount
375 East Raleigh Boulevard
Rocky Mount, NC 27801
Office: 252 972 1490
Mobile: 252 343 3167
Facsimile: 252 972 1670
william.mayo@rockymountnc.gov
rockymountfire.org

Raleigh

From: Max [<mailto:mkasselt@suddenlink.net>]
Sent: Sunday, September 30, 2012 9:32 PM
To: Fish, Nancy
Subject: Public Comment Sought on Shrimp Trawling and Bycatch - Outline for a possible solution

Dear Ms Fish,
I passed my letter to you (below) to a few friends - but only received ome reply. I deleted his name as this info could become public.

----- Original Message -----

From: [Max](#)
To:
Sent: Tuesday, September 25, 2012 7:54 PM
Subject: Re: Public Comment Sought on Shrimp Trawling and Bycatch - Outline for a possible solution

Pay no taxes - Yet they rob the nursery.

----- Original Message -----

From:
To: mkasselt@suddenlink.net
Sent: Tuesday, September 25, 2012 9:22 AM
Subject: RE: Public Comment Sought on Shrimp Trawling and Bycatch - Outline for a possible solution

The fishermen never declare profits thus pay no income taxes....if this was put into law you would all of a sudden have huge declared incomes and tax return revisions.

From: Max [<mailto:mkasselt@suddenlink.net>]
Sent: Monday, September 24, 2012 11:00 PM
To: ;
Subject: Public Comment Sought on Shrimp Trawling and Bycatch - Outline for a possible solution

This is what I want to to send. Please make suggestions. Please send in your own comments as well

nancy.fish@ncdenr.gov

Dear Ms Fish,

The damage done to our fish resources by commercial and "recreational" shrimp trawling in internal waters is a known fact.
That we allow it is a tragedy.

My rough outline for a possible solution:

1. Ban all trawling for shrimp in internal waters
2. Have the state buy all the operational boats of fisherman who claim they cannot survive - then resell them - even at a loss.

Sellers of boats, may not buy or have an interest in any sold boat for 5 years.

3. Reimburse all active fishermen (who sold their boat and/or turned in their commercial license) 10 or 20 times their highest declared income (from fishing) in the last 10 years.

The rewards of creating an angler's paradise will more than pay for the above.

Max Kasselt

From: Jim Parsons [<mailto:jparsons5@ec.rr.com>]

Sent: Tuesday, October 02, 2012 10:31 AM

To: Fish, Nancy

Subject: Shellfish/Crustacean Meeting

I am also unable to attend the meeting tonight due to a previous appointment. I concur with the analysis of Dr. Danzler which is reprinted below. I have seen the carnage first hand. I have owned a shrimp trawl and the terrible damage it was inflicting upon the resource is why I no longer own it. The skimmer rigs do a lot of damage, but the trawls have done, and do, immense damage to the finfish resource.

I urge the MFC to amend the Shrimp FMP to eliminate trawls in the inland waters. It is the right thing to do for the finfish resource!

**James C Parsons
210 Sumter Court
Havelock, NC 28532**

From: Lee [<mailto:lee.dantzler@adamscreek.com>]

Sent: Tuesday, October 02, 2012 9:30 AM

To: Fish, Nancy

Cc: Jim Parsons; Tony Michalek

Subject: Unable to attend tonight's Shellfish/Crustacean Meeting New Bern meeting on shrimping by-catch etc, BUT ... written comments provided

Due to prior business plans I am unable to attend the **Shellfish/Crustacean Meeting** at the Craven County Cooperative Extension Office in New Bern. I understand you accept written comments, which I am providing below:

The latest studies by the DMF, the by catch ratio was typically 4 to 5: 1 and indicates that an estimated 24 million pounds of finfish by-catch has occurred each year for the last five years. The study, which is considered valid and solid by DMF scientists, also determined that the finfish killed in the trawlers were small, averaging around 20 finfish per pound. Multiplying 20 fish per pound by 24 million pounds allows us to get a rough estimate of 480 million juvenile finfish that are killed annually by trawlers operating in North Carolina waters. This is unacceptable to the long-term health of our finfish populations.

I implore the MFC to amend the Shrimp FMP and address the issue of by catch by shrimp trawlers operating in our in shore waters.

I urge you to consider issuing an emergency order to suspend all shrimping and powered boat netting in inshore waters (including all our sounds) until an effective emergency by-catch reduction plan can be developed, vetted and approved.

Dr. H. Lee Dantzler
Ph.D, Oceanography
538 Joyner Drive
Havelock, NC 28532

Sent: Friday, October 05, 2012 3:22 PM

To: Fish, Nancy

Subject: Shrimp trawls

To whom it may concern:

Something has got to change in NC about trawling for Shrimp and the by-catch kill it is causing. To do nothing for the next five years borders on being criminal. At this rate no fishery can sustain itself. I don't understand how how Mr. Stryron can sit chair this board and be impartial. It is the same old story of " Having the Fox watch the Chickens " It's no wonder this State is so far behind our neighbors when it comes to fish management.

I would implore you to take action to stop this needless killing of our small fish, which will eventually lead to the decimation of all our inshore fish species. We need to act now !

Thank you,

David L. Griffith
New Bern NC

From: Larry Mize [<mailto:jmize2@triad.rr.com>]

Sent: Sunday, October 07, 2012 4:08 PM

To: Fish, Nancy

Subject: Shrimp Fisheries Management Plan

Larry Mize

701 N. Main St.

Graham N.C. 27253

336-227-1592

E-Mail jmize2@triad.rr.com

Date; October, 7, 2012

To: North Carolina Commission of Marine Fisheries

Subject; Shrimp Fisheries Management Plan

Dear Sirs,

I would like to recommend that the Marine Fisheries Commission amend the current Shrimp Fisheries Management Plan.

While I wouldn't want to see the shrimping industry in North Carolina shut down, I do believe we have a responsibility to properly and immediately manage all of our saltwater resources.

Sincerely,

Larry Mize

From: Ron Zielinski [<mailto:reellucky@centurylink.net>]

Sent: Wednesday, October 10, 2012 8:42 AM

To: Fish, Nancy

Subject: Comments on proposed 2012 shrimp fisheries management plan

Dear Ms. Nancy Fish,-----I attended the Oct. 2nd Shellfish/Crustacean meeting in New Bern to gather printed information and hear those who chose to speak during the comment period. After reviewing my notes and reading all material gathered, I wish to offer my thoughts on the future of N.C. coastal water's commercial shrimping and fish management. My past experiences include working with an organization to help clean up the Hudson River in New York, guest speaker at many fishing clubs and seminars, promoting the use of circle hooks to reduce by-catch mortality, spearheading the rebuilding of the local AR-396 Artificial Reef in conjunction with the NCDMF Artificial Reef program, occasionally writing articles for magazines, and being a licensed charter captain for 14 years. I am devoted to protecting our local natural resources for future generations' enjoyment and use. Before moving to North Carolina, many visits were made to experience the quality of local fishing. Talking to long time residents about their past fishing experiences made me realize there had been a reduction of sport fish in the Neuse River and Pamlico Sound. In the past 14 years, since first starting to fish the local waters, this reduction seems to be continuing. Several reasons have been offered for this development: lack of water quality, hurricanes, oxygen depletion leading to fish kills, commercial netting in nursery areas of local waters, the influx of commercial fishermen from states that have banned netting in inland waters, and the shrimp trawler's fin fish by-catch mortality. We cannot control Mother Nature but can work on modifying what we as individuals can do. Being we have the possibility of changing the N.C. commercial netting practices with the upcoming Shrimp Fisheries Management Plan, I wish to offer some changes for consideration. With an estimated annual 24 million pounds of juvenile finfish, (mostly spot, croaker, and grey trout), being netted and wasted as by-catch in our inshore waters, we should think of ways to protect these needed " seeds " to increase the numbers of these depleted species. From what has been printed, these figures were gathered from recent trawler studies conducted by the N.C. Dept. of Marine Fisheries. This summer and early fall, while piloting on the lower Neuse River and the Pamlico Sound, there were days when 30-40 shrimp trawlers were actively working in small areas from the mouth of Broad Creek (west of the NR buoy) and into the sound while chasing the migrating shrimp. This has also happened in previous years. Discussing this practice with other recreational fishermen and local commercial netters, the idea of permitting shrimp trawling only in the eastern part of the Pamlico Sound seemed an acceptable alternative. This compromise would reduce the juvenile by-catch mortality in the

Neuse River and near other feeder creeks and rivers used as nursery areas that flow into the west part of the sound. This would continue allowing the shrimp harvest in somewhat weather-protected inland waters behind the Outer Banks. A north/south longitude line or something similar like 2 marker buoys or points of land could be used for a defined separation. An additional benefit would be the elimination of the destruction of bottom habitat and structure which is needed for fin fish habitat. This could also stop the movement for the total elimination of trawling in the N.C. inland waters, as most other coastal states have already done. A reduction of the total head rope while in the Pamlico Sound was brought up by several speakers and sounded like a good idea. There did not seem to be a strong negative reaction to this proposal. In the end we are all in North Carolina to enjoy what it has to offer for work and recreation. We have to consider what's best for the environment, the economy and wellness for our state.-----Best Regards, Ron Zielinski

14.3 APPENDIX 3 – BYCATCH REDUCTION DEVICES AND TURTLE EXCLUDER DEVICES



FOR INFORMATION CONTACT:

Michael Barnette, Michael.Barnette@noaa.gov
2012
(727) 551-5794, FAX (727) 824-5309

May 21,

FB12-037

NOAA Certifies Additional Designs and Materials for Fishermen Currently Required to Use Turtle Excluder Devices

NOAA Fisheries Service announces a final rule adding allowable Turtle Excluder Device (TED) modifications and additional certified TED designs for the shrimp and summer flounder trawl fisheries. The additional designs and modifications may enhance TED effectiveness in reducing sea turtle mortality, promote catch retention, and increase vessel fuel efficiency. The allowable modifications are not mandatory, but provide additional options for fishermen currently required to utilize TEDs.

Provisions of the rule include:

- The addition of flat bar, box pipe, and oval pipe as construction material in currently-approved TED grids.
- An increase in the maximum mesh size on TED escape flaps from 1-5/8 to 2 inches (4.1 to 5.1 cm).
- The addition of the Boone Big Boy TED and Boone Wedge Cut TED escape opening for use in the shrimp fishery.
- The addition of three large TED escape openings.
- The addition of a brace bar as an allowable modification to hard TEDs.
- The addition of the Chauvin Shrimp Deflector to improve shrimp retention.
- The addition of a new TED for use in the summer flounder fishery.

There is also a correction to the TED regulations to rectify an oversight regarding the maximum size chain that can be used on the Parker TED escape opening flap.

NOTE: This final rule is not related to the proposed rule published in the *Federal Register* on May 10, 2012, that would, if implemented, require all skimmer trawls, pusher-head trawls, and wing nets (butterfly trawls) to use TEDs in their nets.

TEDs incorporate an escape opening, usually covered by a webbing flap, which allows sea turtles to escape from trawl nets. To be approved by NOAA Fisheries, a TED design must be shown to be 97 percent effective in excluding sea turtles during testing based upon specific testing protocols.

These latest modifications were developed in coordination with the commercial trawl industry, the Southeast Fisheries Science Center, and the Southeast Regional Office's Protected Resources Division. The modifications and TED designs were developed and tested by NOAA Fisheries gear specialists. Results from a study conducted by the Gulf and South Atlantic Fisheries Foundation were utilized in the development of these allowable modifications.


If you would like to receive these fishery bulletins via e-mail as soon as they are published, e-mail us at: SERO.Communications.Comments@noaa.gov . You will still receive a hard copy of these bulletins through the mail.

This bulletin provides only a summary of the information regarding the existing regulations. Any discrepancies between this bulletin and the regulations as published in the *Federal Register* will be resolved in favor of the *Federal Register*.

Recommended Construction and Installation Instructions

for the Extended Funnel Bycatch Reduction Device

NOAA Fisheries Service, Mississippi Laboratories
P.O. Drawer 1207, Pascagoula, Ms 39568-1207
March, 2008



The Extended Funnel Bycatch Reduction Device (BRD) has been certified for use in the Atlantic and provisionally certified for use in the Gulf of Mexico shrimp fishery. A **provisional certification** applies to an experimental BRD not quite meeting the criteria for certification, but deemed likely to meet the criteria with further testing. A provisional certification of a BRD is effective for 2 years from the date of publication in the Federal Register. This time period allows additional wide scale industry evaluation of the BRD candidate. The intent is to further refine the design or application of the experimental BRD so it could eventually meet the certification criterion.

Extended Funnel Description

The extended funnel BRD consists of an extension with large mesh webbing in the center and small mesh webbing on each end held open by a semi-rigid hoop. A funnel of small mesh webbing is placed inside the large mesh section to form a passage for shrimp to the codend. It also creates an area of reduced water flow to allow for fish escapement through the large mesh.

One side of the funnel is extended vertically to form a lead panel and area of reduced water flow.
Minimum Construction and Installation Requirements

Figure 1. Extension Material

The small mesh used on both sides of the large mesh escape section is constructed from #30, 1-5/8 inch (41 mm) stretch mesh nylon webbing. The front section is 120 meshes around by 6-1/2 meshes deep. The back section is 120 meshes around by 23 meshes deep.

Figure 2. Large Mesh Section

The large mesh escape section is constructed of 8 to 10 inch (20-25 cm) stretch mesh webbing. This section is cut on the bar to form a section that is 15 inches (38 cm) long, 95 inches (241 cm) in circumference. The leading edge is attached to the 6-1/2 mesh extension section and the rear edge is attached to the 23 mesh extension section.

Figure 3. Semi-Rigid Hoop

A 30 inch (76 cm) diameter hoop constructed of plastic coated trawl cable installed evenly 5 meshes behind the trailing edge of the large mesh section. The hoop is constructed using a 94-1/2 inch x 1/2 inch (240cm x 12.7mm) plastic coated cable. The ends are joined using a 3/8 inch micropress sleeve.

Figure 4. Funnel

The funnel is constructed of 1-1/2 inch (38 mm) stretch mesh #30 depth stretched and heat set polyethylene webbing. The circumference of the leading edge is 120 meshes and the back edge is 104 meshes. The short side of the funnel is 34 to 36 inches (86-91 cm) long and half of the opposite side of the funnel extends an additional 22 to 24 inches (5661 cm).

Figure 5. Funnel Attachment


The leading edge of the funnel is attached 3 meshes forward of the large mesh leading edge. Seven meshes of the short side of the funnel is attached to the back section of extension webbing on the top and bottom, 8 meshes back from the trailing edge of the large mesh section. The extended side of the funnel is attached on a slight angle to the top and bottom of the back extension webbing.

Installation of the Extended Funnel BRD

The Extended Funnel BRD is attached behind a hard TED 8 inches (20 cm) behind the posterior edge with the codend attached to the trailing edge of the BRD. If a soft TED is used a second hoop must be installed in the front section of the BRD extension webbing at the leading edge of the funnel.

The Extended Funnel BRD is designed to be installed immediately behind the Turtle Excluder Device (TED). To install the BRD, first remove the rear portion of the TED extension by cutting the TED extension on an even row of meshes 4 meshes behind the posterior edge of

the TED grid (a). Next, join the leading edge of the BRD extension evenly to the TED extension directly behind the TED (b). When attached, the BRD extension should be oriented so that the BRD extension seam is located on top of the trawl when towing. Complete the installation by attaching the codend (bag) to the trailing edge of the BRD extension.



This document was prepared for general informational purposes in March 2008 and has no legal force or effect. Please refer to the federal BRD regulations, 50 CFR part 622 and 622 Appendix D and the Federal Register for specific and controlling BRD requirements.

For more information contact:
NOAA Fisheries
Harvesting Systems and Engineering Branch
P.O. Drawer 1207
Pascagoula, MS 39568
(228) 762-4591
Robert.D.Stevens@noaa.gov or
Daniel.G.Foster@noaa.gov

Figure 1.







Figure 2.



Large Mesh
10" x 10"mm polyester


Figure 3.



1/2" plastic coated cable
ring circumference 94-1/4"

Figure 4.

Funnel






Side

Top

Side


Bottom



Webbing Panels

Figure 5.

Top View




Note: The 8 meshes should be on the funnel's long side of the extension's center seam.

99-FEG-33

Dennis Sawyer
Pamlico County Schools

CERTIFICATION OF BYCATCH REDUCTION DEVICES IN NORTH CAROLINA



99-FEG-33

November, 2001

Bruce Morris, Captain
F/V Miss PCHS
Pamlico County Schools
507 Anderson Drive
Bayboro, North Carolina 28515
Ph. (252) 745-4171 Fax (252) 745-4172

**CERTIFICATION OF
BYCATCH REDUCTION DEVICES
IN NORTH CAROLINA**

By

Bruce Morris

Pamlico County Schools

507 Anderson Drive
Bayboro, NC 28515
(252)745-4171
FAX
(252)745-4172

The research in this report was financed by the North Carolina Sea Grant administered Fisheries Resource Grant Program and conducted by the North Carolina Department of Public Instruction and Pamlico County Schools.

The contents of this publication do not necessarily reflect the views or policies of any of the above agencies or entities.

Grant 99-FEG-33

November, 2001

ACKNOWLEDGMENTS

The researcher wishes to thank the following individuals for their assistance during the course of this grant.

- Owen Lupton, Pamlico County Schools-retired, computer analysis, statistics, and report writing
- Dr. Julia Mobley, Hank Waters, Steve Curtis, Caroline Willis, and Jean Abbott from the Pamlico County Board of Education
- Allen Buck, Ryan Cuthrell, Travis Hardison, Tyler Hardison, Jason Harris, Justin Harris, Ricky Henries, Brad Newman, Scott Paul, Todd Powers, Daniel Rios, and Robbie Simpson, –students in the Marine Occupations II class of Pamlico County High School
- Sean McKenna, North Carolina Division of Marine Fisheries, technical assistance
- Greg Faulkner, net designer from Louisiana
- Tom Jamir, Gulf and South Atlantic Fisheries Foundation, Inc.

EXECUTIVE SUMMARY

The purpose of Marine Resources Grant 99-FEG-33 awarded to Pamlico County Schools was to help the North Carolina Division of Marine Fisheries certify bycatch reduction devices in the shrimp trawl fishery. The Atlantic States Fisheries Management Council (ASFMC) has adopted a protocol which Atlantic states within their jurisdiction must follow for the certification of bycatch reduction devices. Before any bycatch reduction device is approved for use in North Carolina, it must be certified according to this adopted protocol. Bycatch reduction devices can substantially reduce the bycatch of most species of finfish without causing shrimp trawling to become economically impossible. A second primary objective was to ascertain the percentage of shrimp loss in each bycatch reduction device.

When all tows for the control net were combined the total number of weakfish in the control was 2448. The total number of weakfish in all bycatch reduction devices was 1724, and total reduction by number was 30%. The total weight of all weakfish in the control was 161 kg. Total weight of weakfish in all bycatch reduction devices was 108 kg, and total reduction by weight was 33%. Total number of Spanish mackerel was 23 in the control, and 3 in all bycatch reduction devices for a reduction of 87%. Total weight of all Spanish mackerel in the control was 1.56 kg, and .18 kg in all bycatch reduction devices for a reduction of 89%.

The control net had a combined shrimp catch of 613 kg. The bycatch reduction devices had a combined shrimp catch of 566 kg. The overall percentage of shrimp loss was 8% for all bycatch reduction devices. Total tows for the entire research project was 126. Total useable tows was 108.

Results from subsamples showed most of the finfish in the tows were overwhelmingly Atlantic croaker and spot with a few miscellaneous species.

The ASFMC has adopted a protocol to use in certifying bycatch reduction devices and is primarily concerned about reaching a 40% reduction in numbers of weakfish and Spanish mackerel. North Carolina must have adequate data to meet the protocol for reduction of weakfish and Spanish mackerel in the shrimp trawl fishery. Bycatch reduction continues to be a major concern in all fisheries. Shrimp trawling takes place in a very fragile ecosystem. Inshore shrimp trawling in particular is adjacent to primary and secondary nursery areas where large numbers of the finfish are young of the year or juveniles that are non-marketable and therefore wasted other than as cull scrap returned to the water. The use of trawling gear is bottom disturbing, but the impact, if any, has not yet been proven to do permanent damage. The bottoms of shallow water sounds in North Carolina are just as easily disturbed by storms and wave action. The shrimp trawl industry is working in areas that are already under stress because of human and animal waste. The trawling industry must help preserve the limited reproductive and growing capacities of our waters, because human population is ever increasing which constantly results in loss of habitat. Less volume in the bycatch translates into more profit and less work for the shrimpers.

TABLE OF CONTENTS

	Page
Acknowledgments.....	ii
Executive Summary.....	iii
Table of Contents.....	iv
List of Tables.....	v
List of Figures.....	vi
Introduction.....	1
Gear Specifications.....	1
Data Analysis and Collection.....	4
Results	
Fish Slot II.....	5
Lake Arthur BRD.....	6
Kelly/Girouard Grid.....	6
Conclusions.....	7
Recommendations.....	8
Appendices	
References	

LIST OF TABLES

	Page
Table 1 Fish Slot II- Total weights.....	10
Table 2 Lake Arthur BRD- Total weights	14
Table 3 Kelly/Girourard Grid- Total weights.....	17
Table 4 Fish Slot II - Finfish average length and average weight	21
Table 5 Lake Arthur BRD - Finfish average length and average weight	22
Table 6 K/G Grid - Finfish average length and average weight	23

LIST OF FIGURES

	Page
Figure 1 Fish Slot II	2
Figure 2 Lake Arthur BRD.....	3
Figure 3 Kelly/Girouard Grid	4
Figure 4 Fish Slot II Total Weights.....	11
Figure 5 Weakfish Length Frequency Distribution- Fish Slot II	12
Figure 6 Spanish mackerel Length Frequency Distribution-Fish Slot II.....	13
Figure 7 Total Weights Lake Arthur BRD.....	15
Figure 8 Weakfish Length Frequency Distribution- Lake Arthur BRD	16
Figure 9 Total Weights K/G Grid.....	18
Figure 10 Weakfish Length Frequency Distribution- K/G Grid	19
Figure 11 Spanish mackerel Length Frequency Distribution- K/G Grid.....	20

INTRODUCTION

The purpose of Fishery Resource Grant 99-FEG-33 awarded to the Pamlico County Board of Education was to gather data to help the Division of Marine Fisheries certify bycatch reduction devices for the shrimp trawl fishery. North Carolina is following a testing protocol established by SAFMC (South Atlantic Fisheries Management Council) and adopted by ASFMC (Atlantic States Fisheries Management Council) for bycatch reduction of weakfish and Spanish mackerel in shrimp trawls. This grant gathered data for three bycatch reduction devices being tested for certification in North Carolina. They were the Fish Slot II, the Lake Arthur BRD, and the Kelly/Girourard Grid. This is the final report on the results.

The paired tow method of testing was used for certification. This method employs a control trawl without a bycatch reduction device pulled alongside a trawl using the test bycatch reduction device. The testing protocol requires a minimum of 30 tows for certification of a BRD with at least 5 or more weakfish or 1 Spanish mackerel in one of the trawls. Bycatch reduction in the shrimp trawl fishery is important from a conservation and economic standpoint. This research determined the numbers of weakfish and Spanish Mackerel caught, reduction of both species, and loss of shrimp with each bycatch reduction device.

The research for the grant began on June 26, 2000, and ended on September 11, 2001. Ten 60-minute tows calibrated the trawls to within 10% of each other prior to testing. Tow times were for 60 minutes unless otherwise noted. The BRD was switched to the opposite side of the vessel after 5 tows to minimize any difference in the gear. The completed summer research during 2000 yielded 41 tows for the Fish Slot II, but 3 were fouled and were discarded. The summer research during 2001 yielded 38 tows with the Lake Arthur BRD, but 4 were fouled and discarded. The 2001 summer research also yielded 37 tows for the Kelly/Girourard Grid with 1 fouled and discarded.


GEAR SPECIFICATIONS

The shrimp trawls were 32 foot (headrope) two-seam nets built by Harris Net Shop at Atlantic with 1.5 inch stretch mesh bodies cut on a 4/1 taper and constructed of #12 nylon webbing. Hanging was on 5/16-inch combination cable and the leg lines were 4 ft. The footrope rope had 1/4 inch galvanized chain. The wings had seven loops of chain (*16 links per loop*) and the belly had three loops. Three additional loops of chain were located between the wings and belly at the jib corners. The 3/16-inch tickler chain was two feet shorter than the footrope. The trawl doors were 5 and 1/2 feet long by 28 inches high and had 4 inch by 3/8-inch shoes on the bottom. The towing bridles were 3/8-inch diameter by 150 feet long. Both trawls were equipped with 120 mesh mini super-shooter TED's with bottom exit and accelerator funnels of stretch poly. The 1.5 inch


stretch mesh tailbags were constructed of #36 nylon webbing, 120 meshes around, and 120 meshes long with tie-off up 15 meshes from the bottom. All tailbags had elephant ears attached in place of choke straps to avoid the normal clogging that occurs with choke straps. Both trawls had YKK zippers about 10 feet in circumference sewn in between the TED and tailbag to facilitate BRD changes. The zippers worked well and did not come apart on any tows. Tailbag changes took less than 15 minutes.

The research vessel was the F/V Miss PCHS. The Miss PCHS is 38 feet long, powered by a CAT 3208 diesel with a 2:1 gear, and double rigged. All tows were at a vessel ground speed of 2.5 kts. as measured by GPS and Loran. Additional information for each tow included the date of tow, time of tow, and starting/stopping coordinates for the tow. Other recorded information included water temperature, salinity, water depth, wind direction and velocity, and tow number. As much as possible, all tows were in a straight line to prevent one net from fishing more ground than the other. No trynet was used during the research tows.


The Fish Slot II (Figure 1) is an aluminum rod-framed structure similar in design to the NMFS-approved Fish Eye. The Fish Slot II, however has a wider curvature and front-deflector. The Fish Slot II is installed at the 12 o'clock position in the tailbag, up 56 meshes from the tailbag tie-off. The Fish Slot II's trapezoid-shaped, bottom-floating trap door, made of extruded mesh, floats up and closes the escape hole to block shrimp from swimming out. This allows for a slightly longer period of time for fish to escape before shutting off as compared to similar devices like the Sea Eagle II, (National Fisherman, April 1999). The Fish Slot II testing yielded 38 useable tows from July 3, 2000 to August 3, 2000.



The Lake Arthur BRD (Figure 2) was originally used in the Lake Arthur area of western Louisiana. It was in use on single-rigged vessels for 20 years before the term BRD was ever coined. The Lake Arthur BRD doesn't depend on any type of framework. There is a 12-inch slit in the top of the tailbag at the 12 o'clock position. This slit was cut 56 meshes ahead of the tailbag tie-off. A chain is affixed to the front side of the slit, and a football float to the rear side. The combination of the float rising and the chain pulling down keeps the slit open while trawling. As the net slows down, the float pulls the net to the surface and bunches up webbing, which keeps the shrimp inside the tailbag (National Fisherman, April 1999). Testing yielded 34 useable tows from June 12, 2001 to July 10, 2001.



The Kelly/Girourard Grid is a grid-like aperture placed directly behind a TED. It can be used in the top or bottom of the cod end. The device has evenly spaced lateral bars that allow the fish to swim out of the trawl. When the BRD is fitted with a floating ramp, placed directly beneath the grid, fish swim out until the forward speed of the net slows down enough for the ramp to float upward and close off the escapement hole (National Fisherman, April 1999). The Kelly/Girourard Grid was tested from July 16, 2001 until September 11, 2001.



DATA ANALYSIS AND COLLECTION

The catch from each tailbag was examined in the following manner. All weakfish and Spanish mackerel were picked up. The numbers were counted from each tailbag and a random subsample of up to 30 individuals was taken. The subsample yielded an average length to the nearest millimeter, total length (TL), and an average weight to the nearest gram. The subsample also yielded a length frequency distribution of the weakfish caught with each BRD. The percentage of difference in total catch weight between the control and test BRD used the formula $[(\text{control-test})/\text{control} \times 100]$. The paired t-test method compared mean catches between the gear. A significance level of $P \leq 0.05$ was used for all tests. All data was recorded and calculated on Microsoft Excel 97. All the research data has been submitted to NCDMF for analysis and certification of the individual BRD's.

A subsample of the fish species caught was taken periodically to determine the types of finfish taken with the trawls. This subsample yielded a percentage of reduction for the different species. This subsample was taken by filling up one shrimp basket from each net and sorting the contents.

All shrimp were picked up and weighed. Then a sample of 3 pounds yielded a count per pound with heads on. A percentage of loss was calculated with each BRD.

Anything found forward of the elephant ear was excluded from the sample. Shaking out the nets before each tow began cleaned the trawls.

RESULTS

Fish Slot II

The number of weakfish in the control was 1770, and the Fish Slot II caught 1280 for a reduction of 27.7% (Table 1). Total weakfish weight in the control was 83.68 kg and in the Fish Slot II 58.68 for a reduction of 30%, (Figure 4).

The average length of weakfish in the control was 169 mm and the average weight was 54 grams. The average length of weakfish in the Fish Slot II was 167 mm and the average weight was 48 grams (Table 4). The control average was 46.6 weakfish per tow and the Fish Slot II average was 37.6 per tow.

The length frequency distribution range of weakfish was 90 mm to 300 mm. Both the control and BRD had weakfish in the greatest abundance in the 150 mm to 180 mm range with 160 mm predominating (Figure 5).

The control caught 7 Spanish mackerel with an average length of 156 mm and an average weight of 57 grams (Table 4). Length frequency distribution for the control was 150 mm to 165 mm with 3 at 150 mm, 3 at 160 mm, and 1 at 165 mm (Figure 6). The Fish Slot II caught no Spanish mackerel.

Total shrimp catch was 283.41 kg in the control and 267.41 kg in the Fish Slot II for a loss of 5.5% (Figure 4).

Weight reduction of spot in the 13 subsample tows was 19.3%, and reduction of Atlantic Croaker was 26.3%. Total finfish weight including the 13 subsample tows was 304.57 kg in the control and 229.65 kg in the Fish Slot II giving an overall reduction of 24.6% (Table 1).

Lake Arthur

The number of weakfish in the control was 199 and the number in the Lake Arthur was 131 for a reduction of 34.2% (Table 2). The control had a total weakfish weight of 31.46 kg and the Lake Arthur BRD had 20.80 kg for a reduction of 33.9% (Figure 7).

The average length of weakfish in the control was 237 mm and the average weight was 157 grams. The average length of weakfish in the Lake Arthur was 234 mm and the average weight was 160 grams (Table 5). The control average was 5.85 weakfish per tow and the Lake Arthur average was 3.85 per tow.

The length frequency distribution range of weakfish was 80 mm to 350-mm. Both the control and BRD had weakfish in the greatest abundance in the 220 mm to 260 mm range with the peak at about 240 mm (Figure 8).

Only one Spanish mackerel was caught in the control at a length of 150 mm and weight of 50 grams (Table 5). No Spanish were caught in the Lake Arthur.

Total shrimp catch was 156.25 kg in the control and 141.36 kg in the Lake Arthur giving a loss of 9.5% in the Lake Arthur (Figure 7).

Weight increased for spot in the 6 subsamples at 4.8%, and Atlantic Croaker had an 18.8% increase in the subsamples. Finfish weight including the 6 subsample tows was 118.78 kg for the control and 116.71 kg for the Lake Arthur giving a reduction of 1.7%. (Table 2).

Kelly/Girourard Grid

Total number of weakfish was 479 in the control and 313 in the K/G Grid for a reduction of 34.7% (Table 3). The control had a total weakfish weight of 45.55 kg and the Kelly/Girourard Grid had 28.25 kg for a reduction of 38% (Figure 9).

The average length of weakfish in the control was 177 mm and the average weight was 97 grams. The average length of weakfish in the K/G Grid was 171 mm and the average weight was 96 grams (Table 6). The control average was 13.3 weakfish per tow and the K/G Grid average was 8.7 weakfish per tow.

The length frequency distribution range of weakfish was 90 mm to 320 mm. Both the control and BRD had weakfish in the greatest abundance in the 120 mm to 170 mm range (Figure 10).

The control caught 15 Spanish mackerel and the K/G Grid caught 3 for a reduction of 80% by number. The average length of Spanish mackerel in the control was 159 mm and the average weight was 67 grams. The average length of Spanish mackerel in the K/G Grid was 133 mm and the average weight was 58 grams (Table 6).

The length frequency distribution for Spanish mackerel was 120 mm to 210 mm. The control had Spanish in the greatest abundance at 170 mm, while the 3 Spanish in the K/G grid were at the 130 mm to 140 mm range (Figure 11).

Total shrimp catch was 172.84 kg in the control and 156.70 kg in the K/G Grid for a loss of 9.3% (Figure 9).

The 5 subsample tows had a 21% weight increase for spot, and an 11.4% weight increase for Atlantic croaker. Total finfish weight including the 5 subsample tows was 104.86 kg in the control and 95.49 kg in the BRD giving an overall reduction of 8.9% (Table 3).

CONCLUSIONS

Fish Slot II

The Fish Slot II, which originated in the Gulf of Mexico shrimp fishery, did not have enough reduction of weakfish, at 27.7%, to certify in North Carolina. While the reduction of Spanish mackerel was 100%, there simply was not enough catch of Spanish, with only 7, to draw any valid conclusions. The shrimp loss was reasonably low at 5.5%, but there was not a high reduction of our two most abundant fish species with croaker at 26% and spot at 19%. This model is prone to hang on the bumper rails of the vessel's sides and can damage or destroy the tailbag or BRD.

Lake Arthur

The Lake Arthur BRD did not certify where it was installed with a 34.2% weakfish reduction by number. Reduction of Spanish mackerel was 100% but only 1 was caught in all the tows and so that data is not reliable. Shrimp loss was fairly substantial at 9.5%. In an unexpected occurrence, both croaker and spot increased in the Lake Arthur at 18.8% and 4.8% respectively. Since only 6 subsample tows were made during this testing, that could account for the

increase where a decrease should normally be expected. This model, much like the Fish Slot II, is prone to hang on the bumper railings of the vessel during haul-back.

Kelly/Girourard Grid

The Kelly/Girourard Grid had almost a 35% reduction of weakfish by number, but was still short of the 40% needed for certification. Reduction of Spanish mackerel was good at 80%, but again the small number, of only 15 in the control and 3 in the K/G Grid, was not really enough to be sure what it would do on a long-term basis. However, 15 Spanish is far more than would normally be caught in the control, and so the reduction of Spanish alone could possibly be used as a basis for certification, or further testing.

Shrimp loss was significant with this BRD at over 9%. This BRD also gave the same unexpected result as the Lake Arthur. During the 5 subsample tests the weight increased for croaker at 11% and spot at 21%. Obviously this should not be occurring since the reduction of both weakfish and Spanish was a significant amount. The relatively small number of tows (5) where a total subsample was taken could account for the increases in croaker and spot.

With the K/G Grid installed so close to the accelerator funnel of the TED some finfish may not be excluded since the bottom of the tailbag is so far from the release point. The K/G Grid did not present a problem hanging up on the bumper rails like the other two devices.

Sportfish Bycatch

Spanish mackerel do not show as a significant part of the bycatch in Pamlico Sound. With 108 tows completed, only 26 Spanish mackerel were caught. Speckled trout are also insignificant in the bycatch with only 3 caught. No red drum were caught in any tows.

RECOMMENDATIONS

The Fish Slot II is not recommended since it did not certify, and was substantially below the needed 40% for weakfish. Changing the placement in the tailbag could make a significant enough difference that the Fish Slot II might certify. This BRD apparently works well in some states, but did not do very good in Pamlico Sound.

While the concept of the Lake Arthur is very simple, and apparently has worked well at excluding menhaden in western Louisiana, it did not do good

enough to certify or be recommended. Changing the placement in the tailbag could fine-tune this BRD so that it might certify. The almost 10% loss of shrimp would be a problem unless that can be lowered. If changed to a different tailbag location, it could conceivably certify.

The Kelly/Girourard Grid worked without any hang-ups, but did not have enough reduction of weakfish to certify. It was reasonably close, and since it had an 80% reduction on Spanish mackerel, further fine-tuning could find a way to certify it. This BRD is recommended for further testing to see whether the reduction of Spanish mackerel remains constant and the reduction of weakfish can be increased to 40%. The loss of over 9% of shrimp needs to be lowered to receive good acceptance by industry even if the K/G Grid could certify.

Our testing took three devices that have worked well in the Gulf of Mexico and none of them did well enough to certify here in North Carolina. Differences in water clarity, bottom conditions, types of finfish encountered and even the species of shrimp being sought can make significant differences from one state to another. The only way to be sure what any BRD will do in a particular state is to test and see what the results finally turn out to be.

Shrimp trawling will constantly be faced with the need to reduce unwanted bycatch to help preserve as many finfish as possible. While there are no scientific studies available to determine what percentage of finfish die from natural mortality before maturity, everyone can agree that dead fish from trawling looks wasteful even if the effect may be negligible.

At the same time, government must not make it impossible, by over-regulation of bycatch reduction requirements, for the shrimp trawling industry to catch enough shrimp to be economically feasible. What we must work together to achieve is maximum bycatch reduction with minimum loss of shrimp.

Lowering bycatch is not just a good conservation measure, but also makes trawling more efficient and saves time, money, fuel, and effort. While we may never reach 100% bycatch reduction, the only way to improve is to continue to invent, test, and modify as much as humanly possible.

TABLE 1. Total catch weights and reduction for 38 tows with Fish Slot II, tested in western Pamlico Sound, North Carolina, Summer 2000.

	Total Number		Percent	
	Control	Fish Slot II	Difference	
***n=38				
WEAKFISH	1770	1280	-27.7%	
SPANISH MACKEREL	7	0	-100.0%	
SPECKLED TROUT	0	0		

	Total Weight (kg)		Percent		P(T<=t)
	Control	Fish Slot II	Difference		
SHRIMP	283.41	267.73	-5.5%	8.26E-03*	
Summer Shrimp	283.41	267.73	-5.5%		
SOUTHERN FLOUNDER**	4.98	4.27	-14.3%		
sublegal-und 330 mm**	4.98	4.27	-14.3%		
SPOT**	113.18	91.36	-19.3%	6.56E-02	
ATLANTIC CROAKER**	96.82	71.36	-26.3%	6.33E-03*	
WEAKFISH	83.68	58.68	-29.9%	2.52E-08*	
SPANISH MACKEREL	0.40	0.00	-100.0%	1.64E-02*	
ATLANTIC MENHADEN**	0.91	0.80	-12.5%		
PIGFISH**	2.80	2.18	-22.0%		
INSHORE LIZARDFISH**	1.00	1.00	0.0%		
SOUTHERN KINGFISH**	0.68	0.00	-100.0%		
BLACK DRUM**	0.13	0.00	-100.0%		
CRABS & OTHER INVERTEBRATES**	43.18	42.50	-1.6%		
JELLYFISH**	9.77	8.41	-14.0%		
Total Finfish + Subsample	304.57	229.65	-24.6%		
Total Weight + Subsample	640.94	548.29	-14.5%		

***all tows conducted during the daytime

**subsample weights only for 13 tows

*Significant difference at the p<=.05 level or less

Figure 4. Total catch weights for 38 tows with the Fish Slot II, tested in western Pamlico Sound, North Carolina, Summer 2000.





Figure 5 Length Frequency -- Weakfish - Fish Slot II for 38 tows tested in western Pamlico Sound, summer 2000






TABLE 2. Total catch weights and reduction for 34 tows with Lake Arthur BRD, tested in western Pamlico Sound, North Carolina, Summer 2001.

	Total Number		Percent Difference	
	Control	Lake Arthur		
***n=34				
WEAKFISH	199	131	-34.2%	
SPANISH MACKEREL	1	0	-100.0%	
SPECKLED TROUT	0	0		

	Total Weight (kg)		Percent Difference	P(T<=t)
	Control	Lake Arthur		
SHRIMP	156.25	141.36	-9.5%	6.25E-04*
Summer Shrimp	156.25	141.36	-9.5%	
SOUTHERN FLOUNDER**	0.91	0.00	-100.0%	
sublegal-und 330 mm**	0.91	0.00	-100.0%	
SPOT**	47.73	50.00	4.8%	8.13E-01
ATLANTIC CROAKER**	38.64	45.91	18.8%	3.61E-01
WEAKFISH	31.46	20.80	-33.9%	5.92E-05*
SPANISH MACKEREL	0.05	0.00	-100.0%	3.24E-01
CRABS & OTHER INVERTEBRATES**	3.86	3.64	-5.9%	
JELLYFISH**	26.36	22.73	-13.8%	
Total Finfish + Subsample	118.78	116.71	-1.7%	
Total Weight + Subsample	305.26	284.44	-6.8%	

***all tows conducted during the daytime

**subsample weights 6 tows only

*Significant difference at the $p < .05$ level or less

Figure 7. Total catch weights for 34 tows with the Lake Arthur BRD, tested in western Pamlico Sound, North Carolina, Summer 2001.




Figure 8. Length Frequency -- Weakfish -- Lake/Arthur BRD for 34 tows tested in western Pamlico Sound, summer 2001.




TABLE 3. Total catch weights and reduction for 36 tows with Kelly/Girourard Grid, tested in western Pamlico Sound, North Carolina, Summer 2001.

	Total Number		Percent Difference	
	Control	K/G Grid		
***n=36				
WEAKFISH	479	313	-34.7%	
SPANISH MACKEREL	15	3	-80.0%	
SPECKLED TROUT	2	1	-50.0%	

	Total Weight (kg)		Percent Difference	P(T<=t)
	Control	K/G Grid		
SHRIMP	172.84	156.70	-9.3%	1.24E-03*
Summer Shrimp	172.84	156.70	-9.3%	
SPOT**	25.91	31.36	21.1%	5.81E-01
ATLANTIC CROAKER**	31.82	35.45	11.4%	3.74E-01
WEAKFISH	45.55	28.25	-38.0%	1.06E-09*
SPANISH MACKEREL	1.11	0.18	-84.2%	8.82E-03*
SOUTHERN KINGFISH**	0.08	0.00	-100.0%	
SPOTTED SEATROUT	0.40	0.25		
CRABS**	2.73	2.73	0.0%	
JELLYFISH**	0.45	0.45	0.0%	
Total Finfish + Subsample	104.86	95.49	-8.9%	
Total Weight + Subsample	280.89	255.37	-9.1%	

***all tows conducted during the daytime

**subsample weights for 5 tows only

*Significant difference at the p<=.05 level or less

Figure 9. Total catch weights for 36 tows with the Kelly/Girouard Grid, tested in western Pamlico Sound, North Carolina, Summer 2001.





Figure 10: Length Frequency -- Weakfish - Kelly/Girouard Grid for 36 tows tested in western Pamlico Sound, summer 2001






Table 4. Average weight and average length of finfish for 38 tows with the Fish Slot II, tested in western Pamlico Sound, summer, 2000.

	Weakfish		Speckled Trout		Spanish Mackerel	
	Control	Fish Slot II	Control	Fish Slot II	Control	Fish Slot II
Ave. LT.	169	167	0	0	156	0
Agg. WT.	57925	45940	0	0	400	0
Ave. WT.	54	48	0	0	57	0

Table 5. Average length and average weight of finfish for 34 tows with the Lake Arthur BRD, tested in western Pamlico Sound, Summer, 2001.

	Weakfish		Speckled Trout		Spanish mackerel	
	Control	Lake Arthur	Control	Lake Arthur	Control	Lake Arthur
Ave. LT.	237	234	0	0	100	0
Agg. WT.	31460	20800	0	0	50	0
Ave. WT..	157	160	0	0	50	0

Table 6. Average length and average weight of finfish for 36 tows with the Kelly/Girourard Grid tested in western Pamlico Sound, summer, 2001.

	WEAKFISH		Speckled Trout		Spanish Mackerel	
	CONTROL	K/G GRID	Control	K/G GRID	Control	K/G GRID
Ave. LT.	177	171	180	110	159	133
Agg. WT.	43150	26925	425	25	1000	175
Ave. WT.	97	96	213	25	67	58

REFERENCES

- Cunningham, P., R. Curry, R. Pratt, S. Stichter, K. West, L. Mercer, P. Phalen, S. Sherman, B. Burns, and S. Winslow. 1992. Watershed planning in the Albemarle-Pamlico estuarine system: fishing practices mapping. Albemarle Pamlico Estuarine Study Rep. No. 92-05.
- Faulkner, Greg. April, 1999. BRD's of Paradise, National Fisherman, Vol. 79, No. 12, Diversified Business Communications, Portland, Maine.
- Knopf, A. 1983. The Audobon Society Field Guide to North American Fishes, Whales, and Dolphins. Chanticleer Press, Inc., New York.
- Lupton, Owen A. 1996. Bycatch reduction in the estuarine crab trawl industry through manipulating tailbag sizes. Marine Fisheries Grant FRG-94-11, Pamlico County Schools, Bayboro, North Carolina.
- Lupton, Owen A. 1997. Bycatch reduction in the inshore shrimp trawl fishery. Marine Fisheries Grant FRG 95-15, Pamlico County Schools, Bayboro, North Carolina.
- Lupton, Owen A. 1998. Certification of Bycatch Reduction Devices for North Carolina. Marine Fisheries Grant FEG 97-40, Pamlico County Schools, Bayboro, North Carolina.
- McKenna, S., and J.T. Camp. 1992. An examination of the blue crab fishery in the Pamlico River Estuary. Albemarle-Pamlico Estuarine Study Rep. No. 92-08.
- McKenna, S., and A.H. Clark 1993. An examination of alternative fishing devices for the estuarine shrimp and crab trawl fisheries. Albemarle-Pamlico Estuarine Study Rep. No. 93-11.
- Morris, Bruce 1999. North Carolina Bycatch Reduction Device Certification. North Carolina Marine Fisheries Grant 98-FEG-23, Pamlico County Schools, Bayboro, North Carolina.
- Tyler, J. and M. McKenzie. 1969. *To Catch a Million Fish: North Carolina Commercial Fishing Gear and Methods*. North Carolina Department of Conservation and Development, Division of Commercial and Sport Fishing, Raleigh, North Carolina.



Southeast Fishery Bulletin

National Marine Fisheries Service, Southeast Regional Office, 263 13th Avenue South, St. Petersburg, FL 33701

FOR INFORMATION CONTACT:

Michael Barnette, Michael.Barnette@noaa.gov
(727) 551-5794, FAX (727) 824-5309

May 21, 2012

FB12-037

NOAA Certifies Additional Designs and Materials for Fishermen Currently Required to Use Turtle Excluder Devices

NOAA Fisheries Service announces a final rule adding allowable Turtle Excluder Device (TED) modifications and additional certified TED designs for the shrimp and summer flounder trawl fisheries. The additional designs and modifications may enhance TED effectiveness in reducing sea turtle mortality, promote catch retention, and increase vessel fuel efficiency. The allowable modifications are not mandatory, but provide additional options for fishermen currently required to utilize TEDs.

Provisions of the rule include:

- VI. The addition of flat bar, box pipe, and oval pipe as construction material in currently-approved TED grids.
- VII. An increase in the maximum mesh size on TED escape flaps from 1-5/8 to 2 inches (4.1 to 5.1 cm).
- VIII. The addition of the Boone Big Boy TED and Boone Wedge Cut TED escape opening for use in the shrimp fishery.
- IX. The addition of three large TED escape openings.
- X. The addition of a brace bar as an allowable modification to hard TEDs.
- XI. The addition of the Chauvin Shrimp Deflector to improve shrimp retention.
- XII. The addition of a new TED for use in the summer flounder fishery.

There is also a correction to the TED regulations to rectify an oversight regarding the maximum size chain that can be used on the Parker TED escape opening flap.

NOTE: This final rule is not related to the proposed rule published in the *Federal Register* on May 10, 2012, that would, if implemented, require all skimmer trawls, pusher-head trawls, and wing nets (butterfly trawls) to use TEDs in their nets.

TEDs incorporate an escape opening, usually covered by a webbing flap, which allows sea turtles to escape from trawl nets. To be approved by NOAA Fisheries, a TED design must be shown to be 97 percent effective in excluding sea turtles during testing based upon specific testing protocols.

These latest modifications were developed in coordination with the commercial trawl industry, the Southeast Fisheries Science Center, and the Southeast Regional Office's Protected Resources Division. The modifications and TED designs were developed and tested by NOAA Fisheries gear specialists. Results from a study conducted by the Gulf and South Atlantic Fisheries Foundation were utilized in the development of these allowable modifications.

If you would like to receive these fishery bulletins via e-mail as soon as they are published, e-mail us at: SERO.Communications.Comments@noaa.gov . You will still receive a hard copy of these bulletins through the mail.

This bulletin provides only a summary of the information regarding the existing regulations. Any discrepancies between this bulletin and the regulations as published in the *Federal Register* will be resolved in favor of the *Federal Register*.