

SEDAR4 Data Workshop Reference Documents List

- SEDAR4-RD01: Bohnsack, J. A. and A. Woodhead. 1995. Proceedings of the 1987 SEAMAP passive gear assessment workshop at Mayaguez, Puerto Rico. NOAA Tech. Mem. NMFS SEFSC 365.
- SEDAR4-RD02: Bullis, H. R. Jr. and A. C. Jones, *ed.* 1976. Proceedings: Colloquium on snapper-grouper fishery resources of the Western Central Atlantic Ocean. FL SeaGrant Report No. 17.
- SEDAR4-RD03: Bullock, L. H., M. F. Godcharles, and R. E. Crabtree. 1996. Reproduction of yellowedge grouper, *Epinephelus flavolimbatus*, from the Eastern Gulf of Mexico. Bull. Mar. Sci. 59(1) 224-228.
- SEDAR4-RD04: Erickson, D. E. and G. D. Grossman. 1986. Reproductive demography of tilefish from the South Atlantic Bight with a test for the presence of protogynous hermaphroditism. Trans. Am. Fish. Soc. 115:279-285.
- SEDAR4-RD05: Grimes, C. B. and S. C. Turner. 1999. The complex life history of tilefish *Lopholatilus chamaeleonticeps* and vulnerability to exploitation. Am. Fish. Soc. Symp. 23:17-26.
- SEDAR4-RD06: Grossman, G. D., M. J. Harris, and J. E. Hightower. 1985. The relationship between tilefish, *Lopholatilus chamaeleonticeps*, abundance and sediment composition off Georgia. Fish. Bull. 83(3):443-447.
- SEDAR4-RD07: Harris, M. J. and G. D. Grossman. 1985. Growth, mortality, and age composition of a lightly exploited tilefish substock off Georgia. Trans. Am. Fish. Soc. 114:837-846.
- SEDAR4-RD08: Hightower, J. E., and G. D. Grossman. 1989. Status of the tilefish, *Lopholatilus chamaeleonticeps*, fishery off South Carolina and Georgia and recommendations for management. Fish. Bull. 87:177-188.
- SEDAR4-RD09: Low, B., G. Ulrich, and F. Blum. 1982. The fishery for tilefish, *Lopholatilus chamaeleonticeps*, off South Carolina and Georgia. SC Wildl. and Mar. Res. Div. Charleston, SC.
- SEDAR4-RD10: Low, R. A. Jr., G. F. Ulrich, and F. Blum. 1983. Tilefish off South Carolina and Georgia. Mar. Fish. Rev. 45(4-6)16-26.
- SEDAR4-RD11: Manooch, C. S., and D. L. Mason. 1987. Age and growth of the Warsaw grouper and black grouper from the Southeast region of the United States. Northeast Gulf Sci. 9(2):65-75.
- SEDAR4-RD12: Matheson, R. H. and G. R. Huntsman. 1984. Growth, mortality, and yield-per-recruit models for speckled hind and snowy grouper from the United States South Atlantic Bight. Trans. Am. Fish. Soc. 115:607-616.
- SEDAR4-RD13: Parker, R. O. Jr. and R. W. Mays. 1998. Southeastern U. S. deepwater reef fish assemblages, habitat characteristics, catches, and life history summaries. NOAA Tech. Report. NMFS-138.
- SEDAR4-RD14: Ross, J. L. 1982. Feeding habits of the gray tilefish *Caulolatilus microps* (Goode and Bean, 1878) from North Carolina and South Carolina waters. Bull. Mar. Sci. 32(2):448-454.

SEDAR4-RD15: Ross, J. L. and G. R. Huntsman. 1982. Age, growth, and mortality of blueline tilefish from North Carolina and South Carolina. Trans. Am. Fish. Soc. 111:585-592.

SEDAR4-RD16: Russel, G. M., E. J. Gutherz, and C. A. Barans. 1988. Evaluation of demersal longline gear off South Carolina and Puerto Rico with emphasis on deep-water reef fish stocks. Mar. Fish. Rev. 50(1):26-31.

SEDAR4-RD17: Tester, P. A., C. A. Wolfe, R. L. Dixon, and G. R. Huntsman. 1983. Reef fish distributions off North Carolina and South Carolina as revealed by headboat catches. NOAA Tech. Mem. NMFS-SEFC-115.

SEDAR4-RD18: Wyanski, D. M., D. B. White, and C. A. Barans. 2000. Growth, population age structure, and aspects of the reproductive biology of snowy grouper, *Epinephelus niveatus*, off North Carolina and South Carolina. Fish. Bull. 98:199-218.