































- Clarke, T. A., and L. A. Privitera.  
1995. Reproductive biology of two Hawaiian pelagic carangid fishes, the bigeye scad, *Selar crumenophthalmus*, and the round scad, *Decapturus macarellus*. *Bull. Mar. Sci.* 56:33-47.
- Dawson, C. E.  
1971. Occurrence and description of prejuvenile and early juvenile Gulf of Mexico cobia, *Rachycentron canadum*. *Copeia* 1971:65-71.
- Ditty, J. G., and R. F. Shaw.  
1992. Larval development, distribution, and ecology of cobia, *Rachycentron canadum*, (Family: Rachycentridae) in the northern Gulf of Mexico. *Fish. Bull.* 90:668-677.
- Farley, J. H. and T. L. O. Davis.  
1998. Reproductive dynamics of southern bluefin tuna, *Thunnus maccoyii*. *Fish. Bull.* 96:223-236.
- Fitzhugh, G. R., B. A. Thompson, and T. G. Snider III.  
1993. Ovarian development, fecundity and spawning frequency of black drum *Pogonias cromis* in Louisiana. *Fish. Bull.* 91:244-253.
- Franks, J. S., J. R. Warren, and M. V. Buchanan.  
1999. Age and growth of cobia, *Rachycentron canadum*, from the northeastern Gulf of Mexico. *Fish. Bull.* 97:459-471.
- Grier, H. J., and M. Abraham.  
1983. A model for testicular recrudescence in *Oreochromis aureus*. In *International symposium on tilapia in aquaculture* (L. Fishelson and Z. Yaron, eds.), p. 200-209. Tel Aviv Univ. Press, Tel Aviv.
- Grier, H. J., and R. G. Taylor.  
1998. Testicular maturation and regression in the common snook. *J. Fish Biol.* 53:521-542.
- Grier, H. J., R. G. Taylor, and R. O. Reese.  
1987. The mechanism of tubule elongation during testicular recrudescence in the redfish, *Sciaenops ocellatus* (Perciformes). In *Proceedings of the V congress of European ichthyologists* (S. O. Kullander and B. Fernholm, eds.), p. 285-291. Swedish Museum of Natural History, Stockholm.
- Hassler, W. W., and R. P. Rainville.  
1975. Techniques for hatching and rearing cobia, *Rachycentron canadum*, through larval and juvenile stages. Publ. UNC-SC-75-30, Univ. N.C. Sea Grant Coll. Prog., Raleigh, NC, 26 p.
- Hay, D. E., and J. R. Brett.  
1988. Maturation and fecundity of Pacific herring (*Clupea harengus pallasii*): an experimental study with comparisons to natural populations. *Can. J. Fish. Aquat. Sci.* 45:399-406.
- Hrinkevich, A. W.  
1993. Mitochondrial DNA analysis of cobia *Rachycentron canadum* population structure using restriction fragment length polymorphisms and cytochrome B sequence variation. M.S. thesis, Univ. Southern Mississippi, Hattiesburg, MS, 91 p.
- Hunter, J. R., and B. J. Macewicz.  
1985a. Rates of atresia in the ovary of captive and wild northern anchovy, *Engraulis mordax*. *Fish. Bull.* 83:119-136.  
1985b. Measurement of spawning frequency in multiple spawning fishes. In *An egg production method for estimating spawning biomass of pelagic fish: application to the northern anchovy, *Engraulis mordax** (R. Lasker, ed.), p. 79-94. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 36.
- Hunter, J. R., B. J. Macewicz, and H. R. Sibert.  
1986. The spawning frequency of skipjack tuna, *Katsuwonus pelamis*, from the south Pacific. *Fish. Bull.* 84:895-903.
- International Game Fish Association.  
1998. World record game fishes. International Game Fish Association, Pompano Beach, FL, 352 p.
- Jons, G. D., and L. E. Miranda.  
1997. Ovarian weight as an index of fecundity, maturity and spawning periodicity. *J. Fish Biol.* 50:150-156.
- Joseph, E. B., J. J. Norcross, and W. H. Massman.  
1964. Spawning of the cobia, *Rachycentron canadum*, in the Chesapeake Bay area, with observations of juvenile specimens. *Chesapeake Sci.* 5:67-71.
- Livingston, R. J., X. Niu, F. G. Lewis III, and G. C. Woodsum.  
1997. Freshwater input to gulf estuaries: long-term control of trophic organization. *Ecol. Appl.* 7:277-299.
- Lotz, J. M., R. M. Overstreet, and J. S. Franks.  
1996. Gonadal maturation in the cobia, *Rachycentron canadum*, from the northcentral Gulf of Mexico. *Gulf Res. Reports* 9:147-159.
- Lowerre-Barbieri, S. K., M. E. Chittenden Jr., and L. R. Barbieri.  
1996. The multiple spawning pattern of weakfish in the Chesapeake Bay and Middle Atlantic Bight. *J. Fish. Biol.* 48:1139-1163.
- McPherson, G. R.  
1993. Reproductive biology of the narrow barred Spanish mackerel (*Scomberomorus commerson* Lacepede, 1800) in Queensland waters. *Asian Fish. Sci.* 6:169-182.
- Richards, C. E.  
1967. Age, growth and fecundity of the cobia, *Rachycentron canadum*, from the Chesapeake Bay and adjacent mid-Atlantic waters. *Trans. Am. Fish. Soc.* 96:343-350.
- Schaefer, K. M.  
1996. Spawning time, frequency, and batch fecundity of yellowfin tuna, *Thunnus albacares*, near Clipperton Atoll in the eastern Pacific Ocean. *Fish. Bull.* 94:66-76.
- Shaffer, R. V., and E. L. Nakamura.  
1989. Synopsis of biological data on the cobia, *Rachycentron canadum*, (Pisces: Rachycentridae). U.S. Dep. Commer., NOAA Tech. Rep. NMFS 82 [FAO Fish Synop. 153], 21 p.
- Smith, J. W.  
1996. Life history of cobia, *Rachycentron canadum* (Osteichthyes: Rachycentridae), in North Carolina waters. *Brimleyana* 23:1-23.
- SPSS, Inc.  
1997. SPSS PC version 7.5. SPSS, Inc., Chicago, IL, 628 p.  
1998. SYSTAT 8.0: statistics. SPSS, Inc, Chicago, IL, 1086 p.
- Taylor, R. G., H. J. Grier, and J. A. Whittington.  
1998. Spawning rhythms of common snook in Florida. *J. Fish Biol.* 53:502-520.
- Watson, J. J. I. G. Priede, P. R. Witthames, and A. Owari-Wadunde.  
1992. Batch fecundity of Atlantic mackerel, *Scomber scombrus* L. *J. Fish Biol.* 40:591-598.
- Wilson, C. A., and D. L. Nieland.  
1994. Reproductive biology of red drum, *Sciaenops ocellatus*, from the neritic waters of the northern Gulf of Mexico. *Fish. Bull.* 92:841-850.