SEDAR 28 (South Atlantic and Gulf of Mexico Cobia and Spanish Mackerel) Document List

All documents available on the SEDAR website

SEDAR58-RD03

30 January 2018



SEDAR 28 Gulf and South Atlantic -- Spanish Mackerel and Cobia Workshop Document List

| Document # | Title | Authors | |
|--|--|---|--|
| Documents Prepared for the Data Workshop | | | |
| SEDAR28-DW01 | Cobia preliminary data analyses – US Atlantic and GOM genetic population structure | Darden 2012 | |
| SEDAR28-DW02 | South Carolina experimental stocking of cobia Rachycentron canadum | Denson 2012 | |
| SEDAR28-DW03 | Spanish Mackerel and Cobia Abundance Indices from SEAMAP Groundfish Surveys in the Northern Gulf of Mexico | Pollack and Ingram, 2012 | |
| SEDAR28-DW04 | Calculated discards of Spanish mackerel and cobia from commercial fishing vessels in the Gulf of Mexico and US South Atlantic | K. McCarthy | |
| SEDAR28-DW05 | Evaluation of cobia movement and distribution using tagging data from the Gulf of Mexico and South Atlantic coast of the United States | M. Perkinson and M. Denson 2012 | |
| SEDAR28-DW06 | Methods for Estimating Shrimp Bycatch of Gulf of Mexico Spanish Mackerel and Cobia | B. Linton 2012 | |
| SEDAR28-DW07 | Size Frequency Distribution of Spanish Mackerel from Dockside Sampling of Recreational and Commercial Landings in the Gulf of Mexico 1981-2011 | N.Cummings, J. Isely | |
| SEDAR28-DW08 | Size Frequency Distribution of Cobia from Dockside Sampling of Recreational and Commercial Landings in the Gulf of Mexico 1986-2011 | J. Isely and N. Cummings | |
| SEDAR28-DW09 | Texas Parks and Wildlife Catch Per unit of Effort Abundance Information for Spanish mackerel | N. Cummings, J. Isely | |
| SEDAR28-DW10 | Texas Parks and Wildlife Catch Per unit of Effort Abundance Information for cobia | J. Isely, N. Cummings | |
| SEDAR28-DW11 | Size Frequency Distribution of Cobia and Spanish Mackerel from the Galveston, Texas, Reef Fish Observer Program 2006-2011 | J Isely and N Cummings | |
| SEDAR28-DW12 | Estimated conversion factors for calibrating MRFSS charterboat landings and effort estimates for the South Atlantic and Gulf of Mexico in 1981-1985 with For Hire Survey estimates with application to Spanish mackerel and cobia landings | V. Matter, N Cummings, J Isely, K Brennen, and K Fitzpatrick | |
| SEDAR28-DW13 | Constituent based tagging of cobia in the Atlantic | E. Orbesen | |

| | and Gulf of Mexico waters | | |
|------------------|---|--------------------|--|
| CEDADO DW14 | | V Motton | |
| SEDAR28-DW14 | Recreational Survey Data for Spanish Mackerel | V. Matter | |
| | and Cobia in the Atlantic and the Gulf of Mexico | | |
| GED A DOO DIVING | from the MRFSS and TPWD Surveys | N. D. d. i. IV | |
| SEDAR28-DW15 | Commercial Vertical Line and Gillnet Vessel | N. Baertlein, K. | |
| | Standardized Catch Rates of Spanish Mackerel in | McCarthy | |
| | the US Gulf of Mexico, 1998-2010 | | |
| SEDAR28-DW16 | Commercial Vertical Line Vessel Standardized | K. McCarthy | |
| | Catch Rates of Cobia in the US Gulf of Mexico, | | |
| | 1993-2010 | | |
| SEDAR28-DW17 | Standardized Catch Rates of Spanish Mackerel | K. McCarthy | |
| | from Commercial Handline, Trolling and Gillnet | | |
| | Fishing Vessels in the US South Atlantic, | | |
| | 1998-2010 | | |
| SEDAR28-DW18 | Standardized catch rates of cobia from commercial | K. McCarthy | |
| | handline and trolling fishing vessels in the US | | |
| | South Atlantic, 1993-2010 | | |
| SEDAR28-DW19 | MRFSS Index for Atlantic Spanish mackerel and | Drew et al. | |
| | cobia | | |
| SEDAR28-DW20 | Preliminary standardized catch rates of Southeast | NMFS Beaufort | |
| | US Atlantic cobia (Rachycentron canadum) from | | |
| | headboat data. | | |
| SEDAR28-DW21 | Spanish mackerel preliminary data summary: | Boylan and Webster | |
| | SEAMAP-SA Coastal Survey | | |
| SEDAR28-DW22 | Recreational indices for cobia and Spanish | Bryan and Saul | |
| | mackerel in the Gulf of Mexico | | |
| SEDAR28-DW23 | A review of Gulf of Mexico and Atlantic Spanish | Palmer, DeVries, | |
| | mackerel (Scomberomorus maculatus) age data, | and Fioramonti | |
| | 1987-2011, from the Panama City Laboratory, | una i ioiumona | |
| | Southeast Fisheries Science Center, NOAA | | |
| | Fisheries Service | | |
| SEDAR28-DW24 | SCDNR Charterboat Logbook Program Data, | Errigo, Hiltz, and | |
| | 1993 - 2010 | Byrd | |
| | | Dylu | |
| SEDAR28-DW25 | South Carolina Department of Natural Resources | Hiltz and Byrd | |
| | State Finfish Survey (SFS) | J ii | |
| | | | |
| SEDAR28-DW26 | Cobia bycatch on the VIMS elasmobranch | Parsons et al. | |
| | longline survey:1989-2011 | | |
| | Documents Prepared for the Assessment Workshop |) | |
| · | | | |
| SEDAR28-AW01 | Florida Trip Tickets | S. Brown | |
| GED A DOC ANYOG | GED ID 30 G | NI (FG D | |
| SEDAR28-AW02 | SEDAR 28 Spanish mackerel bycatch estimates | NMFS Beaufort | |
| | | | |

| | from US Atlantic coast shrimp trawls | |
|--------------|---|---------------------------------|
| | Documents Prepared for the Review Workshop | |
| SEDAR28-RW01 | The Beaufort Assessment Model (BAM) with application to cobia: mathematical description, implementation details, and computer code | Craig |
| SEDAR28-RW02 | Development and diagnostics of the Beaufort assessment model applied to Cobia | Craig |
| SEDAR28-RW03 | The Beaufort Assessment Model (BAM) with application to Spanish mackerel: mathematical description, implementation details, and computer code | Andrews |
| SEDAR28-RW04 | Development and diagnostics of the Beaufort assessment model applied to Spanish mackerel | Andrews |
| | Final Assessment Reports | |
| SEDAR28-SAR1 | Assessment of Spanish mackerel in the US South Atlantic | To be prepared by SEDAR 28 |
| SEDAR28-SAR2 | Assessment of Spanish mackerel in the US Gulf of Mexico | To be prepared by SEDAR 28 |
| SEDAR28-SAR3 | Assessment of cobia in the US South Atlantic | To be prepared by SEDAR 28 |
| SEDAR28-SAR4 | Assessment of cobia in the US Gulf of Mexico | To be prepared by SEDAR 28 |
| | Reference Documents | I |
| SEDAR28-RD01 | List of documents and working papers for SEDAR 17 (South Atlantic Spanish mackerel) – all documents available on the SEDAR website | SEDAR 17 |
| SEDAR28-RD02 | 2003 Report of the mackerel Stock Assessment Panel | GMFMC and SAFMC, 2003 |
| SEDAR28-RD03 | Assessment of cobia, <i>Rachycentron canadum</i> , in the waters of the U.S. Gulf of Mexico | Williams, 2001 |
| SEDAR28-RD04 | Biological-statistical census of the species entering fisheries in the Cape Canaveral area | Anderson and Gehringer, 1965 |
| SEDAR28-RD05 | A survey of offshore fishing in Florida | Moe 1963 |

| SEDAR28-RD06 | Age, growth, maturity, and spawning of Spanish mackerel, <i>Scomberomorus maculates</i> (Mitchill), from the Atlantic Coast of the southeastern United States | Schmidt et al. 1993 |
|--------------|---|-------------------------------------|
| SEDAR28-RD07 | Omnibus amendment to the Interstate Fishery Management Plans for Spanish mackerel, spot, and spotted seatrout | ASMFC 2011 |
| SEDAR28-RD08 | Life history of Cobia, <i>Rachycentron canadum</i> (Osteichthyes: Rachycentridae), in North Carolina waters | Smith 1995 |
| SEDAR28-RD09 | Population genetics of cobia <i>Rachycentron</i> canadum: Management implications along the Southeastern US coast | Darden et al, 2012 |
| SEDAR28-RD10 | Inshore spawning of cobia (Rachycentron canadum) in South Carolina | Lefebvre and Denson, 2012 |
| SEDAR28-RD11 | A review of age, growth, and reproduction of cobia <i>Rachycentron canadum</i> , from US water of the Gulf of Mexico and Atlantic ocean | Franks and Brown- Peterson, 2002 |
| SEDAR28-RD12 | An assessment of cobia in Southeast US waters | Thompson 1995 |
| SEDAR28-RD13 | Reproductive biology of cobia, <i>Rachycentron</i> canadum, from coastal waters of the southern United States | Brown-Peterson et al. 2001 |
| SEDAR28-RD14 | Larval development, distribution, and ecology of cobia <i>Rachycentron canadum</i> (Family: Rachycentridae) in the northern Gulf of Mexico | Ditty and Shaw 1992 |
| SEDAR28-RD15 | Age and growth of cobia, <i>Rachycentron canadum</i> , from the northeastern Gulf of Mexico | Franks et al 1999 |
| SEDAR28-RD16 | Age and growth of Spanish mackerel, Scomberomorus maculates, in the Chesapeake Bay region | Gaichas, 1997 |
| SEDAR28-RD17 | Status of the South Carolina fisheries for cobia | Hammond, 2001 |
| SEDAR28-RD18 | Age, growth and fecundity of the cobia, Rachycentron canadum, from Chesapeake Bay and adjacent Mid-Atlantic waters | Richards 1967 |
| SEDAR28-RD19 | Cobia (<i>Rachycentron canadum</i>) tagging within Cheasapeake Bay and updating of growth equations | Richards 1977 |
| SEDAR28-RD20 | Synopsis of biological data on the cobia Rachycentron canadum (Pisces: Rachycentridae) | Shaffer and Nakamura 1989 |
| SEDAR28-RD21 | South Carolina marine game fish tagging program 1978-2009 | Wiggers, 2010 |
| SEDAR28-RD22 | Cobia (Rachycentron canadum), amberjack | MARFIN 1992 |

| | (Seriola dumerili), and dolphin (Coryphaena hipurus) migration and life history study off the southwest coast of Florida | |
|--------------|--|---------------------------|
| SEDAR28-RD23 | Sport fish tag and release in Mississippi coastal water and the adjacent Gulf of Mexico | Hendon and Franks 2010 |
| SEDAR28-RD24 | VMRC Cobia otolith preparation protocol | VMRC |
| SEDAR28-RD25 | VMRC Cobia otolith ageing protocol | VMRC |
| SEDAR28-RD26 | Age, growth, and reproductive biology of greater amberjack and cobia from Louisiana waters | Thompson et al. 1991 |
| SEDAR28-RD27 | Gonadal maturation in the cobia, <i>Rachycentron</i> canadum, from the northcentral Gulf of Mexico | Lotz et al. 1996 |
| SEDAR28-RD28 | Cobia (<i>Rachycentron canadum</i>) stock assessment study in the Gulf of Mexico and in the South Atlantic | Burns et al. 1998 |
| SEDAR28-RD29 | Total mortality estimates for Spanish mackerel captured in the Gulf of Mexico commercial and recreational fisheries 1983 to 2011 | Bryan 2012 |
| | | |