

Final Report: Patterns of habitat utilization by reef fish on St. Croix  
F-7-17, Study 3  
Period: 1 October 2001 to 30 September 2002

**Quantitative estimates of species composition  
and abundance of fishes, and fish  
species/habitat associations in  
St. Croix, U.S. Virgin Islands**

Dr. Wesley Toller  
Fisheries Biologist II

Division of Fish and Wildlife  
Department of Planning and Natural Resources  
US Virgin Islands

November 2002

## ABSTRACT

Reef fish communities around St. Croix, U.S. Virgin Islands were studied by visual census methods in order to identify fisheries-habitat links, to document the role of local coral reefs as essential fish habitat, and to provide a baseline for monitoring studies. Seven reef sites were surveyed by stationary fish censuses during spring and fall of 2002. In general, the fish communities were similar across sites and survey periods: all reefs supported a diverse and abundant fish assemblage that was largely replicated between spring and fall surveys. Planktivorous fish (primarily labrids and pomacentrids) dominated the fauna numerically. Herbivorous fish (acanthurids and scarids) were common and abundant at all sites. Piscivores and other specialized feeders were least abundant. Serranids were common and relatively diverse, however most observed fish were either diminutive or small species (e.g. harlequin bass, hamlets, coney, and graysby). Lutjanids were of larger size, but less common, less abundant, and less diverse than serranids. Very few large reef-associated piscivorous species were seen, and the significance of this observation is discussed.

Some variation in fish assemblages was observed among reefs. Comparison of species richness and diversity ( $H'$ ) among reefs showed a suggestive positive relationship to vertical relief, but no obvious relation to percentage live coral cover or percentage turf alga cover. Planktivorous pomacentrids predominated at the two reef sites with the highest coral cover. Acanthurids and scarids, while both nearly ubiquitous, showed a reciprocal density relationship, with scarids predominating on well-developed reefs. Chaetodontids were more abundant and diverse near a submarine canyon (Salt River). Lutjanids were not observed at Lang Bank – an offshore platform with sparse coral cover.

These results imply that at least some members of reef fish communities respond to presently unidentified biotic or abiotic differences among the coral reef habitats of St. Croix, and that the source of this variation warrants additional investigation. Additional studies are recommended to monitor changes in these reef fish communities over time, to improve assessments of rare (but recreationally and commercially important) species, and to further elucidate critical fish-habitat links.

## Table of Contents

	Page No.
Abstract	ii
Table of Contents	iii
List of Tables	iv
List of Figures	iv
INTRODUCTION	1
METHODS	1
Description of Study Sites	1
Visual Census Methods	2
RESULTS	3
DISCUSSION	5
Planktivores	6
Herbivores	6
Piscivores, Carnivores and Specialized Invertebrate Feeders	7
Recommendations	8
ACKNOWLEDGEMENTS	8
LITERATURE CITED	8
APPENDICES	
1. Distribution of observations on 85 fish species across seven St. Croix reef sites.	19-23
2. A. Fish Census Data from Lang Bank	24-25
B. Fish Census Data from Jacks & Isaacs Bay	26-28
C. Fish Census Data from East Wall, Salt River Canyon	29-31
D. Fish Census Data from West Wall, Salt River Canyon	32-34
E. Fish Census Data from Eagle Ray	35-38
F. Fish Census Data from Cane Bay	39-41
G. Fish Census Data from Sprat Hole	42-44

	<b>List of Tables</b>	Page No.
1	Description of St. Croix reef sites for fish census study.	15
2	Fish census effort	15
3	Total number of fish and species observed during stationary fish censuses	16
4	The 25 most abundant fish species observed in this study.	17-18

	<b>List of Figures</b>	Page No.
1	Map of St. Croix showing the seven reef sites where fish visual censuses were conducted.	11
2	Summary of fish community structure at seven St. Croix reef sites. A. Diversity ( $H'$ ) B. Species Richness C. Fish Abundance	12 12 12
3	Fish abundance by family at seven St. Croix reef sites.	12
4	Abundance of fishes representing 8 families at 7 St. Croix reef sites. A. Labridae (wrasses) B. Pomacentridae (damselfishes) C. Acanthuridae (surgeonfishes) D. Scaridae (parrotfishes) E. Serranidae (groupers) F. Lutjanidae (snappers) G. Haemulidae (grunts) H. Chaetodontidae (butterflyfishes) & Pomacanthidae (anglefish)	13 13 13 13 14 14 14 14

## INTRODUCTION

Coral reefs of the U.S. Virgin Islands support a diverse assemblage of fishes. In addition to their ecological roles on reefs, these fish have economic importance to local recreational and commercial fisheries (Appledoorn et al. 1992).

Management of reef fish resources depends upon sound information about the status of local populations, and knowledge of critical links between habitat and fish life-history (i.e. essential fish habitats). Thus a first step is to simply establish the fine-scale patterns of fish distribution among habitats. A descriptive study, such as this one, can provide valuable information about habitat-species associations, which may ultimately lead to identification of critical links.

A second objective of management must be to monitor resources, so that the trajectory of populations through time can be evaluated. A thorough and descriptive baseline survey, whether it represents a natural or disturbed community (Jackson 1997), is the essential starting point for any monitoring program. On St. Croix, for example a proposed Marine Protected Area (MPA) will encompass the nearshore, eastern end of the island. Implementation of a broad "No-Take" zone may conceivably influence the structure of entire reef fish communities located in this area. Monitoring will be instrumental in gauging the effects of this proposed MPA on fish communities. Alternatively, on a scale of individual species, monitoring may allow managers to determine whether specific fisheries regulations, such as localized closures to protect spawning aggregations, are effective at restoring depleted stocks (sensu Tobias et al. 1988).

## METHODS

### Description of Study Sites

Seven reef sites around the island of St. Croix, U. S. Virgin Islands were selected for this study (see Figure 1): Lang Bank (LB), Jacks and Isaacs Bay (JI), East Wall (EW) of Salt River Canyon, West Wall (WW) of Salt River Canyon, Eagle Ray (ER) on Long Reef, Cane Bay (CB), and Sprat Hole (SH). Each of these reefs is surveyed annually by the Center for Marine and Environmental Studies, University of the Virgin Islands, as part of their coral reef monitoring program. Those studies provide much of the descriptive information on sessile reef biota, as presented in Table 1 (Nemeth and Herzlieb 2002). Detailed geological, biological, and spatial descriptions of these reef sites have been made previously, either in a general sense (Hubbard 1989a), in detail (at CB - Hubbard 1989b; at EW and WW - Hubbard 1989c) or at comparable nearby reefs (at JI - Hubbard 1989d). Briefly, the seven reef sites all represent shallow water (< 15 m depth) hard bottom substrates situated on the insular shelf of St. Croix. Despite geographic proximity, there is considerable variation among reefs in terms of benthic flora and fauna, physical structure, exposure to prevailing currents and wave forces, and adjacent habitats (to name but a few of the variables). For example, the development of scleractinian coral communities varies considerably among sites, from about 5% living coral cover at LB to >24% at SH (Nemeth and Herzlieb 2002). Some of the reefs have previously been classified (Hubbard 1989a-d) as barrier-type reefs (ER), fringing reefs (JI), shelf edge reefs (CB, SH), or submarine

hardgrounds (LB). In the present study, only those attributes presented in Table 1 were considered in relation to reef fish community structure.

### Visual Census Methods

The stationary fish census method (Bohnsack and Bannerot 1986) was used to assess reef fish community structure. Compared to other visual census methods (belt transects, timed random swim methods), the stationary count offers the advantage of estimating fish community structure as a function of density per unit area for each observed species, while enabling size (length) estimates of individuals. Generally, results from stationary point counts are comparable to results obtained from belt transects (e.g. Bortone et al. 1989), and the two methods differ primarily in duration (belt transects are faster) and area surveyed per replicate. A typical belt transect of 30 m x 2 m yields 60 m<sup>2</sup> whereas a 15 m wide stationary census (used in this study) yields 176 m<sup>2</sup> per replicate. However neither of these two methods is likely to enumerate all of the species present within a given area (i.e. to generate a species list) - that purpose is best served using replicated timed random diver swims (Kimmel 1985) or related methods (see Rogers et al 2001).

The census protocol is only slightly modified from Bohnsack and Bannerot (1986), and a brief description is provided here. A 15 m diameter census "cylinder" was defined by transect tape with the observer positioned in the center (7.5 m mark). Fish within this cylinder were censused as follows. During an initial 5-minute "listing" period, the names of all observed fish species were recorded onto pre-printed data forms. At the end of the listing period, the observer began enumerating all individuals of each species, working from the bottom of the list upward and making one 360° sweep. Strict adherence was made to the 5-minute listing period. New species observed during the enumeration period (e.g. fish that swam into the cylinder after 5 minutes of listing) were not included in the stationary count data. Divers estimated fish total length to the nearest cm, using a measuring "T-bar" for underwater visual reference. Maximum, minimum, and average size was recorded for each species. Divers also recorded a brief description/sketch of habitat features within the census area. All fish were identified to species, however gobies (Gobiidae) and blennies (Blenniidae) were excluded from census counts. To insure that our fish census areas coincided with ongoing UVI coral surveys, but that replicates did not cover overlapping areas, census stations were marked with a small piece of rebar driven into nonliving reef substrate.

Surveys were conducted during two sampling periods: Spring (May – June, 2002) and Fall (September, 2002). For each reef site, six replicate censuses were made [this is a level of replication thought to adequately sample the species representing > 90% of all individuals at a site (Bohnsack and Bannerot 1986)] except for LB (Spring) where only four replicate censuses were conducted due to unfavorable weather conditions. Six replicate censuses surveyed a total area of 1,056 m<sup>2</sup> per reef site. A summary of census dates, number of replicates, and census duration is presented in Table 2.

Data from this study were analyzed using Microsoft Excel software.

## RESULTS

A total of 35,322 fish were sighted during the surveys (Table 3). More fish were observed in fall (16,859) than spring (18,463). The number of fish observed at each site ranged from 1,385 (JI, fall) to 5,443 (CB, fall). A total of 85 fish species representing 27 families were sighted (Table 3; Appendix 1). More species were seen during fall (77 species) than during spring (72 species; Table 3). The number of species observed at each site during spring or fall ranged from 24 (LB, spring) to 42 (WW, ER, CB; all fall) and the cumulative number of species observed at each reef site ranged from 32 (LB) to 52 (ER).

In general, fish assemblages present at each reef site were similar between spring and fall. Average fish abundance (Figure 2A) remained relatively constant at 6 of the reef sites. At CB, an increase in average abundance in fall was due to a large influx of small (~3 to 5 cm TL) creole wrasses (*Clepticus parrae*; see Appendix 2F). Average species richness also remained relatively constant between spring and fall (Figure 2B). However the Shannon-Weiner diversity index ( $H'$ ), computed as an average per census, showed greater variability between survey periods (Figure 2C; LB and ER), suggesting that community composition had changed in some cases. At LB, change was attributed to a large (30%) decrease in abundance of the numerically dominant species, the bluehead wrasse, *Thalassoma bifasciatum*, and a 3-fold increase in bicolor damselfish (*Stegastes partitus*) abundance. At ER, the fall increase in diversity corresponded to a 40% decrease in abundance of the bluehead wrasse and the appearance of large schools of brown chromis, *Chromis multilineata* (an increase from 1.7 to 43.3 fish/census; Appendix 2E).

In contrast to temporal comparisons (i.e. spring vs. fall), comparisons across reef sites showed greater dissimilarity in fish assemblages (Figure 2). Average abundance (Figure 2A) was highest at CB and SH, sites with the highest coral cover, but also at LB where coral cover is sparse. Average species richness varied from ~15 to ~22 species observed/census (Figure 2B) and LB had the fewest species while ER, CB and SH had the most. Diversity indices showed a similar trend (Figure 2C).

The 25 most abundant species observed in this study are present in Table 4. The bluehead wrasse alone accounted for 38.6% of all fish sightings. The five most abundant species were the bluehead wrasse, creole wrasse, bicolor damselfish, brown chromis, and blue chromis (*Chromis cyanaea*) - all primarily planktivorous species. Together, they accounted for 87.2% of all fish sightings. Despite comparably high abundance levels shared among these five species, their distribution and frequency of occurrence were different (Table 4). While the strongly reef-associated bluehead wrasse and bicolor damselfish were ubiquitous (present at each reef site) and common (frequency > 97% of censuses), the schooling creole wrasse and brown chromis were more patchy in distribution (Appendix 1) and less common (frequency < 45% of censuses).

A detailed account of species is beyond the scope of the present work (see Appendix 2A to G). Here, an analysis of reef fish community composition is presented based upon taxonomic divisions (families) that delineate eight commercially, recreationally or ecologically important fish groups. These families are: Labridae (wrasses), Pomacentridae (damselfishes), Acanthuridae (surgeonfishes), Scaridae (parrotfishes), Serranidae (groupers), Lutjanidae

(snappers), Haemulidae (grunts), Chaetodontidae (butterflyfishes) and Pomacanthidae (angelfishes). The latter two families were combined as a single group (butterflyfish and angelfish) for these analyses. Use of these taxonomic divisions is justified because: (1) each family loosely represents a trophic (ecological) group, and (2) the families collectively represented over 95% (usually >98%) of fish observed at each reef site (e.g. Table 4; Appendices 2A to G).

Fish abundance by family is summarized in Figure 3, where data were pooled from all seven reef sites. Average fish abundance was similar between spring and fall, however abundance varied considerably among families. Labrids and Pomacentrids were over 10 times more abundant than all other families (~ 100 fish observed/census). Acanthurids and scarids had similar abundance levels of ~ 10 fish/census. The abundance of serranids, lutjanids, haemulids, chaetodontids and pomacanthids was generally low (1 to 4 fish/census) and typically quite variable within and among sites (see below).

To examine fish community structure across the seven reef sites, the same eight fish families were utilized as a basis for analyses (Figure 4). Labrids were abundant at all sites (average 242 fish/census) but highly variable within and among sites (Figure 4a) [as noted above, an exceptionally high abundance of creole wrasses was recorded at CB in spring]. Pomacentrids were also abundant (average 148 fish/census), and showed the greatest variation among sites. The highest abundances were seen at CB and SH, where > 300 fish/census were recorded in spring and fall (Figure 4B). Blue chromis and brown chromis were the most abundant pomacentrids at CB and SH (Appendix 2F and G). Herbivorous fish were less abundant than planktivores. The abundance of acanthurids (average 10.7 fish/census) and scarids (average 9.6 fish/census) was also variable across the reef sites (Figures 4C and D). Acanthurids had their highest abundances at WW, EW, JI, and LB and were least abundant at CB and SH. Scarids had their highest abundance at SH.

Predatory fishes (piscivores, carnivores) are represented by the serranids, lutjanids and haemulids. As a family, serranids had low abundance (average 4 fish/census) that showed little variation among reef sites (Figure 4E). Serranid diversity was variable among reef sites. The coney, *Cephalopholis fulvus*, was uniformly distributed across reefs (Appendix 1) and relatively common (Appendix 2A-G), however the graysby, *C. cruentatus*, was more frequent in habitats of high topographic complexity (ER, CB, SH). The red hind, *Epinephelus guttatus*, was observed only infrequently at 4 sites (Appendices 1 and 2). Hamlets (genus *Hypoplectrus*) also showed highest diversity levels at SH, ER, EW and WW but not CB. In contrast to serranids, lutjanids showed greater variation in abundance and distribution. The yellowtail snapper, *Ocyurus chrysurus*, and the mahogany snapper, *Lutjanus mahogoni*, were the most common and abundant snappers, however the schoolmaster, *L. apodus*, and the mutton snapper, *L. analis*, were also observed (Appendix 1). No lutjanids were recorded at LB. Haemulid abundance was relatively uniform (average 2.4 fish/census) across reef sites (Figure 4G).

The combined group of Chaetodontidae and Pomacanthidae represent long-lived reef residents with specialized feeding habits (e.g. feeding on cnidarians, sponges). Abundance and diversity of this group varied among reef sites (Figure 4H) and was generally low (2.3 fish/census).

Chaetodontids were most frequent and abundant at EW, WW and ER. It is interesting that EW and WW had the highest diversity of butterflyfish (4 and 3 species, respectively) whereas other reef sites had 2 or 1 species. Only the banded butterflyfish, *Chaetodon striatus*, was recorded from LB and JI, whereas only the foureye butterflyfish, *C. capistratus*, was recorded from CB and SH (Appendix 2).

## DISCUSSION

The reefs investigated in this study supported diverse and abundant fish communities. This is not surprising - previous studies have documented well over 200 reef-associated species in St. Croix waters (Clavijo et al. 1980), and over 100 species may occur on a single patch reef (Gladfelter and Gladfelter 1978). For the following discussion, it is important to note that the stationary visual census method does not capture total reef fish diversity (i.e. Appendices 2A to G do not represent complete species lists). Rather, these are quantitative data on the most conspicuous and abundant components of each fish community. The absence of a given species from our observations should be considered "relative" absence. Nonetheless, the average number of species observed per census is a quantitative estimate of community species richness, and a useful measure for comparing reef sites.

The goal of the present study was to document, in an exploratory manner, species-habitat links of coral reef fish populations. This experimental design allows only a preliminary analysis of the relationship between individual fish species (or fish communities) and specific reef attributes (e.g. % live coral, vertical relief, benthic invertebrate community composition, wave exposure, prevailing currents, etc). When fish communities were compared across the seven reef sites, there was no obvious relation of average fish abundance to percentage live coral nor percentage turf algal cover (not shown). Planktivore abundance (blue chromis and brown chromis), on the other hand, may be positively related to percentage live coral cover (e.g. CB and SH; Figure 4B). Species richness showed a more suggestive positive relation to coral cover, although ER represents an important exception. At ER, coral cover is relatively low, but vertical relief is high (a spur and groove area formed of *Montastraea* skeletons). A relatively high number of fish species was observed at ER. This suggests that the richness of St. Croix reef fish communities may be influenced more by three-dimensional architecture (topographic complexity) of the reef than by the presence of living (versus dead) corals. This idea is not new (Roberts and Ormond 1987), and it has been suggested by results from long-term studies of fish populations at Buck Island, St. Croix (Gladfelter et al. 1992), where hurricane damage and disease permanently reduced live coral cover (*Acropora palmata*), but left habitat structural complexity relatively intact. Despite the loss of living coral, resident fish populations recovered to pre-hurricane densities.

Nemeth and Herzlieb (2002) suggested that reef sites along St. Croix's northern coast harbor a greater abundance of reef fish than the other reef sites. Our results did not corroborate this finding (see Figure 2 and Figure 4). Neither that study nor this one included sufficient replication of reef sites, adequate spatial coverage of insular reefs, nor control over confounding variables (e.g. topographic complexity) to reach a definitive conclusion. Nonetheless, previous

studies indicate that recruitment patterns are highest on St. Croix's northwestern and southeastern shores (Caselle and Warner 1996, and Sweare 2002). Recruitment studies, in conjunction with community assessments, may clarify a postulated relationship between larval supply and resident reef fish abundance.

### **Planktivores:**

The reef fish communities were numerically dominated by labrids (bluehead wrasse, creole wrasse) and planktivorous pomacentrids (bicolor damselfish, blue chromis, and brown chromis), which were collectively 10 to 100-fold more abundant than all other species. The bluehead wrasse, perhaps the most abundant of all West Indian reef fishes, is a facultative planktivore, and the remaining species feed primarily on zooplankton (Randall 1968).

Large populations of planktivores form a major trophic link between coral reef habitats and open-water communities (e.g. Hobson 1991). Undoubtedly, a large number of these fish fall prey to resident and reef associated predatory fishes. Between spring and fall, bluehead wrasse abundance decreased by ~ 20%, which is suggestive of heavy predation. However we found no relation between planktivore abundance and piscivore abundance (compare Figure 4A and B with 4E and F). Some authors have argued that, rather than predation, the primary trophic link between planktivores and coral reefs is through production of feces, which is then consumed by a wide variety of reef fish and invertebrates (Robertson 1982). Nonetheless, the role of planktivores as forage for commercially and recreationally important species should be investigated further.

### **Herbivores:**

Herbivorous fish were common, moderately abundant, and relatively diverse on all seven reefs. For acanthurids and scarids, average abundance was similar (9.6 fish/176 m<sup>2</sup> census and 10.7 fish /176 m<sup>2</sup> census, respectively). In 2001, Nemeth and Herzlieb (2002) censused fish from the same sites using belt transects. When normalized to 100m<sup>2</sup>, the density of acanthurids observed in our study was slightly lower than in theirs (6.0 vs 9.0 fish/100 m<sup>2</sup>) and the density of scarids observed in this study was substantially lower (5.5 vs 13 fish/100m<sup>2</sup>). The apparent decline in parrotfish populations is unexplained. Methodological differences between the studies may account for this discrepancy. Alternatively, scarid populations may be in decline due to overfishing – parrotfishes are actively sought by a local gillnet fishery. Given the important trophic role that scarids play in maintaining scleractinian coral growth and diversity (e.g. Pennings 1997), local parrotfish populations should be monitored carefully in the future.

In this study, parrotfishes (especially the redband parrotfish, *Sparisoma aurofrenatum*, and the princess parrotfish, *Scarus taeniopterus*) and surgeonfishes (ocean surgeonfish, *Acanthurus bahianus*, and blue tang, *A. coeruleus*) were nearly ubiquitous among the reef sites. Analysis of their abundance patterns (Figure 4C and D) suggests that their densities were reciprocal. Acanthurids were more abundant on low relief, low coral cover sites (LB, JI, EW, WW) while Scarids were more abundant on reefs with greater topographic complexity (ER, CB, SH). This

pattern may arise from differences in feeding mode: acanthurids tend to browse over soft bottoms while scarids are scrapers of hardbottom substrates (e.g. Choat 1991).

### **Piscivores, Carnivores and Specialized Invertebrate Feeders:**

A diversity of predatory fishes was recorded during censuses. The observed species varied in their degree of reef association. For example carangids (bar jack, *Carynx ruber*, and blue runner, *C. crysos*) and scombrids (cero mackerel, *Scomberomorus regalis*) were infrequently observed, in part due to their highly mobility and their loose association with reefs. On the other hand, the coney appears to represent a strongly reef-associated resident species that displayed no habitat preference. This species was the most abundant serranid, occurring with moderate to high frequency at all studied reef sites. The coney is important to both the recreational and commercial fishery of the USVI (Appledoorn et al. 1992; Bolden 1994). The graysby, another frequently observed serranid, showed habitat preference for reefs with a high degree of topographic complexity.

As adults, lutjanids are largely residents of reef habitats. However, the juveniles of many lutjanid species, as well as haemulids and chaetodontids, depend upon "nursery habitats" during their early post-settlement life. On St. Croix for example, Adams and Tobias (1993) documented the importance of an estuarine environment (Salt River) for juvenile schoolmasters (*L. apodus*), french grunts (*Haemulon flavolineatum*), and foureye butterflyfish (*C. capistratus*). Likewise, St. Croix embayments (comprised of patch reef, sand and seagrass habitats) appear to function as important nursery habitats for French grunts and yellowtail snapper (Mateo and Tobias 2001). In our study, adults of these species (but not juveniles) were commonly observed on reefs. For example, adult French grunts were among the most frequently observed fish (~ 80% of all censuses), but no juveniles were observed at these reef sites. For such species, these results serve to underscore the connectivity between adult populations on reefs and juvenile populations residing in nursery habitats.

Nonetheless, it is clear that many of the long-lived reef species were relatively rare. Commercially and recreationally important species such as the red hind (*E. guttatus*) or mutton snapper (*L. analis*) were observed only infrequently (6.1% and 3.7% of censuses, respectively). Some of the larger non-piscivorous reef species such as the french angelfish (*Pomacanthus paru*), queen angelfish (*Holacanthus ciliaris*), or puddingwife (*Halichoeres radiatus*) were also rare in our studies. Formerly, all of the above species were considered "common" on St. Croix reefs (Clavijo et al 1980). Overfishing is the likely explanation for the relative rarity of these species. Establishment of a marine protected area should augment populations of long-lived fish species with high susceptibility to fishing pressures (Bohnsack 1993), such as those discussed above.

Crosby and Reese (1996) have suggested that butterflyfish density is an indicator of reef health. In our study, butterflyfish reached their greatest abundance and diversity at Salt River canyon. This distribution pattern was unique among fishes in our study. Localized abundance at Salt River Canyon may be due to reduced fishing pressure in the immediate vicinity - chaetodontids

are readily caught in fish traps (DFW, unpubl. SEAMAP study) or to the proximity of nursery habitat (Salt River Estuary; see Adams and Tobias 1993), which may augment local butterflyfish populations. Alternatively, the chaetodontids species may share a habitat preference for some unidentified feature of the submarine canyon itself.

### **Recommendations:**

Continued monitoring of these seven reef sites should be conducted semi-annually. One or two additional reef sites should be added to include reef fish communities along the southern insular shelf of St. Croix. The visual census method used here is robust and reproducible and should therefore be used in future monitoring efforts. However, a second assessment method, such as the timed random diver swim (TRDS) method, should also be included in the protocol. Sufficient TRDS replicates (performed on an annual or biannual basis) would establish a species lists for each reef site and enable more powerful inferences regarding rare species.

Population assessments of large, resident reef fishes were compromised by the rarity of these fish. This is particularly troublesome because many are commercially and recreationally important species. However, for species such as the Nassau grouper (*Epinephelus striata*), goliath grouper (*E. itajara*), tiger grouper (*Mycteroperca tigris*), black grouper (*M. bonaci*), yellowfin grouper (*M. venenosa*), mutton snapper (*Lutjanus analis*), and cubera snapper (*L. cyanopterus*), populations are probably too low to adequately assess them by any of the aforementioned visual census methods. Instead, future monitoring efforts should be directed towards censusing these species during their annual spawning aggregations (e.g. Colin 1996). Aggregation sites are already known for at least two species on St. Croix (*L. analis* and *E. guttatus*) and these could be monitored using diver surveys or ROV.

### **ACKNOWLEDGMENTS**

William Tobias provided technical guidance and assisted in many of the field surveys. Hector Rivera and Willy Ventura were boat operators. A special thanks goes to Dianne "Dee" Osinski and Robert Daniel Stone for their participation in (and enthusiasm for) the fish counts.

### **LITERATURE CITED**

- Adams, A.J., and W.J. Tobias. 1993. Red mangrove prop-root habitat as a finfish nursery area. A case study of Salt River, St. Croix, U.S.V.I. *Proc. Gulf Carib. Fish. Inst.* 46:22-47
- Appledoorn, R., J. Beets, J. Bohnsack, S. Bolden, D. Matos, S. Meyers, A. Rosario, Y. Sadovy, and W. Tobias. 1992. Shallow water reef fish stock assessment for the U.S. Caribbean. NOAA Tech. Memo. NMFS:SEFSC-304, 70 pp.
- Bohnsack, J.A. 1993. Marine reserves: they enhance fisheries, reduce conflicts, and protect resources. *Oceanus* 36(3): 63-71

- Bohnsack, J.A., and S.P. Bannerot. 1986. A stationary visual census technique for quantitatively assessing community structure of coral reef fishes. NOAA Technical Report NMFS 41, U.S. Dept of Commerce, 15 pp.
- Bortone, S.A., J.J. Kimmel, and C.M. Bundrick. 1989. A comparison of three methods for visually assessing reef fish communities: time and area compensated. *Northeast Gulf Sci.* 10(2):85-96.
- Caselle, J.E., and R.R. Warner. 1996. Variability in recruitment of coral reef fishes: the importance of habitat at two spatial scales. *Ecology* 77: 2488-2504.
- Choat, J.H. 1991. The biology of herbivorous fishes on coral reefs. Pp. 120-155 In: P. F. Sale (ed.) *The Ecology of Fishes on Coral Reefs*. Academic Press, Inc., New York, 754 pp.
- Clavijo, I.E., J.A. Yntema, and J.C. Ogden. 1980. An annotated checklist of the fishes of St. Croix U. S. Virgin Islands. West Indies Laboratory, St. Croix, USVI, 49 pp.
- Colin, P.L. 1996. Longevity of some coral reef fish spawning aggregations. *Copeia* 1996(1):189-2.
- Crosby, M.P., and E.S. Reese. 1996. A manual for monitoring coral reefs with indicator species: butterflyfishes as indicators of change on Indo Pacific reefs. Office of Ocean and Coastal Resource Management, NOAA, Silver Springs MD, 45 pp.
- Gladfelter, W.B., and E.H. Gladfelter. 1978. Fish community structure as a function of habitat structure on West Indian patch reefs. *Rev. Biol. Trop.* (suppl. 1):65-84.
- Gladfelter, E.H., J.C. Bythell, and Z.M. Hillis. 1992. Changes in fish assemblage structure at Buck Island, St. Croix, U. S. Virgin Islands from 1980-1990: an indication of predictability in coral reef fish assemblages based on known habitat changes? Pages 14-29 In: *Ecological Studies of Buck Island Reef National Monument, St. Croix, U. S. Virgin Islands. Part II*. Report prepared for U.S. Department of the Interior, National Park Service.
- Hobson, E.S. 1991. Trophic relationships of fishes specialized to feed on zooplankters above coral reefs. Pp. 69-95 In: P. F. Sale (ed.) *The Ecology of Fishes on Coral Reefs*. Academic Press, Inc., New York, 754 pp.
- Hubbard, D. 1989a. Modern carbonate environments of St. Croix and the Caribbean: A general overview. Pages 85-94 In: D. Hubbard (ed.) *Terrestrial and marine geology of St. Croix, U. S. Virgin Islands. Sp. Publ. No. 8*, West Indies Laboratory, St. Croix, USVI, 213 pp.
- Hubbard, D. 1989b. The shelf-edge reefs of Davis and Cane Bays, Northwestern St. Croix, U.S.V.I. Pages 167-179 In: D. Hubbard (ed.) *Terrestrial and marine geology of St. Croix, U. S. Virgin Islands. Sp. Publ. No. 8*, West Indies Laboratory, St. Croix, USVI, 213 pp.

- Hubbard, D. 1989c. Depositional environments of Salt River Estuary and submarine canyon, St. Croix, U.S.V.I. Pages 181-196 In: D. Hubbard (ed.) Terrestrial and marine geology of St. Croix, U. S. Virgin Islands. Sp. Publ. No. 8, West Indies Laboratory, St. Croix, USVI, 213 pp.
- Hubbard, D. 1989d. Depositional environments of the Turner Hole reef complex. Pages 155-160 In: D. Hubbard (ed.) Terrestrial and marine geology of St. Croix, U. S. Virgin Islands. Sp. Publ. No. 8, West Indies Laboratory, St. Croix, USVI, 213 pp.
- Jackson, J.B.C. 1997. Reefs since Columbus. *Coral Reefs* 16:S23-S32.
- Kimmel, J.J. 1985. A new species-time method for visual assessment of fishes and its comparison with established methods. *Env. Biol. Fishes* 12:23-32.
- Mateo, I. and W.J. Tobias. 2001. The role of nearshore habitats as nursery grounds for juvenile fishes on the northeast coast of St. Croix. *Proc. Gulf Carib. Fish. Inst.* 52:512-530.
- Nemeth, R.S., and S. Herzlieb. 2002. Video monitoring assessment of coral reefs in proposed marine parks St. Croix, United States Virgin Islands. Center for Marine and Environmental Studies, University of the Virgin Islands, St. Thomas, USVI. In: U. S. Virgin Islands Coral Reef Monitoring Project, Year 1 Final Report. USVI Department of Planning and Natural Resources, Division of Coastal Zone Management.
- Pennings, S.C. 1997. Indirect interactions on coral reefs. Pp. 249-272 In: C. Birkeland (ed) Life and Death of Coral Reefs. Chapman & Hall, New York, 536 pp.
- Randall, J.E. 1968. Caribbean Reef Fishes. T.F.H. Publications, Inc., Jersey City, NJ, 318 pp.
- Roberts, C.M. and R.F.G. Ormond. 1987. Habitat complexity and coral reef fish diversity and abundance on Red Sea fringing reefs. *Mar. Ecol. Prog. Ser.* 41:1-8.
- Robertson, D.R. 1982. Fish feces as food on a Pacific coral reef. *Mar. Ecol. Prog. Ser.* 7:253-265.
- Rogers, C.S., G. Garrison, R. Grober, Z. Hillis, and M.A. Franke. 2001. Coral Reef Monitoring Manual for the Caribbean and Western Atlantic. 2<sup>nd</sup> Ed., Virgin Islands National Park.
- Swearer, S.E. 2000. Self-recruitment in coral reef fish populations. Ph.D. dissertation, University of California, Santa Barbara, 151 pp.
- Tobias, W., E. Telemaque, and M. Davis. 1988. Buck Island fish and shellfish populations. Biosphere Reserve Research Report No. 26, Virgin Islands National Park, 27 pp.

Figure 1. Map of St. Croix showing the seven reef sites where fish visual censuses were conducted. Abbreviations are as follows: Lang Bank (LB), Jacks and Isaacs Bay (JI), East Wall (EW) of Salt River Canyon, West Wall (WW) of Salt River Canyon, Eagle Ray (ER) on Long Reef, Cane Bay (CB), and Sprat Hole (SH).






Figure 2. Summary of fish community structure at seven St. Croix reef sites.  
 A.



C.

Fish Abundance




Figure 3. Fish abundance by family at seven St. Croix reef sites. Error bars show range of average abundance values across sites.

Summary of Fish Abundance






Figure 4. Abundance of fishes representing 8 families at 7 St. Croix reef sites. Families are: A. Labridae (wrasses), B. Pomacentridae (damselfishes), C. Acanthuridae (surgeonfishes), D. Scaridae (parrotfishes), E. Serranidae (groupers), F. Lutjanidae (snappers), G. Haemulidae (grunts), and H. Chaetodontidae (butterflyfishes) and Pomacanthidae (anglefishes). Error bars represent standard deviation.


A



B.



C.



D.






Figure 4 (continued). Abundance of fishes representing 8 families at 7 St. Croix reef sites.  
 Families are: A. Labridae (wrasses), B. Pomacentridae (damselfishes), C. Acanthuridae (surgeonfishes), D. Scaridae (parrotfishes), E. Serranidae (groupers), F. Lutjanidae (snappers), G. Haemulidae (grunts), and H. Chaetodontidae (butterflyfishes) and Pomacanthidae (anglefishes). Error bars represent standard deviation.


E.



F.



G.



H.




Table 1. Description of St. Croix reef sites for fish census study.

Reef Site	Lang Bank	Jack's & Isaac's	East Wall (Salt River)	West Wall (Salt River)	Eagle Ray (Long Reef)	Cane Bay	Sprat Hole
Abbreviation	LB	JI	EW	WW	ER	CB	SH
Location	East	Southeast	North	North	North	Northwest	West
Latitude	N 17° 44.448'	N 17° 44.586'	N 17° 47.221'	N 17° 47.116'	N 17° 45.688'	N 17° 46.433'	N 17° 44.038'
Longitude	W 64° 32.186'	W 64° 34.310'	W 64° 45.445'	W 64° 45.564'	W 64° 41.929'	W 64° 48.810'	W 64° 53.722'
Min. Depth	44	36	32	24	22	24	30
Max. Depth	48	49	45	27	35	36	35
Avg. Depth	45.8	41.7	36.8	25.3	28.0	28.8	32.3
Reef Description <sup>1</sup>	Submarine Hardgounds	Fringing Reef	Sloping Cobble, Canyon Wall	Steep Canyon Wall	Barrier Reef	Shelf-Edge Reef	Shelf-Edge Reef
Vertical Relief <sup>2</sup>	Low	Moderate	Moderate	Low & High	High	High	High
% Cover Live Coral <sup>3</sup>	~ 5%	~ 9%	~ 12 %	~ 11%	~ 7%	~ 23%	~ 25%
% Cover Turf Algae <sup>3</sup>	~ 80%	~ 50%	~ 72%	~ 78%	~ 70%	~ 60%	~ 60%

\*notes:

1. After Hubbard (1989a-d)
2. From diver observations (this study)
3. From Nemeth and Herzlieb (2002)

Table 2. Fish census effort

	LB	JI	EW	WW	ER	CB	SH
Spring 2002							
Date of Censuses	29-May	11-Jun	23, 28-May	22, 23-May	3-May	15-May	16-May
No. of censuses	4	6	6	6	6	6	6
Total Census Time (min)	91	195	187	190	233	242	255
Avg Time / Census (min)	22.8	32.5	31.2	31.7	38.8	40.3	42.5
Fall 2002							
Date of Censuses	3-Sept	4-Sept	6-Sept	12-Sept	4 & 6-Sept	10-Sept	11-Sept
No. of censuses	6	6	6	6	6	6	6
Total Census Time (min)	197	179	225	254	309	294	285
Avg Time / Census (min)	32.8	29.8	37.5	42.3	51.5	49.0	47.5

Table 3. Total number of fish and species observed during stationary fish censuses<sup>\*1</sup>

Reef Site	Survey	LB	JI	EW	WW	ER	CB	SH	Total
No. of Fish	Spring '02	1,659 <sup>*2</sup>	1,921	1,755	1,922	2,156	3,120	4,326	16,859
Observed	Fall '02	2,491	1,385	1,456	2,221	2,148	5,443	3,319	18,463
	Total	4,150	3,306	3,211	4,143	4,304	8,563	7,645	35,322
No. of Species	Spring '02	24	39	37	34	40	32	40	72
Observed	Fall '02	28	37	38	42	42	42	40	77
	Cumulative	32	47	47	48	52	44	49	85

Notes:

1. Values reported for Numbers of species are not additive because there is substantial overlap in species compositions among different reef sites and between sampling periods.
2. Total number of fish from 4 replicate censuses.

Table 4. The 25 most abundant fish species observed in this study.

Common Name	Species	Family	Spring '02			Fall '02			Total No. of Fish Observed	Avg No. Fish per Census	% Freq (n=40)	Total No. of Fish Observed	Avg No. Fish per Census	% Freq (n=42)	Total No. of Fish	% of All Obs*
			Total No. of Fish Observed	Avg No. Fish per Census	% Freq (n=40)	Total No. of Fish Observed	Avg No. Fish per Census	% Freq (n=42)								
bluehead wrasse	<i>Thalassoma bifasciatum</i>	Labridae	7,790	194.8	97.5	5,830	138.8	97.6	13,620	38.6						
creole wrasse bicolor	<i>Clepticus parrae</i>	Labridae	2,010	50.3	45	3,635	86.5	38.1	5,645	16.0						
damselfish	<i>Stegastes partitus</i>	Pomacentridae	1,466	36.7	100	2,485	59.2	100.0	3,951	11.2						
brown chromis	<i>Chromis multilineata</i>	Pomacentridae	1,661	41.5	40	2,172	51.7	40.5	3,833	10.9						
blue chromis ocean	<i>Chromis cyanus</i>	Pomacentridae	1,864	46.6	75	1,880	44.8	76.2	3,744	10.6						
surgeonfish yellowhead wrasse	<i>Acanthurus bahianus</i>	Acanthuridae	367	9.2	97.5	287	6.8	97.6	654	1.9						
black durgon redband parrotfish	<i>Halichoeres garnoti</i>	Labridae	162	4.1	72.5	149	3.5	81.0	311	0.9						
parrotfish princess	<i>Melichthys niger</i>	Balistidae	94	2.4	50	201	4.8	52.4	295	0.8						
dusky damselfish threespot damselfish coney sergeant major blue tang french grunt yellow goatfish soureye butterflyfish stoplight parrotfish yellowtail damselfish	<i>Sparisoma aurofrenatum</i>	Scaridae	144	3.6	90	139	3.3	83.3	283	0.8						
	<i>Scarus taeniopterus</i>	Scaridae	138	3.5	87.5	138	3.3	85.7	276	0.8						
	<i>Stegastes adustus</i> (S. fuscus)	Pomacentridae	150	3.8	60	100	2.4	50.0	250	0.7						
	<i>Stegastes planifrons</i>	Pomacentridae	111	2.8	22.5	94	2.2	21.4	205	0.6						
	<i>Cephalopholis fulvis</i>	Serranidae	97	2.4	72.5	108	2.6	81.0	205	0.6						
	<i>Abudefduf saxatilis</i>	Pomacentridae	47	1.2	32.5	157	3.7	42.9	204	0.6						
	<i>Acanthurus coeruleus</i>	Acanthuridae	89	2.2	87.5	106	2.5	90.5	195	0.6						
	<i>Haemulon flavolineatum</i>	Haemulidae	93	2.3	82.5	85	2.0	78.6	178	0.5						
	<i>Mulloidichthys martinicus</i>	Mullidae	14	0.4	17.5	158	3.8	23.8	172	0.5						
	<i>Chaetodon capistratus</i>	Chaetodontidae	59	1.5	57.5	61	1.5	54.8	120	0.3						
	<i>Sparisoma viride</i>	Scaridae	36	0.9	55	62	1.5	64.3	98	0.3						
	<i>Microspathodon chrysurus</i>	Pomacentridae	46	1.2	60	48	1.1	52.4	94	0.3						

Table 4 (continued). The 25 most abundant fish species observed in this study.

Common Name	Species	Family	Spring '02			Fall '02			Total No. of Fish	% of All Obs*
			Total No.	Avg No.		Total No.	Avg No.			
			of Fish Observed	Fish per Census	% Freq (n=40)	of Fish Observed	Fish per Census	% Freq (n=42)		
striped parrotfish	<i>Scarus iseri</i> (S. croicensis)	Scaridae	19	0.5	22.5	73	1.7	40.5	92	0.3
blackbar soldierfish	<i>Myripris jacobus</i>	Holocentridae	45	1.1	35	43	1.0	33.3	88	0.2
yellowtail snapper	<i>Ocyurus chrysururus</i>	Lutjanidae	49	1.2	32.5	23	0.5	16.7	72	0.2
mahogany snapper	<i>Lutjanus mahogoni</i>	Lutjanidae	23	0.6	12.5	27	0.6	23.8	50	0.1
fairy basslet	<i>Gramma loreto</i>	Grammatidae	11	0.3	10	37	0.9	16.7	48	0.1
			Subtotal = 16,585		Subtotal = 18,098				Subtotal = 34,683	98.2

\* percent of all observations (n = 35,322)

Appendix 1. Distribution of observations on 85 fish species across seven St. Croix reef sites.



**Appendix 1 (continued).** Distribution of observations on 85 fish species across seven St. Croix reef sites.

Appendix 1 (continued). Distribution of observations on 85 fish species across seven St. Croix reef sites.

Common Name	Species	Family	Distribution of observations on 85 fish species across seven St. Croix reef sites.											
			Lang Bank		Jacks		West Wall		Eagle Ray		Cane Bay		Sprat Hole	
			Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall
threespot damselfish	<i>Stegastes planifrons</i>	Pomacentridae									1	1	1	1
cocoa damselfish	<i>Stegastes variabilis</i>	Pomacentridae									1	1		1
striped parrotfish	<i>Scarus iserti</i> (S. croicensis)	Scaridae	1	1	1	1	1	1	1	1	1	1	1	1
princess parrotfish	<i>Scarus taeniopterus</i>	Scaridae	1	1	1	1	1	1	1	1	1	1	1	1
queen parrotfish	<i>Scarus vetula</i>	Scaridae												
greenblotch parrotfish	<i>Sparisoma atomarium</i>	Scaridae												
redband parrotfish	<i>Sparisoma aurofrenatum</i>	Scaridae	1	1	1	1	1	1	1	1	1	1	1	1
redtail parrotfish	<i>Sparisoma chrysopterum</i>	Scaridae												
yellowtail parrotfish	<i>Sparisoma rubripinne</i>	Scaridae												
stoplight parrotfish	<i>Sparisoma viride</i>	Scaridae	1	1	1	1	1	1	1	1	1	1	1	1
highhat	<i>Pareques acuminatus</i>	Sciaenidae												
cero mackerel	<i>Scomberomorus regalis</i>	Scombridae												
spotted scorpionfish	<i>Scorpaena plumieri</i>	Scorpaenidae			1						1	1	1	1
graysby	<i>Cephalopholis cinctatus</i>	Serranidae									1	1	1	1
coney	<i>Cephalopholis fulvus</i>	Serranidae	1	1	1	1	1	1	1	1	1	1	1	1
red hind	<i>Epinephelus guttatus</i>	Serranidae												
yellowtail hamlet	<i>Hoploplectrus chlorurus</i>	Serranidae									1		1	
shy hamlet	<i>Hoploplectrus guttavarius</i>	Serranidae									1			
black hamlet	<i>Hoploplectrus nigricans</i>	Serranidae									1		1	

**Appendix 1** (continued). Distribution of observations on 85 fish species across seven St. Croix reef sites.

Common Name	Species	Family	Distribution of observations on 85 fish species across seven St. Croix reef sites.												
			Lang Bank		Jacks		East Wall		West Wall		Eagle Ray		Cane Bay		
			Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	Spring	Fall	
barred hamlet	<i>Hoploplectrus puella</i>	Serranidae													
butter hamlet	<i>Hoploplectrus unicolor</i>	Serranidae													
greater soapfish	<i>Rypticus saponaceus</i>	Serranidae													
tobaccofish	<i>Serranus cabrilla</i>	Serranidae													
harlequin bass	<i>Serranus tigrinus</i>	Serranidae													
jolthead porgy	<i>Caesio bajonado</i>	Sparidae													
sand diver	<i>Synodus intermedius</i>	Synodontidae													
sharpnose puffer	<i>Canthigaster rostrata</i>	Tetraodontidae													
porcupinefish	<i>Diodon hystrix</i>	Tetraodontidae													
Grand Total =	85 species		Subtotal =	24	28	39	37	37	38	34	42	40	42	40	40

Appendix 2A. Fish census data from Lang Bank

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=4)	Total	Avg.	StDev	% Freq (n=6)	Total	Avg.	StDev	% Freq	No. of Fish	Avg	Min	Max	
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	100.0	30	7.5	2.6	14.5	11	17	100.0	64	10.7	2.8	14.8	10	20
doctorfish	<i>Acanthurus chirurgus</i>	Acanthuridae	50.0	2	0.5	0.6	18.0	17	19	33.3	7	1.2	2.0	20.5	17	23
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	100.0	11	2.8	1.7	16.3	14	19	100.0	24	4.0	1.5	14.7	10	19
queen triggerfish	<i>Balistes vetula</i>	Balistidae	25.0	1	0.3	0.5	27.0	27	27	16.7	1	0.2	0.4	20.0	20	20
black durgon	<i>Melichthys niger</i>	Balistidae	100.0	15	3.8	4.2	22.0	20	26	83.3	58	9.7	11.1	21.0	15	30
banded butterflyfish	<i>Chaetodon striatus</i>	Chaetodontidae	25.0	2	0.5	1.0	10.5	10	11	33.3	4	0.7	1.0	12.3	11	14
caesar grunt	<i>Haemulon carbonarium</i>	Haemulidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	20.0	20	20
french grunt bluesstriped grunt	<i>Haemulon flavolineatum</i>	Haemulidae	100.0	9	2.3	1.3	15.8	15	18	66.7	13	2.2	2.2	16.5	14	20
squirrelfish longspine squirrelfish spanish hogfish	<i>Holocentrus acensionis</i>	Holocentridae	25.0	1	0.3	0.5	26.0	26	26	0.0	0	0.0	0.0	-	-	-
slippery dick yellowcheek wrasse yellowhead wrasse clown wrasse	<i>Holocentrus rufus</i>	Holocentridae	100.0	5	1.3	0.5	17.5	17	20	50.0	5	0.8	1.2	16.5	16	17
	<i>Bodianus rufus</i>	Labridae	0.0	0	0.0	0.0	-	-	-	50.0	3	0.5	0.5	14.3	9	17
	<i>Halichoeres bivittatus</i>	Labridae	50.0	8	2.0	2.4	6.0	5	9	0.0	0	0.0	0.0	-	-	-
	<i>Halichoeres cyanophthalmus</i>	Labridae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	18.5	12	25
	<i>Halichoeres garnoti</i>	Labridae	100.0	11	2.8	1.0	10.5	8	16	83.3	25	4.2	3.1	9.1	4	18
	<i>Halichoeres maculipinna</i>	Labridae	25.0	2	0.5	1.0	9.5	8	11	33.3	15	2.5	4.0	5.5	3	9

Appendix 2A (continued). Fish census data from Lang Bank

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=4)		No. of Fish		Size		% Freq (n=6)		No. of Fish		Size			
			Total	Avg.	Total	Avg.	StDev	Min	Max	Total	Avg.	StDev	Min	Max		
bluehead wrasse	<i>Thalassoma bifasciatum</i>	Labridae	100.0	1340	335.0	102.5	3.0	2	9	100.0	1405	234.2	86.2	4.3	2	10
orange-spotted filefish	<i>Cantherhines pullus</i>	Monacanthidae	0.0	0	0.0	0.0	-	-	-	33.3	3	0.5	0.8	21.8	17	25
scrawled cowfish	<i>Acanthostracion quadricornis</i>	Ostraciidae	25.0	1	0.3	0.5	22.0	22	22	0.0	0	0.0	0.0	-	-	-
queen angelfish	<i>Holacanthus ciliaris</i>	Pomacanthidae	25.0	1	0.3	0.5	25.0	25	25	0.0	0	0.0	0.0	-	-	-
rock beauty	<i>Holacanthus tricolor</i>	Pomacanthidae	75.0	3	0.8	0.5	17.7	15	22	33.3	4	0.7	1.0	9.0	3	20
blue chromis	<i>Chromis cyanus</i>	Pomacentridae	75.0	7	1.8	1.5	4.0	3	7	83.3	45	7.5	4.4	4.8	2	8
yellowtail damselfish bicolor	<i>Microspathodon chrysurus</i>	Pomacentridae	0	0	0	0	-	-	-	16.7	1	0.2	0.4	12.0	12	12
damsel fish striped parrotfish	<i>Stegastes partitus</i>	Pomacentridae	100.0	171	42.8	18.6	4.0	2	6	100.0	721	120.2	25.6	3.8	2	6
princess parrotfish	<i>Scarus iserti (S. croicensis)</i>	Scaridae	25.0	1	0.3	0.5	24.0	24	24	16.7	1	0.2	0.4	30.0	30	30
redband parrotfish	<i>Scarus taeniopterus</i>	Scaridae	100.0	11	2.8	1.3	16.8	12	21	100.0	28	4.7	1.9	17.2	10	30
redtail parrotfish	<i>Sparisoma aurofrenatum</i>	Scaridae	75.0	7	1.8	1.3	14.7	12	18	83.3	16	2.7	1.8	17.0	10	20
stoplight parrotfish	<i>Sparisoma chrysopurpureum</i>	Scaridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	22.0	22	22
coney	<i>Sparisoma viride</i>	Scaridae	50.0	3	0.8	1.0	27.8	25	30	50.0	7	1.2	1.5	30.7	25	40
red hind harlequin bass	<i>Cephalopholis fulvus</i>	Serranidae	100.0	14	3.5	2.4	16.6	14	20	100.0	29	4.8	2.4	15.2	10	22
	<i>Epinephelus guttatus</i>	Serranidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	25.0	25	25
	<i>Serranus tigrinus</i>	Serranidae	50.0	3	0.8	1.0	8.3	6	10	16.7	1	0.2	0.4	7.0	7	7
			Total=	1659						Total=	2491					

Final Report: Patterns of habitat utilization by reef fish on St. Croix

F-7-17, Study 3

Period: 1 October 2001 to 30 September 2002

Appendix 2B. Fish census data from Jacks & Isaacs Bay

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)			No. of Fish			% Freq (n=6)			No. of Fish				
			Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev		
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	100.0	74	12.3	5.5	15.0	7	20	100.0	32	5.3	1.2	13.7	6	20
doctorfish	<i>Acanthurus chirurgus</i>	Acanthuridae	50.0	5	0.8	1.2	19.0	1.7	21	16.7	1	0.2	0.4	18.0	18	18
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	100.0	12	2.0	0.9	14.7	8	20	100.0	17	2.8	1.0	11.2	3	17
black durgon	<i>Melichthys niger</i>	Balistidae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	17.5	15	20
blue runner	<i>Caranx cryos</i>	Carangidae	16.7	1	0.2	0.4	30.0	30	0.0	0.0	0	0.0	0.0	-	-	-
bar jack	<i>(C. fuscus)</i>	Carangidae	16.7	2	0.3	0.8	22.5	21	24	16.7	2	0.3	0.8	20.0	20	20
banded butterflyfish	<i>Chaetodon striatus</i>	Chaetodontidae	16.7	4	0.7	1.6	11.0	10	12	16.7	2	0.3	0.8	9.0	8	10
tomtate	<i>Haemulon aurolineatum</i>	Haemulidae	16.7	1	0.2	0.4	19.0	19	19	16.7	2	0.3	0.8	14.5	14	15
caesar grunt	<i>Haemulon carbonarium</i>	Haemulidae	16.7	1	0.2	0.4	19.0	19	19	16.7	1	0.2	0.4	16.0	16	16
smallmouth grunt	<i>Haemulon chrysargyreum</i>	Haemulidae	16.7	3	0.5	1.2	12.0	12	12	0.0	0	0.0	0.0	-	-	-
french grunt bluesstriped grunt	<i>Haemulon flavolineatum</i>	Haemulidae	83.3	19	3.2	2.9	18.0	14	23	66.7	9	1.5	1.6	15.5	13	16
squirrelfish longpine	<i>Holocentrus acensionis</i>	Holocentridae	0.0	0	0.0	0.0	-	-	-	16.7	2	0.3	0.8	19.0	18	20
squirrelfish blackbar soldierfish spanish hogfish creole wrasse	<i>Holocentrus rufus</i>	Holocentridae	83.3	7	1.2	0.8	19.6	18	22	33.3	3	0.5	0.8	19.3	19	20
	<i>Myripristis jacobus</i>	Holocentridae	66.7	7	1.2	1.2	16.4	16	18	33.3	6	1.0	2.0	13.5	12	15
	<i>Bodianus rufus</i>	Labridae	33.3	2	0.3	0.5	18.5	17	20	16.7	1	0.2	0.4	6.0	6	6
	<i>Clepticus parrae</i>	Labridae	0.0	0	0.0	0.0	-	-	-	33.3	13	2.2	4.8	10.0	4	15

## Appendix 2B (continued). Fish census data from Jacks & Isaacs Bay

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)			No. of Fish			% Freq (n=6)			No. of Fish				
			Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev		
yellowhead wrasse	<i>Halichoeres garnotii</i>	Labridae	83.3	13	2.2	1.3	9.8	6	1.7	83.3	16	2.7	2.6	9.0	4	15
clown wrasse	<i>Halichoeres maculipinna</i>	Labridae	0.0	0	0.0	0.0	-	-	-	16.7	20	3.3	8.2	3.5	3	5
puddingwife bluehead wrasse	<i>Halichoeres radiatus</i>	Labridae	16.7	1	0.2	0.4	8.0	8	8	0.0	0	0.0	0.0	-	-	-
schoolmaster mahogany snapper	<i>Thalassoma bifasciatum</i>	Labridae	100.0	99.5	165.8	78.3	5.3	2	12	100.0	640	106.7	59.6	4.0	2	10
sand tilefish scrawled filefish	<i>Lutjanus apodus</i>	Lutjanidae	16.7	1	0.2	0.4	17.0	17	17	16.7	1	0.2	0.4	20.0	20	20
yellow goatfish honeycomb cowfish	<i>Lutjanus mahogoni</i>	Lutjanidae	16.7	3	0.5	1.2	22.0	20	22	33.3	3	0.5	0.8	18.0	15	20
smooth trunkfish	<i>Matacanthus plumieri</i>	Malacanthidae	50.0	3	0.5	0.5	27.3	26	30	16.7	1	0.2	0.4	25.0	25	25
rock beauty sergeant major	<i>Aluterus scriptus</i>	Monacanthidae	16.7	1	0.2	0.4	25.0	25	25	0.0	0	0.0	0.0	-	-	-
bluehead wrasse	<i>Mulloidichthys martinicus</i>	Mullidae	0.0	0	0.0	0.0	-	-	-	16.7	5	0.8	2.0	18.0	16	20
yellowtail damselfish	<i>Acanthostracion polyacanthus</i>	Ostraciidae	33.3	2	0.3	0.5	24.0	22	26	16.7	1	0.2	0.4	20.0	20	20
yellowtail damselfish	<i>Lactophrys triqueter</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	-	33.3	3	0.5	0.8	17.0	15	20
yellowtail damselfish	<i>Holacanthus tricolor</i>	Pomacentridae	33.3	2	0.3	0.5	11.5	8	15	16.7	1	0.2	0.4	18.0	18	18
blue chromis brown chromis	<i>Abudefduf saxatilis</i>	Pomacentridae	16.7	1	0.2	0.4	13.0	13	13	0.0	0	0.0	0.0	-	-	-
yellowtail damselfish	<i>Chromis cyanus</i>	Pomacentridae	100.0	197	32.8	32.0	5.3	2	8	100.0	139	23.2	19.3	4.3	2	7
dusky damselfish	<i>Chromis multilineata</i>	Pomacentridae	16.7	8	1.3	3.3	5.5	5	6	16.7	2	0.3	0.8	5.5	5	6
yellowtail damselfish	<i>Microspathodon chrysurus</i>	Pomacentridae	50.0	6	1.0	1.1	12.2	11	13	66.7	6	1.0	0.9	12.3	12	13
dusky damselfish	<i>Siganus austrius</i>	Pomacentridae	33.3	3	0.5	0.8	7.8	6	10	16.7	1	0.2	0.4	8.0	8	8
beaugregory leucostictus	<i>Siganus fusca</i>	Pomacentridae	33.3	3	0.5	0.8	7.0	6	8	0.0	0	0.0	0.0	-	-	-

## Appendix 2B (continued). Fish census data from Jacks & Isaacs Bay

Appendix 2B (continued). Fish census data from Jacks & Isaacs Bay																
Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq		No. of Fish		Size		% Freq		No. of Fish		Size			
			(n=6)	Total	Avg.	StDev	(n=6)	Total	Avg.	StDev	(n=6)	Total	Avg.	StDev		
bicolor damselfish	<i>Siegestes partitus</i>	Pomacentridae	100.0	446	74.3	22.5	5.1	3	7	100.0	400	66.7	31.1	3.8	2	6
striped parrotfish	<i>Scarus iserti</i> (S. <i>croicensis</i> )	Scaridae	16.7	2	0.3	0.8	13.0	12	14	16.7	3	0.5	1.2	10.0	6	18
princess parrotfish	<i>Scarus</i> <i>teeniophterus</i>	Scaridae	83.3	17	2.8	2.8	15.4	8	22	66.7	5	0.8	0.8	21.5	16	30
queen parrotfish	<i>Scarus vetula</i>	Scaridae	16.7	2	0.3	0.8	21.0	20	22	0.0	0	0.0	0.0	-	-	-
redband parrotfish	<i>Sparisoma</i> <i>aurofrenatum</i>	Scaridae	100.0	24	4.0	1.3	14.2	5	25	100.0	17	2.8	1.6	16.8	10	22
redtail parrotfish	<i>Sparisoma</i> <i>chrysopurum</i>	Scaridae	16.7	12	2.0	4.9	23.0	18	24	16.7	1	0.2	0.4	22.0	22	22
stoplight parrotfish	<i>Sparisoma viride</i>	Scaridae	66.7	6	1.0	0.9	19.5	15	24	50.0	4	0.7	0.8	22.7	16	30
highhat	<i>Pareques</i>	Sciaenidae	0.0	0	0.0	-	-	-	-	16.7	2	0.3	0.8	11.5	8	15
spotted scorpionfish	<i>Scorpaena</i> <i>acuminatus</i>	Scorpaenidae	0.0	0	0.0	-	-	-	-	16.7	2	0.3	0.8	-	-	-
coneys	<i>Cephalopholis</i> <i>flavilatus</i>	Serranidae	16.7	1	0.2	0.4	27.0	27	27	0.0	0	0.0	0.0	-	-	-
red hind sharknose puffer	<i>Epinephelus</i> <i>gruttatus</i> <i>Canthigaster</i> <i>rostrata</i>	Serranidae	16.7	1	0.2	0.4	22.0	22	22	0.0	0	0.0	0.0	-	-	-
		Tetraodontidae	16.7	1	0.2	0.4	5.0	5	5	0.0	0	0.0	0.0	-	-	-
														Total = 1921	Total = 1385	

Appendix 2C. Fish census data from East Wall, Salt River Canyon

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)			No. of Fish			% Freq (n=6)			No. of Fish				
			Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev		
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	100.0	73	12.2	4.4	14.5	1.2	18	100.0	62	10.3	2.8	13.3	8	18
doctorfish	<i>Acanthurus chirurgus</i>	Acanthuridae	33.3	3	0.5	0.8	20.3	1.9	23	0.0	0	0.0	0.0	-	-	-
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	83.3	18	3.0	0.9	15.4	1.4	18	100.0	20	3.3	1.6	15.0	10	20
trumpetfish	<i>Aulostomus maculatus</i>	Aulostomidae	16.7	2	0.3	0.8	27.5	2.5	30	33.3	3	0.5	0.8	42.5	30	50
black durgon	<i>Melichthys niger</i>	Balistidae	83.3	11	1.8	1.5	23.7	2.0	25	83.3	31	5.2	7.4	21.2	15	25
bar jack	<i>Caranx ruber</i>	Carangidae	33.3	4	0.7	1.0	21.0	14	20	16.7	2	0.3	0.8	19.0	18	20
longsnout butterflyfish	<i>Chaetodon aculeatus</i>	Chaetodontidae	16.7	1	0.2	0.4	8.0	8	8	16.7	3	0.5	1.2	9.0	8	10
soureye butterflyfish	<i>Chaetodon capistratus</i>	Chaetodontidae	100.0	16	2.7	1.8	9.3	7	11	83.3	18	3.0	2.4	9.0	5	11
spottin butterflyfish	<i>Chaetodon ocellatus</i>	Chaetodontidae	16.7	1	0.2	0.4	16.0	16	16	16.7	1	0.2	0.4	13.0	13	13
banded butterflyfish	<i>Chaetodon striatus</i>	Chaetodontidae	50.0	6	1.0	1.1	10.5	8	12	16.7	2	0.3	0.8	11.0	11	11
french grunt	<i>Haemulon flavolineatum</i>	Haemulidae	100.0	11	1.8	0.8	15.3	12	17	100.0	17	2.8	1.2	15.6	10	20
white grunt	<i>Haemulon plumieri</i>	Haemulidae	50.0	3	0.5	0.5	22.3	20	25	16.7	1	0.2	0.4	20.0	20	20
bluestriped grunt	<i>Haemulon sciurus</i>	Haemulidae	16.7	1	0.2	0.4	22.0	22	22	0.0	0	0.0	0.0	-	-	-
squirlfish longspine	<i>Holocentrus acensionis</i>	Holocentridae	0.0	0	0.0	-	-	-	16.7	3	0.5	1.2	19.0	18	20	
squirlfish blackbar	<i>Holocentrus rufus</i>	Holocentridae	50.0	5	0.8	1.0	16.2	15	17	0.0	0	0.0	0.0	-	-	-
soldierfish spanish hogfish	<i>Myripristis jacobus</i>	Holocentridae	0.0	0	0.0	-	-	-	16.7	1	0.2	0.4	18.0	18	18	
	<i>Bodianus rufus</i>	Labridae	16.7	2	0.3	0.8	11.5	3	20	0.0	0	0.0	0.0	-	-	-

## Appendix 2C

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			%Freq (n=6)			No. of Fish			%Freq (n=6)			No. of Fish				
			Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev		
yellowhead wrasse	<i>Halichoeres garnoti</i>	Labridae	50.0	8	1.3	1.5	9.8	5	14	100.0	22	3.7	2.6	8.6	4	14
bluehead wrasse	<i>Thalassoma bifasciatum</i>	Labridae	100.0	1290	215.0	114.8	3.2	2	9	100.0	820	136.7	61.5	4.8	2	12
schoolmaster snapper	<i>Lutjanus apodus</i>	Lutjanidae	0.0	0	0.0	0.0	-	-	-	16.7	5	0.8	2.0	13.0	10	15
yellowtail snapper	<i>Ocyurus chrysurus</i>	Lutjanidae	100.0	19	3.2	2.1	22.6	17	28	16.7	3	0.5	1.2	21.0	18	24
orange-spotted filefish	<i>Cauferhinus pullus</i>	Monacanthidae	16.7	1	0.2	0.4	17.0	17	17	16.7	1	0.2	0.4	16.0	16	16
yellow goatfish	<i>Mulloidichthys martinicus</i>	Mullidae	16.7	1	0.2	0.4	20.0	20	20	16.7	3	0.5	1.2	19.0	18	20
spotted goatfish	<i>Pseudupeneus maculatus</i>	Mullidae	50.0	3	0.5	0.5	18.7	16	20	0.0	0	0.0	0.0	-	-	-
honeycomb cowfish	<i>Acanthostracion polyacanthus</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	26.0	26	26
smooth trunkfish	<i>Iactophryss trispinosus</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	-	66.7	4	0.7	0.5	15.8	15	17
rock beauty	<i>Holacanthus tricolor</i>	Pomacanthidae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	17.5	13	22
french angelfish	<i>Pomacanthus paru</i>	Pomacanthidae	33.3	2	0.3	0.5	24.5	23	26	0.0	0	0.0	0.0	-	-	-
sergeant major	<i>Abudefduf saxatilis</i>	Pomacentridae	16.7	2	0.3	0.8	13.0	12	14	83.3	67	11.2	10.7	11.8	8	15
blue chromis	<i>Chromis cyanus</i>	Pomacentridae	33.3	71	11.8	24.0	3.0	2	4	50.0	65	10.8	13.9	4.3	3	7
yellowtail damselfish	<i>Microspathodon chrysurus</i>	Pomacentridae	83.3	5	0.8	0.4	12.4	12	13	66.7	7	1.2	1.0	11.6	10	13
bicolor damselfish	<i>Siganus partitus</i>	Pomacentridae	100.0	143	23.8	7.8	4.0	2	7	100.0	211	35.2	8.1	4.0	2	6
striped parrotfish	<i>Scarus isertii (S. croicensis)</i>	Scaridae	16.7	2	0.3	0.8	8.0	4	12	16.7	6	1.0	2.4	5.0	4	5
princess parrotfish	<i>Scarus taeniopterus</i>	Scaridae	83.3	15	2.5	2.1	17.0	14	25	83.3	21	3.5	2.3	17.1	7	25

Appendix 2C

(continued). Fish census data from East Wall, Salt River Canyon

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq			No. of Fish			% Freq			No. of Fish				
			Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev		
greenblotch parrotfish	<i>Spurisoma atomarium</i>	Scaridae	0.0	0	0.0	0.0	-	-	16.7	4	0.7	1.6	14.5	14	15	
redband parrotfish	<i>Spurisoma aurofrenatum</i>	Scaridae	83.3	13	2.2	1.7	16.9	12	21	83.3	21	3.5	1.9	17.2	11	23
stoplight parrotfish	<i>Spurisoma viride</i>	Scaridae	33.3	2	0.3	0.5	22.5	15	30	66.7	6	1.0	0.9	19.8	6	32
graysby	<i>Cephalopholis crenatus</i>	Serranidae	16.7	1	0.2	0.4	17.0	17	17	16.7	1	0.2	0.4	18.0	18	18
coney	<i>Cephalopholis fulvus</i>	Serranidae	100.0	15	2.5	1.6	16.4	14	19	83.3	15	2.5	1.9	17.8	8	24
red hind	<i>Epinephelus guttatus</i>	Serranidae	16.7	1	0.2	0.4	25.0	25	25	16.7	1	0.2	0.4	20.0	20	20
shy hamlet	<i>Hypoplectrus guittavrinii</i>	Serranidae	16.7	1	0.2	0.4	13.0	13	13	0.0	0	0.0	0.0	-	-	-
black hamlet	<i>Hypoplectrus nigricans</i>	Serranidae	16.7	1	0.2	0.4	14.0	14	14	0.0	0	0.0	0.0	-	-	-
butter hamlet	<i>Hypoplectrus unicolor</i>	Serranidae	16.7	1	0.2	0.4	10.0	10	10	0.0	0	0.0	0.0	-	-	-
greater soapfish	<i>Rypticus saponaceus</i>	Serranidae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	15.0	15	15	
tobacofish	<i>Serranus tabacarius</i>	Serranidae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	5.0	5	5	
harlequin bass	<i>Serranus tigrinus</i>	Serranidae	16.7	1	0.2	0.4	11.0	11	11	50.0	3	0.5	0.5	7.0	5	8
sharpnose puffer	<i>Canthigaster rostrata</i>	Tetraodontidae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	6.0	6	6	
			Total=	1755									Total=	1456		

Final Report: Patterns of habitat utilization by reef fish on St. Croix

F-7-17, Study 3

Period: 1 October 2001 to 30 September 2002

Final Report: Patterns of habitat utilization by reef fish on St. Croix

F-7-17, Study 3

Period: 1 October 2001 to 30 September 2002

Common Name	Species Name	Family	Spring '02 Census						Fall '02 Census						Size		
			% Freq	No. of Fish	Size	% Freq	No. of Fish	Avg	Min	Max							
			(n=6)	Total	Avg.	StDev	No. of Fish	Avg.	Min	Max	(n=6)	Total	Avg.	StDev	Total	Avg.	Min
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	100.0	96	16.0	6.3	13.6	9	17	100.0	49	8.2	4.3	14.8	7	22	
doctorfish	<i>Acanthurus chirurgus</i>	Acanthuridae	0.0	0	0.0	0.0	-	-	-	33.3	3	0.5	0.8	16.5	12	21	
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	83.3	14	2.3	1.6	16.4	14	20	100.0	12	2.0	0.6	16.8	14	20	
black durgon	<i>Melichthys niger</i>	Balistidae	66.7	27	4.5	6.3	23.1	20	26	66.7	44	7.3	8.9	23.0	12	25	
peacock flounder	<i>Bothus lunatus</i>	Bothidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	12.0	12	12	
bar jack	<i>Caranx ruber</i>	Carangidae	16.7	1	0.2	0.4	22.0	22	22	33.3	3	0.5	0.8	29.5	18	40	
longsnout butterflyfish	<i>Chaetodon aculeatus</i>	Chaetodontidae	33.3	3	0.5	0.8	6.3	6	7	33.3	3	0.5	0.8	8.0	7	10	
foureye butterflyfish	<i>Chaetodon capistratus</i>	Chaetodontidae	100.0	20	3.3	2.3	9.5	4	11	83.3	15	2.5	1.6	9.2	7	12	
banded butterflyfish	<i>Chaetodon striatus</i>	Chaetodontidae	50.0	5	0.8	1.0	11.5	11	12	0.0	0	0.00	0.0	-	-	-	
yellowfin mojarra	<i>Gerres cinereus</i>	Gerieidae	0.0	0	0.0	0.0	-	-	-	16.7	2	0.3	0.8	25.0	24	26	
french grunt	<i>Haemulon flavolineatum</i>	Haemulidae	100.0	17	2.8	1.0	13.8	9	17	83.3	15	2.5	2.4	17.2	15	20	
white grunt	<i>Haemulon plumieri</i>	Haemulidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	22.0	22	22	
squirmelfish	<i>Holocentrus adscensionis</i>	Holocentridae	33.3	3	0.5	0.8	17.3	15	20	16.7	1	0.2	0.4	18.0	18	18	
longspine squirrelfish	<i>Holocentrus rufus</i>	Holocentridae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	16.5	16	17	
blackbar soldierfish	<i>Myripristis jacobus</i>	Holocentridae	16.7	1	0.2	0.4	14.0	14	14	16.7	1	0.2	0.4	15.0	15	15	
creole wrasse	<i>Clepticus parrae</i>	Labridae	66.7	136	22.7	27.0	10.5	8	15	66.7	345	57.5	58.8	16.3	10	20	

Appendix 2D (continued). Fish census data from West Wall, Salt River Canyon

Common Name	Species Name	Family	Spring '02 Census						Fall '02 Census						
			% Freq (n=6)			No. of Fish			% Freq (n=6)			No. of Fish			
			Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	Total	Avg.	StDev	
slippery dick	<i>Halichoeres bivittatus</i>	Labridae	16.7	2	0.3	0.8	2.5	2	3	0.0	0	0.0	0.0	-	-
yellowhead wrasse	<i>Halichoeres garnoti</i>	Labridae	33.3	2	0.3	0.5	10.0	10	10	83.3	14	2.3	1.2	9.4	5
bluehead wrasse	<i>Thalassoma bifasciatum</i>	Labridae	100.0	133.3	222.2	41.8	3.0	1.5	10	100.0	123.5	205.8	130.2	4.6	2
schoolmaster	<i>Lutjanus apodus</i>	Lutjanidae	33.3	3	0.5	0.8	22.0	18	25	33.3	2	0.3	0.5	30.0	30
yellowtail snapper	<i>Ocyurus chrysuris</i>	Lutjanidae	33.3	20	3.3	5.2	25.0	20	30	50.0	15	2.5	2.8	26.0	21
scrawled filefish	<i>Ahuterus scriptus</i>	Monacanthidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	35.0	35
orange-spotted filefish	<i>Cantherhines pullus</i>	Monacanthidae	0.0	0	0.0	0.0	-	-	-	50.0	3	0.5	0.5	15.7	15
yellow goatfish	<i>Mulloidichthys martinicus</i>	Mullidae	0.0	0	0.0	0.0	-	-	-	16.7	4	0.7	1.6	17.0	16
spotted goatfish	<i>Pseudupeneus maculatus</i>	Mullidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	20.0	20
rock beauty	<i>Holacanthus tricolor</i>	Pomacentridae	16.7	1	0.2	0.4	17.0	17	17	16.7	1	0.2	0.4	6.0	6
sergeant major blue chromis	<i>Abudefduf saxatilis</i>	Pomacentridae	50.0	20	3.3	5.9	11.3	10	14	50.0	20	3.3	4.3	12.8	10
brown chromis yellowtail damselfish	<i>Chromis cyanus</i>	Pomacentridae	16.7	3	0.5	1.2	4.0	3	5	0.0	0	0.0	0.0	-	-
dusky damselfish	<i>Microspathodon chrysurus</i>	Pomacentridae	33.3	41	6.8	14.0	5.3	4	7	33.3	90	15.0	23.5	6.0	4
beaugregory	<i>Siganus adustus</i> (S. fuscus)	Pomacentridae	100.0	14	2.3	0.8	10.0	5	12	100.0	17	2.8	1.2	12.1	2.5
	<i>Siganus leucostictus</i>	Pomacentridae	16.7	1	0.2	0.4	9.0	9	9	83.3	15	2.5	1.8	8.9	7
										0	0.0	0.0	0.0	-	-

Appendix 2D (continued). Fish census data from West Wall, Salt River Canyon

Common Name	Species Name	Family	Spring '02 Census						Fall '02 Census							
			%&Freq (n=6)	Total	Avg.	No. of Fish	Size	Avg.	%&Freq (n=6)	Total	Avg.	StDev	No. of Fish	Size		
bicolor damselfish	<i>Stegastes partitus</i>	Pomacentridae	100.0	52	8.7	8.7	5.3	1	12	100.0	220	36.7	10.8	4.4	2	8
striped parrotfish	<i>Scarus iseri</i> (S. croicensis)	Scaridae	33.3	2	0.3	0.5	24.5	24	25	50.0	6	1.0	1.3	20.2	15	23
princess parrotfish	<i>Scarus taeniopterus</i>	Scaridae	66.7	16	2.7	2.9	17.8	10	22	100.0	16	2.7	1.6	20.9	14	30
queen parrotfish	<i>Scarus vetula</i>	Scaridae	16.7	3	0.5	1.2	24.0	18	31	16.7	1	0.2	0.4	40.0	40	40
redband parrotfish	<i>Sparisoma aurifrenatum</i>	Scaridae	100.0	18	3.0	1.9	15.5	6	20	83.3	20	3.3	2.8	15.6	5	23
redtail parrotfish	<i>Sparisoma chrysopterum</i>	Scaridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	23.0	23	23
yellowtail parrotfish	<i>Sparisoma rubripinne</i>	Scaridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	25.0	25	25
stoplight parrotfish	<i>Sparisoma viride</i>	Scaridae	16.7	2	0.3	0.8	15.0	10	20	50.0	5	0.8	1.0	26.0	13	32
cero mackerel	<i>Scomberomorus regalis</i>	Scombridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	30.0	30	30
coney yellowtail hamlet	<i>Cephalopholis fulva</i>	Serranidae	100.0	19	3.2	1.5	11.5	5	18	100.0	20	3.3	1.0	18.7	12	28
shy hamlet barred hamlet harlequin bass	<i>Hypoplectrus chlorurus</i>	Serranidae	16.7	2	0.3	0.8	12.5	10	15	0.0	0	0.0	0.0	-	-	-
sand diver sharpnose puffer	<i>Hypoplectrus guatavarius</i>	Serranidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	11.0	11	11
	<i>Hypoplectrus puella</i>	Serranidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	10.0	10	10
	<i>Serranus tigrinus</i>	Serranidae	66.7	4	0.7	0.5	8.0	6	9	66.7	10	1.7	1.6	8.9	7	10
	<i>Synodus intermedius</i>	Synodontidae	16.7	1	0.2	0.4	22.0	22	22	0.0	0	0.0	0.0	-	-	-
	<i>Canthigaster rostrata</i>	Tetraodontidae	16.7	1	0.2	0.4	4.0	4	4	50.0	3	0.5	0.5	7.7	4	13
			Total =	1922						Total =	2221					

Appendix 2E. Fish census data from Eagle Ray

Common Name	Species	Family	Spring '02 Census						Fall '02 Census						Size		
			% Freq	(n=6)	Total	Avg.	StDev	No. of Fish	Size	Min	Max	% Freq	(n=6)	Total	Avg.	StDev	No. of Fish
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	100.0	54	9.0	1.8	-	-	-	100.0	37	6.2	2.6	12.8	7	16	
doctorfish	<i>Acanthurus chirurgus</i>	Acanthuridae	0.0	0	0.0	0.0	-	-	-	16.7	3	0.5	1.2	18.0	16	20	
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	66.7	6	1.0	0.9	15.625	12	18	83.3	14	2.3	1.8	13.7	8	18	
trumpetfish	<i>Aulostomus maculatus</i>	Aulostomidae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	41.0	32	50	
Melichthys	<i>Melichthys niger</i>	Balistidae	16.7	2	0.3	0.8	23	22	24	0.0	0	0.0	0.0	-	-	-	
black durgon	<i>Caranx ruber</i>	Carangidae	0.0	0	0.00	0.00	-	-	-	33.3	7	1.2	2.0	22.5	20	30	
bar jack	<i>Caranx ignobilis</i>	Carangidae	0.0	0	0.00	0.00	-	-	-	33.3	7	1.2	2.0	22.5	20	30	
foureye	<i>Chaetodon capistratus</i>	Chaetodontidae	83.3	11	1.8	1.0	9.2	7	12	83.3	13	2.2	1.2	10.1	9	12	
butterflyfish	<i>Chaetodon striatus</i>	Chaetodontidae	0.0	0	0.00	0.00	-	-	-	66.7	8	1.3	1.0	11.5	10	15	
banded butterflyfish	<i>Haemulon carbonarium</i>	Haemulidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	30.0	30	30	
caesar grunt	<i>Haemulon flavolineatum</i>	Haemulidae	83.3	10	1.7	1.8	15.8	15	17	100.0	13	2.2	1.0	14.8	12	18	
french grunt	<i>Haemulon plumieri</i>	Haemulidae	33.3	2	0.3	0.5	23	21	25	0.0	0	0.0	0.0	-	-	-	
white grunt	<i>Holocentrus acensionis</i>	Holocentridae	0.0	0	0.0	0.0	-	-	-	50.0	6	1.0	1.3	19.0	17	31	
squirrelfish	<i>Holocentrus rufus</i>	Holocentridae	50.0	4	0.7	0.8	18.167	17	20	33.3	2	0.3	0.5	16.0	16	16	
longspine squirrelfish	<i>Myripristis jacobus</i>	Holocentridae	50.0	19	3.2	4.8	15.167	12	17	83.3	18	3.0	3.6	13.8	10	18	
blackbar soldierfish	<i>Bodianus rufus</i>	Labridae	16.7	1	0.2	0.4	20	20	20	66.7	5	0.8	0.8	11.4	5	25	
spanish hogfish	<i>Clepticus parrae</i>	Labridae	66.7	96	16.0	12.9	7	3	12	50.0	260	43.3	70.9	6.3	2	15	

Appendix 2E (continued). Fish census data from Eagle Ray

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq	(n=6)	Total	Avg.	StDev	No. of Fish	Size	% Freq	(n=6)	Total	Avg.	StDev	No. of Fish	Size
yellowcheek wrasse	<i>Halichoeres cyanopephalus</i>	Labridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	-	15.0	15
yellowhead wrasse	<i>Halichoeres garnoti</i>	Labridae	100.0	18	3.0	1.4	8.3333	3	15	83.3	29	4.8	3.3	8.6	4	16
bluehead wrasse	<i>Thalassoma bifasciatum</i>	Labridae	100.0	1460	243.3	36.1	4.25	2	10	100.0	805	134.2	75.7	4.6	2	12
mutton snapper	<i>Lutjanus analis</i>	Lutjanidae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	35.0	35	35
schoolmaster	<i>Lutjanus apodus</i>	Lutjanidae	16.7	2	0.3	0.8	17	17	17	16.7	1	0.2	0.4	17.0	17	17
yellowtail snapper	<i>Ocyurus chrysuris</i>	Lutjanidae	83.3	10	1.7	1.8	20.1	15	25	50.0	5	0.8	1.2	20.7	20	25
yellow goatfish	<i>Mulloidichthys martinicus</i>	Mullidae	33.3	2	0.3	0.5	23.5	22	25	16.7	7	1.2	2.9	17.0	0	19
spotted goatfish	<i>Pseudupeneus maculatus</i>	Mullidae	50.0	3	0.5	0.5	16.333	14	19	83.3	6	1.0	0.6	17.2	15	20
honeycomb cowfish	<i>Acanthostracion polyacanthus</i>	Ostraciidae	16.7	1	0.2	0.4	24	24	24	0.0	0	0.0	0.0	-	-	-
spotted trunkfish	<i>Lactophrys bicaudalis</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	13.0	13	13
smooth trunkfish	<i>Lactophrys triqueter</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	14.0	14	14
rock beauty	<i>Holacanthus tricolor</i>	Pomacanthidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	17.0	17	17
sergeant major blue chromis	<i>Abudefduf saxatilis</i>	Pomacentridae	33.3	12	2.0	3.1	11.5	10	14	83.3	57	9.5	7.4	11.7	8	14
brown chromis	<i>Chromis cyanus</i>	Pomacentridae	100.0	104	17.3	12.9	4.6667	2	8	100.0	200	33.3	11.1	5.0	3	8
yellowtail damselfish	<i>Microspathodon chrysurus</i>	Pomacentridae	16.7	10	1.7	4.1	3	3	4	66.7	260	43.3	68.4	3.6	3	6
			66.7	7	1.2	1.5	11.25	6	12	33.3	4	0.7	1.2	11.0	8	12

## Appendix 2E (continued). Fish census data from Eagle Ray

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			No. of Fish			Size			No. of Fish			Size				
			% Freq (n=6)	Total	Avg.	StDev	Avg	Min	Max	Total	Avg.	StDev	Avg.	Min	Max	
dusky damselfish	<i>Stegastes adustus</i> (S. fuscus)	Pomacentridae	83.3	31	5.2	4.0	7.2	5	10	83.3	22	3.7	2.2	8.2	6	12
beaugregory bicolor damselfish	<i>Stegastes leucostictus</i>	Pomacentridae	66.7	9	1.5	1.5	6.5	4	9	33.3	7	1.2	1.8	9.5	10	12
cocoa damselfish	<i>Stegastes partitus</i>	Pomacentridae	100.0	192	32.0	8.1	4.3333	2	7	100.0	251	41.8	19.6	3.8	2	6
striped parrotfish	<i>Stegastes variabilis</i>	Pomacentridae	16.7	5	0.8	2.0	6	5	8	33.3	6	1.0	1.7	9.0	5	12
princess parrotfish	<i>Scarus iserti</i> (S. croicensis)	Scaridae	50.0	8	1.3	2.0	16.833	10	21	50.0	20	3.3	5.6	14.0	7	21
queen parrotfish	<i>Scarus taeniopterus</i>	Scaridae	83.3	15	2.5	1.9	17.7	10	26	66.7	13	2.2	2.2	15.5	10	22
redband parrotfish	<i>Scarus vetula</i>	Scaridae	16.7	3	0.5	1.2	30	25	35	0.0	0	0.0	0.0	-	-	-
parrotfish redtail	<i>Spurisoma aurofrenatum</i>	Scaridae	83.3	21	3.5	2.7	16.9	8	24	83.3	26	4.3	3.1	16.9	10	22
parrotfish stoplight	<i>Spurisoma chrysopurum</i>	Scaridae	50.0	4	0.7	0.8	26.5	20	32	0.0	0	0.0	0.0	-	-	-
parrotfish stoplight	<i>Spurisoma viride</i>	Scaridae	66.7	5	0.8	0.8	23	5	35	83.3	9	1.5	1.0	25.9	7	40
graysby coney	<i>Cephalopholis crenatus</i>	Serranidae	66.7	9	1.5	1.6	16.5	14	25	50.0	4	0.7	0.8	18.7	17	20
red hind yellowtail hamlet	<i>Cephalopholis fulvus</i>	Serranidae	50.0	7	1.2	1.3	14.667	10	17	66.7	11	1.8	1.6	17.6	12	21
black hamlet butter hamlet	<i>Epinephelus guttatus</i>	Serranidae	0.0	0	0.0	-	-	-	16.7	1	0.2	0.4	18.0	18	18	
	<i>Hoploplectrus chlorurus</i>	Serranidae	16.7	1	0.2	0.4	12	12	0.0	0	0.0	0.0	-	-	-	
	<i>Hoploplectrus nigricans</i>	Serranidae	16.7	1	0.2	0.4	13	13	0.0	0	0.0	0.0	-	-	-	
	<i>Hoploplectrus unicolor</i>	Serranidae	16.7	1	0.2	0.4	11	11	0.0	0	0.0	0.0	-	-	-	

Appendix 2E (continued). Fish census data from Eagle Ray

Common Name	Species	Family	Spring '02 Census						Fall '02 Census					
			% Freq (n=6)		No. of Fish		Size		% Freq (n=6)		No. of Fish		Size	
			Total	Avg.	Total	Avg.	StDev	Min	Max	Total	Avg.	StDev	Min	Max
tobacofish	<i>Serranus tabacarius</i>	Serranidae	16.7	2	0.3	0.8	4	4	4	0.0	0	0.0	0.0	-
harlequin bass	<i>Serranus tigrinus</i>	Serranidae	16.7	3	0.5	1.2	8	7	9	50.0	6	1.0	1.3	6.3
jolthead porgy	<i>Calamus bajonado</i>	Sparidae	33.3	2	0.3	0.5	25.5	25	26	0.0	0	0.0	0.0	-
sharpnose puffer	<i>Canthigaster rostrata</i>	Tetraodontidae	33.3	3	0.5	0.8	5.75	5	6	33.3	3	0.5	0.8	5.5
			Total =		2156					Total =		2148		

Appendix 2F. Fish census data from Cane Bay

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)		No. of Fish		Size		% Freq (n=6)		No. of Fish		Size			
			Total	Avg.	StDev	Avg.	Min	Max	Total	No.	Avg.	StDev	Avg.	Min		
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	100.0	24	4.0	1.1	14.7	10	20	100.0	29	4.8	2.2	16.0	11	20
doctorfish	<i>Acanthurus chirurgus</i>	Acanthuridae	16.7	1	0.2	0.4	15.0	15	15	50.0	5	0.8	1.0	17.0	15	20
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	100.0	13	2.2	0.4	15.8	10	20	66.7	10	1.7	1.6	15.3	11	18
trumpetfish	<i>Aulostomus maculatus</i>	Aulostomidae	16.7	1	0.2	0.4	35.0	35	35	0.0	0	0.0	0.0	-	-	-
orange-spotted filefish	<i>Cantherhines pullus</i>	Monacanthidae	33.3	3	0.5	0.8	12.8	12	14	33.3	3	0.5	0.9	15.5	14	16
black durgon	<i>Melichthys niger</i>	Balistidae	100.0	39	6.5	6.4	22.0	12	28	100.0	66	11.0	6.7	22.7	13	30
bar jack	<i>Caranx ruber</i>	Carangidae	16.7	1	0.2	0.4	17.0	17	17	83.3	11	1.8	1.5	22.0	12	30
soureye	<i>Chaetodon capistratus</i>	Chaetodontidae	66.7	8	1.3	1.0	11.3	8	15	50.0	5	0.8	1.0	9.7	8	10
butterflyfish	<i>Gramma loreto</i>	Grammatidae	50.0	10	1.7	2.1	2.7	1	3	66.7	32	5.3	4.6	4.0	2	5
fairy basslet	<i>Haemulon carbonarium</i>	Haenulidae	0.0	0	0.0	0.0	-	-	-	33.3	2	0.3	0.5	21.5	18	25
caesar grunt	<i>Haemulon chrysargyreum</i>	Haenulidae	0.0	0	0.0	0.0	-	-	-	16.7	2	0.3	0.9	14.5	14	15
smallmouth grunt	<i>Haemulon flavolineatum</i>	Haenulidae	50.0	20	3.3	6.7	13.5	12	15	66.7	13	2.2	2.2	16.0	12	20
french grunt	<i>Haemulon plumieri</i>	Haenulidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	18.0	18	18
white grunt	<i>Haemulon sciurus</i>	Haenulidae	16.7	3	0.5	1.2	22.0	20	25	16.7	4	0.7	1.6	24.0	20	25
bluestriped grunt	<i>Holocentrus adscensionis</i>	Holocentridae	16.7	1	0.2	0.4	13.0	13	13	16.7	3	0.5	1.3	18.0	16	20
squirrelfish longspine	<i>Holocentrus rufus</i>	Holocentridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	15.0	15	15
squirrelfish blackbar	<i>Myripristis jacobus</i>	Holocentridae	16.7	2	0.3	0.8	15.0	15	15	33.3	5	0.8	1.7	17.0	15	20

i

Appendix 2F (continued). Fish census data from Cane Bay

Common Name	Species	Family	Spring '02 Census						Fall '02 Census						
			% Freq (n=6)	Total	Avg.	StDev	No. of Fish	Size	% Freq (n=6)	Total No.	Avg. No.	StDev	No. of Fish	Size	
longjaw squirrelfish	<i>Neoniphon mariannus</i>	Holocentridae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	13.0	13	
spanish hogfish	<i>Bodianus rufus</i>	Labridae	50.0	3	0.5	0.5	19.7	17	22	33.3	2	0.3	0.5	25.0	20
creole wrasse	<i>Clepticus parrae</i>	Labridae	66.7	311	51.8	84.9	7.0	1	16	50.0	2332	388.7	463.4	4.0	2
yellowhead wrasse	<i>Halichoeres garnoti</i>	Labridae	66.7	10	1.7	1.6	8.0	5	12	66.7	24	4.0	4.3	6.1	4
bluehead wrasse	<i>Thalassoma bifasciatum</i>	Labridae	100.0	722	120.3	43.5	2.3	1	6	100.0	600	100.0	91.0	4.4	2
schoolmaster	<i>Lutjanus apodus</i>	Lutjanidae	0.0	0	0.0	0.0	-	-	-	50.0	5	0.8	1.0	36.2	30
mahogany snapper	<i>Lutjanus maculatus</i>	Lutjanidae	0.0	0	0.0	0.0	-	-	-	33.3	4	0.7	1.3	19.5	18
yellow goatfish	<i>Mulloidichthys martinicus</i>	Mullidae	33.3	5	0.8	1.3	16.8	12	22	50.0	12	2.0	3.4	17.0	12
spotted trunkfish	<i>Lactophrys bicaudalis</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	18.0	18
sergeant major	<i>Abudefduf saxatilis</i>	Pomacentridae	83.3	10	1.7	1.0	11.8	10	15	83.3	13	2.2	2.1	13.0	10
blue chromis brown chromis	<i>Chromis cyanus</i>	Pomacentridae	100.0	682	113.7	79.0	3.7	1	10	100.0	642	107.0	62.3	5.0	2
yellowtail damselfish	<i>Microspathodon chrysurus</i>	Pomacentridae	83.3	12	2.0	1.1	9.9	4	14	83.3	13	2.2	1.2	13.3	4
dusky damselfish	<i>Stegastes partitus</i>	Pomacentridae	100.0	227	37.8	14.0	3.7	1	7	100.0	285	47.5	26.0	4.5	2
beaugregory bicolor damselfish	<i>Steagastes leucostictus</i>	Pomacentridae	100.0	48	8.0	5.3	8.2	5	12	100.0	36	6.0	2.8	9.8	5
	<i>Steagastes fuscus</i>	Pomacentridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	7.0	7

Appendix 2F (continued). Fish census data from Cane Bay

Common Name	Species	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)		No. of Fish		Size		% Freq (n=6)		No. of Fish		Size			
			Total	Avg.	SDDev	Avg	Min	Max	Total	Avg.	No.	StDev	Avg	Min		
threespot damselfish	<i>Steagastes planifrons</i>	Pomacentridae	50.0	10	1.7	2.0	5.7	3	10	66.7	24	4.0	5.4	7.3	3	11
striped parrotfish	<i>Scarus iserti (S. eroicensis)</i>	Scaridae	0.0	0	0.0	0.0	-	-	-	50.0	11	1.8	2.4	8.7	5	25
princess parrotfish	<i>Scarus taeniopterus</i>	Scaridae	100.0	18	3.0	1.1	16.9	10	22	83.3	20	3.3	3.2	19.3	5	30
queen parrotfish	<i>Scarus venula</i>	Scaridae	16.7	1	0.2	0.4	32.0	32	32	0.0	0	0.0	0.0	-	-	-
redband parrotfish	<i>Sparisoma aurofrenatum</i>	Scaridae	83.3	20	3.3	2.3	17.4	7	24	66.7	14	2.3	1.6	16.8	10	30
stoplight parrotfish	<i>Sparisoma viride</i>	Scaridae	50.0	5	0.8	1.2	20.7	6	35	83.3	17	2.8	2.3	27.0	5	35
graysby coney	<i>Cephalopholis cinctus</i>	Serranidae	33.3	2	0.3	0.5	20.0	18	22	83.3	11	1.8	0.8	18.3	15	23
black hamlet	<i>Cephalopholis fulvus</i>	Serranidae	50.0	6	1.0	1.1	14.8	12	18	50.0	5	0.8	1.2	17.0	15	20
harlequin bass	<i>Hopplectrus nigricans</i>	Serranidae	0.0	0	0.0	0.0	-	-	-	50.0	3	0.5	0.5	9.0	6	12
sharpnose puffer	<i>Serranus tigrinus</i>	Serranidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	6.0	6	6
	<i>Canthigaster rostrata</i>	Tetraodontidae	16.7	1	0.2	0.4	6.0	6	6	33.3	4	0.7	1.1	5.5	4	7
			Total =	3120										Total =	5443	

Appendix 2G. Fish census data from Sprat Hole

Common Name	Species Name	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)		No. of Fish		Size		% Freq (n=6)		No. of Fish		Size			
			Total	Avg.	StDev	Avg.	Min	Max	Total	Avg.	StDev	Avg.	Min	Max		
ocean surgeonfish	<i>Acanthurus bahianus</i>	Acanthuridae	83.3	16	2.7	2.1	15.1	12	18	83.3	14	2.3	2.3	15.1	12	20
blue tang	<i>Acanthurus coeruleus</i>	Acanthuridae	83.3	15	2.5	1.5	14.4	10	18	83.3	9	1.5	1.0	16.8	15	20
trumpetfish	<i>Autostomus maculatus</i>	Autostomidae	16.7	1	0.2	0.4	50.0	50	50	50.0	3	0.5	0.5	24.7	15	34
blue runner	<i>Caranx cryos</i>	Carangidae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	30.0	30	30
bar jack	<i>C. fuscus</i>	Carangidae	33.3	5	0.8	1.6	28.0	26	30	16.7	2	0.3	0.8	21.0	20	22
foureye butterflyfish	<i>Caranx ruber</i>	Carangidae	33.3	5	0.8	1.6	28.0	26	30	16.7	2	0.3	0.8	21.0	20	22
yellowfin mojarra	<i>Chaetodon capistratus</i>	Chaetodontidae	33.3	4	0.7	1.0	9.0	7	11	83.3	10	1.7	0.8	9.6	7	12
fairy basslet	<i>Gerres cinereus</i>	Gerridae	16.7	2	0.3	0.8	18.5	17	20	0.0	0	0.0	0.0	-	-	-
caesar grunt	<i>Gramma loreto</i>	Grammatidae	16.7	1	0.2	0.4	2.0	2	2	50.0	5	0.8	1.0	4.0	3	5
french grunt	<i>Haemulon carbonarium</i>	Haemulidae	16.7	2	0.3	0.8	17.5	15	20	0.0	0	0.0	0.0	-	-	-
squirrelfish	<i>H. flavolineatum</i>	Haemulidae	66.7	7	1.2	1.2	15.5	14	17	66.7	5	0.8	0.8	17.0	15	20
blackbar soldierfish	<i>H. adamsi</i>	Holocentridae	16.7	1	0.2	0.4	17.0	17	17	0.0	0	0.0	0.0	-	-	-
spanish hogfish	<i>Myripristis jacobus</i>	Holocentridae	83.3	16	2.7	2.9	14.2	10	17	50.0	12	2.0	2.3	15.7	14	21
creole wrasse	<i>Bodianus rufus</i>	Labridae	16.7	1	0.2	0.4	22.0	22	22	50.0	6	1.0	1.3	15.2	5	25
yellowhead wrasse	<i>Clepticus parviceps</i>	Labridae	100.0	1467	244.5	252.7	5.5	1	14	66.7	685	114.2	240.1	9.0	2	16
clown wrasse	<i>Halichoeres garnoti</i>	Labridae	83.3	100	16.7	31.3	6.3	2	14	66.7	19	3.2	3.0	7.6	4	12
bluehead wrasse	<i>Halichoeres maculipinnis</i>	Labridae	0.0	0	0.0	0.0	-	-	-	16.7	1	0.2	0.4	11.0	11	11
	<i>Thalassoma bifasciatum</i>	Labridae	83.3	650	108.3	79.6	2.7	1	10	83.3	325	54.2	31.8	4.0	2	10

Appendix 2G (continued). Fish census data from Sprat Hole

Common Name	Species Name	Family	Spring '02 Census						Fall '02 Census							
			% Freq (n=6)		No. of Fish		Size		% Freq (n=6)		No. of Fish		Size			
			Total	Avg.	StDev	Avg.	Min	Max	Total	Avg.	StDev	Avg.	Min	Max		
nutton snapper	<i>Lutjanus analis</i>	Lutjanidae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	38.0	38	38	
	<i>Lutjanus apodus</i>	Lutjanidae	16.7	2	0.3	0.8	47.0	40	55	16.7	1	0.2	0.4	40.0	40	40
schoolmaster mahogany snapper	<i>Lutjanus mahogoni</i>	Lutjanidae	66.7	20	3.3	5.0	16.3	10	20	100.0	20	3.3	1.6	20.3	15	25
sand tilefish yellow goatfish spotted	<i>Malacanthus plumieri</i>	Malacanthidae	16.7	1	0.2	0.4	40.0	40	40	0.0	0	0.0	0.0	-	-	-
moray honeycomb cowfish smooth trunkfish	<i>Mulloidichthys martinicus</i>	Mullidae	33.3	6	1.0	2.0	14.0	12	16	50.0	127	21.2	31.2	14.3	10	24
rock beauty sergeant major blue chromis brown chromis yellowtail damselfish	<i>Gymnothorax moringa</i>	Muraenidae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	60.0	60	60	
dusky damselsfish bicolor damselsfish threespot damselsfish	<i>Acanthostracion polygonia</i>	Ostraciidae	16.7	1	0.2	0.4	30.0	30	30	0.0	0	0.0	0.0	-	-	-
	<i>Iactophrys trispinosus</i>	Ostraciidae	0.0	0	0.0	0.0	-	-	16.7	1	0.2	0.4	10.0	10	10	
	<i>Holacanthus tricolor</i>	Pomacentridae	33.3	2	0.3	0.5	13.5	12	15	50.0	4	0.7	0.8	18.7	15	22
	<i>Abudefduf saxatilis</i>	Pomacentridae	16.7	2	0.3	0.8	13.0	13	13	0.0	0	0.0	0.0	-	-	-
	<i>Chromis cyanus</i>	Pomacentridae	100.0	800	133.3	48.9	2.8	1	7	100.0	789	131.5	61.9	4.7	2	9
	<i>Chromis multilineata</i>	Pomacentridae	100.0	701	116.8	32.0	4.5	1	7	83.3	650	108.3	70.3	4.8	2	8
	<i>Microspathodon chrysurus</i>	Pomacentridae	16.7	2	0.3	0.8	10.0	10	0.0	0	0	0.0	0.0	-	-	-
	<i>Sleagases adustus</i> (S. fuscus)	Pomacentridae	83.3	29	4.8	3.8	6.4	4	10	66.7	26	4.3	4.4	10.3	7	15
	<i>Sleagases parvitus</i>	Pomacentridae	100.0	235	39.2	13.6	4.1	1	6	100.0	397	66.2	29.6	4.5	2	10
	<i>Sleagases planifrons</i>	Pomacentridae	100.0	101	16.8	5.7	5.5	3	8	83.3	70	11.7	9.2	8.2	4	15

Appendix 2G (continued). Fish census data from Sprat Hole

Common Name	Species Name	Family	Spring '02 Census						Fall '02 Census						
			% Freq (n=6)	Total	No. of Fish	Avg	Size Min	Max	% Freq (n=6)	Total	No. of Fish	Avg	Size Min	Max	
cocoa damselfish	<i>Stegastes variabilis</i>	Pomacentridae	16.7	1	0.2	0.4	5.0	5	5	0.0	0	0.0	0.0	-	-
striped parrotfish	<i>Scarus iseri (S. croicensis)</i>	Scaridae	16.7	4	0.7	1.6	5.0	4	6	83.3	26	4.3	2.7	11.6	4
princess parrotfish	<i>Scarus taeniopterus</i>	Scaridae	100.0	46	7.7	5.0	17.2	6	25	100.0	35	5.8	4.0	18.6	5
queen parrotfish	<i>Scarus velula</i>	Scaridae	66.7	10	1.7	1.5	26.8	15	45	33.3	2	0.3	0.5	27.5	30
redband parrotfish	<i>Sparisoma aurofrenatum</i>	Scaridae	100.0	41	6.8	1.9	16.2	10	25	83.3	25	4.2	2.7	16.0	3
stoplight parrotfish	<i>Sparisoma viride</i>	Scaridae	100.0	13	2.2	0.8	21.6	6	40	66.7	14	2.3	2.6	13.5	4
grayshy Cephalopholis crenatus	<i>Cephalopholis crenatus</i>	Serranidae	66.7	9	1.5	1.4	14.9	12	19	66.7	7	1.2	1.0	16.5	10
coney yellowtail hamlet	<i>Hoploplectrus fulvus chlorurus</i>	Serranidae	16.7	4	0.7	1.6	17.0	15	20	66.7	9	1.5	1.5	16.1	8
shy hamlet black hamlet	<i>Hoploplectrus gutatarius</i>	Serranidae	0	0	0.0	0.0	-	-	-	16.7	2	0.3	0.8	9.0	8
barred hamlet harlequin bass	<i>Hoploplectrus phaelia</i>	Serranidae	16.7	1	0.2	0.4	8.0	8	8	0.0	0	0.0	0.0	-	-
sand diver sharpnose puffer porcupinefish	<i>Serranus tigrinus intermedius</i>	Serranidae	0.0	0	0.0	0.0	-	-	-	50.0	3	0.5	0.5	10.0	9
	<i>Synodus canthigaster</i>	Synodontidae	33.3	3	0.5	0.8	13.5	10	15	16.7	1	0.2	0.4	10.0	10
	<i>Canthigaster rostrata</i>	Tetraodontidae	33.3	3	0.5	0.8	4.2	3	5	50.0	6	1.0	1.3	4.5	6
	<i>Diodon hystrix</i>	Tetraodontidae	0.0	0	0.0	0.0	-	-	-	16.7	2	0.3	0.8	35.0	35
			Total =	4326						Total =	3319				