

SEDAR

Southeast Data, Assessment, and Review

SEDAR 30

Stock Assessment Report

U.S. Caribbean Queen Triggerfish

April 2013

SEDAR


4055 Faber Place Drive, Suite 201

North Charleston, SC 29405

Table of Contents

Section I. Introduction	PDF Page 3
Section II. Assessment Workshop Report	PDF Page 30
Section III. Review Report	PDF Page 106
Section IV. Research Recommendations	PDF Page 186

SEDAR



Southeast Data, Assessment, and Review

SEDAR 30

U.S. Caribbean Queen Triggerfish

SECTION I: Introduction

SEDAR
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405

1. SEDAR PROCESS DESCRIPTION

SouthEast Data, Assessment, and Review (**SEDAR**) is a cooperative Fishery Management Council process initiated in 2002 to improve the quality and reliability of fishery stock assessments in the South Atlantic, Gulf of Mexico, and US Caribbean. SEDAR seeks improvements in the scientific quality of stock assessments and the relevance of information available to address fishery management issues. SEDAR emphasizes constituent and stakeholder participation in assessment development, transparency in the assessment process, and a rigorous and independent scientific review of completed stock assessments.

SEDAR is managed by the Caribbean, Gulf of Mexico, and South Atlantic Regional Fishery Management Councils in coordination with NOAA Fisheries and the Atlantic and Gulf States Marine Fisheries Commissions. Oversight is provided by a Steering Committee composed of NOAA Fisheries representatives: Southeast Fisheries Science Center Director and the Southeast Regional Administrator; Regional Council representatives: Executive Directors and Chairs of the South Atlantic, Gulf of Mexico, and Caribbean Fishery Management Councils; and Interstate Commission representatives: Executive Directors of the Atlantic States and Gulf States Marine Fisheries Commissions.

SEDAR is normally organized around three workshops and a series of webinars. First is the Data Workshop, during which fisheries, monitoring, and life history data are reviewed and compiled. The second stage is the Assessment Process, which is conducted via a workshop and a series of webinars, during which assessment models are developed and population parameters are estimated using the information provided from the Data Workshop. Third and final is the Review Workshop, during which independent experts review the input data, assessment methods, and assessment products. The completed assessment, including the reports of all 3 workshops and all supporting documentation, is then forwarded to the Council SSC for certification as ‘appropriate for management’ and development of specific management recommendations.

SEDAR workshops are public meetings organized by SEDAR staff and the lead Cooperator. Workshop participants are drawn from state and federal agencies, non-government organizations, Council members, Council advisors, and the fishing industry with a goal of including a broad range of disciplines and perspectives. All participants are expected to contribute to the process by preparing working papers, contributing, providing assessment analyses, and completing the workshop report.

SEDAR 30 was organized differently from the normal SEDAR process, due to the unique data and cultural issues for the U.S. Caribbean. Given the limited amount of data available for use in the assessment, and the assessment approach being used, a formal Data Workshop was not conducted. Instead a series of public meetings was held (one each on Puerto Rico, Saint Thomas, and Saint Croix) in which the data was discussed with local scientists and fisherman.

An Assessment Workshop was held, as per standard SEDAR procedures, but an in-person Review Workshop was not conducted and replaced by several CIE Desk Reviews. This modification of the Review Process was also related to the limited data availability and assessment methods used during this process.

2. MANAGEMENT OVERVIEW ((Prepared by Graciela García-Moliner and Bill Arnold)

2.1. Fishery Management Plan, Plan Amendments, and Local Regulations

Queen triggerfish (*Belistes vetula*) is one of several species of triggerfish within the triggerfish and filefish fishery management unit (CFMC 2005). The regulatory overview describes the management measures that directly or indirectly impact queen triggerfish. There are no regulations specific to any of the species of triggerfish and filefish in the US Caribbean.

The U.S. Caribbean includes the islands of Puerto Rico and the U.S. Virgin Islands (USVI) including St. Thomas, St. John, and St. Croix. The state waters of Puerto Rico extend 9 nm from the shore and the state waters of the USVI extend 3 nm from shore. The following summary applies to these jurisdictions separately.

The following is a summary of the management measures that directly or indirectly have impacted the queen triggerfish fishery in the U.S. Caribbean. The Fishery Management Plan for the Shallow-water Reef Fish Fishery of Puerto Rico and the U.S. Virgin Islands (1985) included 4 species of triggerfish (Balistidae) in the FMU, queen triggerfish, ocean triggerfish, black durgon, and Sargassum triggerfish (*Balistes vetula*, *Canthidermis sufflamen*, *Melichthys niger*, *Xanthichthys ringens*), it did not include the orange spotted or gray triggerfish (*Catherines pullus* or *Balistes capriscus*) and did not include size limits, seasonal closures or other management measures directed at the triggerfish fishery in the EEZ. However, regulations on the mesh size of traps that were implemented at the time had a direct impact on the triggerfish fishery. The orange spotted triggerfish (*C. pullus*) was incorporated into Reef Fish Fishery Management Plan in 1993 along with the scrawled and whitespotted filefish (Monacanthidae: *Aluterus scriptus* and *Cantherhines macrocerus*). The 1993 Amendment changes the name of the FMP formerly known as the Fishery Management Plan for the Shallow-water Reef Fish Fishery of Puerto Rico and the U.S. Virgin Islands; Federal Register, Vol. 50, No. 167: 34850-34855). .

Measures in the original FMP, and in the follow-on amendments, that affect triggerfish include changes to requirements for the constructions of traps (in both the Spiny Lobster and Shallow-water Reef Fish FMPs) as well as seasonal and/or area closures established through amendments to the Reef Fish FMP and the Coral FMP:

Description of Action	FMP/Amendment	Effective Date
Traps: construction and requirement for degradable panel	Spiny Lobster FMP	1985
Traps: construction and requirement for degradable panel; changes to mesh size.	Reef Fish FMP/Amen. 1/ Reg. Amen./Amen. 2/SFA	1985; 1990;1993;2005
Seasonal area closure	Reef Fish FMP/Amen. 2; Amen. 3/Interim Rule/ SFA	1993,1996, 1999,2005
Closed area	Coral FMP Amen. 1	1999
Seasonal closure for reef fish	SFA	2005

In 2005, the Council ratified the Sustainable Fisheries Act (SFA) Amendment, which categorized triggerfish and filefish (along with groupers and other species) into fishery management units (FMUs). Triggerfish were all included in the Triggerfish and Filefish FMU (species included are: queen triggerfish (in the USVI is known as old wife) (*Balistes vetula*), ocean triggerfish (*Canthidermis sufflamen*), black durgon (*Melichthys niger*), Sargassum triggerfish (*Xanthichthys ringens*), scrawled filefish (*Aluterus scriptus*) and whitespotted filefish (*Cantherhines macrocerus*) [NOTE: The orange spotted triggerfish, *Catherines pullus*, is not included in the latest Triggerfish and Filefish FMU.]. Measures were included in the SFA Amendment to institute a total ban on the use of nets in the EEZ and a prohibition on the bottom tending gear in all seasonally closed areas (HAPCs). A number of closed seasons for various FMUs during the peak spawning months of each species were also implemented and indirectly impact triggerfish. The implementation took place on November 28, 2005, and continues to the present. At the time of implementation these measures applied only to U.S. Caribbean EEZ waters.

Compatible regulations on the ban of nets were implemented by the Government of the Territory of the USVI for their local water on July 5, 2006. Also, the prohibition on the use of bottom tending gear (including traps) from the HAPCs was implemented by the USVI Government at the time.

Puerto Rico followed a slightly different strategy by implementing a ban on the use of SCUBA and nets fished together in 2004 (Regulation 6768 with an implementation date of March 12, 2004), establishing a mesh size for gillnets of more than 2 inches (51 mm) (as it was since 1936) and various restrictions on the construction of beach seines (beach seines were subsequently banned and re-established since 2004 to present). Although, as noted above, the implementation of Regulation 6768 was to begin in March 2004, a subsequent Administrative Order (Number 2004-12) delayed enforcement until January 1, 2005, in an effort to provide adequate opportunity to educate fishers on the new regulations. The Administrative Order included a 2-year grace period for these restrictions to be implemented.

Regulation 6768 included a quota for the species collected for the aquarium trade. A total of 561 Sargassum triggerfish (also known as redbtail triggerfish, *Xanthichthys ringes*) were allowed to be harvested per year. All other triggerfish not included in the list of species permitted for capture and exportation were prohibited for the aquarium trade industry. In 2010, the amendment to the regulations (Reglamento de Pesca de Puerto Rico -2010 Numero 7949) listed the species permitted in the aquarium trade (capture and exportation) but did not set any quotas. Reglamento 7949 also allowed importation of the species in the family Balistidae.

Note that the closed seasons in all three island groups, were designed to benefit other species but not triggerfish (i.e., SU1 including silk snapper (October – December), Grouper Unit 4 (February-April, SU3 (mutton and lane snapper April-June), GU3 (red hind, December-February) and might have limited impact on triggerfish. However, the seasonal area closures described in Tables 2.1.1, 2.1.2, and 2.1.3 although not designed specifically for the protection of triggerfish could indirectly benefit triggerfish because the closure areas encompass habitats occupied by triggerfish species including queen triggerfish. The SFA Amendment also prohibited placement of bottom tending gear (e.g., traps, bottom longlines, nets) within Habitat Areas of Particular Concern (HAPCs).

The changes in trap construction and mesh size, the ban on bottom tending gear from HAPCs, the regulations on mesh and length of nets, the prohibition of nets in the EEZ and the USVI are all regulations that impact queen triggerfish (mostly caught by traps, but also with hook and line and spear) and the other species of trigger and filefish.

Table 2.1.1. Annual Commercial/Recreational Triggerfish Regulatory Summary: St. Croix

Year	Fishing Year	Minimum size limit			Trip limit			Closed season		Closed Area		
		Size	Start date	End date	Amount	Start date	End date	Start date	End date	Area/Seasonal ⁴	Start date	End date
1993										Lang Bank ¹	11/15	12/31
1994										Lang Bank ¹	1/1; 12/1	2/28 ² ; 12/31
										Mutton Snapper ³	3/1	6/30
1995										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
1996										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
1997										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
1998										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
1999										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2000										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2001										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2002										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2003										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2004										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30

2005										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2006										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2007										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2008										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2009										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30
2010										Lang Bank ¹	1/1; 12/1	2/28; 12/31
										Mutton Snapper ³	3/1	6/30

¹EEZ waters only; ²2/29 during leap years; ³1993 territorial area closure; 1994 EEZ and territorial area closure; 1996 boundary change to make EEZ compatible with state; ⁴Bottom tending gear (e.g., traps, nets, bottom longlines) prohibited from seasonally closed areas (i.e., HAPCs); ⁵Boundary change to Tourmaline Bank closed area; ⁶Closure extended to six months (October 1 through March 31) beginning with 2011 calendar year.

Table 2.1.2. Annual Commercial/Recreational Triggerfish Regulatory Summary: St. Thomas

Year	Fishing Year	Minimum size limit			Trip limit			Closed season		Closed Area		
		Size	Start date	End date	Amount (lbs)	Start date	End date	Start date	End date	Area/Seasonal ⁴	Start date	End date
1990										Hind Bank(MCD)	12/1	12/31
1991										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1992										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1993										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1994										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1995										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1996										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1997										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1998										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
1999										Hind Bank(MCD)	1/1; 12/1	2/28; 12/31
2000										Hind Bank(MCD)	1/1	12/31
2001										Hind Bank(MCD)	1/1	12/31
2002										Hind Bank(MCD)	1/1	12/31
2003										Hind Bank(MCD)	1/1	12/31
2004										Hind Bank(MCD)	1/1	12/31
2005										Hind Bank(MCD)	1/1	12/31
2006										Grammanik Bank	2/1	4/30
										Hind Bank(MCD)	1/1	12/31
2007										Hind Bank(MCD)	1/1	12/31
										Grammanik Bank	2/1	4/30
2008										Hind Bank(MCD)	1/1	12/31

										Grammanik Bank	2/1	4/30
2009										Hind Bank(MCD)	1/1	12/31
										Grammanik Bank	2/1	4/30
2010										Hind Bank(MCD)	1/1	12/31
										Grammanik Bank	2/1	4/30

¹EEZ waters only; ²1993 territorial area closure; 1994 EEZ and territorial area closure; 1996 boundary change to make EEZ compatible with state; ³2/29 during leap years; ⁴Bottom tending gear (e.g., traps, nets, bottom longlines) prohibited from seasonally closed areas (i.e., HAPCs); ⁵Boundary change to Tourmaline Bank closed area; ⁶Closure extended to six months (October 1 through March 31) beginning with 2011 calendar year.

Table 2.1.3. Annual Commercial/Recreational Triggerfish Regulatory Summary: Puerto Rico

Year	Fishing Year	Minimum size limit			Trip limit			Closed season		Closed Area		
		Size	Start date	End date	Amount (lbs)	Start date	End date	Start date	End date	Area/Seasonal ⁴	Start date	End date
1983												
1984												
1985												
1986												
1987												
1988												
1989												
1990												
1991												
1992												
1993										Tourmaline Bank	11/15	12/31
1994										Tourmaline Bank	1/1; 12/1	2/28; 12/31
1995										Tourmaline Bank	1/1; 12/1	2/28; 12/31
1996										Tourmaline Bank ⁵	1/1; 12/1	2/28; 12/31
1997										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
1998										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
1999										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31

2000										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2001										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2002										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2003										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2004										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2005										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2006										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31

										Bajo de Sico	1/1; 12/1	2/28; 12/31
2007										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2008										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2009										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2010										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico	1/1; 12/1	2/28; 12/31
2011										Tourmaline Bank	1/1; 12/1	2/28; 12/31
										Abrir La Sierra	1/1; 12/1	2/28; 12/31
										Bajo de Sico ⁶	1/1; 12/1	2/28; 12/31

¹EEZ waters only; ²1993 territorial area closure; 1994 EEZ and territorial area closure; 1996 boundary change to make EEZ compatible with state; ³2/29 during leap years; ⁴Bottom tending gear (e.g., traps, nets, bottom longlines) prohibited from seasonally closed areas (i.e., HAPCs); ⁵Boundary change to Tourmaline Bank closed area; ⁶Closure extended to six months (October 1 through March 31) beginning with 2011 calendar year.

2.2 Emergency and Interim Rules:

No Emergency or Interim Rules have been developed specific to the management of triggerfishes.

2.3. Secretarial Amendments

None.

2.4. Control Date Notices

The CFMC at its 130th meeting on March 24-26, 2009, established a control date of March 24, 2009, for every fishery managed by the Council, including SU1.

2.5. Management Program Specifications

The following is a summary of general information regarding management of triggerfish in the U.S. Caribbean:

Species	Queen triggerfish
Management Unit	Triggerfishes
Management Unit Definition	Includes queen triggerfish (<i>Balistes vetula</i>), ocean triggerfish (<i>Canthidermis sufflamen</i>), black durgon (<i>Melichthys niger</i>), Sargassum triggerfish (<i>Xanthichthys ringens</i>), scrawled filefish (<i>Aluterus scriptus</i>) and whitespotted filefish (<i>Cantherhines macrocerus</i>)
Management Entity	Caribbean Fishery Management Council
Management Contacts	William Arnold - SERO
SERO / Council	Graciela García-Moliner - CFMC
Current stock exploitation status	Unknown (Not overfished)
Current stock biomass status	Unknown

As described in the following table, the 2005 SFA Amendment established reference points for triggerfishes. This fishery unit includes 6 species of trigger and filefish. The 2011 ACL Amendment established ACLs for the triggerfish and filefish for the commercial and recreational sectors separate for Puerto Rico and the commercial and recreational sectors together in the USVI. No additional measures such as bag limits for the recreational catch or size limits were implemented. This ACL 2011 Amendment also redefines management reference points based on average current catch for each geographically distinct area (i.e., Puerto Rico, St. Thomas/St. John (STT/STJ), and St. Croix (STX).

Note that reference points were based upon commercial and recreational landings only. Although discards may occur in these fisheries, there has been no available method for estimating the extent of those discards.

Criteria	Current		Proposed	
	Definition	Value	Definition	Value
MSST	$MSST = [(1-M) \text{ or } 0.5 \text{ whichever is greater}] * B_{MSY}$	686,000	$MSST = [(1-M) \text{ or } 0.5 \text{ whichever is greater}] * B_{MSY}$	SEDAR 30
MFMT	F_{MSY} Proxy	0.32	F_{MSY}	SEDAR 30
MSY	Yield at F_{MSY} Proxy	196,000	Yield at F_{MSY}	SEDAR 30
F_{MSY}	M	0.32	F_{MAX}	SEDAR 30
OY	Yield at F_{OY}	184,000	Yield at F_{OY}	SEDAR 30
F_{OY}	$F_{OY} = 0.75 * F_{MSY}$ Proxy	Not specified	$F_{OY} = 50\%, 75\%, 85\% F_{MSY}$	SEDAR 30
M		0.32		SEDAR 30

Stock Rebuilding Information

According to NOAA’s Fish Stock Sustainability Index (http://www.nmfs.noaa.gov/sfa/statusoffisheries/2011/first/FSSInonFSSIstockstatusQ1_2011.pdf), the triggerfish and filefish FMU is not considered to be undergoing overfishing and not overfished. Thus, no rebuilding plan is required.

Stock Projection Information

The 2011 ACL Amendment to the FMPs proposes the following criteria for applying AMs in the management of trigger and file fishes:

Requested Information	Value
First Year of Management	2012
Projection Criteria during interim years should be based on (e.g., exploitation or harvest)	Commercial + Recreational Landings in Puerto Rico, Commercial Landings in the USVI
Projection criteria values for interim years should be determined from (e.g., terminal year, avg of X years)	2011 landings for 2012; average of 2011-2012 landings for 2013, and average 3 years of landings for 2014 and forward

The 2011 Caribbean ACL Amendment was approved and established ACL for Triggerfish and Filefish fishes in Puerto Rico (commercial and recreational) and in the USVI:

Current Quota Value ACL (pounds)	80,404 (PR); 74,447 (STT/STJ) 24,980 (STX)
Next Scheduled Quota Change	TBD
Annual or averaged quota	Averaged
If averaged, number of years to average	6-7 years ¹
Does the quota include bycatch/discard ?	No

¹6 years for PR recreational and STT commercial; 7 years for PR and STX commercial.

How is the quota calculated - conditioned upon exploitation or average landings?

Average landings.

Does the quota include bycatch/discard estimates? If so, what is the source of the bycatch/discard values? What are the bycatch/discard allowances?

No.

Are there additional details of which the analysts should be aware to properly determine quotas for this stock?

The CFMC recognizes the limitations of the data. Improvements in data collection are anticipated resulting from more fisherman-friendly reporting forms and enhanced data collection and effort monitoring. For most effective management of the fisheries, the CFMC will need timely in-season data which is currently lacking for the U.S. Caribbean.

2.6. Management and Regulatory Timeline

Tables 2.1.1, 2.1.2, and 2.1.3 provide event timelines of State and federal EEZ management actions by island group. Additional details regarding regulatory history for Puerto Rico, USVI, and EEZ waters follow.

The principal gears used in targeting triggerfish are traps and nets (including herding with SCUBA gear (note: needs to be confirmed)) and spear. Trap and pots configurations have been regulated since 1936 in Puerto Rico, since 1972 in the USVI, and since 1985 in the EEZ. Nets (beach seines) have been regulated in Puerto Rico since 1936 and not until 2005 in the EEZ and in 2006 in the USVI.

Trap Regulations

Puerto Rico's Fishing Law of 1936 required that all traps have a self-destruct panel but made no mention of mesh size. In 1972, Act 3330 of the USVI regulated the mesh size of traps, establishing a minimum size of 1.25" in the smallest dimension. By convention rather than regulation, trap construction was of wood and hexagonal (i.e., chicken wire) wire mesh.

Starting in 1985 regulations associated with spiny lobster management established that, in all three jurisdictions, self-destruct panels must be configured into traps. Also in 1985, traps to be deployed in the EEZ had to have a minimum mesh size of 1.25" in the smallest dimension, resulting in compatible regulations between the USVI and the EEZ. However, the regulations were not implemented in federal waters until 1988. These regulations were established under the Fishery Management Plan for the Shallow-water Reef Fish Fishery of Puerto Rico and the USVI (i.e., Reef Fish FMP).

In the 1990's, amendments to the Reef Fish FMP (1990, 1991, 1993) increased the minimum mesh size for traps to 1.5" hexagonal or 1.5" square until September 13, 1993, and to 2" square thereafter. These regulations also required that the degradable panel be fastened with degradable material and established specifications for the composition of the degradable material. The intention of the CFMC was to have only one panel but as the regulations were written there was a requirement for two degradable panels

plus a degradable door. This regulation was revised through a technical amendment, resulting in a requirement for only a single biodegradable panel.

In the EEZ by 1994, regulations for traps required that traps have a minimum mesh size of 1.5” hexagonal or 2” square mesh. The documentation for this requirement is available at the Council’s office.

Act 3330 of the USVI regulates fishing and fishing gear. This 1972 regulation required that a minimum mesh size in traps be of 1.25” hexagonal in the smallest dimension and that an escape panel with degradable material also be incorporated into the traps.

In the USVI, each District imposes different requirements for trap construction but all traps have to be inspected and certified. In STT/STX beginning in 1996, all new traps must have a minimum mesh size identical to that required for the EEZ: 1.5” hexagonal or 2” square in the smallest dimension. The requirement for an escape panel with degradable fasteners is compatible in all jurisdictions.

However, in STX beginning in 1996, the minimum mesh size required for new traps was 1.5” hexagonal or 1.5” square in the smallest dimension.

Beginning in 1998, the EEZ, STT/STJ, and STX required the same mesh size of 1.5” hexagonal (but not for the square mesh) for all new traps. Also beginning in 1998, all traps in the USVI had to comply with the 1.5” minimum hexagonal mesh size. The difference between the Districts remained with STT/STJ continuing to allow a square mesh of 2” whereas STX still required a 1.5” minimum mesh.

Beginning in 2000, STT/STJ required that all traps had a minimum mesh size of 2” hexagonal or 2” square. All traps had to be inspected and tagged. This is the most restrictive minimum mesh size requirement in the U.S. Caribbean.

Beginning in 2004, Puerto Rico regulated the construction of traps and required, but did not implement, a minimum mesh size of 1.5” hexagonal or 2” square. Implementation of these mesh size requirements did not occur until 2007.

In summary, the time periods during which mesh size regulations were stable are as follows:

- 1) For the EEZ, (a) 1.25” minimum mesh from 1988 to 1993 (total of 6 years), and (b) 1.5” hexagonal or 2” square from 1994 to present (total of 17 years through 2010).
- 2) For STT/STJ, (a) 1.25” minimum mesh from 1972 to 1996 (total of 25 years), (b) 1.5” hexagonal or 2” square from 1997 to 2000 (total of 4 years), and (c) 2” square or 2” hexagonal from 2001 to present (total of 10 years through 2010).
- 3) For STX, (a) 1.25” minimum mesh from 1972 to 1996 (total of 25 years), (b) 1.5” hexagonal or 1.5” square from 1996 (new traps) to 2009 (total of 14 years) or from 1998 (all traps) to 2009 (total of 12 years), (c) 1.5” hexagonal during 2009 (total of 1 year), and (d) 1.5” hexagonal or 2” square beginning in 2010 (total of 1 year to present).

Compatibility in the mesh size regulation was achieved as follows:

(a) 1.25” mesh in both USVI and EEZ waters between 1985 and 1988 (total of 4 years), (b) 1.5” hexagonal in both STT/STJ and EEZ waters between 1996 and 2000 (total of 5 years), and (c) 2” square from 1996 to present for both STT/STJ and EEZ waters (total of 16 years through 2010).

The missing years correspond to periods when the regulations included a mix of 1.25” minimum mesh, 1.5” hexagonal or square mesh, and 2” square mesh.

Compatibility of EEZ and STX regulations included (a) 1.25” between 1988 and 1994 (total of 7 years), (b) 1.5” hexagonal between 1996 and present (total of 16 years through 2010), and (c) for 2” square in 2010 (total of 1 year through 2010).

Tables 2.6.1, 2.6.2, and 2.6.3 summarize the number of years during which the mesh size was the same for the EEZ and each of the islands or island groups. Figure 2.7.1 provides a visual timeline for changes in regulations, clearly showing the predominance of regulations regarding mesh size.

Table 2.6.1. EEZ and STT/STJ years of compatible mesh size for traps.

		EEZ			
		1.25"	1.5"	2" square	2" hex
		1988	1994	1994	NO
STT	1.25"	1972	7 (1988-1996)		
	1.5"	1996	5 (1996-2000)		
	2" square	1996		5 (1996-2010)	
	2" hex	2000			

Table 2.6.2. EEZ and STX years of compatible mesh size in traps.

		EEZ			
		1.25"	1.5"	2" square	2" hex
		1988	1994	1994	NO
STX	1.25"	1972	7 (1988-1994)		
	1.5"	1996	5 (1996-2010)		
	2" square	2010		1 (2010)	
	2" hex				

Table 2.6.3. EEZ and Puerto Rico (this table needs to be reviewed).

		EEZ			
		1.25"	1.5"	2" square	2" hex
		1988	1994	1994	NO
PR	1.25"				
	1.5"	2007	4 (2007-2010)		
	2" square	2007	4 (2007-2010)		
	2" hex				

Table 2.6.4. Periods of time during which the same regulation of mesh size was maintained in each of the areas or jurisdictions.

Years during which the same mesh size was maintain in each jurisdiction				
Mesh/Area	EEZ	USVI	STT	STX
1.25"	1988-1993	1972-1996	1972-1998	1972-1998
1.5"hex/square	1993-1994			1996-1998
1.5" hex	1994-2010		1996-2000	1996-2010
2"square	1994-2010		1996-2010	2010
2" hex			2000-2010	




Figure 2.6.1. Schematic representation of the various FMPs and regulations that changed the way in which traps were constructed. The most significant changes are for mesh size.

2.7 References

Caribbean Fishery Management Council (CFMC). 1981. Fishery management plan, final environmental impact statement, and regulatory impact review for the spiny lobster fishery of Puerto Rico and the U.S. Virgin Islands. Caribbean Fishery Management Council, San Juan, Puerto Rico. 43 pp. + Appendices.

Caribbean Fishery Management Council (CFMC). 1985. Fishery management plan, final environmental impact statement, and draft regulatory impact review for the shallow-water reef fish fishery of Puerto Rico and the U.S. Virgin Islands. Caribbean Fishery Management Council, San Juan, Puerto Rico. 69 pp. + Appendices.

Caribbean Fishery Management Council (CFMC). 1990a. Amendment number 1 to the fishery management plan for the shallow-water reef fish fishery, preliminary environmental assessment and regulatory impact review. Caribbean Fishery Management Council, San Juan, Puerto Rico. 51 pp. + Appendices.

- Caribbean Fishery Management Council (CFMC). 1990b. Amendment number 1 to the fishery management plan for the spiny lobster fishery of Puerto Rico and the U.S. Virgin Islands. Caribbean Fishery Management Council, San Juan, Puerto Rico. 24 pp.
- Caribbean Fishery Management Council (CFMC). 1991. Regulatory amendment to the shallow water reef fish fishery management plan. Caribbean Fishery Management Council, San Juan, Puerto Rico. 24 pp. + Appendix.
- Caribbean Fishery Management Council (CFMC). 1993. Amendment 2 to the fishery management plan for the shallow-water reef fish fishery of Puerto Rico and the U.S. Virgin Islands. Caribbean Fishery Management Council, San Juan, Puerto Rico. 29 pp. + Appendices.
- Caribbean Fishery Management Council (CFMC). 1998. Essential fish habitat (EFH) generic amendment to the fishery management plans (FMPs) of the U.S. Caribbean including a draft environmental assessment. Caribbean Fishery Management Council, San Juan, Puerto Rico. 169 pp + Appendices.
- Caribbean Fishery Management Council (CFMC). 1999. Amendment number 1 to the fishery management plan for corals and reef associated plants and invertebrates of Puerto Rico and the United States Virgin Islands for establishing a marine conservation district, including regulatory impact review and initial regulatory flexibility analysis and a final supplemental environmental impact statement. Caribbean Fishery Management Council, San Juan, Puerto Rico. 47 pp. + Appendices.
- Caribbean Fishery Management Council (CFMC). 2004. Final environmental impact statement for the generic essential fish habitat amendment to: Spiny lobster fishery management plan (FMP), queen conch FMP, reef fish FMP, and coral FMP for the U.S. Caribbean, Vol. I and II. Caribbean Fishery Management Council, San Juan, Puerto Rico.
- Caribbean Fishery Management Council (CFMC). 2005. Comprehensive amendment to the fishery management plans (FMPs) of the U.S. Caribbean to address required provisions of the Magnuson-Stevens Fishery Conservation and Management Act: Amendment 2 to the FMP for the spiny lobster fishery of Puerto Rico and the U.S. Virgin Islands; Amendment 1 to the FMP for queen conch resources of Puerto Rico and the U.S. Virgin Islands; Amendment 3 to the FMP for the reef fish fishery of Puerto Rico and the U.S. Virgin Islands; Amendment 2 to the FMP for the corals and reef associated invertebrates of Puerto Rico and the U.S. Virgin Islands. Caribbean Fishery Management Council, San Juan, Puerto Rico. 533 pp. + Appendices. Obtained online at: <http://www.caribbeanfmc.com/SCANNED%20FMPS/06%20FINAL%20SFA%20-%20MAY%2003,2005/SFA-FMP.htm>
- Matos, D. 1992. Annual report to the National Marine Fisheries Service Interjurisdictional Fisheries Program. P.R. Dept. Nat. Res. Fish. Res. Lab. 55 pp

3. ASSESSMENT HISTORY AND REVIEW

Caribbean queen triggerfish have not been formally assessed prior to SEDAR 30.

4. REGIONAL MAPS




Figure 4.1. Caribbean management region including Council and EEZ Boundaries.

5. ASSESSMENT SUMMARY

The Summary Report provides a broad but concise view of the salient aspects of the stock assessment. It recapitulates: (a) the information available to and prepared by the Data Workshop; (b) the application of those data, development and execution of one or more assessment models, and identification of the most reliable model configuration as the base run by the Assessment Process (AP); and (c) the findings and advice determined during the Review Workshop.

U.S. CARIBBEAN QUEEN TRIGGERFISH ASSESSMENT SUMMARY

The Summary Report provides a broad but concise view of the salient aspects of the stock assessment. It recapitulates: (a) the information available to and prepared by the Data Workshop; (b) the application of those data, development and execution of one or more assessment models, and

identification of the most reliable model configuration as the base run by the Assessment Process; and (c) the findings and advice determined during the Review Workshop.

Executive Summary

The SEDAR 30 Caribbean Queen triggerfish assessment process consisted of a data scoping process, with one public data scoping meeting held on Puerto Rico, St. Thomas, and St. Croix, and an assessment process which consisted of an in-person workshop held in Miami, FL and a series of webinars. The SEDAR 30 Peer Review Process was conducted via a CIE (Center for Independent Experts) Desk Review in lieu of a Panel Review Workshop. Three reviewers were selected by provided the CIE and provided with the assessment report and background materials. Each reviewer conducted a review of the material and produced an independent review report.

No clear status determination can be made from the assessment as the independent reviewers differed on the appropriateness of the assessment for making such determinations.

Stock Status and Determination Criteria

Data limitations in the US Caribbean preclude the use of advanced quantitative analyses that provide measures of uncertainty. However, the following conclusions can be drawn based on the data-poor methods employed in this assessment, the fundamental principles of population dynamics, and an overall interpretation of the raw data.

Given the available information for this assessment, it is difficult to interpret the sustainability of the estimated current exploitation rates; therefore the overfishing status for queen triggerfish in the U.S. Caribbean is unknown. The overfished status is also unknown.

Stock Identification and Management Unit

- Queen triggerfish occur along the eastern and western gulfs of the Atlantic Ocean (Carpenter 2002). In the western Atlantic, queen triggerfish range from Massachusetts to Brazil (including the Caribbean and Gulf of Mexico). Adults of this species are found on coral reef habitats as well as sandy and grassy areas.
- Queen triggerfish frequently occupy depths fewer than 100m, but they can also occur in depths up to 275m (Aiken 1975; Tortonese 1986). A
- The queen triggerfish management areas within the U.S. Caribbean include the islands of Puerto Rico and the U.S. Virgin Islands (USVI) including St. Thomas, St. John, and St. Croix. The state waters of Puerto Rico extend 9 nm from the shore and the state waters of the USVI extend 3 nm from shore.

Assessment Data

- Species-specific self-reported commercial landings are available from 1983 – 2011 for Puerto Rico

- Commercial landings in St. Croix could only be provided for triggerfish (all species combined) due to a lack of species-species specific reporting by commercial fishers. In St. Croix, landings data were available for the years 1998-2011.
- Commercial landings of queen triggerfish in St. Thomas and St. John were the self-reported logbook records from 2000-2011. Landings could only be provided for triggerfish (all species combined) due to a lack of species-specific reporting by commercial fishers.
- Recreational landings and discard estimates are available for Puerto Rico for the years 2000-2011 via MRFSS/MRIP however data were not used in quantitative or qualitative analyses to determine overfishing status. No recreational information was available for the USVI.
- Commercial length data were available from the Trip Interview Program:
 - Pots and Traps data from St. Croix for years 1983-2011
 - Pots and Traps data from St. Thomas/St. John for years 1983-1988, 1991-1996, 2002-2011
 - Pots and Traps data from Puerto Rico for years 1983-2009, 2011
- The reported ranges for age and growth parameters were:
 - L_{inf} : 415mm-441mm FL
 - K: 0.14-0.57 per year
- The following life history parameter inputs were used in the length-frequency analysis:
 - Lower and upper values for K= 0.165 and 0.435 per year used in sensitivity.
 - Lower and upper values (in mm) used in sensitivity analyses for L_{inf} were 373 and 456.

Release Mortality

No data on release mortality for queen triggerfish in the U.S. Caribbean exists. Release mortality information is not required for the length-based approach attempted in this assessment.

Assessment Methods

- The length frequency analysis for queen triggerfish pot and trap fishery in Puerto Rico focused on time series analyses and relative differences in total mortality estimates rather than on absolute values of total mortality due to considerable uncertainty in age-growth parameters.
- Total mortality (Z) estimates and the ability to detect changes in mortality were explored using a variant of the Beverton-Holt length-based mortality estimator.

Fishing Mortality Trends

Estimates of total mortality can be translated to fishing mortality (F) by subtracting natural mortality (M). Lacking direct estimates of natural mortality, life history invariant relationships were used and given the uncertainty in total mortality estimates this was not pursued.

Stock Abundance and Biomass Trends

Given the data limitations, accurate estimates of stock abundance or biomass could not be developed.

Key Sources of Scientific Uncertainty

- The calculation of traditional benchmarks based on MSY theory using the mean length mortality estimation method were not derived due to considerable uncertainty in the available life-history parameters. Lack of current, species-specific life history information greatly hindered the assessment.
- Small sample size for the length data in recent years complicated the interpretation of the results.

Projections:

Given the data limitations, projections for future status could not be constructed. Furthermore, the AW panel acknowledged that projections were not appropriate for the queen triggerfish stocks due to violations of the selectivity assumption and concerns about life history parameters.

Figures




Figure 1. Puerto Rico yearly commercial fishery landings of queen triggerfish, expansion factors applied, by gear and year. (Figure 1 in the Assessment Workshop Report)




Figure 2. Yearly commercial landings of triggerfishes as reported (no expansion factors applied) on fisher logbooks from St. Croix by gear and year. (Figure 6 from the Assessment Workshop Report)





Figure 3. Yearly commercial landings of triggerfishes as reported (no expansion factors applied) on fisher logbooks from St. Thomas and St. John by gear and year. (Figure 4 from the Assessment Workshop Report)

6. SEDAR ABBREVIATIONS

ABC	Allowable Biological Catch
ACCSP	Atlantic Coastal Cooperative Statistics Program
ADMB	AD Model Builder software program
ALS	Accumulated Landings System; SEFSC fisheries data collection program
ASMFC	Atlantic States Marine Fisheries Commission
B	stock biomass level
BMSY	value of B capable of producing MSY on a continuing basis
CFMC	Caribbean Fishery Management Council
CIE	Center for Independent Experts
CPUE	catch per unit of effort
F	fishing mortality (instantaneous)
F_{MAX}	fishing mortality that maximizes the average weight yield per fish recruited to the fishery
F_{MSY}	fishing mortality to produce MSY under equilibrium conditions
F_{OY}	fishing mortality rate to produce Optimum Yield under equilibrium
$F_{XX\% SPR}$	fishing mortality rate that will result in retaining XX% of the maximum spawning production under equilibrium conditions
F_0	a fishing mortality close to, but slightly less than, F_{max}
FL FWCC	Florida Fish and Wildlife Conservation Commission
FWRI	(State of) Florida Fisheries and Wildlife Research Institute
GA DNR	Georgia Department of Natural Resources
GLM	general linear model
GMFMC	Gulf of Mexico Fishery Management Council
GSMFC	Gulf States Marine Fisheries Commission
GULF FIN	GSMFC Fisheries Information Network
M	natural mortality (instantaneous)
MARMAP	Marine Resources Monitoring, Assessment, and Prediction
MFMT	maximum fishing mortality threshold, a value of F above which overfishing is deemed to be occurring
MRFSS	Marine Recreational Fisheries Statistics Survey; combines a telephone survey of households to estimate number of trips with creel surveys to estimate catch and effort per trip
MRIP	Marine Recreational Information Program
MSST	minimum stock size threshold, a value of B below which the stock is deemed to be overfished

MSY	maximum sustainable yield
NC DMF	North Carolina Division of Marine Fisheries
NMFS	National Marine Fisheries Service
NOAA	National Oceanographic and Atmospheric Administration
OY	optimum yield
SAFMC	South Atlantic Fishery Management Council
SAS	Statistical Analysis Software, SAS Corporation
SC DNR	South Carolina Department of Natural Resources
SEDAR	Southeast Data, Assessment and Review
SEFSC	Fisheries Southeast Fisheries Science Center, National Marine Fisheries Service
SERO	Fisheries Southeast Regional Office, National Marine Fisheries Service
SPR	spawning potential ratio, stock biomass relative to an unfished state of the stock
SSB	Spawning Stock Biomass
SSC	Science and Statistics Committee
TIP	Trip Incident Program; biological data collection program of the SEFSC and Southeast States.
Z	total mortality, the sum of M and F



SEDAR

Southeast Data, Assessment, and Review

SEDAR 30

U.S. Caribbean Queen Triggerfish

SECTION II: Assessment Process Report

January 2013

SEDAR
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405

Table of Contents

1. WORKSHOP PROCEEDINGS 4

1.1. INTRODUCTION..... 4

1.1.1 Workshop time and Place 4

1.1.2 Terms of Reference 4

1.1.3. List of Participants 6

1.1.4. List of Assessment Process Working and Reference Papers 6

1.2. PANEL RECOMMENDATIONS AND COMMENT 7

1.2.1. Term of Reference 1 7

1.2.2. Term of Reference 2 7

1.2.3. Term of Reference 3 8

1.2.4. Term of Reference 4 8

1.2.5. Term of Reference 5 8

1.2.6. Term of Reference 6 8

1.2.7. Term of Reference 7 9

1.2.8. Term of Reference 8 9

1.2.9. Term of Reference 9 9

1.2.10. Term of Reference 10..... 10

2. DATA REVIEW..... 10

2.1. Commercial landings..... 10

2.2. Recreational data 11

2.3. Life history 12

2.4. Length-frequency data..... 12

3. Length-Based Mortality Estimator Methods 14

3.1. Overview 14

3.2. Data sources 14

3.3. Model configuration and equations..... 14

3.4. Estimated parameters 16

3.5. Uncertainty and measures of precision 16

4. Model Results 16

4.1. Puerto Rico..... 16

4.2. St. Thomas and St. John 18

4.3. St. Croix 19

5. Discussion 20

5.1. Puerto Rico 20

5.2. USVI 21

6. Stock status and general conclusions 21

7. References 22

8. Tables 24

9. Figures 47

1. WORKSHOP PROCEEDINGS

1.1. INTRODUCTION

1.1.1 Workshop time and Place

The SEDAR 30 Assessment Workshop was held October 16-18, 2012 in Miami, Florida.

1.1.2 Terms of Reference

1. Review any changes in data following the data scoping and any analyses suggested by the data scoping. Summarize data as used in each assessment model.
2. Develop population assessment models that are compatible with available data and document input data, model assumptions and configuration, and equations for each model considered.
3. Provide estimates of stock population parameters, if feasible
 - When available, include fishing mortality, abundance, biomass, selectivity, stock-recruitment relationship, and other parameters as necessary to describe the population
 - Include appropriate and representative measures of precision for parameter estimates.
4. Characterize uncertainty in the assessment and estimated values.
 - Consider uncertainty in input data, modeling approach, and model configuration.
 - Consider other sources as appropriate for this assessment
 - Provide appropriate measures of model performance, reliability, and ‘goodness of fit’
 - Provide measures of uncertainty for estimated parameters
5. Provide evaluations of yield and productivity
 - Include yield-per-recruit, spawner-per-recruit, and stock-recruitment models
6. Provide estimates of population benchmarks or management criteria consistent with the available data, applicable FMPs, proposed FMPs and Amendments, other ongoing or proposed management programs, and National Standards.
 - Evaluating existing or proposed management criteria as specified in the management summary
 - Recommend proxy values when necessary
7. Provide declarations of stock status relative to benchmarks or alternative data-poor approaches if necessary.
8. Perform a probabilistic analysis of proposed reference points, stock status, and yield.
 - Provide the probability of overfishing at various harvest or exploitation levels
 - Provide a probability density function for biological reference point estimates.
 - If the stock is overfished, provide the probability of rebuilding within mandated time periods as described in the management summary or applicable federal regulations.

9. Project future stock conditions (biomass, abundance, and exploitation) and develop rebuilding schedules if warranted; include estimated generation time. Stock projections shall be developed in accordance with the following:
 - A) If stock is overfished:
F=0, F=current, F=Fmsy, Ftarget (OY),
F=Frebuild (max that rebuild in allowed time)
 - B) If stock is overfishing
F=Fcurrent, F=Fmsy, F= Ftarget (OY)
 - C) If stock is neither overfished nor overfishing
F=Fcurrent, F=Fmsy, F=Ftarget (OY)
 - D) If data-limitations preclude classic projections (i.e. A, B, C above), explore alternate models to provide management advice.
10. Provide recommendations for future research and data collection.
 - Be as specific as practicable in describing sampling design and sampling intensity
 - Emphasize items which will improve future assessment capabilities and reliability
 - Consider data, monitoring, and assessment needs
11. Complete the Assessment Workshop Report for Review (Section III of the SEDAR Stock Assessment Report).

1.1.3. List of Participants

Assessment Workshop Panel

Adyan Rios..... NMFS SEFSC Miami
 Daniel Matos PR DNER
 Francisco PagánCaribbean Coral Reef Institute
 Kevin McCarthy..... NMFS/SEFSC/Miami
 Meaghan Bryan.....NMFS/SEFSC
 Nancie Cummings..... NMFS/SEFSC/Miami
 Richard Appeldoorn..... SSC Representative/University of Puerto Rico

Council Representation

Carlos Farchette CFMC

Appointed Observers

Carlos Velazquez Industry Representative/Puerto Rico
 Daryl Bryan..... Industry Representative/St. Thomas
 Gerson Martinez..... Industry Representative/St. Croix
 Gregory Ledee Industry Representative/St. Thomas

Attendees

Shannon Cass-Calay NMFS/SEFSC/Miami
 Clay Porch..... NMFS/SEFSC/Miami

Staff

Andrea Grabman..... SEDAR
 Bill Arnold SERO
 Graciela García-MolinerCFMC Staff
 Julia Byrd SEDAR
 Julie A. Neer SEDAR
 Michael Larkin..... SERO

1.1.4. List of Assessment Process Working and Reference Papers

Document #	Title	Authors
Documents Prepared for the Assessment Workshop		
SEDAR30-AW-01	Summary of recreational catch and effort for blue tang and queen triggerfish caught in	Meaghan Bryan

	Puerto Rico since 2000	
SEDAR30-AW-02	Evaluation of the available length-frequency information in the US Caribbean Trip Interview Program (TIP) data	Meaghan Bryan
SEDAR30-AW-03	A review of the life history characteristics of blue tang and queen triggerfish	Adyan B. Rios
SEDAR30-AW-04	Commercial fishery landings of queen triggerfish and blue tang in the United States Caribbean, 1983-2011	Kevin J. McCarthy
Reference Documents		
SEDAR30-RD01	A pilot program to assess methods of collecting bycatch, discard, and biological data in the commercial fisheries of St. Thomas, U.S. Caribbean	MRAG Americas
SEDAR30-RD02	A pilot program to assess methods of collecting bycatch, discard, and biological data in the commercial fisheries of U.S. Caribbean (Saint Croix)	MRAG Americas

1.2. PANEL RECOMMENDATIONS AND COMMENT

1.2.1. Term of Reference 1

Review any changes in data following the data scoping and any analyses suggested by the data scoping. Summarize data as used in each assessment model.

Commercial landings were presented for Puerto Rico, St. Thomas/St. John, and St. Croix. The recreational intercept data and the length-frequency data from the Trip Interview Program (TIP) were also reviewed. Basic data inputs for this assessment, such as length-frequency data by island and gear, and life history information from published literature, were reviewed in detail.

1.2.2. Term of Reference 2

Develop population assessment models that are compatible with available data and document input data, model assumptions and configuration, and equations for each model considered.

The AW panel recommended that the length-frequency data from TIP were the most consistent species-specific data available for queen triggerfish. As such, the length based total mortality

estimator (Gedamke and Hoenig 2006) was applied to the available length data. This approach was applied to the data from the pot and trap fisheries in Puerto Rico, St. Thomas/St. John, and St. Croix.

1.2.3. Term of Reference 3

Provide estimates of stock population parameters, if feasible

- *When available, include fishing mortality, abundance, biomass, selectivity, stock-recruitment relationship, and other parameters as necessary to describe the population*
- *Include appropriate and representative measures of precision for parameter estimates.*

Given data limitations, population parameters defining abundance, selectivity, and the stock-recruitment relationship of queen triggerfish in the US Caribbean were not estimated. A mean-length analytical approach was used to estimate total mortality. A sensitivity analysis was conducted to describe the uncertainty in the total mortality estimates. Fishing mortality estimates were derived from the range of total mortality estimates and a range of natural mortality estimates. Natural mortality estimates were derived from several estimators, which were dependent on various life history parameters.

1.2.4. Term of Reference 4

Characterize uncertainty in the assessment and estimated values.

- *Consider uncertainty in input data, modeling approach, and model configuration.*
- *Consider other sources as appropriate for this assessment*
- *Provide appropriate measures of model performance, reliability, and ‘goodness of fit’*
- *Provide measures of uncertainty for estimated parameters*

Since considerable uncertainty exists in the absolute estimates of total mortality from the mean length analysis, a comprehensive sensitivity analysis was conducted.

1.2.5. Term of Reference 5

Provide evaluations of yield and productivity

- *Include yield-per-recruit, spawner-per-recruit, and stock-recruitment models*

Due to limited data and concerns regarding life history parameters, the AW panel did not recommend calculations of yield-per-recruit, spawner-per-recruit, and stock-recruitment estimations. The data limitations are discussed throughout the model results and discussion sections.

1.2.6. Term of Reference 6

Provide estimates of population benchmarks or management criteria consistent with the available data, applicable FMPs, proposed FMPs and Amendments, other ongoing or proposed management programs, and National Standards.

- *Evaluating existing or proposed management criteria as specified in the management summary*
- *Recommend proxy values when necessary*

Derived fishing mortality estimates are discussed with regard to natural mortality, which can be used as a proxy for F_{MSY} . However, due to data limitations, concerns regarding the life history parameters and the resulting uncertainty in the model results, the AW panel concluded that useful population benchmarks could not be developed during this assessment.

1.2.7. Term of Reference 7

Provide declarations of stock status relative to benchmarks or alternative data-poor approaches if necessary.

The AW Panel concluded that this was not applicable to this assessment given the data limitations.

1.2.8. Term of Reference 8

Perform a probabilistic analysis of proposed reference points, stock status, and yield.

- *Provide the probability of overfishing at various harvest or exploitation levels*
- *Provide a probability density function for biological reference point estimates.*
- *If the stock is overfished, provide the probability of rebuilding within mandated time periods as described in the management summary or applicable federal regulations.*

The AW Panel concluded that a probabilistic analysis of proposed reference points was not possible given data limitations.

1.2.9. Term of Reference 9

Project future stock conditions (biomass, abundance, and exploitation) and develop rebuilding schedules if warranted; include estimated generation time. Stock projections shall be developed in accordance with the following:

A) If stock is overfished:

*$F=0$, $F=current$, $F=Fmsy$, $F=target$ (OY),
 $F=Frebuild$ (max that rebuild in allowed time)*

B) If stock is overfishing

$F=Fcurrent$, $F=Fmsy$, $F=Ftarget$ (OY)

C) If stock is neither overfished nor overfishing

$F=Fcurrent$, $F=Fmsy$, $F=Ftarget$ (OY)

D) If data-limitations preclude classic projections (i.e. A, B, C above), explore alternate models to provide management advice.

Due to the limited data available, a data poor methodology was attempted that does not include projections of stock dynamics. Therefore, projections were not conducted for this assessment. Furthermore, the AW panel acknowledged that projections were not appropriate for the queen

triggerfish stocks due to violations of the selectivity assumption and concerns about life history parameters.

1.2.10. Term of Reference 10

Provide recommendations for future research and data collection.

- *Be as specific as practicable in describing sampling design and sampling intensity*
- *Emphasize items which will improve future assessment capabilities and reliability*
- *Consider data, monitoring, and assessment needs*

The ability to utilize length-frequency data is contingent upon having reliable estimates of life history parameters (von Bertalanffy parameters in particular). Studies on basic life history (e.g. age-growth relationships and estimating natural mortality) in the US Caribbean will greatly enhance the utility of the existing length-frequency data and should provide the greatest benefit to providing management advice in the short term. This should be placed as a top priority for key species.

Fishery-independent surveys should be considered as a top research priority for additional data collection. Fishery-independent surveys designed using a rigorous statistical framework will allow for the collection of species-specific catch and effort data that can be used to develop indices of abundance. Indices of abundance are used in stock assessments to inform models about how a population may be changing over time. Fishery-independent surveys can also be used to supplement existing programs by collecting age, length, weight, and reproductive data.

It is essential that continued efforts to improve the data collection of fishery-dependent catch and effort statistics be made. More specifically, continued efforts to collect species-specific catch statistics will be important for future assessments.

During the assessment workshop, the fishers from the USVI indicated that the selectivity pattern for queen triggerfish violated the assumption of knife-edge selectivity in the mean-length model. Efforts should be made to expand this model to accommodate other selectivity patterns.

2. DATA REVIEW

2.1. Commercial landings

A detailed description of the methods and results of commercial landings calculations can be found in working paper SEDAR30-AW-04.

Puerto Rico

Commercial fishery landings data for Puerto Rico were available from self-reported fisher logbooks/sales receipts for the years 1983-2011. Data were reported by species, fishing gear, and fishing center where the catch was landed. Puerto Rico commercial landings have been incompletely reported (Caribbean Fisheries Data Evaluation Final Report, 2009) and required expansion factors to estimate total landings. For the years 2003 to 2011, expansion

factors have been coast-specific (north, south, east, west). Estimation of commercial fishery landings of earlier years used a single, island-wide, expansion factor.

Puerto Rico landings were estimated as: year-specific reported landings * year-specific expansion factor. Estimation of landings for the most recent years (2003-2011) included year and coast-specific expansion factors. Reported landings were assigned to coasts based upon the fishing center reported for a trip. Queen triggerfish landings have been reported by species in Puerto Rico throughout the years of available data. Yearly total expanded queen triggerfish landings for Puerto Rico are provided in Table 1. The expanded landings of queen triggerfish by gear and year are shown in Figure 1. The numbers of trips with reported queen triggerfish landings are plotted in Figure 2 (by coast and year) and Figure 3 (by gear and year).

St. Thomas and St. John

In the US Virgin Islands logbook landings data from the islands of St. Thomas and St. John were compiled separately from St. Croix. Logbook reporting began in 1974; however, landings were initially reported by gear type (e.g., net fish, hook fish, pot fish, and spear fish) and as either snapper/grouper or as other fin fish during the period 1974-1995. Beginning in 1997 in St. Thomas/St. John, some landings data were reported by species group; (e.g., snappers, groupers, parrotfishes, surgeonfishes, etc.) and by gear (hook and line, gill net, SCUBA, trap, etc.). All commercial fishery data reports included species group beginning in 2000. Species-specific data were reported in the US Virgin Islands during the 2011-2012 fishing year.

The data available for summing commercial landings of queen triggerfish in St. Thomas and St. John were the self-reported logbook records from commercial fishers. Landings could only be provided for triggerfish (all species combined) due to a lack of species-specific reporting by commercial fishers. Annual landings data, as reported, were summed by species group and fishing gear and are provided in Table 2 and Figure 4. The numbers of commercial fishing trips with landings of triggerfish in St. Thomas and St. John are shown in Figure 5 by gear and year.

St. Croix

See the St. Thomas and St. John section above for a brief description of the available landings data. Landings for St. Croix could only be provided for triggerfish (all species combined) due to a lack of species-specific reporting by commercial fishers. In St. Croix, landings data were available for the years 1998-2011. Yearly landings data, as reported, were summed by species group and fishing gear and are provided in Table 3 and Figure 6. The numbers of commercial fishing trips with landings of triggerfish in St. Croix are shown in Figure 7 by gear and year.

2.2. Recreational data

Recreational data from the Marine Recreational Fisheries Statistics Survey (MRFSS) were evaluated to determine 1) whether the intercept data could be used to develop a relative index of abundance for queen triggerfish and 2) whether the length data was sufficient to use for length-based analyses. The reader can find a detailed summary of the available recreational fishery data collected by the MRFSS in the SEDAR working paper, SEDAR30-AW-01, but a brief summary follows.

In the US Caribbean, MRFSS only collects data in Puerto Rico. Tables 4-6 summarize the available intercept data for queen triggerfish. Table 4 provides a summary of the number of AB1 catch, fish observed or reported as dead, and B2 catch, fish reported as released alive, and the proportion of intercepted trips that caught queen triggerfish relative to all intercepted recreational fishing trips (i.e., proportion positive). The proportion of positive trips was less than one percent in all years. Upon the request of the AW panel, the intercept data was summarized by fishing area and is shown in Table 5. The majority of intercepted trips catching queen triggerfish were in area 3, which is classified as ocean less than 10 miles. Area 3 can be considered an inshore area. Table 6 summarizes the MRFSS intercept data for areas 3 and 5; area 5 was also included because it is defined as an inshore fishing area. The number of total intercepted trips was reduced, but the annual proportion positives were still less than one percent, except in 2009 when the proportion of the positive recreational trips was three percent (Table 6).

The MRFSS length data for queen triggerfish were also evaluated. Sixty queen triggerfish were measured between 2000 and 2011. These length frequency data can be found in Figure 8. The AW panel determined that at this time the intercepted catch, effort, and length data should not be used for the assessment of queen triggerfish given the low proportion positives and the infrequent collection of length data.

2.3. Life history

A detailed summary of queen triggerfish life history can be found in working paper SEDAR30-AW-03. The available literature describing the age-length relationship for queen triggerfish was limited and is summarized in Table 7. Two studies were available and reported similar asymptotic length estimates, 415mm FL and 441mm FL, but differed in their estimates of the growth coefficient, 0.3 and 0.14 (Manooch and Drennon 1987, de Albuquerque et al. 2011). Figure 9 shows the resulting von Bertalanffy growth curves from these studies. Neither growth curve reaches the asymptotic length by the maximum age reported in their respective study (Figure 9). The AW panel expressed concern about these age-length relationships since they estimate disparate growth coefficients. The growth parameters reported by Manooch and Drennan (1987) were calculated from queen triggerfish collected in Puerto Rico and the US Virgin Islands. These parameters were used to develop the range of growth parameters that were explored in the length-based mortality analysis.

2.4. Length-frequency data

Working paper SEDAR30-AW-02 provides a detailed presentation of the available length-frequency data from the NMFS Trip Interview Program (TIP). A summary of the number of annual queen triggerfish length measurements can be found in Appendix B in SEDAR30-AW-02. Table 8 provides a general summary of the TIP data for queen triggerfish including; the total number of queen triggerfish lengths measured, the number of years in which queen triggerfish were measured, and the average number of measured lengths per year. The majority of queen triggerfish lengths in Puerto Rico, St. Thomas and St. John, and St. Croix came from their respective pot and trap fisheries. The AW panel agreed that the data available from the US Caribbean pot and trap fisheries had sufficient annual sample sizes for length-based analyses.

Puerto Rico

The annual length-frequency data for queen triggerfish caught by the pot and trap fishery in Puerto Rico are shown in Figure 10. Measures of central tendency were calculated and are summarized in Table 9. The annual modes and the length-frequency distributions shifted towards larger individuals starting in the late 1990s (Table 9, Figure 10). Prior to 1998 the mode was approximately 20cm and after 1998 the modes were approximately 30cm (Table 9, Figure 10).

Large queen triggerfish were measured throughout the time-series, but in the late 1990s queen triggerfish with lengths greater than 45cm were measured more frequently than earlier in the time series (Figure 10). The AW panel was interested in determining whether the larger queen triggerfish were found on both the east and west coasts of Puerto Rico or whether they were isolated to a particular area, namely the west or east coasts of Puerto Rico. Figures 11 and 12 show the annual length-frequency plots for queen triggerfish measured on the west and east coast, respectively. The coasts were defined according to fishing center and followed the definitions in SEDAR26-DW-05 (see map on page 25 of SEDAR26-DW-05). Although large queen triggerfish were measured on the west coast throughout the time series, in the late 1990s and early 2000s, a greater number of large queen triggerfish were measured on the east coast of Puerto Rico as compared to other years (Figure 12). At the AW, the industry representative from Puerto Rico indicated that fishers from Puerto Rico have always kept queen triggerfish of any size; however, in recent years the larger queen triggerfish have been in higher demand for culinary purposes. This suggests a potential reason for the appearance of larger queen triggerfish later in the time-series. Another potential explanation for the appearance of larger individuals could be spatial expansion to areas that were previously relatively unfished. The AW panel discussed this possible explanation with respect to localized depletion overtime; however, the spatial expansion of the queen triggerfish fishery in Puerto Rico was not verified.

St. Thomas and St. John

Figure 13 shows the annual length-frequency data for queen triggerfish caught by the St. Thomas/St. John pot and trap fishery. The annual length range was generally between 20cm and 45cm. In 1986, there were many queen triggerfish length measurements that were less than 20cm (Figure 13). During the AW, the industry representatives from St. Thomas indicated that they regularly release queen triggerfish larger than 45cm or 50cm unless a client specifically requests larger queen triggerfish. This provides an explanation for the infrequent measurement of queen triggerfish larger than 45cm and for the repeatedly observed right truncation of the length-frequency data. Over time the mode of the length-frequency data varied and ranged between 20cm and 37cm (Table 10). Prior to 1996, the mode was generally less than 30cm and after 1996, the mode was generally larger than 30cm (Table 10).

St. Croix

The length-frequency data for queen triggerfish caught by the St. Croix pot and trap fishery are shown in Figure 14. The number of queen triggerfish lengths measured annually in St. Croix was generally lower than in Puerto Rico or St. Thomas/St. John. The length range was between 20cm and ~40cm in most years (Figure 14). Larger queen triggerfish, with lengths greater than 40cm, were seen periodically throughout the time-series, but were not a prominent feature except in 1989 and 1990 (Figure 14). Similar to the St. Thomas industry representatives,

the St. Croix industry representative at the AW indicated that there is very low demand for large queen triggerfish in St. Croix. The representatives also explained that St. Croix fishers release queen triggerfish larger than ~40-45cm, unless a client specifically requests large queen triggerfish. Table 11 summarizes the measures of central tendency for the annual length-frequency data. The mode varied over time and ranged between 21cm and 31cm (Table 11).

3. Length-Based Mortality Estimator Methods

3.1. Overview

A review of the length frequency data available from the NMFS Trip Interview Program (TIP) database indicated that sample sizes were sufficient to conduct a time-series length analysis for a limited number of island and gear combinations (Table 8). The length data were subset by gear due to concerns about differential selectivity among the gear types. Total mortality (Z) estimates and the ability to detect changes in mortality were explored using a variant of the Beverton-Holt length-based mortality estimator (Beverton and Holt 1956, 1957).

3.2. Data sources

The AW panel reviewed the only available source of length composition data, the TIP database, and recommended that these data be used for analysis. Input values for other parameters populating the model were gathered from the available literature. Preliminary analyses were performed using the values summarized in Table 12. The SEDAR 30 AW panel noted very limited growth information available for queen triggerfish. This will be discussed with regard to the TIP data and the length analysis in later sections of this report.

3.3. Model configuration and equations

The Beverton-Holt mortality estimator has received widespread use, especially in data-limited situations, owing mainly to the minimal parameter inputs, namely the von Bertalanffy growth parameters K and L_∞ , the length of first capture (smallest size at which animals are fully vulnerable to the fishery and to the sampling gear), L_c , and the mean length of the animals (\bar{L}) above the length L_c :

$$Z = \frac{K(L_\infty - \bar{L})}{\bar{L} - L_c}$$

There are six assumptions behind this method:

1. Asymptotic growth with known parameters K and L_∞ , which are constant over time.
2. No individual variability in growth.
3. Constant and continuous recruitment over time.
4. Mortality rate is constant with age for all ages $t > t_c$, where t_c is the age at first capture.
5. Mortality rate is constant over time.

6. Population is in equilibrium (i.e., enough time has passed following any change in mortality that mean length now reflects the new mortality level).

A criticism of this method is that the assumption of equilibrium (6) is very difficult to meet in the real world situations where any change in fishing pressure disrupts the equilibrium age distribution. For example, with increased fishing pressure, there is a delay before larger/older animals are removed from the population and before the mean length decreases in accordance with the current mortality rate. Likewise, when fishing pressure is decreased, equilibrium can only be reestablished once the smaller/younger animals have grown, and the mean length increases in accordance with the reduced mortality rate.

Gedamke and Hoenig (2006) developed an extension of the Beverton-Holt length-based mortality estimator for use in non-equilibrium situations. This method is quantitatively attractive because it requires minimal and commonly available data, and it does not require the assumption that catch rate is proportional to abundance. The method also allows for a broader application of the mean length approach as it does not require an assumption of equilibrium, which is not often met in real world situations. In addition, the transitional form of the model allows mortality estimates to be made within a few years of a change, rather than having to wait for the mean lengths to stabilize at their new equilibrium level. In other words, as soon as a decline in mean lengths is detected, this model can be applied and the trajectory of decline can be used to estimate the new Z and how mean lengths will change over time.

The method is described in detail in Gedamke and Hoenig (2006) and is summarized below. Like the Beverton and Holt estimator, this extension only requires a series of mean length above a user defined minimum size and von Bertalanffy growth parameters. Therefore, it can be applied in many data poor situations. Gedamke and Hoenig (2006) demonstrated the utility of this approach using both simulated data and an application to data for goosefish caught in the NEFSC fall groundfish survey.

The mean length in a population can be calculated d years after a single permanent change in total mortality from Z_1 to Z_2 yr⁻¹ by the following equation:

$$\bar{L} = L_{\infty} - \frac{Z_1 Z_2 (L_{\infty} - L_c) \{Z_1 + K + (Z_2 - Z_1) \exp(-(Z_2 + K)d)\}}{(Z_1 + K)(Z_2 + K)(Z_1 + (Z_2 - Z_1) \exp(-Z_2 d))}$$

This equation has been generalized to allow for multiple changes in mortality rate over time (e.g. one change, two changes, three changes etc.). The algorithm was programmed in AD Model Builder in a maximum likelihood framework and used to estimate mortality rates from the observed mean lengths. A shell program was written in R to conduct a grid search of the potential year(s) of change and also to conduct a sensitivity analysis to input parameters.

Models were run starting with the simplest (i.e. no change in mortality) and then with increasing complexity by sequentially adding additional years of change (Note: each year of change adds two parameters). The Akaike Information Criterion with a correction for small sample size (AIC_c) was calculated for each scenario and will be referred to simply as AIC throughout the remainder of this document. To compare models, the change in AIC (Δ AIC, the difference between AIC and the minimum AIC) was also calculated. When comparing models, a

Δ AIC value less than 2 indicated strong support for the model and a Δ AIC value between 3 and 5 indicated moderate support for the model (Burnham and Anderson, 2002).

The first step in the application of this mean-length approach is to determine the length at which animals become fully vulnerable to the gear, L_c . Annual length-frequency plots were constructed for island-gear combinations for which sufficient sample sizes were available. L_c was selected visually (Thorson and Prager, 2011) and estimates of central tendency were also calculated to help with the selection. The highest L_c value over the time series was chosen as the input for preliminary runs and the central value from which to develop a sensitivity range. Using the highest L_c value avoids the confounding between selectivity and mortality in the calculation of annual mean lengths. The three values of L_c that were identified for the pot and trap fishery in Puerto Rico were 28cm, 29cm, and 31cm (Table 12). For the initial, single run analysis an L_c value equal to 29cm was used. The three values of L_c that were identified for the St. Thomas/St. John pot and trap fishery were 32cm, 34cm, and 35cm for use in the preliminary analyses (Table 12). For the initial, single run analysis an L_c value equal to 34cm was used. The three values of L_c that were identified for the St. Croix pot and trap fishery for use in the preliminary analyses were 27cm, 28cm, and 29cm (Table 12). For the initial, single run analysis an L_c value equal to 28cm was used. For each fishery, the value identified as the lower possible value of L_c was chosen to avoid confounding between changes in selectivity and total mortality. These L_c values were also used to develop of input values for the sensitivity analysis.

3.4. Estimated parameters

The parameters estimated by the non-equilibrium length method, as described above, are the total mortality rates (Z) and the year(s) of change. The total mortality in the most recent time periods is referred to Z_{current} for the remainder of the document.

3.5. Uncertainty and measures of precision

When the assessment workshop (AW) began, preliminary analyses had been done using the von Bertalanffy growth parameters from a single study, Manooch and Drennan (1987). The von Bertalanffy growth coefficient from this study was equal to 0.3 and the asymptotic length was equal to 41.5cm FL. Sensitivity analyses had also been conducted due to the limited published information on the age-length relationship for queen triggerfish. The sensitivity range for the von Bertalanffy growth coefficient was $41.5 \pm 45\%$ in increments of 1.5cm. The sensitivity range for the asymptotic length was $0.03 \pm 10\%$ in increments of 0.1. Table 12 summarizes the sensitivity range used for queen triggerfish for the different island and gear combinations.

4. Model Results

4.1. Puerto Rico

Initial analyses

The AW panel agreed that annual sample sizes were sufficient to conduct the length-based analysis for the pot and trap fishery in Puerto Rico (Table 8).

The AIC results for the initial analysis are summarized in Table 13a. The model with the lowest AIC value predicted one change in total mortality in 1990. Total mortality was predicted

to decline after 1990 from 0.6 to 0.05, a 92% decline in total mortality. Mean length was fairly stable and approximately equal to 33cm between 1983 and 1990 (Figure 15). After 1990, mean length was variable, but in the late 1990s and early 2000s mean length had an increasing trend corresponding to the reduced total mortality estimate. The model predicting two changes in total mortality was also strongly supported by AIC criteria (i.e., $\Delta AIC < 2$). This model also predicted a decline in total mortality after 1990, followed by an increase after 2000 (Table 13, Figure 16). The predicted decline in total mortality corresponded to larger average lengths between 1998 and 1999 and the predicted increase in total mortality corresponded to smaller mean lengths in and after 2000 (Figure 16). Although this model fit the data well, the model predicting a single change in total mortality also fits the mean length data well and was also more parsimonious.

Sensitivity analyses

Since the life history information available for queen triggerfish was limited, a sensitivity analysis was conducted to quantify uncertainty in the total mortality estimates and to evaluate how model choice was influenced by the input parameters.

Figure 17 illustrates the sensitivity in the absolute estimates of current total mortality given the range of input parameters. The total mortality estimates ranged between 0.02 and 1. Assuming that the von Bertalanffy growth parameters are negatively correlated, some parameter combinations were not biologically plausible and the uncertainty in total mortality may have been overestimated. In general, larger values of the von Bertalanffy growth coefficient and larger values of the asymptotic length led to larger values of total mortality. The only exceptions were for those parameter combinations where the length-at-full vulnerability was less than, but within, 2cm of the asymptotic length. In these cases, total mortality was estimated to be close to zero and reached the imposed lower bound of the parameter range. In this situation the resulting mean length was similar to the asymptotic length, thus informing the model that mean size remained close the theoretical maximum and experienced very little mortality.

The majority of sensitivity runs resulted in strong support, based on AIC criteria, for models predicting a single change or two changes in total mortality (Table 14). A greater number of sensitivity runs supported a single change in total mortality than two changes (Table 14). Irrespective of the number of changes predicted, the predicted first year of change was sometime in the late 1980s or early 1990s (Table 15, Table 16). For the models predicting two changes in total mortality, the predicted second year of change was 1999 (Table 15, Table 16). Total mortality was predicted to decline by 75-100% sometime between 1988 and 1991 (Table 17, Table 18). Mean length in the late 1990s, was similar to the asymptotic length input value, informing the model that the queen triggerfish population had experienced very little mortality. This explains the large percent decline in total mortality. For the sensitivity runs strongly supporting two changes in mortality, the percent increase was high (Table 18). The high percent increase predicted after the second change is due to an increase in total mortality from ~ 0.001 to anywhere between 0.3 – 0.6.

Fishing mortality was derived from the minimum and maximum total mortality estimates from the sensitivity analysis and was compared to estimates of natural mortality. A rule of thumb that is sometimes adopted in data poor situations is that the fishing mortality to achieve

maximum sustainable yield (F_{MSY}) is approximately equal to natural mortality. Fishing mortality estimates greater than natural mortality can indicate that a fishery is experiencing overfishing while fishing mortality estimates less than natural mortality can indicate that a fishery is not experiencing overfishing. Our ability to derive appropriate estimates of fishing mortality is also dependent on having a reliable estimate of natural mortality.

Natural mortality estimates derived from several published equations are presented in Table 19. The estimates ranged between 0.21 and 0.74. The estimates of natural mortality were higher when derived from the higher von Bertalanffy growth coefficient input, 0.3 y^{-1} , or the lower maximum age input, 7 years, than the input values from de Albuquerque et al. (2011). The fishing mortality estimates derived from the smallest value of total mortality were less than the natural mortality estimates, irrespective of the natural mortality estimator (Table 19). The relationship between the fishing mortality derived from the larger estimate of total mortality and natural mortality was mainly dependent on the input parameters. Fishing mortality was less than natural mortality for all estimators, except the Hoenig estimators, the Ralston estimator, and the Jensen estimators, when using a maximum age input equal to 14 or the a von Bertalanffy growth coefficient equal to 0.14 (Table 19). The AW panel suggested that the Pauly equation (Pauly 1980) be used as the natural mortality estimator since the von Bertalanffy growth coefficient and the asymptotic length are both included in the equation and this equation may be robust to the uncertainty in these input parameters. The corresponding fishing mortality estimates were less than natural mortality, and in some cases negative indicating that the queen triggerfish experienced minimal fishing mortality.

4.2. St. Thomas and St. John

Initial analyses

The AIC results from the initial analyses are summarized in Table 13b. The model that predicted no change in total mortality had the lowest AIC value and strongest support, based on AIC criteria, as compared to the other models (Table 13b). Figure 18 shows the model fit to the mean length data. The annual sample sizes were variable, but the mean length was fairly stable overtime. During the AW the St. Thomas industry representatives in attendance indicated that they practice a selection process for queen triggerfish, where they release queen triggerfish that are above ~45cm. This provides an explanation for the stability in mean length over time.

Sensitivity analyses

The sensitivity analyses also indicated that the model that predicted no change in total mortality had the strongest support based on AIC criteria (Table 20). All sensitivity runs supported the model that predicted a single, constant total mortality estimate. The individual absolute total mortality estimates were variable and ranged between 0.1 and 1.25, and were sensitive to the input parameters (Figure 19). In general, higher values of the von Bertalanffy growth coefficient and higher values of the asymptotic length led to higher values of total mortality. Assuming that the von Bertalanffy growth parameters are negatively correlated, some of the parameter combinations explored can be considered biologically implausible and therefore, the uncertainty in total mortality may be overestimated.

Estimates of fishing mortality were derived from the minimum and maximum total mortality estimates obtained in the sensitivity analysis and from estimates of natural mortality. The fishing mortality estimates were then compared to estimates of natural mortality, which is used as a proxy for F_{MSY} in data poor situations. Fishing mortality estimates greater than natural mortality can indicate that a fishery is experiencing overfishing while fishing mortality estimates less than natural mortality can indicate that a fishery is not experiencing overfishing. Natural mortalities were derived from several published estimators. The natural mortality estimates varied among the natural mortality equations and ranged between 0.21 and 0.74 (Table 21).

The fishing mortality estimates derived from the smallest estimate of total mortality from the sensitivity analysis were less than natural mortality, irrespective of the natural mortality estimator (Table 21). All estimates were also negative indicating fishing mortality would have been negligible. Fishing mortality derived from the larger estimate of total mortality was greater than natural mortality when maximum age was equal to 14 or the von Bertalanffy growth coefficient was equal to 0.14, regardless of the natural mortality estimator (Table 21). Fishing mortality derived from the larger total mortality estimates was also greater than natural mortality when the maximum age was set to 7 in the Hoenig (1983) estimator or the Hewitt and Hoenig (2005) estimator. This was also the case when the von Bertalanffy growth coefficient was set at 0.3 in the Jensen (1996) estimator (Table 21).

The AW panel suggested that the Pauly equation (Pauly 1980) be used as the natural mortality estimator since the von Bertalanffy growth coefficient and the asymptotic length are both included in the equation and it may be robust to the uncertainty in these input parameters. Fishing mortality derived from the smaller estimate of total mortality was less than natural mortality. This was also the case when fishing mortality was derived from the larger total mortality estimate along with natural mortality estimates derived from all estimators, except for the Hoenig (1983), Ralston (1987), and Hewitt and Hoenig (2005) estimators when using a von Bertalanffy growth coefficient equal to 0.3 or a maximum age input value equal to 7 years (Table 21).

4.3. St. Croix

Initial analyses

The AIC results from the initial analysis suggest that the model predicting a single change in total mortality had the lowest AIC value and strongest support, based on AIC criteria, (Table 13c). The predicted year of change based on the grid search was 1985 and the total mortality was predicted to increase from 0.7 to 1.07 after 1985. The predicted increase in total mortality corresponds to an approximately 2cm decline in mean length in the early 1980s (Figure 20). This perceived change in size also corresponds to a reduction in annual sample size after 1987 (Figure 20).

Sensitivity analyses

The majority of sensitivity runs provided strong support for the model predicting a single change in total mortality and the predicted year of change was either 1985 or 1986 (Table 22, Table 23). In all cases, total mortality was predicted to increase after 1985 or 1986. The current total mortality estimates ranged between 0.4 and 2 and were consistent among the input values

for L_c (Figure 21). Although the absolute estimates of total mortality were uncertain, the percent increase in total mortality ranged between 50% and 90% for almost all sensitivity parameter combinations (Table 24). The estimates of percent change were larger for the lowest L_c input value and may be due to confounding between selectivity and mortality.

The natural mortality and fishing mortality estimates were derived as previously discussed for Puerto Rico and St. Thomas/St. John (Table 25). The fishing mortality estimates derived from the lowest estimate of total mortality from the sensitivity analysis were all less than natural mortality, irrespective of the method used. In contrast, the fishing mortality estimates derived from the highest estimate of total mortality from the sensitivity analysis were all greater than the estimates of natural mortality (Table 25).

5. Discussion

Although it is possible to identify overfishing if natural mortality can be considered a proxy for F_{MSY} , it is not possible to determine stock status relative to biomass based metrics using length-based analytical methods alone. Given the present data limitations, the AW panel concluded that the length-based approach should be applied to the queen triggerfish length data from the TIP database. The panel decided on this approach in order to ascertain whether mortality has changed over time, to identify the direction of change and the relative magnitude of change, and to derive fishing mortality estimates from total mortality and estimates of natural mortality. The implications of these analyses are discussed below.

5.1. Puerto Rico

The sensitivity results from the analysis of the queen triggerfish length data from the pot and trap fishery in Puerto Rico suggest that total mortality declined in the late 1990s. This corresponds to an increase in mean length in 1998 and 1999. During the AW, the industry representative from Puerto Rico indicated that fishers have always kept queen triggerfish of all sizes. In the late 1990s, however, queen triggerfish with lengths ~60cm were measured more frequently than in previous years. Closer inspection of the annual length-frequency data for the west coast and east coasts of Puerto Rico, suggested that these larger animals were being caught mainly on the east coast of Puerto Rico. A shift in mean length towards larger individuals could indicate a reduction in mortality, a change in selectivity by fishers, or spatial expansion into new fishing areas that had been relatively untouched prior to the late 1990s. Given the information provided by the industry representative, it was thought that a change in selectivity did not explain the increase in mean size. Spatial expansion into new fishing areas can mask a change in mortality and the erosion of the age or size structure of a population, however, there is no evidence supporting this conclusion. Without a time-series of detailed spatial catch data, it is difficult to conclude if the increase in mean length was due to a reduction in total mortality or due to spatial expansion into new fishing areas. Improvements to the collection of commercial catch and effort data are on-going in the US Caribbean and are strongly recommended to continue. To that end, it is recommended that species-specific, spatial catch and effort data be collected to study the spatial evolution of the fishery as it relates to local or more spatially expansive population changes.

The uncertainty in the relationship between fishing mortality and natural mortality suggests that current fishing mortality rates may or may not be sustainable even though the

sensitivity analysis indicated that there has been a reduction in total mortality. The resulting mortality estimates were fully dependent on the age and growth parameters used.

5.2. USVI

During the AW, fishers from St. Thomas and St. Croix indicated that they often capture queen triggerfish larger than the maximum size seen in the TIP data. Generally, queen triggerfish larger than ~45cm are released as they do not have market value. This selectivity pattern violates the model assumption that selectivity is knife-edge (i.e., fish above the length-at-full vulnerability are equally vulnerable to fishing). The violation of this assumption may have led to the overestimation of mortality. It should be noted that release mortality is unknown and concerns about release mortality were discussed at the AW. Opinions expressed by industry representatives and AW panelists suggest that queen triggerfish are a hardy species and release mortality was thought to be minimal. If release mortality is truly minimal, the release of larger queen triggerfish may simultaneously act as a conservation measure. It is currently impossible to evaluate the extent of this fishing behavior on the queen triggerfish population in the USVI since the proportion of released to retained queen triggerfish by size group is unknown. In the short term, the collection of discard data will remain unavailable. Efforts should be made to evaluate the feasibility of modifying the Gedamke-Hoenig model to accommodate other selectivity assumptions and implement them when needed. Efforts should also be made to collect discard data from the fisheries in the USVI to obtain a more accurate measure of fishing mortality.

6. Stock status and general conclusions

Determination of stock status with regard to biomass based metrics using length-based analytic approaches alone is not possible. The length-based approach applied to the queen triggerfish length data for this analysis was used to ascertain whether mortality has changed over time and to derive fishing mortality estimates from total mortality. The AW panel decided that given the reasoning provided in the above sections and the limited life history information available it is difficult to interpret the sustainability of the estimated, current exploitation rates and that the absolute estimates of mortality should be interpreted with caution.

Given the limited life history information, a wide range of growth parameter input values were evaluated and led to a wide range of total mortality estimates for the pot and trap fisheries in Puerto Rico and the USVI. The available life history information indicated that queen triggerfish were either shorter lived and faster growing or longer lived and slower growing (Manooch and Drennon 1987, de Albuquerque et al. 2005). The von Bertalanffy growth parameters are important inputs that influence the estimate of total mortality, as was shown in the sensitivity analyses. The growth coefficient was also a key input parameter used to calculate natural mortality, which was used to derive fishing mortality and evaluate the status of the fishery. The disparate estimates of the growth parameters led to considerable uncertainty in the mortality estimates. They also made it difficult to meaningfully interpret stock status in terms of fishing mortality, in the absence of a weighting system giving credence to one life-history strategy over another.

The AW panel evaluated the protocols used, as well as the sample sizes and the size range of sampled queen triggerfish from the available published studies. Manooch and Drennon (1987) and de Albuquerque et al. (2001) aged queen triggerfish using dorsal spines, which is

considered to be an inferior method for ageing as compared to using otoliths. In addition, Manooch and Drennon (1987) indicated that their estimate of maximum age may be low due to the size range of the fish that were aged. The size range of the aged queen triggerfish was thought to reflect gear selectivity of the commercial hook and line and trap fisheries in Puerto Rico and the USVI and not reflect the full size range of the queen triggerfish in the US Caribbean (Manooch and Drennon 1987). The sampled size range of queen triggerfish was 111mm – 425mm FL by Manooch and Drennon (1987) and 180mm FL - 460mm FL by de Albuquerque et al. (2011). The smaller size range provides reasoning for the lower estimate of the von Bertalanffy growth coefficient by de Albuquerque et al. The disparate growth coefficient and maximum age estimates between the two studies highlight the need for well-designed age and growth studies for queen triggerfish to ensure a representative sample of individuals by age/size, region, season etc.

The lack of detailed spatial information and the problems with the underlying size selectivity also make it difficult to interpret the mortality estimates. As such, it is strongly recommended that the implementation of fishery-independent surveys should be considered as a top research priority. Fishery-independent surveys designed using a rigorous statistical framework can be used to supplement existing programs by collecting age, length, weight, and reproductive data that are representative of the entire population.

7. References

- Alverson, D.L., and M.J. Carney. 1975. A graphic review of the growth and decay of population cohorts. *Journal du Conseil international pour l'Exploration de la Mer* 36(2):133–143.
- Beverton, R.J.H. and S.J. Holt. 1956. A review of methods for estimating mortality rates in fish populations, with special reference to sources of bias in catch sampling. *Rapports et Proces-verbaux des Reunions, Conseil International Pour L'Exploration de la Mer* 140:67-83.
- Beverton, R.J.H. and S.J. Holt. 1957. On the dynamics of exploited fish populations. U.K. Ministry of Agriculture, Fisheries, Food, and Fishery Investigations Series II, Vol. XIX.
- Burnham, K. P., and Anderson, D.R. 2002. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*, 2nd ed. Springer-Verlag.
- Cummings, N. 2011. Updated catch per unit abundance indices for silk and queen snapper from the commercial fisheries in Puerto Rico. SEDAR26-DW-05. 37pp.
- de Albuquerque, C.Q., A.S. Martins, L. Junior, N. de Oliveira, J.N.D. Araújo, and A.M. Ribeiro. 2011. Age and growth of the queen triggerfish *Balistes vetula* (tetraodontiformes, balistidae) of the central coast of Brazil. *Brazilian Journal of Oceanography* 59(3): 231-239.
- Gedamke, T. and J.M. Hoenig. 2006. Estimating mortality from mean length data in nonequilibrium situations, with application to the assessment of goosfish. *Transactions of the American Fisheries Society* 135: 476-487.

- Hewitt, D.A., and J.M. Hoenig. 2005. Comparison of two approaches for estimating natural mortality based on longevity. *Fishery Bulletin* 81(4):898–903.
- Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin* 81(4):898–903.
- Jensen, A.L. 1996. Beverton and Holt life history invariants result from optimal trade-off of reproduction and survival. *Canadian Journal of Fisheries and Aquatic Sciences* 53(4):820–822.
- Manooch, C.S and C.L. Drennon. 1987. Age and growth of yellowtail snapper and queen triggerfish collected from the U S. Virgin Islands and Puerto Rico. *Fisheries Research* 6: 53-68.
- Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal du Conseil international pour l'Exploration de la Mer* 39(2):175–192.
- Ralston, S. 1987. Mortality rates of snappers and groupers. Pages 375–404 *In: J.J. Polovina and S. Ralston (eds.), Tropical Snappers and Groupers: Biology and Fisheries Management.* Westview Press, Boulder Colorado. 659 p.
- Thorson, J. T., and M. H. Prager. 2011. Better catch curves: Incorporating age-specific natural mortality and logistic selectivity. *Transactions of the American Fisheries Society* 140: 356–366.

8. Tables

Table 1. Puerto Rico expanded commercial landings of queen triggerfish, 1983-2011.

Year	Queen Triggerfish
1983	147,333
1984	123,607
1985	82,811
1986	41,387
1987	51,140
1988	49,322
1989	64,896
1990	56,042
1991	60,835
1992	46,221
1993	63,605
1994	73,006
1995	97,152
1996	89,521
1997	93,872
1998	82,659
1999	63,607
2000	72,711
2001	87,910
2002	62,187
2003	69,611
2004	97,721
2005	122,423
2006	44,237
2007	33,409
2008	56,715
2009	47,782
2010	48,455
2011	51,032
Total	2,081,209

Table 2. St. Thomas and St. John reported commercial landings of triggerfish (no expansion factors applied), 2000-2011.

Year	Triggerfishes
2000	72,090
2001	82,688
2002	97,543
2003	101,523
2004	87,420
2005	76,462
2006	70,120
2007	72,642
2008	84,131
2009	79,469
2010	79,555
2011	30,555
Total	934,196

Table 3. St. Croix reported commercial landings of triggerfish (no expansion factors applied), 1998-2011.

Year	Triggerfishes
1998	24,940
1999	23,647
2000	22,815
2001	29,522
2002	33,906
2003	26,902
2004	27,334
2005	26,717
2006	25,908
2007	27,440
2008	32,413
2009	38,735
2010	30,511
2011	25,286
Total	396,074

Table 4. Summary of the MRFSS intercept data for queen triggerfish by year. The summary includes the number of AB1 catch, which refers to the number of observed landed queen triggerfish and queen triggerfish reported as dead, number of B2 catch, which refers to the number of queen triggerfish released alive, the number of trips catching queen triggerfish; total number of intercepted trips catching any recreationally caught species in Puerto Rico, the proportion of trips catching queen triggerfish, and the number of angler hours.

Year	AB1	B2	# of positive trips	Total # of trips	Proportion positive	Angler hours
2000	9	0	6	737	0.008	3658
2001	22	0	13	768	0.017	4349
2002	2	0	2	517	0.004	3098
2003	5	0	5	812	0.006	5022
2004	2	1	3	621	0.005	3643
2005	4	1	4	426	0.009	2329
2006	2	0	2	366	0.005	2118
2007	3	0	2	572	0.003	2953
2008	23	2	13	623	0.021	3393
2009	23	3	16	581	0.028	3148
2010	10	0	5	588	0.009	3054
2011	2	0	2	774	0.003	3530
Total	107	7	73	27614	0.009	40295

Table 5. Summary of the MRFSS intercept data for queen triggerfish by area. The summary includes the number of AB1 catch, which refers to the number of observed landed queen triggerfish and queen triggerfish reported as dead, number of B2 catch, which refers to the number of queen triggerfish released alive, the number of trips catching queen triggerfish; total number of intercepted trips catching any recreationally caught species in Puerto Rico, the proportion of trips catching queen triggerfish, and the number of angler hours. The areas are: 2 – ocean greater than 10 miles, 3 – ocean less than 10 miles, 4 – ocean greater than 10 miles, and 5 – inshore.

Areas	AB1	B2	Positive trips	Trips	Proportion positive	Angler hours
2	0	0	0	3	0	15
3	93	6	62	4334	0.014	22447
4	12	0	8	1119	0.007	10039
5	2	1	3	1929	0.002	7797
Total	107	7	73	7385	0.009	40298

Table 6. Summary of the MRFSS intercept data for queen triggerfish by year for areas 3 and 5 in Puerto Rico. Areas: 3 – ocean less than 3 miles, and 5 – inshore. The summary includes the number of AB1 catch, which refers to the number of observed landed queen triggerfish and queen triggerfish reported as dead, number of B2 catch, which refers to the number of queen triggerfish released alive, the number of trips catching queen triggerfish; total number of intercepted trips catching any recreationally caught species in Puerto Rico, the proportion of trips catching queen triggerfish, and the number of angler hours.

Year	AB1	B2	# of positive trips	Total # of trips	Proportion positive	Angler hours
2000	5	0	4	628	0.006	2850
2001	18	0	10	656	0.015	3161
2002	2	0	2	419	0.005	2121
2003	5	0	5	657	0.008	3412.5
2004	2	1	3	520	0.006	2682.5
2005	4	1	4	378	0.011	1854.5
2006	2	0	2	287	0.007	1384
2007	2	0	1	450	0.002	2106.5
2008	22	2	12	547	0.022	2665
2009	23	3	16	513	0.031	2668
2010	8	0	4	485	0.008	2215
2011	2	0	2	723	0.003	3124.5
Total	95	7	65	6263	0.010	30244.5

Table 7. Summary of maximum ages and lengths (FL: fork length; SL: Standard length; TL: total length), and the age-length relationship for queen triggerfish in the western Atlantic. (STT: St. Thomas, STJ: St. John, and PR: Puerto Rico.)

Location	t_{max} (years)	L_{max} (mm)	Length -age				Source
			L_{∞}	K	t_o	n	
Jamaica	-	447 FL	-	0.570	-	-	Aiken (1975)
PR	-	546 FL	-	-	-	-	Bohnsack and Harper (1988)
STT & STJ	-	465 FL	-	-	-	-	Bohnsack and Harper (1988)
PR & USVI	7	-	415 FL	0.300	-0.600	494	Manooch and Drennon (1987)
PR	-	760 FL	-	-	-	-	Pagán 2002
St. John	-	385 FL	-	-	-	-	Randall (1962)
USVI	-	572 FL	-	-	-	-	Randall (1968)
Brazil	14	460 FL	441 FL	0.140	-1.800	476	de Albuquerque et al. (2011)
south Florida	-	425 FL	-	-	-	-	Bohnsack and Harper (1988)
Brazil	-	450 FL	-	-	-	-	Menezes (1979)

Table 8. Summary of the Trip Interview Program (TIP) data available for queen triggerfish. The data include the number of length measurements, the number of years sampled, and the average number of lengths per year by island and gear type. The analysis type for each strata is also noted. TS indicates that a length-based time-series analysis was done and ID indicates that there were insufficient data for analysis.

Island	Gear type	Number of lengths	Number of years sampled	Average number of lengths year ⁻¹	Analysis
-	Pots & traps	122	2	-	-
Puerto Rico	Dive/Spear/By Hand	1011	26	38.8	ID
Puerto Rico	Hook & line	933	28	33.3	ID
Puerto Rico	Nets	1157	26	44.5	ID
Puerto Rico	Pots & traps	4917	28	175.6	TS
St. Thomas/St John	Dive/Spear/By Hand	6	1	6	ID
St. Thomas/St John	Hook & line	297	12	24.75	ID
St. Thomas/St John	Nets	2	2	1	ID
St. Thomas/St John	Pots & traps	7731	21	368.14	TS
St. Croix	Dive/Spear/By Hand	644	12	53.67	ID
St. Croix	Hook & line	313	12	26.08	ID
St. Croix	Nets	37	7	5.28	ID
St. Croix	Pots & traps	8025	28	286.61	TS

Table 9. Estimates of central tendency and skewness of the queen triggerfish length frequency data from the pot and trap fishery in Puerto Rico and collected by TIP. Estimates correspond to the annual length frequency plots shown in Figure 10.

Species	Gear	Island	Year	Mean	Median	Mode	Skewness
Queen triggerfish	Pots and traps	Puerto Rico	1983	27.53	27.00	26.25	0.66
			1984	28.98	28.50	29.25	1.03
			1985	27.33	27.00	23.25	0.66
			1986	27.31	25.50	23.25	0.86
			1987	28.88	28.50	29.25	0.54
			1988	28.01	28.50	27.75	0.36
			1989	27.31	27.00	29.25	0.29
			1990	28.54	28.50	27.75	0.72
			1991	27.99	27.00	26.25	1.30
			1992	27.70	27.00	21.75	1.82
			1993	27.13	27.00	23.25	0.47
			1994	27.08	25.50	23.25	0.85
			1995	26.59	25.50	24.75	0.78
			1996	30.50	28.50	27.75	2.69
			1997	27.89	26.25	24.75	0.31
			1998	31.78	28.50	24.75	1.90
			1999	36.57	30.00	27.75	0.90
			2000	35.12	30.00	32.25	0.90
			2001	29.10	28.50	27.75	0.10
			2002	31.42	30.00	32.25	1.74
			2003	36.77	33.00	32.25	0.93
			2004	34.85	33.00	32.25	1.48
			2005	33.25	33.00	32.25	-0.16
2006	28.94	28.50	30.75	0.52			
2007	34.28	31.50	24.75	1.64			
2008	27.30	27.00	30.75	0.28			
2009	36.38	36.00	26.25	0.01			
2011	29.88	28.50	27.75	0.38			

Table 10. Estimates of central tendency and skewness of the queen triggerfish length frequency data from the St. Thomas and St. John pot and trap fishery collected by TIP. Estimates correspond to the annual length frequency plots shown in Figure 11.

Species	Gear	Island	Year	Mean	Median	Mode	Skewness
Queen triggerfish	Pots and traps	St. Thomas and St. John	1983	28.76	28.50	23.25	0.53
			1984	31.29	31.50	32.25	-0.30
			1985	31.50	31.50	29.25	0.07
			1986	32.94	33.00	36.75	-0.71
			1987	28.53	28.50	24.75	0.27
			1988	31.42	31.50	32.25	0.31
			1991	31.50	31.50	30.75	NaN
			1992	27.84	28.50	27.75	0.49
			1993	29.51	30.00	27.75	0.43
			1994	29.68	28.50	26.25	0.63
			1995	29.44	28.50	27.75	0.50
			1996	26.67	25.50	23.25	1.76
			2002	32.44	32.25	33.75	0.21
			2003	34.13	34.50	35.25	-0.08
			2004	32.64	31.50	30.75	3.25
			2005	31.49	31.50	30.75	-0.18
			2006	29.82	30.00	20.25	0.60
			2008	30.41	31.50	32.25	-0.22
			2009	32.41	33.00	33.75	-0.13
			2010	32.31	33.00	32.25	-0.03
2011	31.72	31.50	35.25	0.20			

Table 11. Estimates of central tendency and skewness of the queen triggerfish length frequency data from the St. Croix pot and trap fishery collected by TIP. Estimates correspond to the annual length frequency plots shown in Figure 12.

Species	Gear	Island	Year	Mean	Median	Mode	Skewness
Queen triggerfish	Pots and traps	St. Croix	1983	30.86	30.00	29.25	0.38
			1984	28.58	28.50	29.25	0.24
			1985	29.21	28.50	26.25	1.57
			1986	29.95	30.00	26.25	0.55
			1987	27.64	27.00	27.75	0.28
			1988	26.96	27.00	26.25	0.95
			1989	26.45	27.00	26.25	0.83
			1990	27.46	27.00	23.25	0.85
			1991	27.22	27.00	29.25	0.72
			1992	27.10	27.00	24.75	0.57
			1993	27.68	27.00	26.25	0.32
			1994	27.32	27.00	24.75	0.62
			1995	26.08	25.50	24.75	0.63
			1996	25.30	25.50	24.75	0.44
			1997	26.90	27.00	24.75	0.09
			1998	26.16	27.00	26.25	0.16
			1999	27.12	27.00	23.25	0.35
			2000	26.31	27.00	26.25	-0.16
			2001	23.43	22.50	20.25	0.71
			2002	28.41	28.50	29.25	0.10
			2003	28.50	28.50	27.75	NaN
2004	26.38	26.25	21.75	0.27			
2005	22.23	21.00	20.25	0.04			
2007	27.86	27.00	24.75	0.51			
2008	25.42	25.50	24.75	-0.06			
2009	28.93	28.50	27.75	0.58			
2010	28.92	28.50	30.75	0.09			
2011	28.30	28.50	27.75	0.17			

Table 12. Input parameter ranges. L_c : length at full vulnerability measured in cm, K : von Bertalanffy growth coefficient, and L_∞ : asymptotic length measured in cm.

Species name	Island	Gear type	L_c (cm FL)	K	L_∞ (cm FL)
Queen triggerfish	Puerto Rico	Pots and traps	28	0.165	37.3
			29	0.3	41.5
			31	0.435	45.6
Queen triggerfish	St. Thomas/St. John	Pots and traps	32	0.165	37.3
			34	0.3	41.5
			35	0.435	45.6
Queen triggerfish	St. Croix	Pots and traps	27	0.165	37.3
			28	0.3	41.5
			29	0.435	45.6

Table 13. The full AIC results from the preliminary analyses using the queen triggerfish length data from the pot and trap fisheries in a) Puerto Rico , b) St. Thomas/St. John, and c) St. Croix. The rows highlighted in gray have the lowest AIC values. Total mortality values that are equal to 0.001 or are greater than or equal to 4.99 indicate the model did not converge and parameter bounds were reached.

L_c	K	L_∞	Npar	Nobs	Nchange	AIC	Δ AIC	LLIKE	Z	Z1	Change Year1	Z2	Change Year2	Z3	Change Year3	Z4
a)																
29	0.3	41.5	0	2	28	165.90	12.27	80.71	0.38	-	-	-	-	-	-	-
			1	4	28	153.63	0.00	71.95	-	0.65	1990	0.05	-	-	-	-
			2	6	28	155.53	1.90	69.76	-	0.65	1990	0.001	1999	0.35	-	-
			3	8	28	162.66	9.03	69.54	-	0.61	1987	1.30	1989	0.001	1999	0.36
b)																
34	0.3	41.5	0	2	20	49.63	0	22.46	0.43	-	-	-	-	-	-	-
			1	4	20	55.15	5.52	22.24	-	0.48	1983	0.42	-	-	-	-
			2	6	20	60.73	11.1	21.14	-	0.65	1983	0.001	1985	0.43	-	-
			3	8	20	70.27	20.64	20.59	-	0.44	2001	0.19	2006	0.73	2009	0.12
c)																
28	0.3	41.5	0	2	27	86.66	14.81	41.08	0.84	-	-	-	-	-	-	-
			1	4	27	71.85	0	31.02	-	0.7	1985	1.07	-	-	-	-
			2	6	27	77.24	5.39	30.52	-	0.7	1985	1.09	2008	0.76	-	-
			3	8	27	79.91	8.06	27.95	-	0.7	1985	1.11	1991	0.54	1993	1.19

Table 14. The number of sensitivity runs predicting zero, one, two, or three changes in total mortality (Z) when using the queen triggerfish length data from the pot and trap fishery in Puerto Rico collected by TIP. Four length-at-full vulnerability (L_c) values were used as inputs. The results highlighted in gray represent the models supported by the majority of sensitivity runs.

Lc	Number of changes in total mortality (Z)			
	No change	1 change	2 changes	3 changes
28	0	22	17	1
29	1	21	18	0
30.5	2	23	15	0
31	4	22	14	0
Total	7	88	64	1

Table 15. The number of sensitivity runs predicting one or two changes in total mortality (Z) and the frequency of the associated predicted first year of change when using the queen triggerfish length data from the pot and trap fishery in Puerto Rico collected by TIP. These results represent all parameter combinations used for the sensitivity analysis.

Year of first change	Number of changes in total mortality (Z)	
	1	2
1983	11	0
1984	4	0
1985	3	0
1986	7	0
1987	8	2
1988	13	1
1989	19	4
1990	12	18
1991	7	15
1992	4	9
1993	0	11
1994	0	4
Total	88	64

Table 16. The number of sensitivity runs predicting two, or three changes in total mortality (*Z*) and the frequency of the associated predicted second year of change when using the queen triggerfish length data from the pot and trap fishery in Puerto Rico collected by TIP. These results represent all parameter combinations used for the sensitivity analysis. The year with support from the majority of sensitivity runs is highlighted in gray.

Year of second change	Number of changes in total mortality (<i>Z</i>)
	2
1998	2
1999	62
Total	64

Table 17. Percent change in total mortality for sensitivity runs strongly supporting the model predicting a single year of change. The data used for these sensitivity runs were the queen triggerfish length data from the pot and trap fishery in Puerto Rico.

		<i>K</i>				
<i>L_c</i>	<i>L_∞</i>	0.165	0.265	0.3	0.365	0.435
28	37	-99.5	-99.7	-99.7	-99.8	-99.8
	38.5	-99.6	-99.7	-99.8	-99.8	-99.8
	40	-99.7	-99.8	-95.0	-92.2	-90.5
	41.5	-99.7				-81.3
	43	-87.6		-78.5	-77.4	
	44.5		-75.4	-75.0		
	45.6					
	46					
29	37		-99.6	-99.7	-99.7	-99.8
	38.5	-99.6	-99.7	-99.8	-99.8	-99.8
	40	-99.7	-99.8	-99.8	-99.8	-99.9
	41.5	-99.7	-96.3			
	43	-97.4		-84.9	-82.5	
	44.5					
	45.6	-82.4				
	46	-81.0				
30.5	37		-99.4	-99.5	-99.7	-99.7
	38.5		-99.7	-99.7	-99.8	-99.8
	40	-99.6	-99.8	-99.8	-99.8	-99.9
	41.5	-99.7	-99.8	-99.8	-99.9	-99.9
	43	-99.7			-89.6	-87.7
	44.5					
	45.6	-87.9				
	46	-86.1				
31	37					-99.7
	38.5	-99.3	-99.7	-99.7	-99.8	-99.8
	40	-99.6	-99.8	-99.8	-99.8	-99.8
	41.5	-99.7	-99.8	-99.8	-99.9	-99.9
	43	-99.8	-99.8	-96.2		
	44.5	-99.8	-89.0	-88.3		
	45.6					
	46					

Table 18. Percent change in total mortality for sensitivity runs strongly supporting the model predicting two years of change. The data used for these sensitivity runs were the queen triggerfish length data from the pot and trap fishery in Puerto Rico.

L_c	L_∞	K									
		0.165		0.265		0.3		0.365		0.435	
28	41.5			-100	34800	-100	36872	-100	41705		
	43			-100	42036					-100	58506
	44.5	-100	34742					-100	60459	-100	67033
	45.6	-100	33019	-100	50871	-100	57079	-100	64259	-100	75865
	46	-100	34541	-100	53016	-100	59449	-100	66977		
	41.5					-100	35333	-100	39749	-100	46038
29	43			-100	41374					-100	56223
	44.5	-100	35412	-100	48041	-100	53162	-100	59275	-100	65198
	45.6			-100	54311	-100	57041	-100	63653	-100	73764
	46			-100	53598	-100	59409	-100	66330	-100	76909
	43			-100	40287	-100	44415				
	44.5	-100	38074	-100	50363	-100	52796	-100	58342	-100	66141
30.5	45.6			-100	54836	-100	59921	-100	66161	-100	72277
	46			-100	57194	-100	62437	-100	68957	-100	75407
	43							-100	45346	-100	52203
	44.5							-100	58178	-100	63685
	45.6	-100	40948	-100	52594	-100	57918	-100	64272	-100	73223
31	46	-100	40986	-100	55215	-100	60688	-100	67308	-100	73733

Table 19. Natural and fishing mortality estimates for queen triggerfish caught by the pot and trap fishery in Puerto Rico. Input parameters include: asymptotic length (L_{∞} , FL mm), the von Bertalanffy growth coefficient (K), maximum age ($tmax$), the proportion of population at the maximum age (P), and temperature (the average for Puerto Rico: www.nodc.noaa.gov/dsdt/cwtg/oatl.html). Footnotes indicate the equation associated with each publication. Fishing mortality estimates ($F = Z - M$) result from two estimates of total mortality from the sensitivity analysis and the natural mortality estimates. Cells heightened in gray indicate F is greater than M .

Input parameters					Source of natural mortality equation							
L_{∞}	K	$tmax$	P	Temp	Alverson and Carney 1975 ¹	Pauly 1980 ²	Hoening 1983 ³ (regression)	Hoening 1983 ⁴	Ralston 1987 ⁵	Jensen 1996 ⁶ (theoretical)	Jensen 1996 ⁷ (derived from Pauly 1980)	Hewitt and Hoening 2005 ⁸
415	0.3	7	0.05	26.6	0.74	0.72	0.62	0.43	0.64	0.45	0.48	0.60
441	0.14	14	0.05	26.6	0.38	0.43	0.32	0.21	0.31	0.21	0.22	0.30
Total mortality assumed equal to 0.02 (lower estimate from sensitivity analysis)												
L_{∞}	K	$tmax$	P	Temp	Fishing mortality estimates							
415	0.3	7	0.05	26.6	-0.72	-0.70	-0.60	-0.41	-0.62	-0.43	-0.46	-0.58
441	0.14	14	0.05	26.6	-0.36	-0.41	-0.30	-0.19	-0.29	-0.19	-0.20	-0.28
Total mortality assumed equal to 0.75 (upper estimate from sensitivity analysis)												
L_{∞}	K	$tmax$	P	Temp	Fishing mortality estimates							
415	0.3	7	0.05	26.6	0.01	0.03	0.13	0.32	0.11	0.30	0.27	0.15
441	0.14	14	0.05	26.6	0.37	0.32	0.43	0.54	0.44	0.54	0.53	0.45

¹ $M = 3K/(\exp[0.38*K*tmax] - 1)$, ² $M = \exp[-0.0152 + 0.6543*\ln(K) - 0.279*\ln(L_{\infty}/10) + 0.4634*\ln(Temp)]$

³ $M = \exp[1.44 - 0.982*\ln(tmax)]$, ⁴ $M = -\ln(P)/tmax$, ⁵ $M = 0.0189 + 2.06K$, ⁶ $M = 1.5K$, ⁷ $M = 1.6K$, ⁸ $M = 4.22/tmax$

Table 20. The frequency of sensitivity runs predicting a change or lack of change in total mortality (Z) when using the queen triggerfish length data from the St. Thomas/St. John pot and trap fishery collected by TIP. Four values of the length-at-full vulnerability (L_c) were used as part of the sensitivity analysis.

L_c	Number of changes in Z	
	0 changes	
32	40	
33.5	40	
34	40	
35	40	
Total	160	

Table 21. Natural and fishing mortality estimates for queen triggerfish caught by the St. Thomas/St. John pot and trap fishery. Input parameters include: asymptotic length (L_{∞} , FL mm), the von Bertalanffy growth coefficient (K), maximum age ($tmax$), the proportion of population at the maximum age (P), and temperature (the average for Puerto Rico: www.nodc.noaa.gov/dsdt/cwtg/oatl.html). Footnotes indicate the equation associated with each publication. Fishing mortality estimates ($F = Z - M$) result from two estimates of total mortality from the sensitivity analysis and the natural mortality estimates. Cells heightened in gray indicate F is greater than M .

Input parameters					Source of natural mortality equation							
L_{∞}	K	$tmax$	P	Temp	Alverson and Carney 1975 ¹	Pauly 1980 ²	Hoening 1983 ³ (regression)	Hoening 1983 ⁴	Ralston 1987 ⁵	Jensen 1996 ⁶ (theoretical)	Jensen 1996 ⁷ (derived from Pauly 1980)	Hewitt and Hoening 2005 ⁸
415	0.3	7	0.05	26.6	0.74	0.72	0.62	0.43	0.64	0.45	0.48	0.60
441	0.14	14	0.05	26.6	0.38	0.43	0.32	0.21	0.31	0.21	0.22	0.30
Total mortality assumed equal to 0.2 (lower estimate from sensitivity analysis)												
L_{∞}	K	$tmax$	P	Temp	Fishing mortality estimates							
415	0.3	7	0.05	26.6	-0.54	-0.52	-0.42	-0.23	-0.44	-0.25	-0.28	-0.40
441	0.14	14	0.05	26.6	-0.18	-0.23	-0.12	-0.01	-0.11	-0.01	-0.02	-0.10
Total mortality assumed equal to 1 (upper estimate from sensitivity analysis)												
L_{∞}	K	$tmax$	P	Temp	Fishing mortality estimates							
415	0.3	7	0.05	26.6	0.26	0.28	0.38	0.57	0.36	0.55	0.52	0.40
441	0.14	14	0.05	26.6	0.62	0.57	0.68	0.79	0.69	0.79	0.78	0.70

¹ $M = 3K/(\exp[0.38*K*tmax] - 1)$, ² $M = \exp[-0.0152 + 0.6543*\ln(K) - 0.279*\ln(L_{\infty}/10) + 0.4634*\ln(Temp)]$

³ $M = \exp[1.44 - 0.982*\ln(tmax)]$, ⁴ $M = -\ln(P)/tmax$, ⁵ $M = 0.0189 + 2.06K$, ⁶ $M = 1.5K$, ⁷ $M = 1.6K$, ⁸ $M = 4.22/tmax$

Table 22. The predicted number of changes in total mortality and the associated frequency of sensitivity runs when using the TIP length data from the St. Croix pot and trap fishery. These results represent all parameter combinations used for the sensitivity analysis. The number of changes in total mortality supported by the majority of sensitivity runs is highlighted in gray.

Lc	Number of changes in total mortality (Z)	
	1	2
27	26	14
28	40	0
28.5	40	0
29	40	0
Total	146	14

Table 23. The predicted year of change for the sensitivity runs that resulted in strong support for the model predicting one change in total mortality when using the length data from the St. Croix pot and trap fishery collected by TIP. The year(s) supported by the majority of sensitivity runs is highlighted in gray.

Change year	One change in total mortality
1983	4
1984	8
1985	92
1986	42
Total	146

Table 24. Percent change in total mortality for sensitivity runs strongly supporting the model predicting a single year of change. The data used for these sensitivity runs were the queen triggerfish length data from the pot and trap fishery in St. Croix.

		K				
L_c	L_∞	0.165	0.265	0.3	0.365	0.435
27	37			84.90		82.93
	38.5		78.50		75.03	72.39
	40				67.78	65.19
	41.5		66.86	65.20	62.44	60.00
	43	65.12	62.71	61.02	58.36	
	44.5	64.73	59.33	57.68	55.16	57.58
	45.6		57.24	55.64		55.90
	46		56.55	54.96		55.34
28	37	80.28	71.30	71.60	70.47	68.74
	38.5	70.21	64.04	63.19	61.31	59.32
	40	60.39	58.33	57.17	55.08	53.14
	41.5	55.46	53.95	52.69	50.62	48.79
	43	54.36	50.53	49.26	47.27	45.58
	44.5	52.02	47.79	46.56	44.68	45.73
	45.6	50.49	46.12	44.92	43.12	44.39
	46	49.97	45.57	44.39	42.61	43.95
28.5	37	83.02	75.87	75.19	73.20	70.61
	38.5	68.33	65.91	64.53	61.75	58.91
	40	62.58	58.81	57.12	54.18	57.38
	41.5	58.47	53.50	51.73	54.35	53.00
	43	55.07	49.39	52.47	51.09	49.71
	44.5	52.15	50.79	49.97	48.50	47.16
	45.6	50.26	49.28	48.41	46.93	45.62
	46	49.62	48.77	47.89	46.42	45.13
29	37	89.89	79.17	77.81	75.00	71.97
	38.5	72.05	67.28	65.50	62.35	59.43
	40	63.20	59.35	57.46	54.44	56.56
	41.5	59.90	53.68	51.86	53.56	51.82
	43	55.89	49.46	51.87	50.00	48.36
	44.5	52.59	50.21	49.07	47.26	45.74
	45.6	50.51	48.50	47.37	45.63	44.20
	46	49.82	47.94	46.81	45.10	43.69

Table 25. Natural and fishing mortality estimates for queen triggerfish caught by the St. Croix pot and trap fishery. Input parameters include: asymptotic length (L_{∞} , FL mm), the von Bertalanffy growth coefficient (K), maximum age (t_{max}), the proportion of population at the maximum age (P), and temperature (the average for Puerto Rico: www.nodc.noaa.gov/dsdt/cwtg/oatl.html). Footnotes indicate the equation associated with each publication. Fishing mortality estimates ($F = Z - M$) result from two estimates of total mortality from the sensitivity analysis and the natural mortality estimates. Cells highlighted in gray indicate F is greater than M .

Input parameters					Source of natural mortality equation							
L_{∞}	K	t_{max}	P	Temp	Alverson and Carney 1975 ¹	Pauly 1980 ²	Hoenig 1983 ³ (regression)	Hoenig 1983 ⁴	Ralston 1987 ⁵	Jensen 1996 ⁶ (theoretical)	Jensen 1996 ⁷ (derived from Pauly 1980)	Hewitt and Hoenig 2005 ⁸
415	0.3	7	0.05	26.6	0.74	0.72	0.62	0.43	0.64	0.45	0.48	0.60
441	0.14	14	0.05	26.6	0.38	0.43	0.32	0.21	0.31	0.21	0.22	0.30
Total mortality assumed equal to 0.4 (lower estimate from sensitivity analysis)												
L_{∞}	K	t_{max}	P	Temp	Fishing mortality estimates							
415	0.3	7	0.05	26.6	-0.34	-0.32	-0.22	-0.03	-0.24	-0.05	-0.08	-0.20
441	0.14	14	0.05	26.6	0.02	-0.03	0.08	0.19	0.09	0.19	0.18	0.10
Total mortality assumed equal to 1.5 (lower estimate from sensitivity analysis)												
L_{∞}	K	t_{max}	P	Temp	Fishing mortality estimates							
415	0.3	7	0.05	26.6	0.76	0.78	0.88	1.07	0.86	1.05	1.02	0.90
441	0.14	14	0.05	26.6	1.12	1.07	1.18	1.29	1.19	1.29	1.28	1.20

¹ $M = 3K/(\exp[0.38*K*t_{max}] - 1)$, ² $M = \exp[-0.0152 + 0.6543*\ln(K) - 0.279*\ln(L_{\infty}/10) + 0.4634*\ln(Temp)]$

³ $M = \exp[1.44 - 0.982*\ln(t_{max})]$, ⁴ $M = -\ln(P)/t_{max}$, ⁵ $M = 0.0189 + 2.06K$, ⁶ $M = 1.5K$, ⁷ $M = 1.6K$, ⁸ $M = 4.22/t_{max}$

9. Figures




Figure 1. Puerto Rico yearly commercial fishery landings of queen triggerfish, expansion factors applied, by gear and year.




Figure 2. Puerto Rico reported yearly commercial fishing trips with queen triggerfish landings by coast and year.




Figure 3. Puerto Rico reported yearly commercial fishing trips with queen triggerfish landings by gear and year.




Figure 4. Yearly commercial landings of triggerfishes as reported (no expansion factors applied) on fisher logbooks from St. Thomas and St. John by gear and year.




Figure 5. St. Thomas and St. John reported commercial fishing trips with triggerfish landings by gear and year.




Figure 6. Yearly commercial landings of triggerfishes as reported (no expansion factors applied) on fisher logbooks from St. Croix by gear and year.




Figure 7. St. Croix reported commercial fishing trips with triggerfish landings by gear and year.




Figure 8. Length-frequency plot for queen triggerfish caught in Puerto Rico and intercepted by MRFSS between 2000 and 2011. N = 60.




Figure 9. The von Bertalanffy growth curves for queen triggerfish as defined in Table 7.




Figure 10. Annual length frequency plots of queen triggerfish from the Puerto Rican pot and trap fishery. Data are from the TIP database. The length bin size is 1.5cm. The solid red line is the mode, the dashed red line is the mean, and the dotted line is the median. In some instances there is more than one mode because the frequency of more than one length bin is equal. N indicates the number of measured lengths.




Figure 10. continued




Figure 10. continued




Figure 10. continued




Figure 11. Annual length frequency plots of queen triggerfish caught by the pot and trap fishery on the west coast of the Puerto Rico. The west coast includes: Aguada, Agudilla, Arecibo, Cabo Rojo, Mayaguez, and Rincon. The bin size is 1.5cm. The solid red line is the mode, the dashed red line is the mean, and the dotted line is the median. In some instances there is more than one mode because the frequency of more than one length bin is equal. N indicates the number of measured lengths.




Figure 11. continued




Figure 11. continued




Figure 12. Annual length frequency plots of queen triggerfish caught by the pot and trap fishery on the east coast of the Puerto Rico. The east coast includes: Ceiba, Culebra, Fajardo, Humacoa, Naguabo, Vieques, and Yabucoa. The bin size is 1.5cm. The solid red line is the mode, the dashed red line is the mean, and the dotted line is the median. In some instances there is more than one mode because the frequency of more than one length bin is equal. N indicates the number of measured lengths.




Figure 12. continued




Figure 12. continued




Figure 13. Annual length frequency plots of queen triggerfish lengths from the St. Thomas/St. John pot and trap fishery. The data are from the TIP database. The bin size is 1.5cm. The solid red line is the mode, the dashed red line is the mean, and the dotted line is the median. In some instances there is more than one mode because the frequency of more than one length bin is equal. N indicates the number of measured lengths.




Figure 13. continued




Figure 13. continued




Figure 14. Annual length frequency plots of queen triggerfish caught by the St. Croix pot and trap fishery. The data are from the TIP database. The bin size is 1.5cm. The solid red line is the mode, the dashed red line is the mean, and the dotted line is the median. In some instances there is more than one mode because the frequency of more than one length bin is equal. N indicates the number of measured lengths.




Figure 14. continued





Figure 14. continued



pots_traps

STX

Fork length (cm)

Figure 14. continued




Figure 15. The fit to the queen triggerfish mean length data from the Puerto Rican pot and trap fishery. The model fit shown predicts one change in total mortality and had the lowest AIC value. The input parameter values used were $L_c = 29\text{cm}$, $L_\infty = 41.5\text{cm}$, and $K = 0.3\text{y}^{-1}$. Bubble size represents the sample size scaled with respect to other years.




Figure 16. The fit to the queen triggerfish mean length data from the Puerto Rican pot and trap fishery. The model fit shown predicts two changes in total mortality was strongly supported by AIC criteria (i.e., $\Delta AIC = 1.9$). The input parameter values used were $L_c = 29\text{cm}$, $L_\infty = 41.5\text{cm}$, and $K = 0.3\text{y}^{-1}$. The bubble size represents the sample size scaled with respect to other years.




Figure 17. The estimates of the current total mortality, $Z_{current}$, from the sensitivity analysis carried out on the queen triggerfish length data from the pot and trap fishery in Puerto Rico. Each panel represents a different length-at-full vulnerability value, L_c , and is indicated above the panel. The colored points represent different values of L_{∞} , the values are indicated in the figure legend.




Figure 18. The model fit to the queen triggerfish mean length data from the St. Thomas/St. John pot and trap fishery. The model fit shown predicts no change in total mortality and had the lowest AIC value. The input parameter values used were $L_c = 34\text{cm}$, $L_\infty = 41.5\text{cm}$, and $K = 0.3\text{y}^{-1}$. Bubble size represents the sample size scaled with respect to other years.




Figure 19. The estimates of the current total mortality, $Z_{current}$, from the sensitivity analysis carried out on the queen triggerfish data from the St. Thomas/St. John pot and trap fishery. Each panel represents a different length-at-full vulnerability value, L_c , and is indicated above the panel. The colored points represent different values of L_{∞} , the values are indicated in the figure legend.




Figure 20. The model fit to the queen triggerfish mean length data from the St. Croix pot and trap fishery. The model fit predicts a single change in total mortality and had the lowest AIC value. The input parameter values used were $L_c = 28\text{cm}$, $L_\infty = 41.5\text{cm}$, and $K = 0.3\text{y}^{-1}$. Bubble size represents the sample size scaled with respect to other years.





Figure 21. The estimates of the current total mortality, $Z_{current}$, from the sensitivity analysis carried out on the queen triggerfish data from the St. Croix pot and trap fishery. Each panel represents a different length-at-full vulnerability value, L_c , and is indicated above the panel. The colored points represent different values of L_∞ , the values are indicated in the figure legend.



SEDAR

Southeast Data, Assessment, and Review

SEDAR 30

U.S. Caribbean Queen Triggerfish

SECTION III: Review Report

March 2013

SEDAR
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405

Table of Contents

Table of Contents 2

1. INTRODUCTION 2

 1.1 WORKSHOP TIME AND PLACE 2

 1.2 TERMS OF REFERENCE..... 2

 1.3 LIST OF PARTICIPANTS 3

2. CIE REVIEWER REPORTS 3

1. INTRODUCTION

1.1 WORKSHOP TIME AND PLACE

The SEDAR 30 Peer Review Process was conducted via a CIE (Center for Independent Experts) Desk Review in lieu of a Panel Review Workshop. Three reviewers were selected by provided the CIE and provided with the assessment report and background materials. Each reviewer conducted a review of the material and produced an independent review report. Those reports are included below.

1.2 TERMS OF REFERENCE

1. Evaluate the data used in the assessment, addressing the following:
 - a) Are data decisions made by the Assessment Workshop sound and robust?
 - b) Are data uncertainties acknowledged, reported, and within normal or expected levels?
 - c) Are data applied properly within the assessment model?
 - d) Are input data series reliable and sufficient to support the assessment approach and findings?
2. Evaluate the methods used to assess the stock, taking into account the available data.
 - a) Are methods scientifically sound and robust?
 - b) Are assessment models configured properly and used consistent with standard practices?
 - c) Are the methods appropriate for the available data?
3. Evaluate the assessment findings with respect to the following:
 - a) Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?
 - b) Is the stock overfished? What information helps you reach this conclusion?

- c) Is the stock undergoing overfishing? What information helps you reach this conclusion?
 - d) Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?
 - e) Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?
4. Evaluate the stock projections, addressing the following:
- a) Are the methods consistent with accepted practices and available data?
 - b) Are the methods appropriate for the assessment model and outputs?
 - c) Are the results informative and robust, and useful to support inferences of probable future conditions?
 - d) Are key uncertainties acknowledged, discussed, and reflected in the projection results ?
5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.
- Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods
 - Ensure that the implications of uncertainty in technical conclusions are clearly stated.
6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.
- Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
 - Provide recommendations on possible ways to improve the SEDAR process.
7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.

1.3 LIST OF PARTICIPANTS

CIE Reviewers

Massimiliano Cardinale	CIE Reviewer
Yong Chen	CIE Reviewer
M. Kurtis Trzcinski.....	CIE Reviewer

2. CIE REVIEWER REPORTS

**Center for Independent Experts (CIE)
Independent Peer Review Report of the SEDAR
30 Caribbean Blue Tang and Queen Triggerfish
assessment review**

Dr. Massimiliano Cardinale

March 2013

Executive Summary

- This document is the individual Center for Independent Experts (CIE) review report of the SEDAR 30 Caribbean blue tang and queen triggerfish assessments conducted during February 2013 and provided at the request of the CIE (see Attachment A).
- This report solely represents the views of the independent reviewer (Dr. Massimiliano Cardinale).
- This reviewer does not completely agree with all of the findings reported in the SEDAR 30 Caribbean queen triggerfish assessment report, while the reviewer is in general agreement concerning the blue tang assessment report. Findings that are reported in the SEDAR 30 Caribbean blue tang and queen triggerfish assessments reports are not necessarily fully repeated in this individual report. This report focuses on clarifications of elements contained in the Summary Report and some additional views of the individual reviewer about how data for queen triggerfish could have been better explored to derive more robust estimates of exploitation rates and thus stock status.
- The assessment team tackled all of the review terms of reference (TORs).
- This reviewer believes that the SEDAR 30 has done a good job in carrying out the assessment, analysing all available source of data, modelling uncertainty and providing a full sensitivity analysis of both the data and the models. However, the reviewer is of the opinion that data for queen triggerfish are underutilised and that the reader is left with the doubt that more could have been done in terms of data analysis to derive estimates of exploitation rates and thus stock status for this species.
- For Caribbean blue tang, the report gives the impression that stability in average length is taken as an indication of a low level of F. The reviewer disagrees with this idea, and considers that given the available information the status of the stock should be considered as unknown.
- Further recommendations aimed at improving the data source used in the Caribbean queen triggerfish and blue tang assessment were made. These are based on additional future research and further re-analysis and modelling of the original data set.

Introduction

SEDAR 30 Caribbean blue tang and queen triggerfish assessments reports and associated background documents containing detailed information on the data used in the assessment were provided to the independent reviewer (Dr. Massimiliano Cardinale) well in advance of the deadline scheduled for the 28th of February 2013. The reports were reviewed at the request of the CIE (see Attachment A).

Description of review activities

This review was undertaken by Dr. Massimiliano Cardinale as desk work during February 2013 at the request of the CIE (see Attachment A).

Relevant documents (see bibliography, Attachment B) were made available four weeks prior to the deadline through email and via a link to an ftp or SEDAR 30 website (https://grunt.sefsc.noaa.gov/sedar/Sedar_Documents.jsp?WorkshopNum=30&FolderType=Assessment). The documentation was reviewed prior to the deadline and the deadline was met. The background information and assessments of Caribbean blue tang and queen triggerfish was presented through two documents (see Attachment A). Background information relevant to this review are presented in a series of appendices, including: CIE Statement of Work (Attachment A); a bibliography (Attachment B), report format (Annex 1); Terms of Reference (Annex 2); Comments included here are provided following the terms of reference (TORs) (Annex 2) and are those of this independent reviewer only. The list of main documents provided as background material is included in Attachment B. Additional presentations and documentations were made available during the meeting and were continuously updated under the ftp or SEDAR 30 website (https://grunt.sefsc.noaa.gov/sedar/Sedar_Documents.jsp?WorkshopNum=30&FolderType=Assessment).

Summary of findings

Recommendations

1. Estimate time series of landings per unit of effort (LPUE) for Puerto Rico queen triggerfish and investigate the possibility to derive the proportion of queen triggerfish within triggerfish and blue tang within surgeonfish from the Trip Interview Program (TIP) data. This would allow estimating the total number of fish landed by size class for the main gear (i.e. traps and pots) for both species, combining landings information with the size frequency data from the TIP (see also comments under ToR2).
2. Explore also the quality of the effort data for both species from the TIP, with the aim to produce an effort standardized time series of length frequency distribution (LFD) for queen triggerfish and blue tang.
3. **Queen triggerfish catch data from Puerto Rico traps and pots:** Estimate the total number of fish landed by size class for the main gear (i.e. traps and pots), combining the landings information with the size frequency data from the TIP; Statistical slicing of the total number of fish landed by size class by the main gear (i.e. traps and pots) to estimate the number of fish landed per age class for years with sufficient length measurements (i.e. for years from 1983 to 1988; a general rule of thumb would be to use years with more than 150 or 200 individuals); Estimation of Z from the catch curve or using a pseudocohort analysis (i.e. VIT when only one or few years of data are available); Conducting a yield-per-recruit (YPR) analysis to estimate F_{01} as proxy for F_{MSY} to be compared against estimate of M
4. **TIP data of queen triggerfish:** Explore the use of effort data from the TIP survey to produce an effort standardized time series of LFD for years with sufficient length measurements (a general rule of thumb would be to use years with more than 150 or 200 individuals); statistical slicing of the total number of fish caught by size class by the main gear (i.e. traps and pots) to estimate the total number of fish per age class; estimation of Z from the catch curve or using a pseudocohort analysis (i.e. VIT when only one or few years of data is available)

5. The reviewer considers that ProdBiom method (see Abella et al.,1997) might be more appropriate for the estimation of M as it combines in a single framework the growth parameters, the length weight relationship and information on the longevity of the species. Or at least it should be used along with the other methodologies presented in the reports.
6. The reviewer is of the opinion that the combination of large L_{inf} and low k are the most plausible set of VBF parameters for queen triggerfish, given what has been presented in SEDAR 30 AW 03 and according to information available in the literature, and therefore they should have been given more weight in the evaluation of the stock status.
7. The reviewer considers $F_{MSY}=M$ as a large oversimplification, which ignores selectivity that has a large impact on F_{MSY} . The assessment team should try to estimate catch at age data from LFD (which is possible for certain combinations of years and gear type) and conduct a VIT and an YPR analysis for queen triggerfish based on selected yearly data to verify how realistic is the $F_{MSY}=M$ assumption.
8. For Caribbean blue tang, there is some implication in the report that stability in average length is taken as an indication of a low level of F. The reviewer disagrees with this idea, and considers that given the available information the status of the stock should be considered as unknown.
9. Selectivity studies should be conducted to estimate the effect of the mesh size of the traps on the amount and size distribution of the catches of Caribbean queen triggerfish and blue tang.

Terms of Reference (ToR)

ToR1: Evaluate the data used in the assessment, addressing the following:

- a) Are data decisions made by the Assessment Workshop sound and robust?
- b) Are data uncertainties acknowledged, reported, and within normal or expected levels?
- c) Are data applied properly within the assessment model?
- d) Are input data series reliable and sufficient to support the assessment approach

Puerto Rico reported landings of Caribbean queen triggerfish and Caribbean blue tang have been adjusted for incomplete reporting using so-called expansion factors to estimate the total real landings. It is however unclear, both from the assessment report and from the background documents, how the expansion factor has been estimated (which is the source of the factors), how large the factors are, and if they vary between years for the different areas.

In general, I feel that the landings and effort data are underutilised, especially for Puerto Rico queen triggerfish, for which landings are reported to the level of species. Even the simple estimation of a LPUE time series for Puerto Rico, would have been an useful addition, especially for evaluating estimated time-changes in mortality derived from the Gedamke and Hoenig (2006) method. Also, simple production models might have been tested as an attempt to validate or corroborate the results from the Gedamke and Hoenig (2006) method.

I accept that it is difficult to utilise landings data for Caribbean blue tang as they are reported within the species-group surgeonfish, but especially for short times series such as St. Thomas and St. John, an assumption of constant proportion of landings of blue tang within the species-group surgeonfish could be made. This would allow building CPUE time series also for the other areas and species. In general, I wonder if the TIP data could provide an estimate of the proportion of both species in the landings, when landings data are provided as a species group instead that at the species level. In other words, it would be a useful addition to know if an estimate of the proportion of queen triggerfish within triggerfish and blue tang within surgeonfish might be derived from the TIP data from which the LFD are also derived.

Although it is reasonable to assume that some form of effort data has been collected during the TIP, it is not clear from the assessment reports if such information exists. This has been specified neither in the assessment reports nor in the background information document (i.e. SEDAR 30 AW 02). Effort data from TIP would give a rather different dimension to the LFD as they could provide information more similar to a survey and thus could be useful for estimating stock parameters such as Z and relative changes in population size, especially for queen triggerfish.

Again, considering the large uncertainty associated with the estimate of Z from the Gedamke and Hoenig (2006) method (i.e. violation of constant selectivity assumption, and uncertainty in the VBF parameters), the assessment team should have tried to produce another source of information concerning the exploitation status of the queen triggerfish stock.

I agree instead with the way the assessment team dealt with the available data for Caribbean blue tang. The large uncertainty in the reported Von Bertalanffy (VBF) parameters and, given the fact that this species presents an initial fast growth but a very high longevity, makes the length data uninformative regarding individual ages after age 5. With such large uncertainty in the basic growth parameters, to which both M and F (and F_{MSY}) depend and, due to the peculiar growth characteristics of the species, I agree with the assessment team that it is not possible to use length data to define the stock status of the species. A further difficulty with the use of length data for Caribbean blue tang is the fact that L_c is almost as large as L_{inf} , which makes most of the age classes for which the age could be in theory derived from length information not fully exploited. In this situation, age data are crucial for a robust assessment of this species.

Recreational data for both species are also presented but they are too sparse for allowing any kind of analysis. In this context, the reviewer agrees with the evaluation made by the assessment team.

Recommendations: Estimate time series of CPUE for Puerto Rico queen triggerfish and investigate the possibility of deriving the proportions of queen triggerfish within triggerfish and blue tang within surgeonfish from the TIP data. This would allow estimating the total number of fish landed by size class for the main gear (i.e. traps and pots) for both species, combining landings information with the size frequency data from the TIP (see also comments under ToR2).

Explore also the quality of the effort data for both species from the TIP, with the aim to produce an effort standardized time series of LFD for queen triggerfish and blue tang.

ToR2: Evaluate the methods used to assess the stock, taking into account the available data.

a) Are methods scientifically sound and robust?

b) Are assessment models configured properly and used consistent with standard practices?

The methodology (i.e. Gedamke and Hoenig (2006) method) used to estimate Z has been applied correctly and I consider it as one that is robust and is an appropriate alternative for deriving estimates of exploitation given the available data. However, as for the landings data, and considering the uncertainty associated with the method used, I consider that the length data for queen triggerfish have been underutilised. Thus, other methods should have been used in conjunction with the Gedamke and Hoenig (2006) method to derive estimate of exploitation rates for this species.

For queen triggerfish from Puerto Rico, length frequency data (LFD) from the trap and pot fisheries between 1983 and 1988 are sufficient to estimate the total number of fish caught by the main gear (i.e. traps and pots) by age class, at least for the first 4-6 age classes, which constitutes the main bulk of the catches (compare for example Figures 9 and 10). A recent method has been developed (statistical slicing; see Kell and Kell 2011; Scott et al., 2011) to generate age-structured data for stock assessment from length frequency data and VBF growth curve parameters. The method is very flexible and offers a sophisticated framework for converting numbers at length to numbers at age as well as estimating the mean length at age assuming different distributions of the length data (i.e. Gaussian, gamma and lognormal).

This would allow the assessment team to obtain another and possibly more robust estimate of Z and F (assuming that M is known) from the same length data and to compare them with those derived from the Gedamke and Hoenig (2006) method. In theory, this would also allow for conducting a yield per recruit (YPR) analysis (at least based on the historical part of the times series) and derive estimates of F_{MSY} (using F_{01} as a proxy), which are independent from the estimates of M and take into account selectivity at size/age. Historical estimates of F would be crucial to evaluate the results from the Gedamke and Hoenig (2006) method as well as YPR would be important to define a more robust estimate of F_{MSY} .

However, I also realise that this is conditional on the standardization of the LFD by fishing effort to make them comparable between years and to allow for the catch curve analysis and estimates of Z . For example, the yearly number of trips from which the LFD are derived would be a reasonable index of the effort and sufficient to make the LFD comparable between years. This would allow the use of the statistical slicing method and the catch curve analysis (see also comments and recommendations under ToR2).

Another method that can be used to derive estimates of mortality is the VIT (Leonart and Salat, 2000), which is even more flexible because it can be used also when a single year of LFD and growth parameters are available, thus no effort standardization of the LDF is needed. The method is extensively used in similar data situations with several Mediterranean stocks (e.g. STECF 2012). VIT conducts a virtual population analysis (VPA) assuming a steady state. This is a rather strong assumption for species such as small pelagic fish species, with highly fluctuating abundance due to both variable recruitment and relatively low number of age classes, but it is a much more likely assumption for demersal fish species such as triggerfish for which the population is made up of several age classes. As it requires knowledge of the catches over one year only (Leonart and Salat, 2000) it might be used for years, areas and species for which the data allow for such an analysis. In addition to the above mentioned data, VIT requires a number of biological parameters as growth, length-weight relationship, natural mortalities and percentage mature by size or age, and proportions caught

by each fishing gear (when available, but these parameters are not necessary). These parameters are all available for queen triggerfish and reported in SEDAR 30 AW 02 and thus they might be used.

For several years, the sample size of queen triggerfish from Puerto Rico is too low to conduct such kinds of analyses. However, this also applies to the estimation of average length used in the Gedamke and Hoenig (2006) method and constitutes a further argument why the assessment team should have combined different approaches to estimate Z and tried a more thorough utilisation of the available data, especially for those years with large sample size of individual length data.

The situation is different for blue tang due to the large uncertainty in the reported Von Bertalanffy (VBF) parameters and, given the fact that the species presents an initial fast growth but a very high longevity, it makes the length data uninformative regarding individual ages after age 5. Therefore, for blue tang, the exploration of the slicing method and the VIT are not feasible and the reviewer agrees with the assessment team concerning the methodology used for assessing this species.

Another method to estimate M is ProdBiom (Abella et al., 1997). The main advantage of this method is that it combines in a single framework the growth parameters, the length weight relationship and information on the longevity of the species. Also, it is able to derive estimates of M by age class, which are very useful in VIT models. It generally gives values of M which are slightly smaller than other methods, thus also avoiding failure to detect overfishing because of unrealistically high values of M . The reviewer considers that Prodbiom might be more appropriate for the estimate of M or it should be used along with the other methodologies presented.

In general, the reviewer considers that the reader is left with the doubt that much more could have been done if there had been a few more assumptions for the queen triggerfish, or at least the assessment team should have explored the possibility of using different methodologies than the Gedamke and Hoenig (2006) method to derive estimates of exploitation rates and F_{MSY} .

Recommendations:

Queen triggerfish catch data from Puerto Rico trap and pots

- Estimate the total number of fish caught by size class for the main gear (i.e. traps and pots), combining the landings information with the size frequency data from the TIP.
- Statistical slicing of the total number of fish landed by size class by the main gear (i.e. traps and pots) to estimate the number of fish landed per age class for years with sufficient length measurements (i.e. for years from 1983 to 1988; a general rule of thumb would be to use years with more than 150 or 200 individuals).
- Estimation of Z from the catch curve or using a pseudocohort analysis (i.e. VIT when only one or few years of data are available).

- Conducting an YPR analysis to estimate F_{01} as a proxy for F_{MSY} to be compared against using M as a proxy for F_{MSY} .

TIP data of queen triggerfish

- Explore the use of effort data from the TIP survey to produce an effort standardized time series of LFD for years with sufficient length measurements (a general rule of thumb would be to use years with more than 150 or 200 individuals).
- Statistical slicing of the total number of fish caught by size class by the main gear (i.e. traps and pots) to estimate the total number of fish per age class.
- Estimation of Z from the catch curve or using a pseudocohort analysis (i.e. VIT) when only one or few years of data is available.

Use the ProdBiom method (Abella et al., 1997) to estimate M along with the other methodologies presented here.

ToR3: Evaluate the assessment findings with respect to the following:

- a) Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?
- b) Is the stock overfished? What information helps you reach this conclusion?
- c) Is the stock undergoing overfishing? What information helps you reach this conclusion?

Generally, a lot of emphasis is given in estimating the uncertainty, which is obviously fine, but with little critical considerations of the likelihood of each of the tested scenarios. This will automatically overestimate the uncertainty and make the evaluation of the stock status even more complicated. This is a more prominent issue for Caribbean queen triggerfish compared to blue tang. As the assessment team correctly pointed out, the key items here are the VBF parameters, which are used to estimate Z , M and F (and F_{MSY}) for both species. It is clear from Table 7 in the assessment report and Table 2 in SEDAR30 AW 03 that the L_{inf} of queen triggerfish estimated by Manooch and Drennon (1987) is generally lower or much lower than L_{max} estimated by other studies in the same area, although no details are given on the number of individuals analysed in these studies. L_{inf} is assumed to range from 37.3 to 45.6, which is in the lower range of the reported L_{max} . The impression I have is that L_{inf} is likely larger than 46.5 and thus the sensitivity analysis should have included also larger L_{inf} and lower k as L_{inf} and k are generally negatively correlated. This has direct consequences on the estimation of M and F , which are likely to be over- and underestimated, respectively. Moreover, L_{inf} and k are negatively correlated, which makes several of the scenarios tested and presented in figure 17, 19 and 21 unrealistic and also inflates the level of uncertainty in the Z estimates. Interestingly, figure 17, 19 and 21 showed that Z estimated for the combination of extreme range of L_{inf} and k are very similar, again corroborating the idea that uncertainty is largely overestimated by the way the sensitivity analysis is set up.

The reviewer is of the opinion that the combination of large L_{inf} and low k are the most plausible set of VBF parameters, given what has been presented in SEDAR30 AW 03 and therefore they should have been given more weight in the evaluation of stock status.

The situation is different for blue tang due to the large uncertainty in the reported VBF parameters, which, together with the fact that the species presents an initial fast growth but a very high longevity, makes the length data uninformative of individual age after age 5. Thus, the reviewer agrees with the assessment team that it is not possible to precisely define the stock status for the Caribbean blue tang and that age-based data are crucial in the future. Stability in mean length is difficult to interpret in this case, and without a robust estimate of the absolute value of Z it cannot be interpreted as an indication of sustainable fishing. Thus, I consider that the stock status is unknown and age data are needed in the future as also pointed out by the assessment team in their general conclusions.

A lot of emphasis has been given to test the effect of L_c on the Z estimates, which was then revealed by the sensitivity analysis to be very small, instead of critically assigning different likelihood to the different scenarios. The authors correctly stress that the estimates are dependent on the parameters but they fail to give critical support to one or fewer scenario over the others to reduce the number of scenarios and help with the evaluation of the stock status.

The impression is that the assessment team is more prone to consider queen triggerfish as being not subject to overexploitation although they correctly stress the fact that the data are not enough to make firm conclusions on the stock's status. However, from Tables 19 and 21, several scenarios indicated that F was in excess of F_{MSY} , which I would interpret as an indication of overfishing being highly likely but this does not emerge from the text of the report. The reviewer also considers $F_{MSY}=M$ as a large oversimplification, which ignores selectivity that has a large impact on F_{MSY} . I would try to estimate catch at age data from LFD and conduct an YPR analysis based on selected yearly data to have an idea of how realistic is this assumption.

For Caribbean blue tang, there is some implication in the report that stability in average length is viewed as an indication of a low level of F . The reviewer disagrees with this view, and considers that given the available information the status of the stock should be considered as unknown.

Recommendations

The reviewer is of the opinion that the combination of large L_{inf} and low k are the most plausible set of VBF parameters, given what has been presented in SEDAR30 AW 03 and therefore they should have been given more weight in the evaluation of the stock's status.

The reviewer considers $F_{MSY}=M$ as a large oversimplification, which ignores selectivity that has a large impact on F_{MSY} . The assessment team should try to estimate catch at age data from LFD and conduct a VIT and an YPR analysis for queen triggerfish based on selected yearly data to have an idea of how realistic is this assumption.

For Caribbean blue tang, the report gives the impression that stability in average length is taken as an indication of low level of F . The reviewer disagrees with this view, and considers that given the available information the status of the stock should be considered as unknown.

ToR 4. Evaluate the stock projections, addressing the following:

- a) Are the methods consistent with accepted practices and available data?
- b) Are the methods appropriate for the assessment model and outputs?
- c) Are the results informative and robust, and useful to support inferences of probable future conditions?
- d) Are key uncertainties acknowledged, discussed, and reflected in the projection results ?.....

The ToR could not be conducted due to data restrictions.

Recommendations

None.

ToR 5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.

- Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods.
- Ensure that the implications of uncertainty in technical conclusions are clearly stated.

See comments under ToR3.

Recommendations

None.

ToR 6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.

- Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
- Provide recommendations on possible ways to improve the SEDAR process.

The assessment team do provide an exhaustive shopping list for future data to be collected, which would greatly improve the capability of assessing the status of the Caribbean queen triggerfish and blue tang stock. However, I also suggest that effort should be devoted to selectivity experiments aimed to evaluate the theoretical changes in selectivity linked with the historical changes in the mesh size of the traps.

Recommendations

Conduct selectivity studies on the effect of the mesh size of the traps on the amount and size distribution of the catches of queen triggerfish and blue tang.

ToR 7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.

See comments under ToR 2 and 3.

Recommendations

None

The key information contained in the introduction of both the assessment for U.S. Caribbean queen triggerfish and the one for blue tang is the management table and the table with the changes in management regulations. The management table should indicate the unit for the value of MSST, MSY and OY, which are now missing. On the other hand, the table with the changes in management regulations is very detailed but without any information about the selectivity of the different mesh size for the traps. Therefore, the reported information is rather uninformative and it is basically impossible to evaluate how these changes might have affected the selectivity of the fisheries. This is crucial information as violating the assumption of time invariant selectivity would affect directly the model output in this case and makes the utilisation of the landings data more complicated. I suggest that effort should be devoted to selectivity experiments aimed to evaluate the theoretical changes in selectivity linked with the historical changes in the mesh size of the traps (see also recommendations in ToR6).

Conclusions

The assessment team should be commended for their effort, timing and clarity in presenting the results. However, I consider that data are underutilised and the uncertainty overestimated by the sensitivity set up used. Also, the lack of alternative estimates of Z beside those coming from the Gedamke and Hoenig (2006) method makes it difficult to evaluate the results and assess the status of the Caribbean queen triggerfish stock. A series of recommendations on how to improve the data utilisation and provide alternative estimates of the exploitation rates have been given under the specific ToRs.

For Caribbean blue tang, the report gives the impression that stability in average length is an indication of a low level of F . The reviewer disagrees with this view, and considers that given the available information the status of the stock should be considered as unknown.

The basic data and model framework were adequately presented through documents and were circulated well in advance of the review. A possible improvement for the presentation of the result in the report could be the creation of a *Glossary* and an *Acronyms* list at the end of the document. This will greatly facilitate the reading of the report for the public.

Reference list

Abella A., Caddy J., Serena F., 1997. Do natural mortality and availability decline with age? An alternative yield paradigm for juvenile fisheries, illustrated by the hake *Merluccius merluccius* in the Mediterranean. *Aquat. Liv. Res.*, 10: 257-269.

Laurence T. Kell and Alexander Kell. A comparison of age slicing and statistical age estimation for mediterranean sword_sh (xiphias gladius). Collect. Vol. Sci. Pap. ICCAT, 66(4):1522{1534, 2011.

Lleonart, J. & Salat, J., 2000. VIT (version 1. 1): Software for fishery analysis. User's manual. On ligne: <http://www.faocopemed.org/es/activ/infodif/vit.htm>.

Scientific, Technical and Economic Committee for Fisheries (STECF) – Assessment of Mediterranean Sea stocks – part 1 (STECF 12-19). (eds. Cardinale M. (Chairman) Osio C. & Charef A.). 2012. Publications Office of the European Union, Luxembourg, EUR 25602 EN, JRC 76735, 502 pp.

Scott F, Osio G, Cardinale M 2011. Comparison of age slicing methods - Working Document in support to the STECF Expert Working Group 11-12 Assessment of Mediterranean Sea stocks - part II. EUR 25054 EN. Luxembourg.

Attachment A: Statement of Work for Dr. Massimiliano Cardinale

External Independent Peer Review by the Center for Independent Experts

SEDAR 30 Caribbean blue tang and queen triggerfish assessment review

Scope of Work and CIE Process: The National Marine Fisheries Service's (NMFS) Office of Science and Technology coordinates and manages a contract providing external expertise through the Center for Independent Experts (CIE) to conduct independent peer reviews of NMFS scientific projects. The Statement of Work (SoW) described herein was established by the NMFS Project Contact and Contracting Officer's Representative (COR), and reviewed by CIE for compliance with their policy for providing independent expertise that can provide impartial and independent peer review without conflicts of interest. CIE reviewers are selected by the CIE Steering Committee and CIE Coordination Team to conduct the independent peer review of NMFS science in compliance the predetermined Terms of Reference (ToRs) of the peer review. Each CIE reviewer is contracted to deliver an independent peer review report to be approved by the CIE Steering Committee and the report is to be formatted with content requirements as specified in **Annex 1**. This SoW describes the work tasks and deliverables of the CIE reviewer for conducting an independent peer review of the following NMFS project. Further information on the CIE process can be obtained from www.ciereviews.org.

Project Description SEDAR 30 will be a compilation of data, an assessment of the stock, and an assessment review conducted for Caribbean blue tang and queen triggerfish. The CIE peer review is ultimately responsible for ensuring that the best possible assessment has been provided through the SEDAR process. The stocks assessed through SEDAR 30 are within the jurisdiction of the Caribbean Fisheries Management Council and the territorial waters of Puerto Rico and the U.S. Virgin Islands. The Terms of Reference (ToRs) of the peer review are attached in **Annex 2**.

Requirements for CIE Reviewers: Three CIE reviewers shall have the necessary qualifications to complete an impartial and independent peer review in accordance with the tasks and ToRs described in the SoW herein. The CIE reviewers shall have expertise in stock assessment, statistics, fisheries science, and marine biology sufficient to complete the tasks of the scientific peer-review described herein. Each CIE reviewer's duties shall not exceed a maximum of 10 days to complete all work tasks of the peer review described herein.

Location of Peer Review: Each CIE reviewer shall conduct the desk review during 4-7 February 2013, therefore no travel will be required.

Statement of Tasks: Each CIE reviewer shall complete the following tasks in accordance with the SoW and Schedule of Milestones and Deliverables herein.

Prior to the Peer Review: Upon completion of the CIE reviewer selection by the CIE Steering Committee, the CIE shall provide the CIE reviewer information (full name, title, affiliation, country, address, email) to the COR, who forwards this information to the NMFS Project Contact no later the date specified in the Schedule of Milestones and Deliverables. The CIE is responsible for providing the SoW and ToRs to the CIE reviewers. The NMFS Project Contact is responsible for providing the CIE reviewers with the background documents, reports, and other information pertinent to the desk review arrangements. Any changes to the SoW or ToRs must be made through the COR prior to the commencement of the peer review.

Pre-review Background Documents: Two weeks before the peer review, the NMFS Project Contact will send (by electronic mail or make available at an FTP site) to the CIE reviewers the necessary background information and reports for the peer review. In the case where the documents need to be mailed, the NMFS Project Contact will consult with the CIE Lead Coordinator on where to send documents. CIE reviewers are responsible only for the pre-review documents that are delivered to the reviewer in accordance to the SoW scheduled deadlines specified herein. The CIE reviewers shall read all documents in preparation for the peer review.

Desk Review: Each CIE reviewer shall conduct the independent peer review in accordance with the SoW and ToRs, and shall not serve in any other role unless specified herein. **Modifications to the SoW and ToRs shall not be made during the peer review, and any SoW or ToRs modifications prior to the peer review shall be approved by the COR and CIE Lead Coordinator.** The CIE Lead Coordinator can contact the Project Contact to confirm any peer review arrangements.

Contract Deliverables - Independent CIE Peer Review Reports: Each CIE reviewer shall complete an independent peer review report in accordance with the SoW. Each CIE reviewer shall complete the independent peer review according to required format and content as described in Annex 1. Each CIE reviewer shall complete the independent peer review addressing each ToR as described in Annex 2.

Specific Tasks for CIE Reviewers: The following chronological list of tasks shall be completed by each CIE reviewer in a timely manner as specified in the **Schedule of Milestones and Deliverables**.

- 1) Conduct necessary pre-review preparations, including the review of background material and reports provided by the NMFS Project Contact in advance of the peer review.
- 2) During February 4-7, 2013 as specified herein, conduct an independent desk peer review in accordance with the ToRs (**Annex 2**).
- 3) No later than February 21, 2013, each CIE reviewer shall submit an independent peer review report addressed to the “Center for Independent Experts,” and sent to Mr. Manoj Shivlani, CIE Lead Coordinator, via email to shivlanim@bellsouth.net, and CIE Regional Coordinator, via email to Dr. David Sampson david.sampson@oregonstate.edu. Each CIE report shall be written using the format and content requirements specified in Annex 1, and address each ToR in **Annex 2**.

Schedule of Milestones and Deliverables: CIE shall complete the tasks and deliverables described in this SoW in accordance with the following schedule.

14 January 2013	CIE sends reviewer contact information to the COR, who then sends this to the NMFS Project Contact
18 January 2013	NMFS Project Contact sends the stock assessment report and background documents to the CIE reviewers.
4-13 February 2013	Each reviewer conducts an independent desk peer review
19 February 2013	CIE reviewers submit draft CIE independent peer review reports to the CIE Lead Coordinator and CIE Regional Coordinator
7 March 2013	CIE submits CIE independent peer review reports to the COR
14 March 2013	The COR distributes the final CIE reports to the NMFS Project Contact and regional Center Director

Modifications to the Statement of Work: This ‘Time and Materials’ task order may require an update or modification due to possible changes to the terms of reference or schedule of milestones resulting from the fishery management decision process of the NOAA Leadership, Fishery Management Council, and Council’s SSC advisory committee. A request to modify this SoW must be approved by the Contracting Officer at least 15 working days prior to making any permanent changes. The Contracting Officer will notify the COR within 10 working days after receipt of all required information of the decision on changes. The COR can approve changes to the milestone dates, list of pre-review documents, and ToRs within the SoW as long as the role and ability of the CIE reviewers to complete the deliverable in accordance with the SoW is not adversely impacted. The SoW and ToRs shall not be changed once the peer review has begun.

Acceptance of Deliverables: Upon review and acceptance of the CIE independent peer review reports by the CIE Lead Coordinator, Regional Coordinator, and Steering Committee, these reports shall be sent to the COR for final approval as contract deliverables based on compliance with the SoW and ToRs. As specified in the Schedule of Milestones and Deliverables, the CIE shall send via e-mail the contract deliverables (CIE independent peer review reports) to the COR (William Michaels, via William.Michaels@noaa.gov).

Applicable Performance Standards: The contract is successfully completed when the COR provides final approval of the contract deliverables. The acceptance of the contract deliverables shall be based on three performance standards:

- (1) The CIE report shall be completed with the format and content in accordance with **Annex 1**,
- (2) The CIE report shall address each ToR as specified in **Annex 2**,
- (3) The CIE reports shall be delivered in a timely manner as specified in the schedule of milestones and deliverables.

Distribution of Approved Deliverables: Upon acceptance by the COR, the CIE Lead Coordinator shall send via e-mail the final CIE reports in *.PDF format to the COR. The COR will distribute the CIE reports to the NMFS Project Contact and Center Director.

Support Personnel:

William Michaels, Program Manager, COR
NMFS Office of Science and Technology
1315 East West Hwy, SSMC3, F/ST4, Silver Spring, MD 20910
William.Michaels@noaa.gov Phone: 301-427-8155

Manoj Shivlani, CIE Lead Coordinator
Northern Taiga Ventures, Inc.
10600 SW 131st Court, Miami, FL 33186
shivlanim@bellsouth.net Phone: 305-383-4229

Roger W. Peretti, Executive Vice President
Northern Taiga Ventures, Inc. (NTVI)
22375 Broderick Drive, Suite 215, Sterling, VA 20166
RPerretti@ntvifederal.com Phone: 571-223-7717

Key Personnel:

NMFS Project Contact:

Julie Neer, SEDAR Coordinator
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405
julie.neer@safmc.net Phone: 843-571-4366

Annex 1: Format and Contents of CIE Independent Peer Review Report

1. The CIE independent report shall be prefaced with an Executive Summary providing a concise summary of the findings and recommendations, and specify whether the science reviewed is the best scientific information available.
2. The main body of the reviewer report shall consist of a Background, Description of the Individual Reviewer's Role in the Review Activities, Summary of Findings for each ToR in which the weaknesses and strengths are described, and Conclusions and Recommendations in accordance with the ToRs. The CIE independent report shall be a stand-alone document for others to understand the weaknesses and strengths of the science reviewed. The CIE independent report shall be an independent peer review of each ToRs.
3. The reviewer report shall include the following appendices:

Appendix 1: Bibliography of materials provided for review

Appendix 2: A copy of the CIE Statement of Work

Annex 2: Terms of Reference for the Peer Review

SEDAR 30 Caribbean blue tang and queen triggerfish assessment review

1. Evaluate the data used in the assessment, addressing the following:
 - a) Are data decisions made by the Assessment Workshop sound and robust?
 - b) Are data uncertainties acknowledged, reported, and within normal or expected levels?
 - c) Are data applied properly within the assessment model?
 - d) Are input data series reliable and sufficient to support the assessment approach and findings?
2. Evaluate the methods used to assess the stock, taking into account the available data.
 - a) Are methods scientifically sound and robust?
 - b) Are assessment models configured properly and used consistent with standard practices?
 - c) Are the methods appropriate for the available data?
3. Evaluate the assessment findings with respect to the following:
 - a) Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?
 - b) Is the stock overfished? What information helps you reach this conclusion?
 - c) Is the stock undergoing overfishing? What information helps you reach this conclusion?
 - d) Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?
 - e) Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?
4. Evaluate the stock projections, addressing the following:
 - a) Are the methods consistent with accepted practices and available data?
 - b) Are the methods appropriate for the assessment model and outputs?
 - c) Are the results informative and robust, and useful to support inferences of probable future conditions?
 - d) Are key uncertainties acknowledged, discussed, and reflected in the projection results?
5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.
 - Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods

- Ensure that the implications of uncertainty in technical conclusions are clearly stated.
6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.
 - Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
 - Provide recommendations on possible ways to improve the SEDAR process.
 7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.

Attachment B: List of main documents provided as background material

Draft Stock Assessment:

Section1_blue tang_v1.pdf

SectionII_S30_Blue_tang_AW_report_complete_w_watermark.pdf

Section1_queen_triggerfish_v1.pdf

SectionII_S30_Queen_triggerfish_AW_report_w_watermark.pdf

Background Materials:

S30 Doc List.pdf

S30_FTP site instructions.pdf

S30_AW_01_SummaryRecreationalBlueTangQueenTriggerfish.pdf.

S30_AW_02_SummaryTIP.pdf

S30_AW_03_Rios_Life History Review.pdf

S30_AW_04_Caribbean queen triggerfish and blue tang landings.pdf

CIE Independent Peer Review Report

on

SEDAR 30

Caribbean blue tang and queen triggerfish assessment review

Prepared by

Dr. Yong Chen

Professor of Fisheries Science
School of Marine Sciences
University of Maine
Orono, ME 04469

March 20, 2013

Contents

SECTION	Page
I. Executive Summary.....	3
II. Background.....	5
III. Description of the Individual Reviewer’s Role in the Review Activities.....	6
IV. Summary of Findings.....	6
IV-1. Queen triggerfish.....	6
IV-2. Blue Tang.....	14
V. Conclusions and Recommendations.....	23
VI. References.....	26
VII. Appendices.....	28
VII-2. List of Documents	28
VII-1. Statement of Work for Dr. Yong Chen.....	29

I. Executive Summary

Queen triggerfish (*Balistes vetula*) and blue tang (*Acanthurus coeruleus*) are reef dwelling and widely distributed in the Atlantic Ocean. They are common in the Caribbean Sea and support two important fisheries in the Puerto Rico and U.S. Virgin Islands. Commercial landing data of the U.S. Caribbean queen triggerfish and blue tang were included in snapper/grouper landings in the 1970s -1990s and then in species groups in the 2000s, and species-specific landing data were only available in recent years. Limited life history parameters such as von Bertalanffy growth parameters are available. Length-composition data are also available for all locations except for the Puerto Rico blue tang. No fishery-independent and fishery-dependent abundance index data were available for the assessment. Because limited data are available for their stock assessment, they are considered data-poor fisheries and a formal stock assessment model is not applicable to these two fish stocks.

An estimator, which uses length-frequency data and requires no assumption of equilibrium population, was used in the assessment for estimating total mortality rate. The natural mortality was estimated from various methods. The fishing mortality was then estimated as the difference between the total and natural mortality rates. No biological reference points such as F_{MSY} and B_{MSY} were estimated. The natural mortality was used as a proxy of F_{MSY} . The estimated fishing mortality was compared with the natural mortality to determine if the fishery was in the status of “overfishing”. Because no biomass could be estimated, it is impossible to determine if the queen triggerfish and blue tang stocks were “overfished” and no stock projection under different management strategies could be done. A comprehensive sensitivity analysis was conducted to evaluate impacts of uncertainty associated with the key life history parameters. The stock assessment suggests that the choice of growth parameters and estimators of natural mortality influenced the determination of status of the U.S. Caribbean queen triggerfish and blue tang. For certain combinations of growth parameters and natural mortality estimators, the fisheries could be defined as experiencing “overfishing”, but for the other combinations, the fisheries were considered not in the status of “overfishing”. Although large uncertainty existed in the status of the fisheries, it appears that these two fish stocks were *less likely* to be in the status of “overfishing”. *Overall, I conclude this assessment is the best the AW panel could do given the restriction of data availability; however I cannot conclude that the assessment is “sound” and “robust” as the assessment quality and results are affected greatly by large uncertainty in the data quality and estimators of natural mortality.*

I have made the following recommendations for improving the assessment: (1) the expert and background knowledge on species of similar life history patterns be used to exclude biologically unrealistic values of K , L_{∞} , natural mortality, and total mortality; (2) uncertainty associated with K , L_{∞} , natural mortality, total mortality, and subsequently fishing mortality be quantified using a Monte Carlo simulation; (3) a program be developed to interview fishermen to collect the information on temporal and spatial variability of the fishing grounds, target species and sizes, and fishing efforts; (4) the information on the species composition of current landings be used to decompose the historical landings of species group into the species-specific landings; (5) a fishery-independent survey program be developed for the queen triggerfish, blue tang and other reef-dependent species sharing similar habitat to collect samples for estimating basic life history parameters and for driving reliable abundance indices; (6) a simulation study be

conducted to evaluate the performance of the length-based estimator and identify factors that are critical in influencing the performance of the estimator; (7) a yield-per-recruit analysis be conducted with the incorporation of uncertainty in life history parameters to estimate biological reference points such as F_{\max} and $F_{0.1}$; and (8) a spawning stock biomass-per-recruit analysis be done with the incorporation of uncertainty associated with life history parameters to estimate reference points such as $F_{20\%}$ and $F_{40\%}$.

II. Background

Queen triggerfish (*Balistes vetula*) is a reef dwelling triggerfish mainly distributed in the Atlantic Ocean. In the West Atlantic, they are distributed from Canada to southern Brazil and are common in the US Caribbean. They are reef-dependent and typically occur at coral and rocky reefs in shallow waters. However, they sometimes also can be found in relatively deep water (up to 275 m) and in areas with sand or seagrass. Adult queen triggerfish are opportunistic feeders. The species is subject to diurnal movement and tends to be either solitary or aggregate in small groups (Randall 1968; Aiken 1975). The maximum length was observed at 572 mm fork length in the U.S. Virgin Islands (Randall 1968). The oldest age recorded in the U.S. Caribbean was 7 years old (Manooch and Drennon 1987).

Blue tang (*Acanthurus coeruleus*), also known as the Atlantic blue tang surgeonfish or the Atlantic blue tang, is a surgeonfish in the Atlantic Ocean. Blue tang is common in the Caribbean Sea and Gulf of Mexico. They inhabit shallow-water, coral reefs and rocky habitat (Carpenter 2002). Adult blue tang are herbivorous, feeding on various benthic algae (Carpenter 2002). The maximum length was observed at 457 mm total length in St. Thomas (Olsen 2011), and the oldest age was found to be 20 years of age (Mutz 2006). Growth parameters estimated in different studies tend to differ greatly as a result of differences in sampling locations, sample sizes, and ranges of age/size composition of sampled fish (Choat and Robertson 2002; Mutz 2006).

Both the fisheries are data-poor with a limited amount of information/data available to the stock assessment. The historical landing data were aggregated by snapper/grouper earlier in the 1970s -1990s, by species groups in the 2000s, and were only separated by species in recent years (after 2011). The recreational data tend to have few trips of positive catch. No reliable fishing effort and no fishery-independent data were available. Size composition data of commercial catch derived from relatively large sample sizes were available for all locations except for the blue tang stock in Puerto Rico (SEDAR30 2013).

Size composition data were used in the stock assessment for estimating the total mortality rate using a length-based estimator developed by Gedamke and Hoenig (2006). This method improves the traditional Beverton-Holt mortality estimator (Beverton and Holt 1957) with no requirement for the assumption of an equilibrium population. The natural mortality was estimated from various methods (Pauly 1980; Hoenig 1983; Jensen 1996). The fishing mortality was then estimated as the difference between the total and natural mortality rates. Because this is data-poor fishery, no biological reference points such as F_{MSY} and B_{MSY} were estimated. The natural mortality was used as a proxy of F_{MSY} (King 1995). Thus, the estimated fishing mortality was compared with the natural mortality to determine if the fishery was in the status of “overfishing”. Because no biomass could be estimated, it is impossible to determine if the queen triggerfish and blue tang stocks are “overfished” and no stock projection can be done to evaluate impacts of various management strategies on the stocks. A comprehensive sensitivity analysis was conducted to evaluate impacts of uncertainty associated with the estimates of key life history parameters. The stock assessment suggests that the choice of growth parameters and estimators of natural mortality influenced the determination of status of the U.S. Caribbean queen triggerfish and blue tang. For certain combinations of growth parameters and natural mortality

estimators, the fisheries could be defined as experiencing “overfishing”, but for the other combinations, the fisheries were considered not in the status of “overfishing” (SEDAR30 2013).

III. Description of the Individual Reviewer’s Role in the Review Activities

As the SoW states that “*Each CIE reviewer shall conduct the independent peer review in accordance with the SoW and ToRs*”, my role as a CIE independent reviewer is to conduct an impartial and independent peer review of SEDAR 30 “Caribbean blue tang and queen triggerfish assessment” with respect to the pre-defined Terms of Reference.

This is a desk review. Thus, I have no opportunity for face-to-face discussion and questioning. I read the “SEDAR30-SAR1: Final Stock Assessment Report for Caribbean Blue Tang” and “SEDAR30-SAR2: Final Stock Assessment Report for Caribbean Queen Triggerfish” and all other background documents that were sent to me (see the list in the Appendix I). I also read references relevant to the topics covered in the reports and the SoW. I addressed each topic covered in the ToRs, evaluated the strengths and weaknesses of what was done in this assessment, and provided recommendations to improve future assessment. Based on these evaluations and analyses, I made research recommendations for future assessment of Caribbean blue tang and queen triggerfish.

IV: Summary of Findings

IV-1. Queen triggerfish

The following summary of my findings is provided with respect to a set of pre-defined TORs for the U.S. Caribbean queen triggerfish.

IV-1-1. Evaluate the data used in the assessment, addressing the following:

Data available to the assessment include commercial landings, recreational intercept data and length frequency data estimated separately for St. Thomas/St. John, St. Croix, and Puerto Rico.

For Puerto Rico, the commercial landing data of queen triggerfish by gear and fishing center were estimated from self-reported fisher logbooks/sale receipts for the time period from 1983 to 2011. Because the report was incomplete, the total landings were adjusted (SEDAR30, 2013). The number of trips with reported queen triggerfish landings was also estimated by gear and year.

For St. Thomas and St. John, although logbook reporting started in 1974, landings were reported by gear and by either snapper/grouper or other finfish prior to 1997. Landings were reported by species group and gear from 1997 to 1999, all commercial landings were reported by species group from 2000 to 2010, and the landing data have been reported by species since 2011. For the years included in this assessment, queen triggerfish-specific landing data are not available prior to 2011, and landing data are only available for the triggerfish species group.

For St. Croix, similar to St. Thomas and St. John, landing data were only available for the triggerfish species group, not for queen triggerfish. Commercial landing data were only available from 1998 to 2011.

Marine Recreational Fisheries Sampling Survey (MRFSS) collects data from Puerto Rico, but not the US Virgin Islands. The sampling design consists of two complementary components, an angler-site intercept survey for estimating catch and length frequency data and a fishing effort telephone survey to estimate fishing effort. However, the positive intercepted trips (i.e., presence of queen triggerfish) were less than 1% in almost all the years. Only 60 queen triggerfish were measured from 2000-2011.

The VBGF parameters were estimated in two studies, but K differs greatly (Manooch and Drennon 1987; de Albuquerque et al. 2011). The maximum age in the sample is 7 (Manooch and Drennon 1987) and 14 years of age (de Albuquerque et al. 2011), and the growth curves in neither study reached the asymptotic lengths. The VBGF parameters estimated in Manooch and Drennon (1987) were used because the samples were taken in Puerto Rico and the US Virgin Islands.

Length frequency data were estimated from samples taken in pot and trap fisheries in Puerto Rico, St. Thomas and St. John, and St. Croix. However, the temporal changes in length frequency data might be influenced by changes in market demand for large sizes of queen triggerfish and/or possible expansion of the fishery into new areas.

IV-1-1-a. Are data decisions made by the Assessment Workshop sound and robust?

The AW panel decided to use the von Bertalanffy growth parameters estimated in Manooch and Drennon (1987) because the samples were taken in Puerto Rico and the US Virgin Islands. The AW panel also decided to use length-frequency data estimated in the pot and trap fisheries for estimating the total mortality using a length-based mortality estimator. The AW decided not to use the intercepted catch, effort, and length data from MRFSS because of the low proportion of positive trips and sample sizes of length data.

Based on the data available, these decisions are the best the AW panel could make. However, based on the information available I do not have evidence to conclude if the data decisions are “sound and robust”.

IV-1-1-b. Are data uncertainties acknowledged, reported, and within normal or expected levels?

The AW panel did acknowledge potential issues which might influence the quality of the data. The MRFSS data were excluded because of small sample sizes, and good discussions were made on potential causes resulting in large shifts in length compositions over time. However, I do not see distributional quantification of uncertainty associated with the estimates of the von Bertalanffy growth parameters. I think the estimation of these parameters should come with estimates of uncertainty (e.g., bootstrap-estimated confidence intervals). Given there are only 7 age groups available in Manooch and Drennon (1987), the uncertainty associated with the estimated L_{∞} and K could be large. It is also unclear if the variation in size within an age group was considered and if the fitting of the VBGF was weighted by the sample sizes of the different age groups.

IV-1-1-c. Are data applied properly within the assessment model?

This is a data-poor fishery, and there is not enough information for a formal stock assessment. Given the data available, I consider the data are properly applied in the estimation of total, natural and fishing mortality rates in the assessment.

However, I believe the sensitivity analysis could be better designed and justified if the uncertainty associated with K and L_{∞} could be better quantified and if expert and background knowledge could be used to exclude biologically unrealistic combinations of values for the growth parameters and natural mortality.

IV-1-1-d. Are input data series reliable and sufficient to support the assessment approach and findings?

Estimation of the length-frequency data might be influenced by temporal variability in the shifted preference of market demand for large queen triggerfish, which might change the selectivity of the fishery over the time. Thus, the temporal variability in length composition might not reflect changes in fish mortality; rather reflect changes in selectivity. Although potential changes in the total mortality could be incorporated in Gedamke and Hoenig (2006), I did not see how possible changes in selectivity could be incorporated. This may yield biases in the assessment results.

IV-1-2. Evaluate the methods used to assess the stock, taking into account the available data.

Given the limitation of data availability, the Beverton-Holt length-based mortality estimator (Beverton and Holt 1957) is a good option for the assessment of this fishery. However, as the AW panel explicitly described in the assessment report, this method, explicitly and implicitly, requires six assumptions: (1) growth is constant over time and space; (2) there is no variability in growth among individuals; (3) there is constant and continuous recruitment over time; (4) the mortality rate is the same for fish older than the age at recruitment; (5) the mortality rate is constant over time and space; and (6) the population is in equilibrium. Apparently none of these assumptions can be satisfied in the U.S. Caribbean queen triggerfish fishery. Instead of using this traditional approach, the AW panel used a modified Beverton-Holt length-based mortality estimator which requires no assumption of an equilibrium population. However, the other assumptions are still required. There were a number of years when mortality rate changes were identified using the Akaike Information Criterion (AIC) as the model selection measure. A systematic sensitivity analysis was done to evaluate impacts of uncertainty in the growth parameters on the estimation of fish mortality rates.

IV-1-2-a. Are methods scientifically sound and robust?

Given the limitation of data availability, this approach may be the best choice the AW panel can have for the assessment of the U.S. Caribbean queen triggerfish. However, based on what has been reported in the AW report, I cannot conclude that this is scientifically sound and robust because it is difficult to evaluate whether this approach can capture the real fishing mortality rate without knowing the true value. A simulation study, similar to the one in

Gedamke and Hoenig (2006) but based on the queen triggerfish data, should be conducted to evaluate the performance and robustness of this mortality estimator for the queen triggerfish with respect to different assumptions associated with the fishery.

IV-1-2-b. Are assessment models configured properly and used consistent with standard practices?

Given what is available, I believe that the configuration of the assessment models is consistent with standard practices. However, the uncertainty in growth parameters was not estimated and the sensitivity analysis could be better designed if the uncertainty associated with K and L_{∞} was explicitly estimated. The values of K and L_{∞} are usually strongly and negatively correlated, and such negative correlations should be considered in the sensitivity analysis. The sensitivity analysis should focus on one parameter (either K or L_{∞}) with the value of the other parameter drawn from a joint probability distribution with a defined covariance structure for K and L_{∞} . A bootstrap approach can be used to define the joint probability distribution of K and L_{∞} .

The AW panel considered different approaches for estimating natural mortality, and recommended that the M estimated using the Pauly's equation (Pauly 1980) be used because the growth parameters were also used in the estimation. I agree with the AW panel and believe this perhaps is the most robust approach to reduce potential biases in the estimated fishing mortality rate (because $F = Z - M$). However, I think a more appropriate approach for estimating M may be the use of a subset of fish species with similar habitat and life history characteristics (e.g., reef-associated species) to modify Pauly's equation to make the estimation of M more consistent with the life history and habitat characteristics of the Caribbean queen triggerfish.

The use of M as a proxy for F_{MSY} is a common practice for a data-poor fishery (King 1995). This *ad hoc* limit reference point appears to be the best choice given the available data.

IV-1-2-c. Are the methods appropriate for the available data?

Overall, I believe that the method is appropriate for the available data. However, I believe a simulation study should be conducted to evaluate the performance of the method.

IV-1-3. Evaluate the assessment findings with respect to the following:

IV-1-3-a. Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?

Because of data limitations, neither abundance nor biomass was estimated. Fishing mortality was derived from the difference between the total mortality estimated from length-composition data and natural mortality estimated from Pauly's model (Pauly 1980). Uncertainty associated with the fishing mortality was evaluated by considering possible ranges of the total mortality estimated using different growth parameters and natural mortality estimated using different estimators.

I believe that the lack of abundance/biomass estimates is consistent with the limitation of data availability. Large uncertainty associated with fishing mortality estimates is consistent with

possible issues related to the estimates of life history parameters used in estimating the total and natural mortality rates.

IV-1-3-b. Is the stock overfished? What information helps you reach this conclusion?

The stock biomass/abundance and biomass-based limit reference points cannot be estimated reliably in this assessment based on the data available. The AW panel concluded that this assessment did not have enough information to determine if the stock was overfished. Given the available data, I agree with the AW panel.

IV-1-3-c. Is the stock undergoing overfishing? What information helps you reach this conclusion?

Puerto Rico

Large uncertainty is associated with the estimation of fishing mortality as a result of the varying choices of estimators for estimating natural mortality and different growth parameters used in estimating the total mortality. However, for most scenarios tested, the estimated fishing mortality tended to be much lower than natural mortality, suggesting that the queen triggerfish experienced low fishing mortality. If the natural mortality is used as a limit reference point in determining if the fishery is in the status of overfishing, we may conclude that the Puerto Rico queen triggerfish fishery is not in the status of overfishing. The analysis of length composition data from the pot and trap fishery shows that fishing mortality has a declining trend in the late 1990s; however large uncertainty as a result of lack of understanding of possible temporal changes in selectivity and fishing grounds complicates the interpretation of this result.

St. Thomas and St. John

The results of comparing fishing mortality and natural mortality depend on the choices of (1) growth parameters used in the estimation of the total mortality; (2) estimators of natural mortality; and (3) maximum age (i.e., 7 in Manooch and Drennon (1987) or 14 in de Albuquerque et al. 2011). Given such large uncertainty and lack of strong evidence to justify the use of one set of life history parameters over the other, it is difficult to conclude if the fishery is in the status of overfishing.

St. Croix

The results of comparing fishing mortality and natural mortality depend on the choice of growth parameters, which determine the estimates of the total mortality. For the set of life history parameters resulting in a high level for the total mortality estimate, the fishing mortality is higher than natural mortality, suggesting that fishing mortality may be too high. However, for the set of life history parameters resulting in a low level for the total mortality estimate, the estimated fishing mortality is lower than natural mortality, suggesting that the fishing mortality is not too high. We do not have strong evidence favoring one set of the life history parameters over the other, and hence it is difficult to decide if the fishery is in the status of overfishing.

IV-1-3-d. Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?

No stock-recruitment relationship can be developed based on the available data.

IV-1-3-e. Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?

Like many data-poor fisheries, natural mortality was used as a proxy for F_{MSY} in the assessment (King 1995). This effectively treats natural mortality as a limit biological reference point to determine if the fishery is in the status of overfishing. The AW panel explored and evaluated different methods in quantifying the natural mortality and found large uncertainty associated with the natural mortality estimates. Given the information available, I believe that yield-per-recruit (and maybe egg-per-recruit) analysis can be conducted, which can produce estimates of $F_{0.1}$ and F_{max} . The AW panel did mention that they did not do per-recruit analysis because of concerns on the quality of life history parameters. However, given the same life history parameters used in estimating the total fishing mortality and natural mortality (for some methods), I do not see the logic here for not doing a per-recruit analysis. I think the uncertainty associated with life history parameters can be readily incorporated in a per-recruit analysis using a Monte Carlo approach (e.g., Chen and Wilson 2002; Chang et al. 2009).

IV-1-4. Evaluate the stock projections, addressing the following:

No formal stock projection was done in the assessment because of data limitations.

IV-1-4-a. Are the methods consistent with accepted practices and available data?

Stock projections were not done in the assessment because of lack of the information on the dynamics of the fish population.

IV-1-4-b. Are the methods appropriate for the assessment model and outputs?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-1-4-c. Are the results informative and robust, and useful to support inferences of probable future conditions?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-1-4-d. Are key uncertainties acknowledged, discussed, and reflected in the projection results?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-1-5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.

- Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods
- Ensure that the implications of uncertainty in technical conclusions are clearly stated.

The AW panel outlined several sources of uncertainty in the assessment. The uncertainty associated with the quality and quantity of fisheries data (e.g., commercial and recreational catch and size composition data, fishing efforts, and sample sizes) is well discussed to determine which data sets should be used in the assessment. Large variabilities on growth parameters among different studies were identified and their impacts on the estimation of total mortality and fishing mortality were evaluated in a sensitivity analysis. Uncertainty resulting from choices of estimators for natural mortality also was discussed.

Although the AW panel discussed the uncertainty of different sources rather thoroughly and developed sensitivity analyses to evaluate impacts of the uncertainty on the estimation of the total, natural and fishing mortality rates, I believe the uncertainty should be incorporated in the assessment in a more systematic way. I suggest using a Monte Carlo simulation approach to systematically incorporate the uncertainty in life history parameters into the estimation of the fishing mortality rate. For each parameter, a distribution (uniform, multinomial, normal, or log-normal) can be defined based on the type of the data and possible ranges of the values. For each run, the value of a given parameter can be randomly drawn from such a distribution. The correlations between L_{∞} and K should be considered and their values should be drawn from a joint distribution of these two values. One hundred or more runs of Monte Carlo simulation can yield a distribution for the total, natural and fishing mortality rates. Such an approach can better capture and quantify the uncertainty, which can be used directly in comparing probability distributions of natural mortality and fishing mortality to determine the likelihood of overfishing. Before this can be done, however, the range of the growth parameters and natural mortality should be narrowed down based on the expert knowledge and background information on fish species of similar life history and habitat needs.

IV-1- 6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.

- Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
- Provide recommendations on possible ways to improve the SEDAR process.

The AW panel recommends improving the quality of life history parameter estimates; developing a fishery-independent monitoring program; continuing the efforts to improve the collection of species-specific catch and effort data; and modifying the length-based total mortality estimator to account for potential changes in selectivity. I consider these research areas

are important for reducing the uncertainty and improving the quality of the assessment. The AW panel probably needs to prioritize the research recommendations and separate the short-term research plan from the long-term plan.

Given the problems associated with the data, an important research goal should be to improve the data quality and quantity. Short-term and long-term plans should be developed to achieve the goal. Short term research priority may include (1) improvement of life history data estimates and the quantification of their uncertainty in the form of probably distributions; (2) identification of major fishing areas and their spatio-temporal variability via conducting interviews with fishermen involved in the fishery; and (3) identification of potential approaches that can be used to estimate species-specific landing data (e.g., based on species composition of landings that become available in recent years). The long-term research plan should include the development of a fishery-independent monitoring program and continued improvement of the sampling protocol for the collection of fishery-dependent data (catch and effort).

Given the data limitations, I believe another research priority that should be addressed soon is to evaluate the performance of the length-based estimator (Gedamke and Hoenig 2006) for the total mortality. Based on the information available and with some assumptions, a queen triggerfish fishery can be simulated, following the approach used in Gedamke and Hoenig (2006). A simulation study can be conducted with this simulated fishery to evaluate the performance of this length-based estimator for estimating the total mortality. Different scenarios can be developed to identify key factors that may have significant impacts on the performance of the estimator. This can guide the future model development and data collection.

IV-1-7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.

I recommend the following key areas for the improvement when scheduling the next assessment:

- Growth parameters K and L_{∞} should be estimated with uncertainty. A bootstrap approach can be used with the von Bertalanffy growth model to quantify the joint probability distribution of K and L_{∞} , which can be used for quantifying probability distributions for the total, natural and fishing mortality rates;
- More basic biological studies need to be conducted to improve our understanding of key life history processes and estimate key life history parameters such as growth parameters, length/age at maturity, fecundity, and their spatial variability;
- An interviewing-fishermen program should be done to identify major fishing grounds and main size classes of landed catch, and possible changes over time;
- Use the proportion of queen triggerfish in the total catch of all triggerfish species estimated in recent years to estimate the queen triggerfish catch in the past (assuming that the proportion is the same over time);

- A simulation study needs to be conducted to evaluate the performance of the length-based estimator of the total mortality rate and identify assumptions/parameters that can influence greatly the performance of the estimator, which will help us understand the quality of the estimates of the total mortality;
- Uncertainty associated with the natural mortality rate should be quantified in the form of a probability distribution, which can be done with a Monte Carlo simulation approach;
- The Pauly natural mortality estimator was derived from many species with very different life history and habitat needs (Pauly 1983), and a subset of fish species that have life history and habitat needs similar to the focal species may yield a more appropriate natural mortality estimator;
- A yield-per-recruit analysis with the consideration of uncertainty associated with life history parameters (e.g., Chen 1996; Chang et al. 2009) can be done to estimate theoretical biological reference points such as F_{max} and $F_{0.1}$; and
- A spawning stock biomass-per-recruit analysis with the incorporation of uncertainty associated with life history parameters can also be done to estimate reference points such as $F_{20\%}$ and $F_{40\%}$.

IV-2. Blue tang

The following summary of my findings is provided with respect to the set of pre-defined TORs for blue tang.

IV-2-1. Evaluate the data used in the assessment, addressing the following:

Data available to the assessment included commercial landings, recreational intercept data and length-frequency data estimated separately for St. Thomas/St. John, St. Croix, and Puerto Rico. Life history data obtained from published studies were also used in the assessment.

For Puerto Rico, the commercial landing data of blue tang were included in the reported catch of the species group surgeonfishes by gear and fishing center, and the proportion of blue tang within the surgeonfishes species group was unknown. Hence, no separate landing data are available for blue tang. The landing data were reported by gear and fishing center and estimated from self-reported fisher logbooks/sale receipts for the time period from 1983 to 2011. Because the report was incomplete, the total landings were adjusted (Caribbean Fisheries Data Evaluation Final Report, 2009). Length composition data were derived from small sample sizes.

For St. Thomas and St. John, although logbook reporting started in 1974, landings were reported by gear and by either snapper/grouper or other finfish prior to 1997. Some landings were reported by species group and gear from 1997 to 1999, and all reported commercial landings was reported by species group from 2000 to 2010, and the landing data have been reported by species since 2011. For the years included in this assessment, landing data were provided as surgeonfishes with all the species combined, and blue tang-specific landing data are not available.

For St. Croix, similar to St. Thomas and St. John, landing data were only available for surgeonfishes, and no blue tang-specific landing data were available. Commercial landing data were only available from 1998 to 2011.

Marine Recreational Fisheries Sampling Survey (MRFSS) collects data from Puerto Rico, but not the US Virgin Islands. The sampling design consists of two complementary components, an angler-site intercept survey for estimating catch and length frequency data and a fishing effort telephone survey to estimate fishing effort. However, the positive intercepted trips (i.e., presence of blue tang) are too small. The AW panel concluded that this data set was not useful for the blue tang assessment.

The VBGF parameters were estimated for different locations in the U.S. Caribbean in two studies (Choat and Robertson 2002; Mutz 2006). Large differences were found in the estimates between the studies. A sensitivity analysis was done to evaluate the range of possible values for K and L_{∞} .

Length frequency data were estimated from samples taken in the NMFS Trip Interview program for the pot and trap fisheries in Puerto Rico, St. Thomas and St. John, and St. Croix. The number of blue tang measured in Puerto Rico was small compared to St. Thomas/St John or St. Croix.

IV-2-1-a. Are data decisions made by the Assessment Workshop sound and robust?

The AW panel concluded that the sample sizes for the length-frequency data in the pot and trap fisheries in the US Virgin Islands were sufficient for length-based mortality estimation. The growth parameters used in the initial analysis were from Mutz (2006). A sensitivity analysis was conducted to evaluate alternative values and their impacts on the estimation of the total fish mortality. The AW panel decided not to use the intercepted catch, effort, and length data from MRFSS because of the low proportion of positive trips and sample sizes of length data.

The AW panel considered that the sample size for estimating length-composition data was not sufficient in Puerto Rico, and derived length-composition data were not appropriate for length-based mortality estimator.

Based on the data available and limited choices the AW panel had, these decisions were the best one could make. However, because there is no scientific evidence showing the results are robust regarding these decisions, I cannot conclude that the data decisions are “sound and robust”.

IV-2-1-b. Are data uncertainties acknowledged, reported, and within normal or expected levels?

The AW did acknowledge potential issues which might influence the quality and quantity of the data. The MRFSS data were excluded because of small sample sizes, and good discussions were made on potential causes resulting in changes in length compositions over time. However, I do not see quantification of uncertainty associated with the estimates of von Bertalanffy growth parameters (although the differences in the parameters estimated in different studies were shown). I think the estimation of these parameters should come with estimates of uncertainty, which could be derived using an approach such as a bootstrap method.

IV-2-1-c. Are data applied properly within the assessment model?

This is a data-poor fishery, and there is not enough information for a formal stock assessment. Given the data available and limitation of stock assessment model choices, I consider the data are properly applied in the estimation of the fish mortality in this stock assessment.

However, I believe the sensitivity analysis could be better designed and justified if the uncertainty associated with K and L_{∞} could be estimated and quantified. The values of K and L_{∞} in the sensitivity analysis should be drawn from their joint distribution (Chen 1996; Chang et al. 2009) rather than varied independently. The correlation between K and L_{∞} and standard errors associated with L and L_{∞} can be estimated in the Nonlinear Least Squares or their joint probability distribution could be derived using the bootstrap approach.

IV-2-1-d. Are input data series reliable and sufficient to support the assessment approach and findings?

Estimation of the length-frequency data may be influenced by spatio-temporal variability in fishing selectivity. Thus, the temporal variability in length composition may not reflect changes in fish mortality; but rather reflect changes in selectivity and fishing locations. Although potential changes in the total mortality rate can be incorporated in Gedamke and Hoenig (2006), I do not see how changes in selectivity can be incorporated. This may yield biases in the assessment.

IV-2-2. Evaluate the methods used to assess the stock, taking into account the available data.

Given the limitations of data availability, the Beverton-Holt length-based mortality estimator (Beverton and Holt 1957) is a good option for the assessment of this fishery. However, as the AW panel explicitly described in the assessment report, this method, explicitly and implicitly, has six assumptions: (1) growth is constant over time and space; (2) there is no variability in growth among individuals; (3) there is constant and continuous recruitment over time; (4) the mortality rate is the same for fish older than the age at recruitment; (5) the mortality rate is constant over time and space; and (6) the population is in equilibrium. Apparently none of these assumptions can be satisfied in the US Caribbean queen triggerfish fishery. Instead of using this traditional approach, the AW panel used a modified length-based mortality estimator. This method does not need to make the equilibrium assumption, but still needs the other five assumptions. The number of years when mortality rate changes was estimated using the Akaike Information Criterion (AIC) as the performance measure. A systematic sensitivity analysis was done to evaluate impacts of uncertainty in growth parameters on the estimation of fish mortality rates.

IV-2-2-a. Are methods scientifically sound and robust?

Given the limitation of data availability, this approach may be the best the AW panel can have for the assessment of the US Caribbean blue tang. However, based on what has been reported in the AW report, I cannot conclude that this is scientifically sound and robust because there is no evidence showing that this approach can capture the real rate of fishing mortality. A simulation study, similar to the one in Gedamke and Hoenig (2006) but based on the blue tang data, should be conducted to evaluate the performance and robustness of this mortality estimator for blue tang with respect to different assumptions associated with the fishery.

IV-2-2-b. Are assessment models configured properly and used consistent with standard practices?

Given what is available, I believe that the configuration of the assessment models is consistent with standard practices to estimate mortality rates and to evaluate impacts of uncertainty in growth parameters on the mortality estimation. However, the uncertainty in the growth parameters was not estimated and the sensitivity analysis could be better designed. The values of K and L_{∞} are usually strongly and negatively correlated, and such negative correlations should be considered in the sensitivity analysis. The sensitivity analysis should focus one parameter (either K or L_{∞}) with the value of the other parameter drawn from a joint probability distribution with defined correlations of K and L_{∞} . Alternatively, a bootstrap approach can be used to estimate a joint distribution of K and L_{∞} , which can be used to quantify the uncertainty associated with the estimates of the total and natural mortality rates.

The AW panel considered different approaches for estimating natural mortality, and recommended that the M estimated from Pauly equation (Pauly 1980) be used because the growth parameters were also used in the estimation. I agree with the AW panel and believe this perhaps is the most robust approach to reduce potential biases in estimating fishing mortality (because $F = Z - M$). However, I think a more appropriate approach for estimating M may be the use of a subset of fish species with similar habitat and life history (e.g., reef-associated species) to modify the Pauly equation, which can make the estimation of M more consistent with the life history and habitat of the Caribbean blue tang. Biologically unrealistic estimates of natural mortality, judged based on life history theory and knowledge on species of similar life history and habitat need, should be excluded from further consideration in the estimation of fishing mortality.

The use of M as a proxy for F_{MSY} is a common practice for a data-poor fishery (King 1995). This *ad hoc* limit reference point appears to be the best choice given the available data.

IV-2-2-c. Are the methods appropriate for the available data?

Overall, I believe the method is appropriate for the available data. However, I believe a simulation study should be conducted to evaluate the performance of the method.

IV-2-3. Evaluate the assessment findings with respect to the following:

IV-2-3-a. Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?

Because of data limitations, neither abundance nor biomass was estimated. Fishing mortality was derived from the difference between the total mortality rate estimated from length-composition data and natural mortality rate estimated from Pauly's model (Pauly 1980). Uncertainty associated with the fishing mortality rate was evaluated by evaluating possible ranges of the total mortality rate estimated using different growth parameters and natural mortality estimated using different estimators.

I consider the lack of abundance/biomass estimates is consistent with the limitations of data availability. Large uncertainty associated with the fishing mortality estimates is consistent with possible issues related to the estimates of life history parameters used in the estimation of the total and natural mortality rates.

The AW panel did not estimate the total mortality rate for the Puerto Rico blue tang from the length-frequency data because they believed that the sample size was too small. I agree with the AW panel and consider this is consistent with the data available.

IV-2-3-b. Is the stock overfished? What information helps you reach this conclusion?

The stock biomass/abundance and biomass-based limit reference points cannot be estimated reliably in this assessment based on the data available. The AW panel concluded that this assessment did not have enough information to determine if the stock is overfished. Given the available data, I agree with the AW panel.

IV-2-3-c. Is the stock undergoing overfishing? What information helps you reach this conclusion?

Puerto Rico

No length-based analysis was done because the AW panel concluded that the sample size used to derive the length-composition data was too small. Thus, there were no estimates of the total, natural and fishing mortality rates for the Puerto Rico blue tang.

St. Thomas and St. John

The results of comparing the fishing mortality and natural mortality rates depend on the choices of (1) growth parameters used in the estimation of the total mortality; and (2) estimators of natural mortality. The AP panel suggested that Pauly's natural mortality estimator be used because both K and L_{∞} were used, which is consistent with what is used in estimating the total mortality. Based on this approach, the fishing mortality rate, estimated as the difference between the total mortality rate estimated from the length-based estimator (Gedamke and Hoenig 2006) and natural mortality rate estimated using Pauly's equation (Pauly 1980), was much smaller than the natural mortality rate, which is commonly used as limit reference point to determining if a data-poor fishery is in the status of overfishing. This suggests that the fishery was not in the status of overfishing. However, if natural mortality was estimated from age-based data, the results would depend on the choice of growth parameters in estimating the total mortality. For the most scenarios tested in the sensitivity analysis, it appears that the fishing mortality rate was lower than the natural mortality rate, suggesting that the St. Thomas and St. John blue tang were

likely not in the status of overfishing. However, given such large uncertainty and the lack of strong evidence to justify the use of one set of life history parameters over the other, it is difficult to yield a conclusive result regarding the status of the fishery.

St. Croix

Like the assessment for the St. Thomas and St. John blue tang, the results of comparing fishing mortality and natural mortality depend on the choice of growth parameters, which determine the estimates of the total and natural mortality rates. For the set of life history parameters resulting in a high level for the total mortality rate, the fishing mortality rate is higher than the natural mortality rate, suggesting that fishing mortality may be too high. However, for the set of life history parameters resulting in a low level for the total mortality rate, the estimated fishing mortality was lower than natural mortality, suggesting that the fishing mortality was not too high. We do not have strong evidence favoring one set of the life history parameters over the other, and hence it is difficult to decide if the fishery is in the status of overfishing.

IV-2-3-d. Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?

No stock-recruitment relationship can be developed based on the available data.

IV-2-3-e. Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?

Like many data-poor fisheries, natural mortality is used as a proxy for F_{MSY} in the assessment (King 1995). This effectively treats the natural mortality rate as a limit biological reference point to determine if the fishery is in the status of overfishing. The AW panel explored and evaluated different methods for quantifying the natural mortality rate and found large uncertainty associated with the natural mortality rate estimates. Given the information available, I believe that a yield-per-recruit (and maybe egg-per-recruit) analysis can be conducted, which can yield estimates for $F_{0.1}$ and F_{max} . The AW panel did mention that they did not do a per-recruit analysis because of concerns on the quality of life history parameters. However, given the same life history parameters used in estimating the total fishing mortality and natural mortality (for some methods), I do not see the logic for not doing a per-recruit analysis. I think the uncertainty associated with the life history parameters can be readily incorporated in a per-recruit analysis using a Monte Carlo approach (e.g., Chen and Wilson 2002; Chang et al. 2009).

IV-2-4. Evaluate the stock projections, addressing the following:

No formal stock projection was done in the assessment because of data limitations.

IV-2-4-a. Are the methods consistent with accepted practices and available data?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-2-4-b. Are the methods appropriate for the assessment model and outputs?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-2-4-c. Are the results informative and robust, and useful to support inferences of probable future conditions?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-2-4-d. Are key uncertainties acknowledged, discussed, and reflected in the projection results?

Stock projections were not done in the assessment because of lack of information on the dynamics of the fish population.

IV-2-5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.

- Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods
- Ensure that the implications of uncertainty in technical conclusions are clearly stated.

The AW panel outlined several sources of uncertainty in the assessment. The uncertainty associated with the quality and quantity of fisheries data (e.g., commercial and recreational catch and size composition data, fishing efforts, and sample sizes) is well discussed to determine which data set should be used in the assessment. Large variability in growth parameters among different studies was identified and their impact on the estimation of the total mortality rate and fishing mortality rate were evaluated in a sensitivity analysis. Uncertainty resulting from choices of estimators for the natural mortality rate also was discussed.

Although the AW panel discussed the uncertainty of different sources rather thoroughly and developed sensitivity analyses to evaluate impacts of the uncertainty on the estimation of the total, natural and fishing mortality rates, I believe the uncertainty should be incorporated in the assessment in a more systematic way. I suggest using a Monte Carlo simulation approach to systematically incorporate the uncertainty in life history parameters into the estimation of the fishing mortality rate. For each parameter, a distribution (uniform, multinomial, normal, or log-normal) can be defined based on the type of the data and possible ranges of the values. For each run, the value of a given parameter can be randomly drawn from such a distribution. The correlations between L_{∞} and K should be considered and their values should be drawn from a joint distribution of these two values. One hundred or more runs of Monte Carlo simulation can yield a distribution for the total, natural and fishing mortality rates. Such an approach can better capture and quantify the uncertainty, which can be used directly in comparing probability

distributions of the natural mortality rate and fishing mortality rate to determine the likelihood of overfishing. Before this can be done, however, the range of the growth parameters and natural mortality rate should be narrowed down based on expert knowledge and background information on fish species of similar life history and habitat needs.

IV-2- 6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.

- Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
- Provide recommendations on possible ways to improve the SEDAR process.

The AW panel recommended improving the quality of life history parameter estimates; developing a fishery-independent monitoring program; continuing efforts to improve the collection of species-specific catch and effort data; and modifying the length-based total mortality rate estimator to account for potential changes in selectivity. I consider these research areas are important for reducing the uncertainty and improving the quality of the assessment. The AW panel probably needs to prioritize the research recommendations and separate the short-term research plan from the long-term plan.

Given the problems associated with the data, an important goal should be to improve the data quality and quantity. Short-term and long-term plans should be developed to achieve the goal. The short term research priority may include (1) improving life history data estimates and the quantification of their uncertainty in the form of probably distributions; (2) identifying major fishing areas and how the fishing areas vary with time via conducting interviews with fishermen involved in the fishery; and (3) identifying potential approaches that can be used to estimate species-specific landing data (e.g., based on species composition of landings that become available in recent years). The long-term research plan should include the development of fishery-independent monitoring program and continue improving the sampling protocol in the collection of fishery-dependent data (catch and effort).

Given the data limitations, I believe another research priority is to evaluate the performance of the length-based estimator (Gedamke and Hoenig 2006) for the total mortality. Based on the information available and with some assumptions, a queen triggerfish fishery can be simulated, following the approach used in Gedamke and Hoenig (2006). A simulation can be conducted with this simulated fishery to evaluate the performance of this length-based estimator in estimating the total mortality rate. Different scenarios can be developed to identify key factors that may have significant impacts on the performance of the estimator. This can guide the future model development and data collection.

IV-2-7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.

I recommend the following key areas for the improvement when scheduling the next assessment:

- Growth parameters K and L_{∞} should be estimated with uncertainty. A bootstrap approach can be used with the von Bertalanffy growth model to quantify the joint probability distribution of K and L_{∞} , which can be used for quantifying probability distributions for the total, natural and fishing mortality rates;
- More basic biological studies need to be conducted to improve our understanding of key life history processes and estimate key life history parameters such as growth parameters, length/age at maturity, fecundity, and their spatial variability;
- An interviewing-fishermen program should be done to identify major fishing grounds and main size classes of landed catch, and possible changes over time;
- Use the proportion of blue tang in the total catch of all surgeonfish species estimated in recent years to estimate the queen triggerfish catch in the past (assuming that the proportion is the same over time);
- A simulation study needs to be conducted to evaluate the performance of the length-based estimator of the total mortality rate and identify assumptions/parameters that can influence greatly the performance of the estimator, which will help us understand the quality of the estimates of the total mortality rate;
- Uncertainty associated with the natural mortality rate should be quantified in the form of a probability distribution, which can be done with a Monte Carlo simulation approach;
- The Pauly natural mortality rate estimator was derived from many species with very different life history and habitat needs (Pauly 1980), and a subset of fish species that have life history and habitat needs similar to the focal species may yield a more appropriate natural mortality rate estimator;
- A yield-per-recruit analysis with the consideration of uncertainty associated with life history parameters (e.g., Chen 1996; Chang et al. 2009) can be done to estimate theoretical biological reference points such as F_{max} and $F_{0.1}$; and
- A spawning stock biomass-per-recruit analysis with the incorporation of uncertainty associated with life history parameters can also be done to estimate reference points such as $F_{20\%}$ and $F_{40\%}$.

V. Conclusions and Recommendations

Given the data limitations, the assessment appears to be well-planned and structured. Uncertainties in the quality and quantity of data, fisheries (e.g., selectivity) and life history parameters (e.g., von Bertalanffy growth parameters and natural mortality rate), and model structure (different estimators used to estimate the total and natural mortality rates) were carefully evaluated. I would like to commend the efforts of the AW panel in addressing data quality and quantity issues, identifying and evaluating implicit and explicit assumptions associated with methods and data, designing and conducting a rather systematic sensitivity analysis, exploring alternative model configurations and parameterization. However, based on the information I have reviewed, I cannot conclude that this assessment is scientifically sound, and adequately addresses needs for management advice. This mainly results from the data limitations.

I do have concerns for both the Caribbean queen triggerfish and blue tang that I hope the AW panel could address to improve the assessment of the Caribbean queen triggerfish and blue tang. I made the following general comments and specific recommendations.

General comments

The Caribbean queen triggerfish and blue tang are typical data-poor fisheries with no fishery-independent data and limited fishery-dependent data of questionable quality. Their life history processes are not well understood and key life history parameters are not well quantified. The low quality and quantity of the information available makes it extremely difficult to assess the status of the Caribbean queen triggerfish and blue tang stocks. The top priority should be to develop a fishery-independent monitoring program for the reef-dependent species such as queen triggerfish and blue tang in the U.S. Caribbean. Such a program can yield a reliable abundance index and provide samples for basic biological studies to estimate key life history parameters such as von Bertalanffy growth parameters, fecundity, and length/age at maturity and their spatial variability, not only for the Caribbean queen triggerfish and blue tang; but also for other reef-dependent fish species inhabiting the same area.

The quality and quantity of fishery-dependent data should also be improved. This can be done by developing a port or sea sampling program or further improve current reporting system by including the information on spatial locations of catch and conducting some cross-validation studies of fishermen's reported data. The report of species-specific landings in recent years is certainly a good way to improve the data quality and quantity, making the landing data useful in the species-specific stock assessment. The information may be useful to decompose the historical landings of species group into species-specific landings. A program should be developed to interview fishermen on their historical and current fishing areas and the changes in their attitude towards the targeted species and size composition in the fishery (Ames 2004). I believe such an interview program is cost effective to collect some valuable historical information regarding fishing grounds and fishermen's preferences for species and size. Such information will be

valuable to improve the quality of the historical data and improve the understanding of possible temporal changes in fishing effort distribution and selectivity.

Given the data limitations, the choice of stock assessment models is rather limited for the U.S. Caribbean queen triggerfish and blue tang stock assessment. Instead of using a traditional Beverton-Holt method to estimate the total mortality from length-composition data, the AW panel listed six assumptions explicitly and explicitly associated with the method and decided to use the method by Gedamke and Hoenig (2006) which does not require the assumption of an equilibrium population. Given the available data, this may be the best approach available. However, this approach also requires some assumptions in temporal variability in selectivity. Although Gedamke and Hoenig (2006) conducted a simulation study to evaluate the performance of the estimator, their simulation was based on a single species with different biology and fishing intensity. I suggest that the AW panel uses the Caribbean queen triggerfish and blue tang data to design a similar simulation study. The AW panel can design a few scenarios to evaluate the performance of the estimator in retrieving the “true” built in the simulation study and identify key factors that may greatly influence the performance of the estimator for the U.S. Caribbean queen triggerfish and blue tang fisheries.

A rather comprehensive sensitivity analysis was conducted to evaluate possible impacts of uncertainty associated with the growth parameters for the estimation of the total mortality rate in the assessment of the queen triggerfish and blue tang. Although I appreciate the AW panel’s efforts, I believe a better structured Monte Carlo simulation approach may be better in quantifying the uncertainty associated with the estimation of the total mortality rate and natural mortality rate. The AW panel can use the sensitivity analysis to identify the most plausible parameterization of the Gedamke-Hoenig model (Gedamke and Hoenig 2006) and then conduct a Monte Carlo simulation approach with parameters K and L_{∞} randomly drawn from their joint distribution which can be derived from bootstrapped nonlinear least squares in fitting the von Bertalanffy growth model to length-at-age data. similar approach can be used for estimating the natural mortality rate.

I also believe expert and background knowledge about the queen triggerfish and blue tang should be used to reduce the magnitude of the uncertainty on the growth parameters and natural mortality rate. Some values for M , K , and L_{∞} appear to be not biologically realistic for a fish species with a life history process similar to the queen triggerfish and blue tang, and should be excluded in the assessment. Maybe a literature search for fish species of similar life history and habitat should be done to derive a range of values that are biologically realistic for the key life history parameters.

The AW panel did not do a yield-per-recruit analysis and SSB-per-recruit (or egg-per-recruit) analysis because of uncertainty associated with the growth parameters and natural mortality rate. However, these values were used in the estimation of the total mortality rate and natural mortality rate. This is a rather inconsistent argument. I would like to suggest that at least a yield-per-recruit analysis can be done to estimate F_{MAX} and $F_{0.1}$ for possible reference points. The fact that both growth parameters and natural mortality rate are used in a yield-per-recruit analysis and estimation of the current fishing mortality rate using the approach described in the

assessment may reduce the impact of uncertainty associated with the growth parameters and natural mortality rate on the determination of the fishery status.

Specific recommendations

Although I have provided comments and recommendations under each TOR, I would like to re-iterate the following recommendations.

- I recommend that expert and background knowledge/information on species of similar life history patterns and habitat needs be used to exclude biologically unrealistic values of K , L_{∞} , natural mortality rate, and total mortality rate;
- I recommend that uncertainty associated with K , L_{∞} , natural mortality rate, total mortality rate, and subsequently fishing mortality rate be quantified using a Monte Carlo simulation;
- I suggest that a program be developed for interviewing fishermen to have a better understanding of temporal and spatial variability of the fishing ground, target fish species and size (i.e. selectivity), and fishing efforts;
- I recommend that information on the fish species composition of current landings be used to decompose the historical landings of species group into the species-specific landings;
- I recommend that a fishery-independent survey program be developed for the U.S. Caribbean queen triggerfish, blue tang and other reef-dependent species sharing similar habitat to collect samples for estimating basic life history parameters and for driving reliable abundance indices;
- I recommend that a simulation study be conducted to evaluate the performance of the length-based estimator of the total mortality rate and identify assumptions/parameters that can influence greatly the performance of the estimator;
- I suggest that a yield-per-recruit analysis be conducted with the incorporation of uncertainty associated with life history parameters to estimate theoretical biological reference points such as F_{\max} and $F_{0.1}$; and
- I recommend that a spawning stock biomass-per-recruit analysis be done with the incorporation of uncertainty associated with life history parameters to estimate reference points such as $F_{20\%}$ and $F_{40\%}$.

VI. References cited

- Aiken, K.A. 1975. Chapter 15: The biology, ecology and bionomics of the triggerfishes, Balistidae. In: Munro, J.L. (Ed.) Caribbean Coral Reef Fishery Resources. International Center for Living Aquatic Resources Management, Manila, Philippines.
- Ames, T. 2004. Atlantic cod stock structure in the Gulf of Maine. *Fisheries* 29:10-29.
- Beverton, R.J.H. and S.J. Holt. 1957. On the dynamics of exploited fish populations. U.K. Ministry of Agriculture, Fisheries, Food, and Fishery Investigations Series II, Vol. XIX.
- Bryan, M.D. 2012. Summary of recreational catch and effort for blue tang and queen triggerfish caught in Puerto Rico since 2000. SEDAR30-AW-01. SEDAR, North Charleston, SC. 15 pp.
- Carpenter, K.E. 2002. The living marine resources of the Western Central Atlantic. Volume 3: Bony fishes part 2 (Opistognathidae to Molidae), sea turtles and marine mammals. FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Special Publication No. 5. Rome, FAO.
- Chang, Y. J., C.J. Sun, Y. Chen, S.Z. Yeh, and W.C. Chiang. 2009. Incorporating uncertainty into the estimation of biological reference points for a spiny lobster (*Panulirus penicillatus*) fishery. *New Zealand Journal of Marine and Freshwater Research*, 43: 429-442.
- Chen, Y. 1996. A Monte Carlo study on impacts of the size of subsample catch on estimation of fish stock parameters. *Fisheries Research* 26(3-4): 207-223.
- Chen, Y., and C. Wilson. 2002. A simulation study to evaluate impacts of uncertainty on the assessment of American lobster fishery in the Gulf of Maine. *Canadian journal of Fisheries and aquatic Sciences*, 59: 1394-1403.
- Choat, J.H. and D.R. Robertson. 2002. Age-based studies on coral reef fishes. In: Sale P.F. (ed) Coral reef fishes. Dynamics and diversity in a complex ecosystem. Academic Press, New York.
- de Albuquerque, C.Q., A.S. Martins, L. Junior, N. de Oliveira, J.N.D. Araújo, and A.M. Ribeiro. 2011. Age and growth of the queen triggerfish *Balistes vetula* (tetraodontiformes, balistidae) of the central coast of Brazil. *Brazilian Journal of Oceanography*, 59(3), 231-239.
- Gedamke, T. and J.M. Hoenig. 2006. Estimating mortality from mean length data in nonequilibrium situations, with application to the assessment of goosefish. *Transactions of the American Fisheries Society* 135: 476-487.

- Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin* 81(4):898–903.
- Jensen, A.L. 1996. Beverton and Holt life history invariants result from optimal trade-off of reproduction and survival. *Canadian Journal of Fisheries and Aquatic Sciences* 53(4):820–822.
- King, M. 1995. *Fisheries Biology, Assessment and Management*. University of Iowa Press.
- Manooch, C.S and C.L. Drennon. 1987. Age and growth of yellowtail snapper and queen triggerfish collected from the U S. Virgin Islands and Puerto Rico. *Fisheries Research* 6: 53-68.
- Mutz, S.J. 2006. *Comparative growth dynamics of Acanthurid fishes*. School of Marine Biology and Aquaculture, James Cook University.
- Olsen, D.A. 2011. *Acanthurus coeruleus*. In Froese, R. and D. Pauly (eds.). Fishbase. 9 Oct. 2012. <http://www.fishbase.org/photos/PicturesSummary.php?StartRow=5&ID=944&what=species&TotRec=8>
- Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal du Conseil international pour l'Exploration de la Mer* 39(2):175–192.
- Randall, J.E., 1968. *Caribbean reef fishes*. T.F.H. Publications, Inc. Ltd., The British Crown Colony of Hong Kong.
- SEDAR 30. 2013. *Caribbean blue tang and queen triggerfish assessment*. SEDAR, 4055Faber Place Drive, Suite 201, North Charleston, SC 29405.

Appendix 1: Bibliography of materials provided for review for SEDAR 30 Caribbean Blue Tang and Queen Triggerfish

Documents Prepared for the Assessment Workshop

- Bryan, M. 2012. Summary of recreational catch and effort for blue tang and queen triggerfish caught in Puerto Rico since 2000. SEDAR30-AW-01.
- Bryan, M. 2012. Evaluation of the available length-frequency information in the US Caribbean Trip Interview Program (TIP) data. SEDAR30-AW-02
- Rios, A. B. 2012. A review of the life history characteristics of blue tang and queen triggerfish. SEDAR30-AW-03
- McCarthy, K. J. 2012. Commercial fishery landings of queen triggerfish and blue tang in the United States Caribbean, 1983-201. SEDAR30-AW-04

Final Stock Assessment Reports

- SEDAR30-SAR1: Blue Tang Assessment Report
- SEDAR30-SAR2: Queen Triggerfish Assessment Report

Reference Documents

- SEDAR30-RD01: A pilot program to assess methods of collecting bycatch, discard, and biological data in the commercial fisheries of St. Thomas, U.S. Caribbean
- SEDAR30-RD02: A pilot program to assess methods of collecting bycatch, discard, and biological data in the commercial fisheries of U.S. Caribbean (Saint Croix)

Appendix 2: A copy of the CIE Statement of Work

Attachment A: Statement of Work for Dr. Yong Chen

External Independent Peer Review by the Center for Independent Experts

SEDAR 30 Caribbean blue tang and queen triggerfish assessment review

Scope of Work and CIE Process: The National Marine Fisheries Service's (NMFS) Office of Science and Technology coordinates and manages a contract providing external expertise through the Center for Independent Experts (CIE) to conduct independent peer reviews of NMFS scientific projects. The Statement of Work (SoW) described herein was established by the NMFS Project Contact and Contracting Officer's Representative (COR), and reviewed by CIE for compliance with their policy for providing independent expertise that can provide impartial and independent peer review without conflicts of interest. CIE reviewers are selected by the CIE Steering Committee and CIE Coordination Team to conduct the independent peer review of NMFS science in compliance the predetermined Terms of Reference (ToRs) of the peer review. Each CIE reviewer is contracted to deliver an independent peer review report to be approved by the CIE Steering Committee and the report is to be formatted with content requirements as specified in **Annex 1**. This SoW describes the work tasks and deliverables of the CIE reviewer for conducting an independent peer review of the following NMFS project. Further information on the CIE process can be obtained from www.ciereviews.org.

Project Description SEDAR 30 will be a compilation of data, an assessment of the stock, and an assessment review conducted for Caribbean blue tang and queen triggerfish. The CIE peer review is ultimately responsible for ensuring that the best possible assessment has been provided through the SEDAR process. The stocks assessed through SEDAR 30 are within the jurisdiction of the Caribbean Fisheries Management Council and the territorial waters of Puerto Rico and the U.S. Virgin Islands. The Terms of Reference (ToRs) of the peer review are attached in **Annex 2**.

Requirements for CIE Reviewers: Three CIE reviewers shall have the necessary qualifications to complete an impartial and independent peer review in accordance with the tasks and ToRs described in the SoW herein. The CIE reviewers shall have expertise in stock assessment, statistics, fisheries science, and marine biology sufficient to complete the tasks of the scientific peer-review described herein. Each CIE reviewer's duties shall not exceed a maximum of 10 days to complete all work tasks of the peer review described herein.

Location of Peer Review: Each CIE reviewer shall conduct the desk review during 4-7 February 2013, therefore no travel will be required.

Statement of Tasks: Each CIE reviewer shall complete the following tasks in accordance with the SoW and Schedule of Milestones and Deliverables herein.

Prior to the Peer Review: Upon completion of the CIE reviewer selection by the CIE Steering Committee, the CIE shall provide the CIE reviewer information (full name, title, affiliation, country, address, email) to the COR, who forwards this information to the NMFS Project

Contact no later the date specified in the Schedule of Milestones and Deliverables. The CIE is responsible for providing the SoW and ToRs to the CIE reviewers. The NMFS Project Contact is responsible for providing the CIE reviewers with the background documents, reports, and other information pertinent to the desk review arrangements. Any changes to the SoW or ToRs must be made through the COR prior to the commencement of the peer review.

Pre-review Background Documents: Two weeks before the peer review, the NMFS Project Contact will send (by electronic mail or make available at an FTP site) to the CIE reviewers the necessary background information and reports for the peer review. In the case where the documents need to be mailed, the NMFS Project Contact will consult with the CIE Lead Coordinator on where to send documents. CIE reviewers are responsible only for the pre-review documents that are delivered to the reviewer in accordance to the SoW scheduled deadlines specified herein. The CIE reviewers shall read all documents in preparation for the peer review.

Desk Review: Each CIE reviewer shall conduct the independent peer review in accordance with the SoW and ToRs, and shall not serve in any other role unless specified herein. **Modifications to the SoW and ToRs shall not be made during the peer review, and any SoW or ToRs modifications prior to the peer review shall be approved by the COR and CIE Lead Coordinator.** The CIE Lead Coordinator can contact the Project Contact to confirm any peer review arrangements.

Contract Deliverables - Independent CIE Peer Review Reports: Each CIE reviewer shall complete an independent peer review report in accordance with the SoW. Each CIE reviewer shall complete the independent peer review according to required format and content as described in Annex 1. Each CIE reviewer shall complete the independent peer review addressing each ToR as described in Annex 2.

Specific Tasks for CIE Reviewers: The following chronological list of tasks shall be completed by each CIE reviewer in a timely manner as specified in the **Schedule of Milestones and Deliverables**.

- 1) Conduct necessary pre-review preparations, including the review of background material and reports provided by the NMFS Project Contact in advance of the peer review.
- 2) During February 4-7, 2013 as specified herein, conduct an independent desk peer review in accordance with the ToRs (**Annex 2**).
- 3) No later than February 21, 2013, each CIE reviewer shall submit an independent peer review report addressed to the “Center for Independent Experts,” and sent to Mr. Manoj Shivlani, CIE Lead Coordinator, via email to shivlanim@bellsouth.net, and CIE Regional Coordinator, via email to Dr. David Sampson david.sampson@oregonstate.edu. Each CIE report shall be written using the format and content requirements specified in Annex 1, and address each ToR in **Annex 2**.

Schedule of Milestones and Deliverables: CIE shall complete the tasks and deliverables described in this SoW in accordance with the following schedule.

14 January 2013	CIE sends reviewer contact information to the COR, who then sends this to the NMFS Project Contact
18 January 2013	NMFS Project Contact sends the stock assessment report and background documents to the CIE reviewers.
4-13 February 2013	Each reviewer conducts an independent desk peer review
19 February 2013	CIE reviewers submit draft CIE independent peer review reports to the CIE Lead Coordinator and CIE Regional Coordinator
7 March 2013	CIE submits CIE independent peer review reports to the COR
14 March 2013	The COR distributes the final CIE reports to the NMFS Project Contact and regional Center Director

Modifications to the Statement of Work: This ‘Time and Materials’ task order may require an update or modification due to possible changes to the terms of reference or schedule of milestones resulting from the fishery management decision process of the NOAA Leadership, Fishery Management Council, and Council’s SSC advisory committee. A request to modify this SoW must be approved by the Contracting Officer at least 15 working days prior to making any permanent changes. The Contracting Officer will notify the COR within 10 working days after receipt of all required information of the decision on changes. The COR can approve changes to the milestone dates, list of pre-review documents, and ToRs within the SoW as long as the role and ability of the CIE reviewers to complete the deliverable in accordance with the SoW is not adversely impacted. The SoW and ToRs shall not be changed once the peer review has begun.

Acceptance of Deliverables: Upon review and acceptance of the CIE independent peer review reports by the CIE Lead Coordinator, Regional Coordinator, and Steering Committee, these reports shall be sent to the COR for final approval as contract deliverables based on compliance with the SoW and ToRs. As specified in the Schedule of Milestones and Deliverables, the CIE shall send via e-mail the contract deliverables (CIE independent peer review reports) to the COR (William Michaels, via William.Michaels@noaa.gov).

Applicable Performance Standards: The contract is successfully completed when the COR provides final approval of the contract deliverables. The acceptance of the contract deliverables shall be based on three performance standards:

- (1) The CIE report shall be completed with the format and content in accordance with **Annex 1**,
- (2) The CIE report shall address each ToR as specified in **Annex 2**,
- (3) The CIE reports shall be delivered in a timely manner as specified in the schedule of milestones and deliverables.

Distribution of Approved Deliverables: Upon acceptance by the COR, the CIE Lead Coordinator shall send via e-mail the final CIE reports in *.PDF format to the COR. The COR will distribute the CIE reports to the NMFS Project Contact and Center Director.

Support Personnel:

William Michaels, Program Manager, COR
NMFS Office of Science and Technology
1315 East West Hwy, SSMC3, F/ST4, Silver Spring, MD 20910
William.Michaels@noaa.gov Phone: 301-427-8155

Manoj Shivlani, CIE Lead Coordinator
Northern Taiga Ventures, Inc.
10600 SW 131st Court, Miami, FL 33186
shivlanim@bellsouth.net Phone: 305-383-4229

Roger W. Peretti, Executive Vice President
Northern Taiga Ventures, Inc. (NTVI)
22375 Broderick Drive, Suite 215, Sterling, VA 20166
RPeretti@ntvifederal.com Phone: 571-223-7717

Key Personnel:

NMFS Project Contact:

Julie Neer, SEDAR Coordinator
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405
julie.neer@safmc.net Phone: 843-571-4366

Annex 1: Format and Contents of CIE Independent Peer Review Report

1. The CIE independent report shall be prefaced with an Executive Summary providing a concise summary of the findings and recommendations, and specify whether the science reviewed is the best scientific information available.
2. The main body of the reviewer report shall consist of a Background, Description of the Individual Reviewer's Role in the Review Activities, Summary of Findings for each ToR in which the weaknesses and strengths are described, and Conclusions and Recommendations in accordance with the ToRs. The CIE independent report shall be a stand-alone document for others to understand the weaknesses and strengths of the science reviewed. The CIE independent report shall be an independent peer review of each ToRs.
3. The reviewer report shall include the following appendices:

Appendix 1: Bibliography of materials provided for review

Appendix 2: A copy of the CIE Statement of Work

Annex 2: Terms of Reference for the Peer Review

SEDAR 30 Caribbean blue tang and queen triggerfish assessment review

1. Evaluate the data used in the assessment, addressing the following:
 - a) Are data decisions made by the Assessment Workshop sound and robust?
 - b) Are data uncertainties acknowledged, reported, and within normal or expected levels?
 - c) Are data applied properly within the assessment model?
 - d) Are input data series reliable and sufficient to support the assessment approach and findings?
2. Evaluate the methods used to assess the stock, taking into account the available data.
 - a) Are methods scientifically sound and robust?
 - b) Are assessment models configured properly and used consistent with standard practices?
 - c) Are the methods appropriate for the available data?
3. Evaluate the assessment findings with respect to the following:
 - a) Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?
 - b) Is the stock overfished? What information helps you reach this conclusion?
 - c) Is the stock undergoing overfishing? What information helps you reach this conclusion?
 - d) Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?
 - e) Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?
4. Evaluate the stock projections, addressing the following:
 - a) Are the methods consistent with accepted practices and available data?
 - b) Are the methods appropriate for the assessment model and outputs?
 - c) Are the results informative and robust, and useful to support inferences of probable future conditions?
 - d) Are key uncertainties acknowledged, discussed, and reflected in the projection results ?
5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.

- Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods
 - Ensure that the implications of uncertainty in technical conclusions are clearly stated.
6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.
 - Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
 - Provide recommendations on possible ways to improve the SEDAR process.
 7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.

30th Southeast Data, Assessment and Review (SEDAR 30)

**Reviewer Report to the Center for Independent Experts on the U.S.
Caribbean Blue Tang and Queen Triggerfish (SEDAR 30)**

March 20, 2013

Prepared for:
Center for Independent Experts

By:
M. Kurtis Trzcinski
5 1/2 A Quarry Rd
Halifax, Nova Scotia, Canada
B3N 1X1

Executive Summary

This document is an independent review of the activities and findings of the 30th Southeast Data, Assessment and Review (SEDAR 30). The review was a desktop review, that is, assessment documents and supporting material were sent out for review, but there was not a meeting or an opportunity for dialogue regarding the assessment. Two stocks were reviewed: the U.S. Caribbean blue tang and queen triggerfish. While I acknowledge that the science reviewed is the best scientific information available and that considerable effort was made to make the best use of the data available, I do not find that either assessment provides a sound basis for management advice. Several of the ToRs were met, but the most important ones regarding stock status, trends and the impact of fishing were not (ToRs 3 and 4). The failure to meet these ToRs are through no fault of the assessment team, but rather, are due to the lack of data and often the poor quality of the data they do have.

The approach to each assessment was similar. It was in fact so similar, I had trouble seeing the differences and I wonder what differences in methods, if any, are warranted given the life history of each species and any differences in the fishery. Consequently, my comments typically apply to both assessments and I only make distinctions where necessary.

My overall conclusions are that we don't know much about the status of these fish or whether overfishing has occurred. A noble effort has been made, but alas, we have tremendous uncertainty and little basis for management advice. Rather than focusing on the particulars of these assessments, which I do below, I encourage the assessment team to take a strategic approach to the problem. In particular, to formulate a medium to long-term plan to, 1) engage external scientists in a program review where an assessment framework (benchmark) can be agreed upon, and 2) make plans to collect the data to support it.

1.0 Background

This document is an independent review of the findings of the 30th Southeast Data, Assessment and Review (SEDAR 30). The review was a desktop review, that is, assessment documents and supporting material were sent out for review, but there was not a meeting or an opportunity for dialogue regarding the assessment. Two stocks were reviewed: the U.S. Caribbean blue tang and queen triggerfish. Assessment documents (Appendix 1) and background materials were provided via a website two weeks before the review. I was also provided with a Statement of Work (Appendix 2), including the Terms of Reference (ToR).

2.0 Individual Reviewer Activities

I reviewed the assessment and background documents provided for the review. This was a desktop review so there was no dialogue between the assessment team and me and all my comments only pertain to documents provided to me. As outlined in Statement of Work (Appendix 2), these reports should state in the reviewer's own words whether each ToR of the Stock Assessment Workshop was completed successfully, should state whether they accepted or rejected the work that they reviewed, and should include an explanation of their decisions (strengths, weaknesses of the analyses, etc.) and recommendations for each ToR. A key determinant of whether a ToR had been met was the extent to which it provided a scientifically credible basis for developing fishery management advice. The following section contains my review for both assessments.

3.0 Review of U.S. Caribbean Blue Tang and Queen Triggerfish

U.S. Caribbean blue tang and queen triggerfish were assessed using a length-based estimate of total mortality (Beverton and Holt 1957, Gedamke and Hoenig 2006). I presume that this is the first time these stocks have been assessed as there were no citations to previous assessments. If these are the first assessments of these stocks, I think it is important to put in the assessment report as context (as it stands now, the information is presented in section 3 of Section I; I suggest making that more prominent in the introductions to Sections I and II). I felt like a lot of the structure of the report and writing was 'cookie cutter', that is, following a particular formulae. This seems counter productive to me. In my view, what is needed in a first assessment is a review of past work and some soul searching for ways forward. This is better done in a free format. Furthermore, the assessments are chopped up into little pieces (six supporting documents). While the details should remain in the supporting documents, I think a synthesis of these documents is needed in the Assessment Process Report. Maybe that is what is lacking the most: a cohesive synthesis and vision. Blue tang and queen triggerfish have similar data and assessment problems, how can we solve them together? Or for perhaps a larger species group? Step back and take stock of the situation, so to speak. I

am having trouble getting the sense of whether people care and if so exactly what they care about. Compared to some of the world's fisheries the landings of surgeonfish and triggerfish are small (~35, ~80 metric tons / year respectively) although this is a relatively small area as well. Maybe more data and analysis to support management is not needed. If it is, then a stronger case should be made for it. Similarly, if one is going to choose an assessment method, it should be defended and not simply stated that the 'AW panel decided'.

The assessment of blue tag and queen triggerfish is difficult given the quality and limited amount of data. Overall, the data have been assembled with considerable care and diligence. Some issues remain, but it is a good point in the history of these fisheries to step back and evaluate the quality of the data and the most effective means of assessing stock status and the impact of fishing. The assessment team makes good use of the data, given what they have and it is clear that progress has been made and that higher quality data are being collected. Stock status, in terms of a biomass based reference, cannot be determined given the approach used and the impact of fishing can only be estimated using strong and weakly supported assumptions about fishing and natural mortality. In my view, the current monitoring and assessment approach is inadequate and what is needed is an overall program review. I am not a manager, but if I were, I would be uncomfortable managing this fishery with such limited tools to assess stock status and the impact of fishing. So that causes me, as a reviewer, to look to the future. What is the framework or benchmark which will be used to assess these populations? Has there been adequate discussion and review of a proposed framework? Once the framework is decided upon, then the focus can be put on the most important and useful information to collect. The discussion of the assessment approach and data collection needs to proceed hand in hand. I will go so far as to suggest that this assessment team formulate a proposal for a new assessment approach along with a plan to collect the necessary data and submit that for review.

I did find the management history and context presented in Section I: Introduction useful, complicated, but important to be aware of. Actually, I would like to be slightly more complimentary of this work, as I don't see it enough and appreciate the work it takes. What I would like is for the managers and biologists to make stronger statements about if and when these management measures would be expected to affect the data used in the assessment. If a management measure was put in place and it was expected to affect mean length in the catch, did we see it? Did we even have the data to see it? Which of these management measures can be ignored, in terms of the assessment, and which should be explicitly accounted for. If you find a change in total mortality, can it be attributed to a management measure and / or a change in fishing practices? This is good work, but it should be pushed to the next level, if possible (I acknowledge the difficulties, but encourage those involved to try).

I find the explanation of methods in the Section II: Assessment Process Report thin, but what is needed even more is better motivation of the general approach and methods used. The motivation is even more important because this is a desktop review. It is much easier to get a sense of why decisions about the data or methods were made when there is a

presentation and reviewers are afforded the opportunity to ask questions. In the situation of a desktop review, we are left with sentences like 'The AW [assessment workshop] panel determined..., The AW panel agreed...', but I need more written explanation supporting the decision or conclusion in order to decide whether I reach the same conclusion or not.

I see at least four major data categories where you might concentrate your efforts: 1) landings data, 2) fisheries independent survey, 3) life history data, and 4) tagging. As I noted, the priority and amount of effort given to each depends on the assessment framework you plan to use. Obviously, if one wishes to assess the stock with fisheries based catch per unit effort (cpue) data, then emphasis would be placed on collecting high quality effort and landings data. If on the other hand, one wishes to assess the stock using a fisheries independent survey, or calculate yield per recruit then the priorities shift and a different investment is required. I fear I state the obvious to my fellow stock assessment scientists, but in an effort to be as helpful and constructive as possible, I elaborate on these issues below (ToR 7).

ToRs 1,2,5 and 6 were met, but ToRs 3 and 4 were not in both assessments. I provide comments on all the ToRs below.

ToR 1

1. *Evaluate the data used in the assessment, addressing the following:*
 - a) *Are data decisions made by the Assessment Workshop sound and robust?*
 - b) *Are data uncertainties acknowledged, reported, and within normal or expected levels?*
 - c) *Are data applied properly within the assessment model?*
 - d) *Are input data series reliable and sufficient to support the assessment approach and findings?*

This ToR was met. There are limited data for this fishery. The landings are unknown for the species blue tang or queen triggerfish, but are recorded collectively as surgeonfish and triggerfish. There has been some length sampling, but they appear to be uninformative for a length-based cohort analysis as one cannot see cohorts in the length frequency plots (both species). Life history data (growth, age at maturity) for blue tang from the management unit is lacking and data are taken from other areas. The life history data for queen triggerfish is taken from Puerto Rico and the US Virgin Islands but the authors suggest that it needs to be verified by another study using otoliths rather than dorsal spines. Plots of the frequency at length by age and examining the modes and overlap among ages would be useful, to help determine if a length or age based model might be effective. Neither species has a fisheries independent survey. Overall, there is very little to go on. So, the greatest advancements in understanding the dynamics in these stocks will occur by collecting better data.

One big difficulty is not knowing the landings. The landings are aggregated into large species groups and until there is a method to estimate landings at a species level it will be hard to answer the question 'should we care'. I would like to say we should care about every fish we remove from the ocean, but the fact is that we must prioritize our efforts. Certainly some surgeonfish and triggerfish are being removed, but even a rough estimate of proportion by species would be useful. It appears that the reporting regulations have changed and that this will no longer be a problem, but I suggest you take the time to demonstrate that it is not a problem in the future with some biological sampling of the catch. It is stated that species-specific data were reported in the US Virgin Islands during the 2011 - 2012 fishing year. I think it would have been very useful to report the proportion of blue tang and queen triggerfish in your assessment. If there have not been a lot of gear changes or changes in fishing practices, these proportions can be applied with some caution to the older landings data. Do we have enough information to partition the landings in the other areas? If not, what additional data would be needed?

I am not entirely sure if blue tang and queen triggerfish are targeted in the fishery or if they are principally bycatch species. If they are only a small proportion of the surgeonfish and triggerfish maybe this assessment approach is adequate. My confusion, of course, would have been cleared up immediately if this review was done at a meeting. This small example shows how extra effort is required by the assessment team to explain the context of the situation.

In general, I would have appreciated more background on the natural history of these fish and the community and ecosystem in general. This context is important in evaluating the appropriateness of this assessment and avenues for its improvement. I would integrate all the information in SEDAR30-AW-03 'A review of the life history characteristics...' into the Assessment Process Report. One notable gap in your knowledge is the length or age at maturity.

I can appreciate the fact that improved sampling of the catch will be difficult, and I found the pilot studies examining these issues useful (SEDAR30-RD-01, SEDAR30-RD-02). Some sort of improved sampling will need to occur, but the level of investment is proportional to the assessment method. If you want to use a length-based or age-based model then this data stream will be very important and will require further study and planning. If you instead choose to use tagging as an assessment method, sampling the catch is less of a priority.

- a) The decisions about how to use the data appear to be robust and sound.
- b) Data uncertainties are acknowledged and reported. I find the uncertainties larger than 'normal' and these uncertainties severely limit the ability to track the population and to estimate the impact of fishing. Only the most general of statements about total mortality and whether it has changed can be made and given the uncertainties we cannot be confident in these conclusions.
- c) The data appear to be applied properly within the model: the length based mortality estimator by Gedamke and Hoenig (2006).

d) This depends on what the goals are. I think the goals should be on how to move to a better assessment framework. The input data series are not reliable and sufficient to support the assessment of stock status and the impact of fishing, the typical goals of an assessment. If the goal is to estimate whether there has been a change (increase?) in the total mortality rate, then this work makes a contribution, but the conclusions must be very tentative given the quality of the data.

ToR 2

2. *Evaluate the methods used to assess the stock, taking into account the available data.*

a) *Are methods scientifically sound and robust?*

b) *Are assessment models configured properly and used consistent with standard practices?*

c) *Are the methods appropriate for the available data?*

This ToR was met. The methods used for a length-based estimate of the total mortality rate look sound, but maybe not very robust. The methods are appropriate for the data available. As noble as these efforts are, I have trouble viewing these methods as an 'assessment'. With better life history and selectivity data we may have more confidence in an estimate of the total mortality rate and any conclusion about whether the total mortality rate has changed. The signal in any changes in length does not appear to be as strong as in the examples in Gedamke and Hoenig (2006). There is some evidence that the total mortality rate has increased for blue tang and possibly decreased for queen triggerfish, but changes in fishing practices may (probably?) make it difficult to interpret these data. In the AIC results, it is essentially profiling over different life history input parameters *and* changes in Z. The interpretation focuses on the best model for Z within a combination of life history parameters, but do these results also indicate the most likely combination of life history parameters? Can one actually put forth one or two models from Table 18 blue tang and Table 13 queen triggerfish as the best model?

a) The methods are scientifically sound and robust, but they cannot estimate stock status and the impact of fishing. I would like to have seen the profile likelihoods or the Bayesian posteriors for the estimates of total mortality and change year, similar to Figure 3 in Gedamke and Hoenig (2006), but more. These kinds of diagnostics are typically important to present.

b) The model is configured properly.

c) The methods are appropriate for the available data but do not form the basis for strong management advice (objectives 3 and 4 below). I have some trouble with the sensitivity analysis. It seems like a very wide range was chosen and that just has the effect of demonstrating that the estimates of the total mortality rate could be just about anything. How were the ranges of the life history parameters chosen? Were they the 95% credible interval (CI) from a growth study? I also have difficulty tracing back the range of total mortality rates used in the tables estimating F and M (blue tang: Tables 16 and 24, queen

triggerfish: Tables 19, 21, 25). Couldn't that range be taken from the CI of Z estimated from your analysis of mean length?

ToR 3

3. *Evaluate the assessment findings with respect to the following:*
 - a) *Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?*
 - b) *Is the stock overfished? What information helps you reach this conclusion?*
 - c) *Is the stock undergoing overfishing? What information helps you reach this conclusion?*
 - d) *Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?*
 - e) *Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?*

This ToR was not met. By and large it is not possible to reach these objectives given the lack of suitable data. The assessment does not provide abundance or biomass estimates and given the data, it currently is not possible. It does provide an estimate of total mortality (Z) from which fishing mortality (F) can be inferred. However, there are large uncertainties in both total mortality and natural mortality making it exceedingly difficult to estimate fishing mortality ($Z-M=F$). They did not provide a convincing argument that their estimate(s) of mortality are useful to support status inferences. They do a good job of showing the uncertainty, but could do a better job arguing which estimate has the most support. A preferred model or estimate brings focus to the discussion about whether the estimate or model is 'useful to support status inferences'. As noted by the authors in the general discussion of the blue tang assessment:

'The disparate estimates of growth led to considerable uncertainty in the mortality estimates. They also made it difficult to meaningfully interpret stock status in terms of fishing mortality, in the absence of a weighting system giving credence to one life-history strategy over another.'

So given these data, the status of the blue tang fishery cannot be well determined. I agree entirely with this statement. Although, I think sometime we need to stick our neck out and rely on 'expert knowledge', if for nothing else than to push things forward (I guess that is what I am doing with all these comments!).

Similarly for queen triggerfish in the section on stock status and general conclusions the authors state:

'...it is difficult to interpret the sustainability of the estimated, current exploitation rates and that the absolute estimates of mortality should be interpreted with caution.'

I also agree entirely with this statement. So given the data, and a good analysis of what data is available, I conclude that we can not determine if the stock is overfished or if overfishing is occurring. The current data and modeling is light-years away from estimating a stock recruitment relationship, and I don't think this should be a short or medium term goal. Stock status cannot be determined with the current data and there are very few other data, if any, that can be used to inform managers about stock trends and conditions. If there were better life history data from the management unit and better length sampling of the catch then statements about overfishing based on F relative to M potentially could be better substantiated, but it will not solve all your problems and I would not rush to this without a more thorough program review.

ToR 4

4. Evaluate the stock projections, addressing the following:

- a) Are the methods consistent with accepted practices and available data?*
- b) Are the methods appropriate for the assessment model and outputs?*
- c) Are the results informative and robust, and useful to support inferences of probable future conditions?*
- d) Are key uncertainties acknowledged, discussed, and reflected in the projection results?*

This ToR was not met. No projections were done.

ToR 5

- 5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.*
 - Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods.*
 - Ensure that the implications of uncertainty in technical conclusions are clearly stated.*

The uncertainties are well addressed, there are just so many! Most of the uncertainties are either due to the lack of data or to data of poor quality / resolution. The AIC analysis and sensitivity analyses are useful and demonstrate the need for better data, but I think that one could and should put forth (tentatively) the best model. You could work on how to better visualize the output from multiple models. The assessment team points out that the estimates from the length based total mortality estimator 'should be considered with caution', which is clearly and appropriately stated.

ToR 6

6. *Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.*
 - *Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.*
 - *Provide recommendations on possible ways to improve the SEDAR process.*

I set this out more generally below. I cannot prioritize these well because I do not know 1) the species biology, ecosystem and fishery well, 2) the long-term assessment goals or 3) the financial constraints. I think the assessment team would benefit from a meeting to discuss these issues and help set out the overall assessment framework. But if prioritizing is at all useful given my limited knowledge, I would work on getting a fisheries independent survey together, I might even do this over the life history work although that should be done as well.

Presumably some discussion occurred about whether the method used was the best given the available data. I think it is important to review and recapitulate that argument in the introduction to the assessment report.

ToR 7

7. *Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.*

I wouldn't schedule another assessment until an assessment framework is chosen and the appropriate data collected to support it. Below I briefly review the potential value of collecting better data in four areas.

Landings

I encourage to continue to try and improve sampling of the landings. Some changes have already been made. Check to see they accomplish your goals. Stratification and expansion factors will be important issues if you decide to go to a length based or age based model.

Fisheries independent survey

A fisheries independent survey can be the most useful and important piece of data indicating stock status and the impact of fishing. I am showing some of my bias, but I think many fisheries scientists believe this as well. The effectiveness of a survey, however, depends on many things including, the natural variability in the distribution and abundance of the species or community of interest, the survey design, the gear, the catchability, etc.. Designing a good survey is a big task and unless the species is extremely valuable, most surveys are used for an array of species. In your research recommendations you identify a fisheries independent survey as a 'top research priority'. If so, then this is quite an undertaking and probably deserves a suite of studies examining

1) the species which the survey would focus on monitoring, 2) the gear and fishing method, and 3) statistical design including stratification, sample size and power analysis. I would like to encourage this work, but it should be acknowledged that commitment to a survey is a high level decision and that a new survey will take a lot of work and will require some long-term planning.

Life history data

There is quite a bit of uncertainty in the basic biology of this species. In particular, there has not been a growth study done in this area for blue tang, and there is quite a bit of variability in growth when looking across other regions. The authors suggest that the growth study for queen triggerfish be redone. Natural mortality is estimated from these growth studies, but you seem to show with the different methods and your sensitivity analysis that it could be just about anything. Which estimate should go forward and why? The authors argue that the Pauly (1980) method is better because it includes the growth coefficient and asymptotic length and then state that it 'may be robust to their negative correlation', but the real question is whether the extra parameter contains new or different information. No information was presented on the length or age at maturity which I believe is used in Roff's method (1984). It is stated that beyond the age of five, length is not informative about blue tang age. While this may be true, it would be better to support it with plots of the frequency at length for each age group, so we can see how much overlap there is. If the first 4 or 5 ages show distinct modes then a length-based model may still be a good way forward.

Although it appears that the authors have some data to estimate the selectivity of the gear, this has not been done. In the analysis of the total mortality rate, the parameter measuring the length of first capture (L_c) was allowed to vary. I would think about how to better estimate the selectivity of the gear. Tagging? If one ever wanted to do a yield per recruit analysis, this parameter along with natural mortality, and maturity would have to be better estimated.

Tagging

A tagging program can be used to address many questions, as I am sure most of those involved are aware. I just think it is important to bring up because it may be useful in designing a program for assessing these fish. Conventional tags can be used to estimate movement and help determine the appropriateness of the management unit, the selectivity of the gear and fish growth. Tagging can be used to get an estimate of the fishing and natural mortality rates, and an estimate of population size, however population size is more difficult and requires more tags. In my view, an overall assessment and research program benefits from a tagging program. It has the potential to reduce the number of assumptions in an assessment and the uncertainty in some parameters. In some cases it is the best method for an assessment given the natural history of the fish and the nature of the fishery.

4.0 References

Beverton, R. J., & Holt, S. J. (1957). *On the dynamics of exploited fish populations*. Springer.

Pauly, D. (1980). A new methodology for rapidly acquiring basic information on tropical fish stocks: growth, mortality and stock-recruitment relationships. *Stock assessment for tropical small-scale fisheries*, 154-172.

Roff, D. A. (1984). The evolution of life history parameters in teleosts. *Canadian Journal of Fisheries and Aquatic Sciences*, 41(6), 989-1000.

5.0 Appendices

Appendix 1: Bibliography of Materials Provided for Review

Appendix 2: CIE Statement of Work

Appendix 1: Bibliography of Materials Provided for Review

Documents Prepared for the Assessment Workshop

SEDAR30-AW-01. Summary of recreational catch and effort for blue tang and queen triggerfish caught in Puerto Rico since 2000. Meaghan Bryan

SEDAR30-AW-02. Evaluation of the available length-frequency information in the US Caribbean Trip Interview Program (TIP) data. Meaghan Bryan

SEDAR30-AW-03. A review of the life history characteristics of blue tang and queen triggerfish. Adyan B. Rios

SEDAR30-AW-04. Commercial fishery landings of queen triggerfish and blue tang in the United States Caribbean, 1983- 2011. Kevin J. McCarthy

Final Stock Assessment Reports

SEDAR30-SAR1. Blue tang

SEDAR30-SAR2. Queen triggerfish

Reference Documents

SEDAR30-RD01. A pilot program to assess methods of collecting bycatch, discard, and biological data in the commercial fisheries of St. Thomas, U.S. Caribbean. MRAG Americas

SEDAR30-RD02. A pilot program to assess methods of collecting bycatch, discard, and biological data in the commercial fisheries of U.S. Caribbean (Saint Croix). MRAG Americas

Appendix 1: CIE Statement of Work.

Attachment A: Statement of Work for Dr. Kurtis Trzcinski

External Independent Peer Review by the Center for Independent Experts

SEDAR 30 Caribbean blue tang and queen triggerfish assessment review

Scope of Work and CIE Process: The National Marine Fisheries Service's (NMFS) Office of Science and Technology coordinates and manages a contract providing external expertise through the Center for Independent Experts (CIE) to conduct independent peer reviews of NMFS scientific projects. The Statement of Work (SoW) described herein was established by the NMFS Project Contact and Contracting Officer's Representative (COR), and reviewed by CIE for compliance with their policy for providing independent expertise that can provide impartial and independent peer review without conflicts of interest. CIE reviewers are selected by the CIE Steering Committee and CIE Coordination Team to conduct the independent peer review of NMFS science in compliance the predetermined Terms of Reference (ToRs) of the peer review. Each CIE reviewer is contracted to deliver an independent peer review report to be approved by the CIE Steering Committee and the report is to be formatted with content requirements as specified in **Annex 1**. This SoW describes the work tasks and deliverables of the CIE reviewer for conducting an independent peer review of the following NMFS project. Further information on the CIE process can be obtained from www.ciereviews.org.

Project Description SEDAR 30 will be a compilation of data, an assessment of the stock, and an assessment review conducted for Caribbean blue tang and queen triggerfish. The CIE peer review is ultimately responsible for ensuring that the best possible assessment has been provided through the SEDAR process. The stocks assessed through SEDAR 30 are within the jurisdiction of the Caribbean Fisheries Management Council and the territorial waters of Puerto Rico and the U.S. Virgin Islands. The Terms of Reference (ToRs) of the peer review are attached in **Annex 2**.

Requirements for CIE Reviewers: Three CIE reviewers shall have the necessary qualifications to complete an impartial and independent peer review in accordance with the tasks and ToRs described in the SoW herein. The CIE reviewers shall have expertise in stock assessment, statistics, fisheries science, and marine biology sufficient to complete the tasks of the scientific peer-review described herein. Each CIE reviewer's duties shall not exceed a maximum of 10 days to complete all work tasks of the peer review described herein.

Location of Peer Review: Each CIE reviewer shall conduct the desk review during 4-7 February 2013, therefore no travel will be required.

Statement of Tasks: Each CIE reviewer shall complete the following tasks in accordance with the SoW and Schedule of Milestones and Deliverables herein.

Prior to the Peer Review: Upon completion of the CIE reviewer selection by the CIE Steering Committee, the CIE shall provide the CIE reviewer information (full name, title, affiliation, country, address, email) to the COR, who forwards this information to the NMFS Project Contact no later the date specified in the Schedule of Milestones and Deliverables. The CIE is responsible for providing the SoW and ToRs to the CIE reviewers. The NMFS Project Contact is responsible for providing the CIE reviewers with the background documents, reports, and other information pertinent to the desk review arrangements. Any changes to the SoW or ToRs must be made through the COR prior to the commencement of the peer review.

Pre-review Background Documents: Two weeks before the peer review, the NMFS Project Contact will send (by electronic mail or make available at an FTP site) to the CIE reviewers the necessary background information and reports for the peer review. In the case where the documents need to be mailed, the NMFS Project Contact will consult with the CIE Lead Coordinator on where to send documents. CIE reviewers are responsible only for the pre-review documents that are delivered to the reviewer in accordance to the SoW scheduled deadlines specified herein. The CIE reviewers shall read all documents in preparation for the peer review.

Desk Review: Each CIE reviewer shall conduct the independent peer review in accordance with the SoW and ToRs, and shall not serve in any other role unless specified herein. **Modifications to the SoW and ToRs shall not be made during the peer review, and any SoW or ToRs modifications prior to the peer review shall be approved by the COR and CIE Lead Coordinator.** The CIE Lead Coordinator can contact the Project Contact to confirm any peer review arrangements.

Contract Deliverables - Independent CIE Peer Review Reports: Each CIE reviewer shall complete an independent peer review report in accordance with the SoW. Each CIE reviewer shall complete the independent peer review according to required format and content as described in Annex 1. Each CIE reviewer shall complete the independent peer review addressing each ToR as described in Annex 2.

Specific Tasks for CIE Reviewers: The following chronological list of tasks shall be completed by each CIE reviewer in a timely manner as specified in the **Schedule of Milestones and Deliverables**.

- (1) Conduct necessary pre-review preparations, including the review of background material and reports provided by the NMFS Project Contact in advance of the peer review.
- (2) During February 4-7, 2013 as specified herein, conduct an independent desk peer review in accordance with the ToRs (**Annex 2**).
- (3) No later than February 21, 2013, each CIE reviewer shall submit an independent peer review report addressed to the “Center for Independent Experts,” and sent to Mr.

Manoj Shivlani, CIE Lead Coordinator, via email to shivlanim@bellsouth.net, and CIE Regional Coordinator, via email to Dr. David Sampson david.sampson@oregonstate.edu. Each CIE report shall be written using the format and content requirements specified in Annex 1, and address each ToR in **Annex 2**.

Schedule of Milestones and Deliverables: CIE shall complete the tasks and deliverables described in this SoW in accordance with the following schedule.

14 January 2013	CIE sends reviewer contact information to the COR, who then sends this to the NMFS Project Contact
18 January 2013	NMFS Project Contact sends the stock assessment report and background documents to the CIE reviewers.
4-13 February 2013	Each reviewer conducts an independent desk peer review
19 February 2013	CIE reviewers submit draft CIE independent peer review reports to the CIE Lead Coordinator and CIE Regional Coordinator
7 March 2013	CIE submits CIE independent peer review reports to the COR
14 March 2013	The COR distributes the final CIE reports to the NMFS Project Contact and regional Center Director

Modifications to the Statement of Work: This ‘Time and Materials’ task order may require an update or modification due to possible changes to the terms of reference or schedule of milestones resulting from the fishery management decision process of the NOAA Leadership, Fishery Management Council, and Council’s SSC advisory committee. A request to modify this SoW must be approved by the Contracting Officer at least 15 working days prior to making any permanent changes. The Contracting Officer will notify the COR within 10 working days after receipt of all required information of the decision on changes. The COR can approve changes to the milestone dates, list of pre-review documents, and ToRs within the SoW as long as the role and ability of the CIE reviewers to complete the deliverable in accordance with the SoW is not adversely impacted. The SoW and ToRs shall not be changed once the peer review has begun.

Acceptance of Deliverables: Upon review and acceptance of the CIE independent peer review reports by the CIE Lead Coordinator, Regional Coordinator, and Steering Committee, these reports shall be sent to the COR for final approval as contract deliverables based on compliance with the SoW and ToRs. As specified in the Schedule of Milestones and Deliverables, the CIE shall send via e-mail the contract deliverables (CIE independent peer review reports) to the COR (William Michaels, via William.Michaels@noaa.gov).

Applicable Performance Standards: The contract is successfully completed when the COR provides final approval of the contract deliverables. The acceptance of the contract deliverables shall be based on three performance standards:

- (1) The CIE report shall be completed with the format and content in accordance with **Annex 1**,
- (2) The CIE report shall address each ToR as specified in **Annex 2**,

(3) The CIE reports shall be delivered in a timely manner as specified in the schedule of milestones and deliverables.

Distribution of Approved Deliverables: Upon acceptance by the COR, the CIE Lead Coordinator shall send via e-mail the final CIE reports in *.PDF format to the COR. The COR will distribute the CIE reports to the NMFS Project Contact and Center Director.

Support Personnel:

William Michaels, Program Manager, COR
NMFS Office of Science and Technology
1315 East West Hwy, SSMC3, F/ST4, Silver Spring, MD 20910
William.Michaels@noaa.gov Phone: 301-427-8155

Manoj Shivlani, CIE Lead Coordinator
Northern Taiga Ventures, Inc.
10600 SW 131st Court, Miami, FL 33186
shivlanim@bellsouth.net Phone: 305-383-4229

Roger W. Peretti, Executive Vice President
Northern Taiga Ventures, Inc. (NTVI)
22375 Broderick Drive, Suite 215, Sterling, VA 20166
RPeretti@ntvifederal.com Phone: 571-223-7717

Key Personnel:

NMFS Project Contact:

Julie Neer, SEDAR Coordinator
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405
julie.neer@safmc.net Phone: 843-571-4366

Annex 1: Format and Contents of CIE Independent Peer Review Report


1. The CIE independent report shall be prefaced with an Executive Summary providing a concise summary of the findings and recommendations, and specify whether the science reviewed is the best scientific information available.
2. The main body of the reviewer report shall consist of a Background, Description of the Individual Reviewer's Role in the Review Activities, Summary of Findings for each ToR in which the weaknesses and strengths are described, and Conclusions and Recommendations in accordance with the ToRs. The CIE independent report shall be a stand-alone document for others to understand the weaknesses and strengths of the science reviewed. The CIE independent report shall be an independent peer review of each ToRs.
3. The reviewer report shall include the following appendices:
 - Appendix 1: Bibliography of materials provided for review
 - Appendix 2: A copy of the CIE Statement of Work

Annex 2: Terms of Reference for the Peer Review

SEDAR 30 Caribbean blue tang and queen triggerfish assessment review

1. Evaluate the data used in the assessment, addressing the following:
 - a) Are data decisions made by the Assessment Workshop sound and robust?
 - b) Are data uncertainties acknowledged, reported, and within normal or expected levels?
 - c) Are data applied properly within the assessment model?
 - d) Are input data series reliable and sufficient to support the assessment approach and findings?
2. Evaluate the methods used to assess the stock, taking into account the available data.
 1. Are methods scientifically sound and robust?
 2. Are assessment models configured properly and used consistent with standard practices?
 3. Are the methods appropriate for the available data?
3. Evaluate the assessment findings with respect to the following:
 - a) Are abundance, exploitation, and biomass estimates reliable, consistent with input data and population biological characteristics, and useful to support status inferences?
 - b) Is the stock overfished? What information helps you reach this conclusion?
 - c) Is the stock undergoing overfishing? What information helps you reach this conclusion?
 - d) Is there an informative stock recruitment relationship? Is the stock recruitment curve reliable and useful for evaluation of productivity and future stock conditions?
 - e) Are the quantitative estimates of the status determination criteria for this stock reliable? If not, are there other indicators that may be used to inform managers about stock trends and conditions?
4. Evaluate the stock projections, addressing the following:
 - a) Are the methods consistent with accepted practices and available data?
 - b) Are the methods appropriate for the assessment model and outputs?
 - c) Are the results informative and robust, and useful to support inferences of probable future conditions?

- d) Are key uncertainties acknowledged, discussed, and reflected in the projection results ?
5. Consider how uncertainties in the assessment, and their potential consequences, are addressed.
 - Comment on the degree to which methods used to evaluate uncertainty reflect and capture the significant sources of uncertainty in the population, data sources, and assessment methods
 1. Ensure that the implications of uncertainty in technical conclusions are clearly stated.
 6. Consider the research recommendations provided by the Assessment workshop and make any additional recommendations or prioritizations warranted.
 - Clearly denote research and monitoring that could improve the reliability of, and information provided by, future assessments.
 - Provide recommendations on possible ways to improve the SEDAR process.
 7. Provide guidance on key improvements in data or modeling approaches which should be considered when scheduling the next assessment.



SEDAR

Southeast Data, Assessment, and Review

SEDAR 30

U.S. Caribbean Queen Triggerfish

SECTION IV: Research Recommendation

March 2013

SEDAR
4055 Faber Place Drive, Suite 201
North Charleston, SC 29405

Recommendations from the Assessment Process

The ability to utilize length-frequency data is contingent upon having reliable estimates of life history parameters (von Bertalanffy parameters in particular). Studies on basic life history (e.g. age-growth relationships and estimating natural mortality) in the US Caribbean will greatly enhance the utility of the existing length-frequency data and should provide the greatest benefit to providing management advice in the short term. This should be placed as a top priority for key species.

Fishery-independent surveys should be considered as a top research priority for additional data collection. Fishery-independent surveys designed using a rigorous statistical framework will allow for the collection of species-specific catch and effort data that can be used to develop indices of abundance. Indices of abundance are used in stock assessments to inform models about how a population may be changing over time. Fishery-independent surveys can also be used to supplement existing programs by collecting age, length, weight, and reproductive data.

It is essential that continued efforts to improve the data collection of fishery-dependent catch and effort statistics be made. More specifically, continued efforts to collect species-specific catch statistics will be important for future assessments.

During the assessment workshop, the fishers from the USVI indicated that the selectivity pattern for queen triggerfish violated the assumption of knife-edge selectivity in the mean-length model. Efforts should be made to expand this model to accommodate other selectivity patterns.

Research Recommendations from the CIE Reviewer Reports (Term of Reference 6)

Cardinale Recommendations

The assessment team does provide an exhaustive shopping list for future data to be collected, which would greatly improve the capability of assessing the status of the Caribbean queen triggerfish and blue tang stock. However, I also suggest that effort should be devoted to selectivity experiments aimed to evaluate the theoretical changes in selectivity linked with the historical changes in the mesh size of the traps.

Chen Recommendations:

The AW panel recommends improving the quality of life history parameter estimates; developing a fishery-independent monitoring program; continuing the efforts to improve the collection of species-specific catch and effort data; and modifying the length-based total mortality estimator to account for potential changes in selectivity. I consider these research areas are important for reducing the uncertainty and improving the quality of the assessment. The AW panel probably needs to prioritize the research recommendations and separate the short-term research plan from the long-term plan.

Given the problems associated with the data, an important research goal should be to improve the data quality and quantity. Short-term and long-term plans should be developed to achieve the goal. Short term research priority may include (1) improvement of life history data estimates and the quantification of their uncertainty in the form of probably distributions; (2) identification of major fishing areas and their spatio-temporal variability via conducting interviews with fishermen involved in the fishery; and (3) identification of potential approaches that can be used to estimate species-specific landing data (e.g., based on species composition of landings that become available in recent years). The long-term research plan should include the development of a fishery-independent monitoring program and continued improvement of the sampling protocol for the collection of fishery-dependent data (catch and effort).

Given the data limitations, I believe another research priority that should be addressed soon is to evaluate the performance of the length-based estimator (Gedamke and Hoenig 2006) for the total mortality. Based on the information available and with some assumptions, a queen triggerfish fishery can be simulated, following the approach used in Gedamke and Hoenig (2006). A simulation study can be conducted with this simulated fishery to evaluate the performance of this length-based estimator for estimating the total mortality. Different scenarios can be developed to identify key factors that may have significant impacts on the performance of the estimator. This can guide the future model development and data collection.

Trzcinski Recommendations:

I set this out more generally below. I cannot prioritize these well because I do not know 1) the species biology, ecosystem and fishery well, 2) the long-term assessment goals or 3) the financial constraints. I think the assessment team would benefit from a meeting to discuss these issues and help set out the overall assessment framework. But if prioritizing is at all useful given my limited knowledge, I would work on getting a fisheries independent survey together, I might even do this over the life history work although that should be done as well.

Presumably some discussion occurred about whether the method used was the best given the available data. I think it is important to review and recapitulate that argument in the introduction to the assessment report.