


Review of Operating Model Parameters for SEDAR 49: Snowy Grouper

Skyler R. Sagarese, J. Jeffery Isely, and Matthew W. Smith

SEDAR49-AW-09

Submitted: 12 August 2016



This information is distributed solely for the purpose of pre-dissemination peer review. It does not represent and should not be construed to represent any agency determination or policy.

Please cite this document as:

Sagarese, S. R., J. J. Isely, and M. W. Smith. 2016. Review of Operating Model Parameters for SEDAR 49: Snowy Grouper. SEDAR 49-AW-09. SEDAR, North Charleston, SC. 17 pp.

Review of Operating Model Parameters for SEDAR 49: Snowy Grouper

Skyler R. Sagarese, J. Jeffery Isely, and Matthew W. Smith
Southeast Fisheries Science Center
75 Virginia Beach Drive
Miami, FL, USA 33149

The first step in a DLMtool data-limited assessment is the development of an operating model (OM) that describes the “true” simulated population dynamics covering the stock and fishing fleet of interest. Within a simulation framework, the operating model represents the biological components of the system to be managed, fisher behavior in response to management actions, how data are collected from the management system, and environmental conditions as well as interactions (Kell et al. 2007; Carruthers et al. 2014; Punt et al. 2014).

During the SEDAR 49 Data Workshop, multiple working groups were convened to review available data and provide recommendations of appropriate life history, stock dynamics and fleet characterizations to aid in parameterizing the OMs. Substantial efforts were also undertaken prior to the workshop which enabled discussion of available literature, data sources, and their reliability at the workshop. A comprehensive literature review of life history parameters was conducted prior to the workshop, with details provided in Adams et al. (2016). A critical component to the Data Workshop was the participation of federal, state, and industry experts in characterizing both stock and fleet dynamics for each of the eight species selected for SEDAR49. Fishermen provided keen insight into potential issues including species misidentification, selectivity patterns, discard mortality, and ecosystem considerations (see SEDAR 49 DW Report, Section 10).

In this working document we describe the parameters recommended and justifications for representing both stock and fleet dynamics within the operating model for Snowy Grouper (*Epinephelus niveatus*). Although a range of data limitations were discussed at the Data Workshop, our simulation analyses should be able address whether applied data-limited methods perform well under the specified life history type, which is expected to represent Snowy Grouper. It is assumed that the operating model under development, as specified in this document, represents reality and reflects the best available science at this point in time (Table 1). The respective distributions of parameters are provided in Table 2.

Literature Cited:

- Adams, M. S., S. R. Sagarese, and A. B. Rios. 2016. Gulf of Mexico Data-Limited Species Life History Compilation. SEDAR49-DW-05, SEDAR, North Charleston, SC. 36 pp.
- Carruthers, T. R., and A. Hordyk. 2016. Package 'DLMtool', version 3.2. 132 pp. Available from: <https://cran.r-project.org/web/packages/DLMtool/>
- Carruthers, T. R., A. E. Punt, C. J. Walters, A. MacCall, M. K. McAllister, E. J. Dick, and J. Cope. 2014. Evaluating methods for setting catch limits in data-limited fisheries. *Fisheries Research* 153:48-68. doi: 10.1016/j.fishres.2013.12.014
- Kowal, K. 2010. Aspects of the Life History of the Snowy Grouper, *Epinephelus niveatus*, in the Gulf of Mexico. University of South Florida.
- Kell, L. T., I. Mosqueira, P. Grosjean, J.-M. Fromentin, D. Garcia, R. Hillary, E. Jardim, S. Mardle, M. Pastoors, and J. Poos. 2007. FLR: an open-source framework for the evaluation and development of management strategies. *ICES Journal of Marine Science* 64:640-646. doi: 10.1093/icesjms/fsm012
- Punt, A. E., D. S. Butterworth, C. L. Moor, J. A. De Oliveira, and M. Haddon. 2014. Management strategy evaluation: best practices. *Fish and Fisheries*. doi: 10.1111/faf.12104
- SEDAR. 2013. SEDAR 36: South Atlantic Snowy Grouper Stock Assessment Report. SEDAR, North Charleston, SC. 146 pp.
- Then, A. Y., J. M. Hoenig, N. G. Hall, and D. A. Hewitt. 2014. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES Journal of Marine Science: Journal du Conseil*:fsu136.
- Thorson, James T., Jason M. Cope, Trevor A. Branch, and Olaf P. Jensen. 2012. Spawning biomass reference points for exploited marine fishes, incorporating taxonomic and body size information. *Canadian Journal of Fisheries and Aquatic Sciences* 69:1556-1568. doi: 10.1139/f2012-077
- Zhou, S., S. Yin, James T. Thorson, Anthony D. M. Smith, and M. Fuller. 2012. Linking fishing mortality reference points to life history traits: an empirical study. *Canadian Journal of Fisheries and Aquatic Sciences* 69:1292-1301. doi: 10.1139/f2012-060

Table 1. Operating model parameters for Snowy Grouper. Decisions made at the Data Workshop are color coded for ease of interpretation (red = Life History Working Group recommendation; green = Commercial/Recreational Working Group recommendation; blue = Index Working Group recommendation). Parameters with no color indicate decisions made by the analysts.

OM input	Description	Value or CV	Source
Life-history			
MaxAge	Maximum age of individuals that are simulated (no plus group)	35 y	Maximum age observed in meta-analysis (SEDAR36 2013)
R0	Magnitude of unfished recruitment (scaling factor). Normally fixed to some arbitrary value since it simply scales the simulated numbers. Typically can be set to 1000. (Carruthers and Hordyk 2016)	1000	Default value
M	Natural mortality rate	$0.153 - 0.189 \text{ y}^{-1}$	Ranged recommended based on plausible maximum ages (35 - 44 y) in reliable literature (SEDAR 2013; Kowal 2010)
Msd	Inter-annual variability in M expressed as a coefficient of variation	c(0,0)	Turned off for now
Mgrad	Mean temporal trend in M, expressed as a percent change in M per year	c(0,0)	Turned off for now
h	Recruitment compensation (steepness)	0.74 – 0.94	Range considered in SEDAR36 for Snowy Grouper (Adams et al. 2016)
SRrel	Type of stock-recruitment relationship (1=BH, 2=Ricker)	1	Adams et al. 2016
Linf	Von Bertalanffy asymptotic size	106.5 – 108.6 cm TL	Ranged recommended based on reliable literature (converted Kowal (2010) Linf from FL to TL, likely prone to errors)

SEDAR49: Gulf of Mexico Snowy Grouper Operating Model Parameters

Linf _{sd}	Inter-annual variability in Linf expressed as a coefficient of variation	c(0,0)	Turned off for now
Linf _{grad}	Mean temporal trend in Linf, expressed as a percent change in M per year	c(0,0)	Turned off for now
K	Von Bertalanffy maximum growth rate	0.077 – 0.111	95% Confidence interval in other reliable literature (Kowal 2010)
K _{sd}	Inter-annual variability in K expressed as a coefficient of variation	c(0,0)	Turned off for now
K _{grad}	Mean temporal trend in K, expressed as a percent change in M per year	c(0,0)	Turned off for now
vbt0	Von Bertalanffy theoretical age at length zero	-1.88 – -3.19	95% Confidence interval in other reliable literature (Kowal 2010)
a	Length-weight parameter a	3.56E-05	Value from SEDAR49 data analysis for Max TL (cm) to W W _t (lbs)
b	Length-weight parameter b	2.98	Value from SEDAR49 data analysis for Max TL (cm) to W W _t (lbs)
D	Current level of stock depletion (B _{now} / B _{unfished})	c(0.05,0.30)	Estimates of depletion over last three years for Yellowedge Grouper range from 0.30 to 0.30
L50	Length at which individuals are 50% mature	58.0 – 62.0 cm TL	SEDAR36 - length where 50% maturity falls
L50_95	Length increment from 50% to 95% maturity (L95-upper L50,L95-lower L50)	c(13.0, 17.0)	75.0 cm TL (Table 1, SEDAR36 2013)
rec _{grad}	Mean slope in recruitment deviations	c(0,0)	Turned off for now
Perr	Process error in recruitment deviations	0.55 – 0.55	Fixed in SEDAR36 for Snowy Grouper

SEDAR49: Gulf of Mexico Snowy Grouper Operating Model Parameters

			(Adams et al. 2016)
AC	Autocorrelation in recruitment deviations	c(0.27,0.32)	Range derived from SEDAR22 Yellowedge Grouper assessment. Autocorrelation was investigated between: (1) estimated recruits and (2) recruitment deviations. No YOY index available
Frac_area_1	Fraction of unfished biomass in area 1 at start of simulation	c(0.095, 0.105)	Default value
Prob_staying	Probability that individuals in area 1 stay there in year	c(0.5, 0.6)	Default value
Fleet			
nyears	Number of years for historical simulation. Should be set as close as possible to the length of time that the fishery has been exploited.	45	Coincident with ramp-up of deep fishing (in 1970s)
Spat_targ	Distribution of fishing in relation to spatial biomass. 1 = fishers are indiscriminate in where they fish (e.g., bycatch species); > 1 indicates targetting areas of higher biomass	c(1,1)	Default value
LFS	Length at full selectivity (LFS/L50) for representative fleet	c(0.94, 1.00)	58.0 cm; DW Report, Section 8.2
L5	Length at 5% selectivity (LFC/L50) for representative fleet	c(0.68, 0.72)	42.0 cm; DW Report, Section 8.2
Vmaxlen	Vulnerability of oldest age class to representative fleet (controls extent of dome-shaped selectivity)	c(1,1)	Asymptotic (DW Report, Table 8.5.2)

SEDAR49: Gulf of Mexico Snowy Grouper Operating Model Parameters

Fsd	Interannual variability in F - determines how much F fluctuates from year to year	c(0.00,0.14)	Range of interannual variability in annual F for dominant fleet ("representative") from SEDAR22 for Yellowedge Grouper
qinc	Mean percentage change in fishing efficiency	c(0,0)	Turned off for now
qcv	Interannual variability in fishing efficiency	c(0,0)	Turned off for now
EffYears	Index of effort	Commercial Longline	DW Report, Section 6.2.5
Observation			
LenMcv	Bias in length at 50% vulnerability	0.2	Default value for biased, imprecise
Cobs	Log-normal catch observation error	c(0.11, 0.11)	DW Report, Section 5.2.5
Cbiascv	CV controlling the sampling of bias	0.11	DW Report, Section 5.2.5
CAA_nsamp	Number of catch-at-age observations per time step	c(150, 200)	General range of annual composition samples desired; or could use default for biased, imprecise (50,100)
CAA_ESS	Effective sample size	c(10,20)	Default for biased, imprecise
CAL_nsamp	Number of catch-at-length observations per time step	c(150, 200)	General range of annual composition samples desired; or could use default for biased, imprecise (50,100)
CAL_ESS	Effective sample size	c(10,20)	Default for biased, imprecise
CALcv	Lognormal variability in length at age	c(0.29,0.29)	Maximum CV derived from length data for the

SEDAR49: Gulf of Mexico Snowy Grouper Operating Model Parameters

			representative fleet by year
Iobs	Observation error in relative abundance index expressed as a CV	c(1,1)	Not available, assuming high variability
Icv	Observation error in relative abundance index expressed as a CV	1	Not available, assuming high variability
Mcv	Bias in M sampled from a log-normal distribution with a CV	0.32	Cross-validation prediction error of updated Hoenig (Then et al. 2014); Carruthers et al. (2014) used 0.5
Lincv	Bias in Linf sampled from a log-normal distribution with a CV	0.06	SEDAR36 2013: mean = 1065 mm TL, SE = 65.22
Kcv	Bias in K sampled from a log-normal distribution with a CV	0.22	SEDAR36 2013: mean = 0.094, SE = 0.021
t0cv	Bias in t0 sampled from a log-normal distribution with a CV	0.33	SEDAR36 2013: mean = -2.88, SE = 0.951
LFCcv	Bias in length at first capture sampled from a log-normal distribution with a CV	0.03	DW Report, Section 10.4
LFScv	Bias in length at full selection sampled from a log-normal distribution with a CV	0.21	DW Report, Section 10.4
B0cv	Bias in unfished biomass sampled from a log-normal distribution with a CV	4	Default for biased, imprecise
FMSYcv	Bias in FMSY sampled from a log-normal distribution with a CV	0.2	Default for biased, imprecise
FMSY_Mcv	Bias in FMSY/M sampled from a log-normal distribution with a CV	0.11	From meta-analysis (Zhou et al. 2012); Carruthers et al. (2014) used 0.2
BMSY_B0cv	Bias in BMSY/B0 sampled from a log-normal distribution with a CV	0.14	From meta-analysis (Thorson et al. 2012);

SEDAR49: Gulf of Mexico Snowy Grouper Operating Model Parameters

			Carruthers et al. (2014) used 0.2 (default for biased, imprecise)
rcv	Bias in intrinsic rate of increase sampled from a log-normal distribution with a CV	0.5	Default for biased, imprecise (used in Carruthers et al. 2014)
Dbiascv	Bias in stock depletion sampled from a log-normal distribution with a CV	1	Carruthers et al. (2014) used 1.0
Dcv	Imprecision in stock depletion among years expressed as a CV	c(0.05,0.20)	Default for biased, imprecise
Btbias	Bias in current stock biomass sampled from a log-normal distribution with a CV	c(0.2,5.0)	Default for biased, imprecise
Btcv	Imprecision in current stock biomass expressed as a CV	c(0.2,0.5)	Default for biased, imprecise
Fcurbiascv	Bias in current F sampled from a log-normal distribution with a CV	0.75	Default for biased, imprecise
Fcurcv	Imprecision in current F among years expressed as a CV	c(0.5,1.0)	Default for biased, imprecise
hcv	Bias in knowledge of steepness	0.12	Maximum value of: $\text{abs}[(\text{range estimate} - \text{point estimate}) / \text{point estimate}]$
maxagecv	Bias in maximum age (not currently used)	0.26	Maximum value of: $\text{abs}[(\text{range estimate} - \text{point estimate}) / \text{point estimate}]$
Reccv	Bias in recent recruitment strength	c(0.1,0.3)	Default for biased, imprecise
Irefcv	Bias in relative abundance index at BMSY	0.3	Default for biased, imprecise
Crefcv	Bias in MSY	0.3	Default for biased, imprecise
Brefcv	Bias in BMSY	0.5	Default for biased, imprecise
beta	Parameter controlling hyperstability (< 1)/hyperdepletion (>1)	c(0.33, 3.0)	Default for biased, imprecise

Table 2. Summary of distributions assumed for parameters within the stock, fleet, and observation subclasses within the operating model. Input parameters are as defined in Table 1.

Input	Distribution	Input	Distribution
Stock subclass		Observation subclass	
MaxAge	-	LenMcv	Log-normal
R0	-	Cobs	Uniform
M	Uniform	Cbiascv	Log-normal
sd	Uniform	CAA_nsamp	Uniform
grad	Uniform	CAA_ESS	Uniform
h	Uniform	CAL_nsamp	Uniform
SRrel	-	CAL_ESS	Uniform
Linf	Uniform	CAL_cv	-
sd	Uniform	Iobs	Uniform
grad	Uniform	Mcv	Log-normal
K	Uniform	Kcv	Log-normal
sd	Uniform	t0cv	Log-normal
grad	Uniform	Linfcv	Log-normal
t0	Uniform	LFCcv	Log-normal
a	-	LFScv	Log-normal
b	-	B0cv	Log-normal
D	Uniform	FMSYcv	-
L50	Uniform	FMSY_Mcv	Log-normal
L50_95	Uniform	BMSY_B0cv	Log-normal
recgrad	Uniform	rcv	-
Perr	Uniform	Dbiascv	Log-normal
AC	Uniform	Dcv	Uniform
Frac_area_1	Uniform	Btbias	Uniform
Prob_staying	Uniform	Btcv	Uniform
Fleet subclass		Fcurbiascv	-
nyears	-	Fcurecv	-
Spat_targ	Uniform	hcv	-
LFS	Uniform	Icv	-
L5	Uniform	maxagecv	-
Fgrad	-	Reccv	Uniform
qinc	Uniform	Irefcv	Log-normal
qcv	Uniform	Crefcv	Log-normal
Vmaxlen	Uniform	Brefcv	Log-normal
		beta	Uniform

Figure 1. Sampled parameters in the operating model for life history. Parameters include: M = natural mortality, L_{∞} = von Bertalanffy asymptotic size, K = von Bertalanffy growth rate, t_0 = von Bertalanffy theoretical age at length zero, age_M = age at 50% maturity (derived from length at 50% maturity), len_M = length at 50% maturity, len_{95} = length at 95% maturity, Steepness = steepness of the stock recruitment curve, Recr Devs = recruitment deviations.




Figure 2. Sampled parameters in the operating model for the fishery. Parameters include: L5 = shortest length corresponding to 5% vulnerability, LFC = length at first capture, LFS = length at full selection, Frac_area_1 = fraction of individuals found in area 1 of a generic two-area model of the simulation framework, Probab_staying = probability that individuals in area 1 remain in that area, F = fishing mortality, Recr Devs = recruitment deviations.




Figure 3. Sampled parameters in the operating model for the reference parameters. Parameters include: RefY = Reference yield (highest long-term yield obtained from fixed F strategy), MSY = maximum sustainable yield, OFLreal = true simulated overfishing limit, BMSY_B0 = most productive stock size relative to unfished, FMSY = fishing mortality rate at maximum sustainable yield, FMSY_M = fishing mortality rate divided by natural mortality rate.




Figure 4. Sampled parameters in the observation model for life history. Parameters as defined in Figure 1 caption, with exception of hbias = bias in steepness parameter. Bias refers to an inaccuracy in an observed parameter that occurs for the duration of a simulation.




Figure 5. Sampled parameters in the observation model for fishery. Parameters as defined in Figure 2 caption, with exception of Cbias = bias in observed catches. Bias refers to an inaccuracy in an observed parameter that occurs for the duration of a simulation.




Figure 6. Sampled parameters in the observation model for the index of abundance. Betas is a parameter controlling hyperstability/hyperdepletion.




Figure 7. Sampled parameters in the observation model for the size and composition. Parameters include: CAA_nsamp = number of catch-at-age observations per time step, CAA_ESS = effective sample size (independent age draws) of the multinomial catch-at-age observation error model, CAL_nsamp = number of catch-at-length observations per time step, and CAL_ESS = effective sample size (independent age draws) of the multinomial catch-at-length observation error model.




Figure 8. Sampled parameters in the observation model for the reference. Parameters as defined in Figure 3 caption, with exception of D = Depletion, A = abundance, Iref = reference index level, Cref = reference catch level, and Bref = reference biomass level. Bias refers to an inaccuracy in an observed parameter that occurs for the duration of a simulation.

