

Survey of the Florida recreational shark fishery utilizing shark tournament and selected longline data.
Final report to Florida Department of Natural Resources, Mote Marine Laboratory Tech. Rpt. 232A:94pp.

SEDAR65- RD07

Robert Hueter

Received: 11/1/19



SURVEY OF THE FLORIDA
RECREATIONAL SHARK FISHERY
UTILIZING SHARK TOURNAMENT
AND SELECTED LONGLINE DATA

FINAL REPORT
FDNR Grant Agreement 6627

Submitted to:

Florida Department of Natural Resources
100 Eighth Avenue Southeast
St. Petersburg, Florida 33701

Submitted by:

Mote Marine Laboratory
1600 Thompson Parkway
Sarasota, Florida 34236
(813) 388-4441



Robert E. Hueter, Ph.D.
Staff Scientist

MOTE MARINE LABORATORY TECHNICAL REPORT 232A

September 1, 1991

ABSTRACT

A survey of shark tournament activity and a comparative database of inshore longlining operations in the northeastern Gulf of Mexico were used to examine historical trends in the Florida recreational shark fishery over the last two decades. Over 200 shark tournaments involving more than 22,500 anglers at 34 different sites have been held in Florida waters from 1971 to 1991. Catch data from these tournaments in terms of numbers and size of sharks and catch per unit effort were analyzed for trends. The longline data contained catch records on 659 sharks from 13 species collected from 1975 to 1989. Both sets of data strongly indicate significant declines in numbers of large, inshore sharks in Florida waters in the late 1980's.

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	2
TABLE OF CONTENTS	3
LIST OF FIGURES	4
LIST OF TABLES	5
INTRODUCTION	6
Recreational Shark Fisheries	6
Commercial Shark Fisheries	7
The Need for Historical Shark Fishery Information	10
OBJECTIVES OF THE STUDY	13
METHODS	14
Shark Sportfishing Tournaments	14
Crooke Longline Data	16
DATA	18
Shark Sportfishing Tournaments	18
Tournament Overview	18
Temporo-Spatial Distribution of Tournament Activity	23
Catch Data	27
Species Distribution/Abundance	44
Crooke Longline Data	47
Raw Data	47
Species Profiles	47
Historical Trends in Size of Sharks	58
Catch per Unit Effort	67
Summary	67
DISCUSSION	69
Historical Trends in Florida Shark Tournaments	69
Historical Trends in the Crooke Longline Data	71
Implications for Local Shark Populations and the Florida Shark Fishery	72
ACKNOWLEDGMENTS	73
REFERENCES	74
APPENDICES	
APPENDIX 1. Crooke longline data arranged chronologically.	A1.1-10
APPENDIX 2. Crooke longline data arranged by species.	A2.1-10

LIST OF FIGURES

	<u>Page</u>
Figure 1. Commercial landings of sharks in Florida, 1976-89.	9
Figure 2. Distribution of commercial shark landings in Florida, 1989.	11
Figure 3. Location and size of Florida shark tournaments, 1971-91.	24
Figure 4. Number of Florida shark tournaments by year, 1971-91.	26
Figure 5A-G. Distribution of Florida Gulf shark tournaments, 1980-91.	28-34
Figure 6. Locations of 7 shark tournaments and Crooke longlining.	36
Figure 7A-E. Catch data for 7 shark tournaments.	38-42
Figure 8A-G. Number of sharks vs. month in Crooke data.	49-55
Figure 9. Temperature ranges for shark species from Crooke data.	56
Figure 10A-G. Shark length vs. year for Crooke data, by species.	59-65
Figure 11. Shark length vs. year for Crooke data, all species combined.	66
Figure 12. CPUE for successful sets of Crooke longline.	68

LIST OF TABLES

	<u>Page</u>
Table 1. Florida shark sportfishing tournaments, 1971-91.	19-22
Table 2. Shark species of Florida coastal waters.	25
Table 3. Logbook records summary for Florida tournaments.	37
Table 4. Catch by species categories in Florida tournaments.	45-46
Table 5. Crooke longline catch by species and sex.	48

INTRODUCTION

The fisheries biology of sharks inhabiting U.S. coastal waters of the Atlantic Ocean and Gulf of Mexico has received relatively little attention from researchers or fishery agencies in the past. Traditionally treated as undesirable species, sharks have often seemed to "fall between the cracks" of fishery resource management. Recently, however, the exploitation of sharks in waters of the southeast U.S. has skyrocketed. Florida in 1991 is now the home of a major recreational and commercial fishery for sharks. With the lack of historical, fisheries-related studies on sharks, the impact of this increased fishing pressure on shark populations has been difficult to assess.

Recreational Shark Fisheries

Through the Marine Recreational Fishery Statistics Survey (MRFSS), the National Marine Fisheries Service (NMFS) has estimated that in 1979 over 1.2 million sharks were caught by recreational fishermen in southeast U.S. waters from North Carolina to Texas. Of this number, an estimated 37%, or about 459,000 sharks, were landed and the remainder were released. In 1986, NMFS estimated nearly 1.2 million sharks again were caught by recreational fishermen in the same region (NMFS, 1984, 1987). By 1989, the last year for which the MRFSS data are currently available, the catch of sharks was 774,000 from North Carolina to Louisiana (Texas data are excluded from the MRFSS after 1986) (NMFS, 1991).

In Florida waters alone, the estimated number of sharks caught by recreational fishermen increased from 450,000 in 1979 to 733,000 in 1986--a 63% increase in shark fishing activity in the state in just seven years. Nearly all

(96%) of this increase in activity was attributed to anglers fishing in Gulf of Mexico waters. With an estimated 560,000 sharks caught in the Florida Gulf in 1986, and with approximately 800 miles of Gulf coastline, the estimated recreational catch in 1986 averages 2 sharks caught per mile of coastline per day in the Florida Gulf.

The increasing desirability for sharks among U.S. recreational fishermen in the 1980's was not a transient fad. The release of the movie *Jaws* in the summer of 1975 was no doubt partly responsible for this rise in popularity, but the fishing trend outlived the movie. In essence, shark fishermen are not a specialty group but rather fish for the same reasons as other fishermen (Graefe and Ditton, 1976). Recreational fishermen from all economic levels discovered in the late 1970's that sharks are accessible, enjoyable to catch, and good to eat with proper handling. Many anglers switched to sharks as stocks of other finfish were reduced.

After 1986, however, recreational catches of sharks in Florida began to fall off. By 1989, the estimated recreational catch had dropped to 392,000, a decrease of 47% from the 1986 Florida catch (NMFS, 1987, 1991). Although the MRFSS data are estimates based on statistical projections from creel censuses rather than total counted landings, the relative changes indicate two features: a significant increase in recreational fishing activity for sharks in Florida through the early and mid-1980's, followed by a downturn in catch and landings of sharks in the late 1980's.

Commercial Shark Fisheries

After recreational fishing pressure on sharks increased, the commercial fishery for sharks in the southeast U.S. also expanded. In the 1980's, this

fishery grew at an exponential rate to reach an all-time high in the annual landings of shark meat. Adding to the commercial value of sharks for their meat--and propelling this fishery upward--was a substantial increase in the value of shark fins, which are currently worth as much as \$23 per pound of dried fins in the Oriental sharkfin soup market.

The build-up in the Florida commercial shark fishery began after 1979, with exponential growth occurring after 1986 when fin prices doubled (Fig. 1). In the two-year period between 1979 and 1981, commercial landings of sharks in Florida nearly quadrupled. The Florida Department of Natural Resources (FDNR) estimated that 170,740 pounds of sharks of all species were landed in 1979 by commercial fishermen on the Florida Atlantic and Gulf coasts. That figure rose to 658,245 pounds for the same region in 1981 (FDNR, 1979, 1981) and landings in-state continued to increase steadily up through 1986, when commercial landings reached approximately 2.5 million pounds. Then, in 1987, total state landings nearly doubled from the previous year to 4.6 million pounds and approached the 6 million pounds per year level in 1988 (FDNR, 1988), which translates to approximately 120,000 sharks landed in the commercial fishery in Florida alone. Since 1985, these FDNR statistics have been actual landings based on trip ticket information, and therefore can be considered to be a conservative estimate of the level of activity of the commercial shark fishery in Florida.

By 1989, sharks had become the fourth most valuable commercial finfish resource in Florida, exceeded only by the state's grouper, mullet, and snapper fisheries. Commercial shark fisheries in Florida accounted for about \$4.4 million in landed meat and fins in 1989 (swordfish and tuna landings were valued higher, but these fishes are caught in waters far outside of the Florida coastal zone). Total state landings of shark meat in 1989 reached 6.5 million pounds

REPORTED LANDINGS OF SHARKS
IN FLORIDA COMMERCIAL FISHERY
1976 - 1989




Fig. 1. Commercial landings of sharks in Florida, 1976-89. Fin prices doubled after 1986, leading to exponential growth of the fishery.

whole weight (FDNR, 1989), corresponding to over 130,000 sharks. Two-thirds of these landings were brought into Florida Gulf ports (Fig. 2). Landed fin weights in the state exceeded 200,000 pounds (FDNR, 1989), taken from an estimated 50,000 sharks, with 86% of these landed at Gulf ports.

The Need for Historical Shark Fishery Information

The National Marine Fisheries Service recently has determined that the ever-increasing demand for sharks and shark products has exceeded the reproductive capacity of some Gulf and Atlantic coast shark species. Because of a life history that includes very slow growth, late sexual maturity, and small number of offspring, sharks are especially vulnerable to overfishing and present special problems to fishery management (Holden, 1974). NMFS has now proposed, through a Secretarial Shark Fishery Management Plan, to place 38 species of sharks under management in federal waters. Furthermore, the Florida Marine Fisheries Commission (MFC) recently has proposed managing sharks in state waters as well. The federal and state management plans include commercial quotas and other limits, closed seasons, recreational bag limits, prohibition of sale of recreational catch, and a ban on shark "finning" (NMFS, 1989; MFC, 1991).

Historical fishery statistics for the shark fishery in Florida are essential both for the assessment of current fishing pressure on shark stocks and for the formulation of sound management plans. Although the FDNR trip-ticket system has become an effective means of tracking commercial landings of shark fishery products in Florida, similar analyses of the recreational fishery are hampered by a lack of studies other than broad-scale estimates, such as those furnished by the MRFSS. Basic descriptions of the in-state recreational fishery for sharks during the recent period of increased pressure are generally lacking.




Fig. 2. Distribution of commercial landings of shark meat (by pounds whole weight) in Florida ports, 1989. Source: Florida DNR.

To assess completely the shark recreational fishery in Florida in terms of catch/effort, species composition, seasonality, and various socio-economic factors, a fine-scale survey is necessary. To provide meaningful, precise, and accurate information, a very large, labor-intensive effort that is technically and logistically complicated--and very expensive--is required. However, such a study typically provides information only on the present and future fishery. Historical data may be non-existent or may be overlooked with such an approach.

A cost-effective first step to assessing historical trends in this fishery is to analyze data available through organized sportfishing tournaments (also called "derbies" or "rodeos") for sharks. Since the mid-1970's, the number of tournaments that include or are directed solely toward sharks along the U.S. Atlantic and Gulf coasts has steadily increased. Such tournaments in this region have been conservatively estimated by NMFS to number about 65 per year as of 1989 (NMFS, 1989).

Over the past decade, shark tournaments became popular as fishing contests and fund-raisers in many fishing communities of the southeast U.S. In recent years, declining catches of sharks, both in number and size, have been reported in these tournaments via anecdotal sources. Although tournament directors have pointed to years of logbook information containing catch records, no formal attempt has been made previously to compile and analyze these records on a statewide basis. In light of the relative lack of published information on Florida's recreational shark fishery, these tournament records represent important historical data.

In addition, another valuable source of historical information exists in a database of longline catches compiled by Mr. C.F. Crooke, Jr. of Warrington (near Pensacola), Florida. The Crooke data pertain to small but remarkably

consistent longlining operations for large, inshore sharks over 15 years, from 1975 through 1989. The records include species, sex, water temperature, and other information for sharks caught in the waters off Pensacola, an area with a significant recreational interest in sharks. Thus, these data can serve as a comparison to the recreational tournament data over the last two decades. No previous attempt to compile and analyze the Crooke data has been made.

OBJECTIVES OF THE STUDY

The primary objectives of this project were as follows:

- (1) Survey the recreational fishery for sharks in Florida from organized shark tournaments held during the period 1971-1991, by compiling historical and current data on tournament activity and distribution, catch, species/size/sex composition of catch, catch/effort, and other factors.
- (2) Compile and analyze the 15 years of Crooke longline data (1975-1989) in conjunction with the recreational tournament data, and evaluate the two data sets for historical trends in the Florida shark fishery.

METHODS

Shark Sportfishing Tournaments

A contact network of current tournament directors, fishermen's clubs and organizations, outdoor writers, marina operators, fishing tackle dealers and bait shops, and individual anglers was used to survey the state for shark tournament activity. All coastal counties were canvassed in person or by telephone. Individuals with local knowledge of sharkfishing were interviewed for their recollections of tournament activity over the past two decades. When a tournament was identified, every attempt was made to contact the most knowledgeable individual available to provide the best available data on that tournament. Preference to quality of information was generally given, in the following order, to: 1) original tournament director with written records; 2) other individual possessing original records; 3) tournament director or committee member without records; 4) outdoor writer with or without written information; 5) regular tournament participants; and 6) others.

From this survey, a total of 34 sportfishing tournaments devoted strictly to sharks and operating in Florida waters sometime between 1971 and 1991 was identified. (This survey does not include other tournaments that had sharks as one *category* of catch--only dedicated shark tournaments are considered.) In 25 (74%) of these, the tournament director or co-director was identified, located, and interviewed. General information was collected on each tournament as follows:

- Official name of the tournament and its base location.
- Years of operation and current status (active or discontinued).
- Number of participating anglers and/or boats per year.
- Status of tournament records.

If written catch records of a tournament were located, the status of those records fell into one of three categories: 1) records provided to this study and suitable for analysis--9 tournaments (26%); 2) records incomplete and not suitable for analysis--6 (18%); or 3) records may have been suitable but were not provided, due either to passive noncompliance or deliberate noncooperation of record curators--5 (15%). For the rest of the tournaments (14/41%), no existing records were located.

Tournaments with suitable records were analyzed for catch by year. Primary data consisted of species and size (by total weight in pounds) of catch. Unfortunately, data on sex and length of sharks caught were rarely recorded, so these could not be included in the analysis. Given that the information came from competitive tournaments, it was assumed that weight measurements were accurate. On the other hand, it was not necessarily assumed that species identifications were accurate. The record-holders were interviewed to gauge the relative accuracy of the species ID's, and adjustments in the data were made where justified. However, some errors in species identification likely exist in the tournament data set, due to the inherent difficulties that anglers have in distinguishing between closely related species of sharks.

Number of sharks caught, heaviest shark (usually the winning catch), and average weight of sharks were determined and plotted (Sigma-Plot). Catch per unit effort was calculated and plotted where possible. Trends were determined from the plotted data.

Crooke Longline Data

Mr. C.F. Crooke, Jr. of Warrington, Florida, generously provided his data collected from 15 years of longlining operations in the Florida panhandle region. Beginning in 1975 and ending in 1989, Mr. Crooke set a small longline regularly in the waters south of Pensacola, specifically to catch large sharks for supply of shark meat to local restaurants. His typical fishing year ran from late March/early April to late November/early December. Since these activities were incidental to Mr. Crooke's regular line of work, he fished usually on weekends or holidays. Through the 15 years, he averaged just over 18 successful sets of the gear per year ("successful" = at least one shark was caught and recorded), with a range of 9 successful sets in some years to 32 in his most active year (1978).

The Crooke longline was set each time on the bottom in the same location: just inside of 5 miles due south of the western end of Santa Rosa Island, near Pensacola Beach, in a depth of 50 feet of water. The gear consisted of Mustad 14/0 giant tuna hooks, baited with fresh or fresh-frozen bonito, jack crevalle, or mullet, on 5-ft. gangings placed about 20 feet apart on the groundline. From 1975 to 1986, Mr. Crooke set 20 hooks; in 1987, he increased the number of hooks to 30, which he used until he ceased operations in 1989.

The gear was typically set at sundown, preferably on an outgoing tide, and picked up the next morning at sunrise. The catch was hauled in to shore in whole condition and each shark was measured by length and identified by species and sex prior to dressing out the carcass. Notes were taken on the reproductive condition of females, stomach contents, and other aspects. Water temperature for each date was noted. All data were recorded on standardized data sheets and

maintained in chronological order in an organized file in Mr. Crooke's home. One unfortunate drawback of the database was that data were taken only when sharks were caught, i.e. only on successful sets. *Sets that caught no sharks were not recorded*, and there appears to be no way to recover this information accurately. This compromises to some extent the catch/effort data, limiting the quantitative analysis to successful sets only.

Length of each shark was measured in feet/inches by Mr. Crooke in two of three ways. In all cases, Mr. Crooke measured total length of the shark by laying a flexible measuring tape over the entire dorsal surface of the shark's body, from snout to precaudal pit to tip of tail. Since this measurement is not equivalent to "total length" typically reported by other researchers (the straight-line distance down the body axis from snout to tail tip), this type of measurement will be referred to as *total contour length* in this report. Other length measurements taken by Mr. Crooke were either: 1) true, straight-line fork length of sharks caught in 1975-78; or 2) over-the-back, contour measurement from snout to precaudal tip, excluding the tail, on sharks from 1979-89. Since *total contour length* was measured on all sharks, raw data in this length category were used in all analyses.

In all, data on 659 sharks comprised of 13 species were collected over the 15 years of longlining operations. The reliability of the species identifications is very high; beginning in the late 1960's, Mr. Crooke accumulated fishing experience with sharks of the area and educated himself on formal species identification. Consultations with shark biologists Dr. Perry Gilbert and Mr. Stewart Springer of Mote Marine Laboratory, as well as contact with other ichthyologists at nearby shark tournaments, reinforced his knowledge of shark biology.

Data from the original data sheets were transcribed and tabulated for analysis in this report. Data were plotted (Sigma-Plot) and trends were assessed from the plotted data, using simple linear regression and ANOVA procedures where applicable.

DATA

Shark Sportfishing Tournaments

Tournament Overview. A total of 34 separate tournament operations, each one organized and conducted as a distinct, usually multi-year tournament for sharks, was held in Florida state waters between 1971 and 1991 (Table 1). (Two of these are based on the Florida border in Orange Beach, Alabama, but are included in this survey because they involve Florida anglers fishing in state waters.) The average lifespan of these 34 tournaments is 6.2 yrs ($s = 4.8$), with a range of 1 yr of operation (one tournament begun in 1991) to a maximum of 18 yrs of continuous operation (one tournament discontinued in 1989). In all, 209 shark tournaments involving approximately 22,560 participants were held in state waters in the past 20 years.

Unlike the huge events held in northeast U.S. waters, such as the annual Bayshore and Montauk shark tournaments on Long Island in New York, shark

Table 1. Florida shark sportfishing tournaments, 1971-1991.

Format:

Name of tournament, location
No. years of operation, actual years held [Status]
No. boats/anglers per yr average
Status of tournament records
Tournament director/Contact person (phone no.)

I. PANHANDLE REGION

1. Monster Shark Tournament, Orange Beach, AL
3 yrs (89-91) [Active]
10 boats/yr -- 35-45 anglers/yr
Records provided
G. Myers (205-981-9361)
2. Perdido Pass Shark Tournament, Orange Beach, AL
4 yrs (88-91) [Active]
20 boats/yr -- 50-80 anglers/yr
Records provided
K. Eberly (205-981-6481)
3. Pensacola (Biggs) Shark Rodeo, Pensacola, FL
6 yrs (76-81) [Discontinued 81]
180-250 anglers/yr
Incomplete records
C.F. Crooke (904-455-4157)
4. Destin Shark Tournament, Destin, FL
16 yrs (76-91) [Active]
50 boats/yr -- 100-200 anglers/yr
Records provided
M. Eller (904-837-4944)
5. Panama City Shark Tournament, Panama City, FL
12 yrs (80-91) [Active]
50-100 anglers/yr
Records not provided (curator noncooperative)
B.J. Putnam (904-234-2621)

II. CENTRAL GULF COAST REGION (Apalachicola to Tampa Bay)

1. Suwannee Shark Tournament, Suwannee, FL
2 yrs (81,83) [Discontinued 83]
10-20 anglers/yr
No records
R. Riemenschneider (904-542-8380)
2. Shark Hunt, Homosassa, FL
3 yrs (89-91) [Active]
50 anglers/yr
No records
J. Slaght (904-628-2474)

Table 1. (continued)

3. Hernando Beach Shark Tournament, Hernando Beach, FL
4 yrs (88-91) [Active]
100-150 anglers/yr
Records not provided
D. Matusik (904-596-8096)
4. Port of Hudson Shark Tournament, Hudson, FL
7 yrs (85-91) [Active]
100-300 (200 avg.) anglers/yr
Records provided
R. Ufnal (813-868-0512)
5. Sportsman's Marina Shark Tournament, Port Richey, FL
2 yrs (87-88) [Discontinued 88]
100 anglers
No records
H. Smith (813-584-7919)
6. Greater Suncoast Shark Tournament, Clearwater, FL
2 yrs (83-84) [Discontinued 84]
400-500 anglers/yr
No records
E. Levy (813-442-9252)
7. Tampa Bay Sharkers Tournament, Tampa/St. Petersburg, FL
13 yrs (78-90) [Discontinued 90]
200 anglers/yr
Records provided
J. Moreland (813-272-3425)
8. O'Neill's Shark Tournament, St. Petersburg, FL
11 yrs (79-89) [Discontinued 89]
100-200 anglers/yr
No records
A. Phillips (813-867-2585)
9. Inter-Club Tampa Bay Shark Tourn., Tampa Bay-Sarasota, FL
5 yrs (82-86) [Discontinued 86]
5-club competition, approx. 100 anglers/yr
No records
H. Smith (813-584-7919)

III. SOUTHWEST GULF COAST (S. of Tampa Bay to Florida Bay)

1. Sarasota Shark Rodeo, Sarasota, FL
5 yrs (83-87) [Discontinued 87]
50-80 anglers
Incomplete records
B. Lamb (813-921-6006)
2. Gulf Coast Shark Census Tournament, Sarasota, FL
3 yrs (89-91) [Active]
120 anglers/yr
Records provided (100% catch-and-release)
R. Hueter (813-388-4441)
3. Venice Shark Tournament, Venice, FL
10 yrs (76-85) [Discontinued 85]
50 anglers/yr
No records
C. Balsinger (813-484-9044)

Table 1. (continued)

4. Englewood Shark Club Tournament, Englewood, FL
3 yrs (78-80) [Discontinued 80]
20-30 boats (50-100 anglers)
No records
B. Matherly (813-475-7342)
5. Port Charlotte Rotary Shark Tournament, Port Charlotte, FL
2 yrs (89-90) [Discontinued 90]
<50 anglers/yr
No records
B. Clayman (813-624-3993)
6. Fishin' Frank's Shark Tournament, Port Charlotte, FL
(2 parts: open-water in June, Charlotte Harbor in Aug)
6 yrs (86-91) [Active]
150-225 anglers/yr
Records not provided
F. Hommemma (813-625-3888)
7. Riviera Marina Shark Tournament, Punta Gorda, FL
3 yrs (87-89) [Discontinued 89]
<50 anglers/yr
No records
B. Howard (813-639-2008)
8. Pineland Marina Shark Tournament, Pineland, Pine Island, FL
3 yrs (86-88) [Discontinued 88]
15-40 anglers/yr first two yrs; 100 anglers third year
Incomplete records
B. Jackson (813-283-0080)
9. Pine Island Shark Tournament, St. James City, Pine Isl., FL
2 yrs (89-90) [Discontinued 90]
10-20 anglers/yr
No records
T. Walsh (813-283-4446)
10. Great Shark Hunt, Ft. Myers/Ft. Myers Beach, FL
4 yrs (86-88,90) [Discontinued 90]
100 anglers/yr
Incomplete records
D. Eget (813-463-4448)
11. Angler's Answer Shark Tournament, Marco Island/Naples, FL
4 yrs (88-91) [Active]
150 anglers/yr
Records not provided
B. Udell (813-775-7336)
12. St. Matthew's House Shark Tournament, Naples, FL
1 yr (91) [Active]
26 anglers/11 boats
1 shark caught (4 ft./45 lb. hammerhead)
J. Shaver (813-774-0500)
13. Jaws I-VIII, Port of the Islands, Naples, FL
8 yrs (84-91) [Active]
150-175 anglers/yr
Incomplete records
C. Ray (813-394-3101)

Table 1. (continued)

IV. FLORIDA KEYS

1. Round-the-Clock Shark Tournament, Key West, FL
4 yrs (88-91) [Active]
15-20 boats/yr -- 30-50 anglers/yr
Records not provided
R. Bostal (305-292-1961)
2. Jaycees Shark Rodeo, Marathon, FL
18 yrs (72-89) [Discontinued 89]
40 anglers/yr
No records
B. Daniels (305-872-2237)
3. Univ. of Miami/Upper Keys Shark Tournament, Islamorada, FL
4 yrs (87-90) [Discontinued 90]
50-100 anglers/yr
Incomplete records
J. Musick (305-664-1003)

V. FLORIDA EAST COAST

1. Jaycees Shark Tournament, Lake Worth, FL
5 yrs (78-82) [Discontinued 82]
100-200 anglers/yr
No records
B. Mostler (407-746-7568)
2. Port Salerno Shark Fishing Tournament, Port Salerno, FL
16 yrs (76-91) [Active]
50-100 anglers/yr
Records provided
D. Taylor (407-692-0661)
3. Sebastian Inlet Beach Tournament, n. of Vero Beach, FL
3 yrs (85-87) [Discontinued 87]
30-50 anglers/yr
No records
E. Dwyer (407-773-8379)
4. Florida Shark Club Tournament, Jacksonville (Mayport), FL
15 yrs (76-90) [Discontinued 90]
50-100 anglers/yr
Records provided
S. Gale (904-389-5659)

TOTAL NO. FLORIDA TOURNAMENTS DOCUMENTED = 34

tournaments in Florida typically have been smaller events held in late spring or summer months, on both Atlantic and Gulf coasts. These tournaments range in size from a few events with over 200 participants, which draw anglers, sponsors, and media attention from well outside the fishing area, to very small contests involving less than 50 anglers (Fig. 3). Average tournament size in Florida is about 100 participants. The duration of the typical tournament is one weekend, usually a 40 to 48-hr period between Friday evening and Sunday afternoon or evening.

Nearly all tournaments are exclusively for boat fishermen, usually 2-4 registered anglers per boat; at least two tournaments also have involved beach or pier fishermen. Strictly rod and reel fishing under IGFA rules typically applies. Until very recently (the last 2-3 yrs), most tournaments had no restrictions on catch, either by number or size of sharks, except that many disallowed nurse sharks (*Ginglymostoma cirratum*; Table 2) as not being a good "fighting" fish.

[A joint study by Texas A&M University of the Gulf of Mexico recreational shark fishery for the NMFS/MARFIN program has been conducted concurrently with this study. Further socio-economic aspects of shark tournament participants in the Florida Gulf are surveyed in-depth in the Texas A&M study, under the supervision of Dr. R. Ditton of the Department of Wildlife and Fisheries Sciences and in cooperation with this study's principal investigator, Dr. R. Hueter.]

Temporo-Spatial Distribution of Tournament Activity. The number of shark tournaments in the state began building after 1975 and reached a peak in 1989, then dropped in 1990 and 1991 (Fig. 4). Only one tournament (Jaycees Shark Rodeo, Marathon) was found for the years prior to 1975. In the summer of 1975,




Fig. 3. Location and size (by number of registered anglers) of shark sportfishing tournaments in Florida, 1971-91. A total of 34 tournaments in the 20-year period was held in state waters. The Gulf Coast Shark Census is a non-kill, 100% catch-and-release shark tournament involving 100-200 anglers per year; all others are kill tournaments.

Table 2. List of shark species of Florida coastal waters by common and scientific names.

<u>Common Name</u>	<u>Scientific Name</u>	<u>Family</u>
Bignose	<i>Carcharhinus altimus</i>	Carcharhinidae
Blacknose	<i>C. acronotus</i>	
Blacktip	<i>C. limbatus</i>	
Bull	<i>C. leucas</i>	
Caribbean reef	<i>C. perezi</i>	
Dusky	<i>C. obscurus</i>	
Sandbar	<i>C. plumbeus</i>	
Silky	<i>C. falciiformis</i>	
Spinner	<i>C. brevipinna</i>	
Tiger	<i>Galeocerdo cuvieri</i>	
Lemon	<i>Negaprion brevirostris</i>	
Atlantic sharpnose	<i>Rhizoprionodon terraenovae</i>	
Bonnethead	<i>Sphyrna tiburo</i>	Sphyrnidae
Great hammerhead	<i>S. mokarran</i>	
Scalloped hammerhead	<i>S. lewini</i>	
Nurse	<i>Ginglymostoma cirratum</i>	Orectolobidae
Sand tiger	<i>Odontaspis taurus</i>	Odontaspidae
Shortfin mako	<i>Isurus oxyrinchus</i>	Lamnidae




Fig. 4. Number of active shark tournaments in Florida waters by year, 1971-91. The increase in activity beginning in 1976 followed the release of the movie Jaws in the summer of 1975. After the heyday of shark tournament activity in 1987-89, the number of tournaments began to decline.

Jaws changed the way much of the public viewed sharks, and the number of new tournaments began growing: five added in 1976, and 10 by 1980. From 1985 to 1989, the total number of tournaments in the state increased by two per year, reaching a peak of 21 separate shark tournaments held in Florida in 1989. In recent historical terms, 1987-89 was the heyday of shark tournament activity in the state.

Since 1989, there has been a sharp decline in the number of shark tournaments held each year. Only 14 remain active, and of these only seven are large, viable events involving 100 anglers or more. Thus, of the 34 tournaments charted between 1971 and 1991, only 21% remain as significant tournaments today. This temporal pattern mirrors the trends evident in the NMFS/MRFSS data for Florida: a build-up in sharkfishing activity through the early and mid-1980's, and a decline in the late 1980's.

The great majority of shark tournaments in the state (nearly 80%) have been held along the Gulf coast (Fig. 3). Only four organized tournaments have been held along the Atlantic coast (Jacksonville, Sebastian Inlet, Port Salerno, and Lake Worth) and three in the Keys (Key West, Marathon, Islamorada). On the Gulf coast, five tournaments have been run in the panhandle region (Panama City, Destin, Pensacola, and two in Orange Beach, Alabama). The remaining 22 Gulf tournaments have been concentrated from just north of Tampa Bay to Naples (Fig. 3). With the availability of large sharks indigenous to the broad continental shelf of the central Gulf coast (Clark and von Schmidt, 1965), sharkfishing in this region has attracted more tournament activity than in other coastal zones.

The spatial distribution of the Gulf tournaments from 1980 to 1991 is illustrated in Figs. 5A-G. Figs. 5A-C show the increase in tournament activity along the central Gulf coast from 1980 through 1987. A spatial pattern then emerges in the yearly sequence of tournament distribution from 1988 through 1991 (Figs. 5D-G). There appears to have been a steady dispersal of tournament activity away from its geographic core area (Tampa Bay to Charlotte Harbor) as overall tournament activity began to decline.

Catch Data. Logbook records from nine tournaments were located and found suitable for analysis. Of the remaining 25 tournaments, usable records from only five may exist but were not made available to this study. Of these five, the




Fig. 5A. Distribution of shark kill tournaments in the Florida Gulf, 1980. Figs. 5A-G show a build-up in tournament activity, especially along the central Gulf coast, through the 1980's, followed by a decline in activity in the latter 1980's/early 1990's.




Fig. 5B. Distribution of shark kill tournaments in the Florida Gulf, 1984-85.




Fig. 5C. Distribution of shark kill tournaments in the Florida Gulf, 1986-87.




Fig. 5D. Distribution of shark kill tournaments in the Florida Gulf, 1988.




Fig. 5E. Distribution of shark kill tournaments in the Florida Gulf, 1989.




Fig. 5F. Distribution of shark kill tournaments in the Florida Gulf, 1990.




Fig. 5G. Distribution of shark kill tournaments in the Florida Gulf, 1991.

record curator of only one tournament (Panama City) was deliberately noncooperative in providing information. Two of the nine tournaments providing suitable records were eliminated from the analysis: the St. Matthew's tournament in Naples, because 1991 was its first year; and the Gulf Coast Shark Census based in Sarasota, because its three-year-old format of 100% catch-and-release is categorically different from the kill tournaments.

The seven tournaments with catch data for analysis are: 1) Florida Shark Club Tournament in Jacksonville (Mayport); 2) Port Salerno Tournament near Stuart; 3) Tampa Bay Sharkers Tournament in St. Petersburg; 4) Port of Hudson Tournament in Hudson Beach, north of Clearwater; 5) Destin Tournament; 6) Perdido Pass Tournament in Orange Beach, Alabama; and 7) Monster Shark Tournament in Orange Beach, Alabama (Fig. 6).

The usable catch records from these seven tournaments are summarized in Table 3. They range in duration from 16 years of continuous data from the Port Salerno tournament (1976-91) to three years of data from the Orange Beach Monster Shark tournament (1989-91). Four categories of data for each tournament are summarized in Table 3: 1) total number of sharks entered in the tournament; 2) largest shark entered by total whole weight in pounds; 3) average shark size entered, either averaging weights of all sharks entered or, where those data were not available, taking the average of the top three or five heaviest shark weights; and 4) number of tournament participants. The fishing effort in terms of numbers of anglers fishing was very steady within six out of the seven tournaments over the time periods indicated in Table 3, except for the last year of the Jacksonville (1990), Tampa Bay (1990), and Hudson (1991) tournaments. In the case of the Port Salerno tournament, angler registration is variable by year and is shown in Table 3.

Lines 1, 2, and 3 of Table 3 are plotted for the seven tournaments in Figs. 7A-E. Each graph plots shark number or CPUE data against the left axis and shark weight against the right axis. Fig. 7A shows the combined data for the three panhandle tournaments (Destin, Perdido Pass, Orange Beach Monster Shark). Although the records are not complete and cover only 3-4 years of activity, all three categories of catch (number of sharks, largest, average size) appear to be lower in 1990-91 vs. 1988-89. Number of sharks caught per registered angler (CPUE) dropped for the Destin tournament from $71/150 = 0.47$




Fig. 6. Locations of seven shark tournaments and longlining operation for which catch records are analyzed.

- 1: Monster Shark Tournament, Orange Beach, Alabama.
- 2: Perdido Pass Tournament, Orange Beach, Alabama.
- 3: Destin Tournament, Destin, Florida.
- 4: Port of Hudson Tournament, Hudson Beach, Florida.
- 5: Tampa Bay Sharkers Tournament, St. Petersburg, Fl.
- 6: Port Salerno Tournament, Port Salerno, Florida.
- 7: Florida Shark Club Tournament, Mayport, Florida.

C: Area of fishing for Crooke longline operation.

Table 3. Data summary for logbook records from seven Florida shark tournaments.

		1976	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91
Jax	1				67	-	-	-	-	80	-	-	-	-	-	8	
(79-90)	2				827	-	-	-	-	672	716	625	516	638	-	254	
	3				587 ₃	-	-	-	-	542 ₃	627 ₃	497 ₃	426 ₃	373 ₃	-	79	
	4				113a	-	-	-	-	-	-	-	-	-	-	50a	
Port	1	18	56	-	40	43	32	45	-	14	19	40	34	19	11	13	7
Salerno	2	389	670	1090	905	750	540	665	210	420	400	375	270	384	272	328	227
(76-91)	3	362 ₅	374 ₅	689 ₅	613 ₃	447 ₅	355 ₅	-	202 ₃	361 ₅	281 ₅	349 ₅	231 ₅	301 ₅	245 ₅	267 ₅	184 ₅
	4	107a	69a	-	45a	51a	83a	78a	109a	131a	111a	78a	100a	90a	76a	63a	30a
Tampa	1										32	18	5	22	11	2	
Bay	2										607	566	322	615	351	182	
(85-90)	3										243	302	234	176	198	162	
	4										(averages 200a/yr)					44a	
Hudson	1										48	22	26	21	11	4	5
(85-91)	2										715	752	530	860	300	95	63
	3										-	197	217	216	187	73	55
	4										(averages 230a/yr)					60a	
Destin	1													71	28	18	-
(88-91)	2													605	757	387	-
	3													157	234	202 ₈	-
	4													(averages 150a/yr)			
Perdido	1													16	9	-	-
Pass	2													476	617	-	225
(88-91)	3													246	322	-	-
	4													20b	18b	-	-
Orange	1														32	10	2
Beach	2														566	989	177
(89-91)	3														353 ₅	-	-
	4														8b	-	-

- 1 No. sharks: Total number of sharks entered in tournament
- 2 Heaviest shark: Largest shark (lbs whole weight) entered in tournament
- 3 Avg. shark weight: Mean weight (lbs) of sharks entered [subscript denotes no. of sharks used in calculating mean weight when all shark weights were not available, e.g. x₅ = avg. wt. of top 5 heaviest sharks in tournament, etc.]
- 4 No. anglers/boats: Total number of registered participants in tournament
 - a = anglers
 - b = boats




Fig. 7A. Total number of sharks entered (left axis) and shark weights (right axis) for three Panhandle tournaments. Data from Table 3. The Δ in 1991 is heaviest shark weight (225 lbs) for the Perdido tournament.

HUDSON BEACH TOURNAMENT 1985-91




Fig. 7B. Catch data from Table 3 for Hudson tournament, 1985-91.

TAMPA BAY TOURNAMENT 1985-90




Fig. 7C. Catch data from Table 3 for Tampa Bay tournament, 1985-90.

PORT SALERNO TOURNAMENT 1976-91




Fig. 7D. Catch data from Table 3 for Port Salerno tournament, 1976-91. Catch per unit effort (CPUE) is calculated as number of sharks entered in tournament per registered angler.

JACKSONVILLE TOURNAMENT 1979-90




Fig. 7E. Catch data from Table 3 for Jacksonville tournament, 1979-90. The average shark weight for 1990 is for all 8 sharks entered in the tournament that year.

in 1988 to $18/202 = 0.09$ in 1991; the Orange Beach CPUE likewise dropped from approximately 1.00 in 1989 to 0.06 in 1991. But the trends are not clear due to the short time period of coverage and small number of participants in the Perdido and Orange Beach tournaments.

For the Hudson tournament, however, the historical trends are quite clear (Fig. 7B). Over the seven years of this tournament (1985-91), all three categories of catch have shown sharp declines. This is true even though effort remained a steady 200-230 anglers per year until the last year (1991), when it dropped to 60 anglers. This tournament, which in 1985 landed 48 sharks with a 715-lb tiger as the winning shark, brought in only 4 sharks in 1991 with a 94.5-lb nurse shark as the winner. The most noticeable part of this decline began after 1988.

The Tampa Bay tournament (Fig. 7C) shows similar but perhaps less clear-cut trends over recent years. The 1987 tournament was nearly cancelled due to bad weather, and the tournament director reported that the boats could not get offshore that year. Thus, the transient declines in the 1987 catch vs. 1986 and 1988 were most likely due to this factor. But the declines from 1988 to 1990 are clear. In 1990, with good weather, the number of registrants dropped to 44, but the poor nature of their catch was nevertheless significant: a total of two sharks, a 182-lb nurse shark and a 142-lb bull shark. CPUE dropped from approximately 0.16 sharks/angler in 1985, when 32 sharks weighing from 606.5 to 79 lbs were entered by about 200 anglers, to 0.05 in 1990. After the 1990 tournament, the event was discontinued.

For the Port Salerno tournament (Fig. 7D), angler registration has varied, but fortunately the numbers of registrants are known for all but one of the 16 years of records. This allows for the plotting of true CPUE over the life of the tournament, and results in the best data set found for this study. Although there are some breaks in the data, this tournament shows declines in all categories of catch through the 1980's. CPUE dropped from a high of 0.89 in 1979 to a low of 0.14 in 1989, recovering slightly to 0.21 in 1990 and 0.23 in 1991. (The 1991 data are compromised by comparatively low angler registration and poor weather on the second day of the tournament.)

The Jacksonville tournament data are shown in Fig. 7E. Although there are large breaks in these data, recent declines in catch by number and size of

sharks are evident. The 1990 tournament resulted in eight sharks entered by 50 anglers (CPUE = 0.16, vs. 0.59 in 1979) and an average weight of 79 lbs for the eight sharks. Attempts by the Florida Shark Club to find large sharks in the area prior to the 1991 scheduled tournament were unsuccessful, even though two separate dates were tried, and so the tournament was cancelled that year.

In summary, it is clear that there have been sharp declines in the recreational catch of sharks by number, size, and CPUE in Florida coastal waters since the mid-to-late 1980's, as exemplified by shark tournament data. These lower catches are having a significant effect on tournament activity, leading to the demise of a number of previously well-organized and well-attended kill tournaments.

Species Distribution/Abundance. The breakdown of the seven-tournament catch into eight species categories (sandbar, dusky, tiger, hammerhead spp., bull, lemon, blacktip/spinner, and nurse; see Table 2) is shown in Table 4. The species representation is typical of the assemblage of large sharks inhabiting Florida coastal waters in spring and summer months (Clark and von Schmidt, 1965; Castro, 1983). Generally, the decline in total catch in all tournaments, particularly after 1988-89, is reflected across-the-board in declines in catch by species. No particular species appears to escape this trend. Large dusky, tiger, and hammerhead sharks become rare after 1989.

Further interpretations of these data are difficult given the low and irregular sample size and the unreliability of some of the species identifications. For example, the identification of 39 dusky sharks caught in the Destin tournament in the summer of 1988 is questionable. The common name "dusky" is used in that region to describe a number of shark species (C.F. Crooke, pers. com.), and no duskies were reported in the Crooke longline data for all of 1988 (see Crooke longline data, below). However, in previous years, Mr. Crooke reported catching duskies in the month of July (see below), so the Destin tournament data cannot be entirely ruled out.

The apparent lack of sandbar sharks in the 7/4/88 and 7/4/90 Jacksonville tournaments may be due to real declines in stocks of this species, which is heavily targeted by commercial longliners. On the other hand, the low numbers could be due to a mismatch between the timing of the 1988 and 1990 tournaments

Table 4. Representation of 8 species categories in catch from 7 Florida shark tournaments.

	1979	80	81	82	83	84	85	86	87	88	89	90	91
SANDBAR													
Orange Beach											0		
Perdido Pass										0	0		
Destin										0	0	0	
Hudson							0	0	0	1	0	0	
Tampa Bay												0	
Port Salerno											1	0	1
Jacksonville	22					31				7		0	
DUSKY													
Orange Beach											6		
Perdido Pass										0	0		
Destin										39	4	9	
Hudson							0	0	5	0	3	0	
Tampa Bay												0	
Port Salerno											0	0	0
Jacksonville	1					3				0		0	
TIGER													
Orange Beach											1		
Perdido Pass										4	6		
Destin										8	2	1	
Hudson							4	5	10	3	1	1	
Tampa Bay												0	
Port Salerno											1	0	0
Jacksonville	11					14				1		4	
HAMMERHEAD spp.													
Orange Beach											1		
Perdido Pass										3	3		
Destin										9	6	3	
Hudson							1	5	3	2	1	0	
Tampa Bay												0	
Port Salerno											0	1	1
Jacksonville	6					12				0		1	
BULL													
Orange Beach											8		
Perdido Pass										9	0		
Destin										15	12	2	
Hudson							2	3	3	5	2	1	
Tampa Bay												1	
Port Salerno											0	0	0
Jacksonville	7					7				2		0	

Table 4. (continued)

	1979	80	81	82	83	84	85	86	87	88	89	90	91
LEMON													
Orange Beach											0		
Perdido Pass										0	0		
Destin										0	0	0	
Hudson							1	2	2	4	1	0	
Tampa Bay												0	
Port Salerno											3	6	3
Jacksonville	8					4				0		2	
BLACKTIP/SPINNER													
Orange Beach											0		
Perdido Pass										0	0		
Destin										0	3	2	
Hudson							1	1	3	1	0	1	
Tampa Bay												0	
Port Salerno											1	0	0
Jacksonville	8					4				0		0	
NURSE													
Orange Beach											0		
Perdido Pass										0	0		
Destin										0	1	1	
Hudson							1	6	0	5	3	1	
Tampa Bay												1	
Port Salerno											1	4	2
Jacksonville	0					2				0		0	

NOTE: Species ID's not necessarily reliable

Species ID records incomplete for:

Orange Beach '89 (16/32 ID'ed)

Hudson '85 (10/48 ID'ed)

Port Salerno '89 (10/11 ID'ed)

Jacksonville '84 (77/80 ID'ed) & '88 (10/? ID'ed)

and movements of the inshore groups of sandbar sharks off northeast Florida during those two years. As an aggregating, highly migratory species, sandbar sharks can vary in the timing of their local seasonal abundance with climatic variation and other factors.

Nevertheless, with the above considerations, the overall picture of decreasing catch by species is consistent with the declines in the general catch data from the seven tournaments.

Crooke Longline Data

Raw Data. The transcribed Crooke longline data are presented in Appendices 1 and 2. Appendix 1 is the database arranged chronologically and Appendix 2 is sorted by species. The area of fishing south of Pensacola for these data is indicated in Fig. 6.

Table 5 shows the compiled species totals by sex. A total of 659 sharks of 13 species were collected, comprised of 338 nongravid females, 43 gravid females, 245 males, and 33 sharks where sex was not determined (usually because the shark had been damaged on the longline by the predatory activity of other sharks). The species caught in order of abundance, from highest to lowest, were sandbar, blacktip, bull, tiger, dusky, scalloped hammerhead, great hammerhead, blacknose, nurse, spinner, sand tiger, lemon, and sharpnose sharks. The size of the longline gear (14/0 hooks) selected for larger sharks, and smaller species that inhabit the fishing area may be under-represented (e.g. blacknose and sharpnose) or not present (e.g. bonnethead, *Sphyrna tiburo*) in the catch.

Species Profiles. The distributions of catch by month of the year over the 15-year period are plotted for each of the 13 species in Figs. 8A-G. Mr. Crooke did not fish in the months of January and February, and rarely in March and December, so the primary sampling period extends from April through November each year. The water temperature regimes for each species are shown in Fig. 9.

Sandbar sharks, the most abundant species, were found throughout the fishing year off Pensacola but reached a distinct peak in October (Fig. 8A). Sex ratio for sandbars was 1:1.37 males to females. Ten percent of all females

Table 5. Compiled Crooke longline catch (1975-89) by species and sex.

<u>SPECIES</u>	<u># NONGRAVID FEMALES</u>	<u># GRAVID FEMALES</u>	<u># MALES</u>	<u>SEX NOT AVAILABLE</u>	<u>TOTALS</u>
1. Sandbar	90	10	73	1	174
2. Blacktip	70	19	23	15	127
3. Bull	58	4	50	1	113
4. Tiger	67	1	28	3	99
5. Dusky	32	1	27	4	64
6. Scal. Hammerhead	0	0	28	4	32
7. Great Hammerhead	12	2	5	0	19
8. Blacknose	3	5	4	2	14
9. Nurse	0	0	5	2	7
10. Spinner	4	0	1	1	6
11. Sand tiger	2	0	0	0	2
12. Lemon	0	1	0	0	1
13. Sharpnose	0	0	1	0	1
	<u>338</u>	<u>43</u>	<u>245</u>	<u>33</u>	<u>659</u>





Fig. 8A. Number of sandbar and blacktip sharks in Crooke longline data by month caught over all years, 1975-89.





Fig. 8B. Number of bull and tiger sharks in Crooke longline data by month caught over all years, 1975-89.




Fig. 8C. Number of dusky and scalloped hammerhead sharks in Crooke longline data by month caught over all years, 1975-89.




Fig. 8D. Number of great hammerhead and blacknose sharks in Crooke longline data by month caught over all years, 1975-89.




Fig. 8E. Number of spinner and nurse sharks in Crooke longline data by month caught over all years, 1975-89.





Fig. 8F. Number of sand tiger and sharpnose sharks in Crooke longline data by month caught over all years, 1975-89.




Fig. 8G. Number of lemon sharks in Crooke longline data by month caught over all years, 1975-89.




Fig. 9. Seawater temperature ranges and means (filled circles) for shark species from Crooke longline data. (Local water temperature in the fishing area was recorded for each day of catch.)

caught were gravid. Size of sandbars ranged from a 4.3-ft (total contour length-TCL) nongravid female caught 6/16/87 to a 7.8-ft TCL nongravid female caught 10/5/81.

Blacktip sharks were more predominant in the late spring, reaching their peak in May and June (Fig. 8A). Sex ratio was 1:3.87 males to females and 21% of females were gravid. For 12% of the blacktip sharks, sex was not identifiable. This was by far the most molested species of shark on the line, comprising over 50% of the sharks preyed on by other sharks after being caught. Size of blacktips ranged from a 3.9-ft TCL male caught 9/7/85 to a 6.4-ft TCL nongravid female caught 6/11/83.

Bull sharks usually appeared in the area in May at their peak abundance, and were caught with decreasing frequency throughout the summer into September (Fig. 8B). One bull was caught in November, a 6.6-ft TCL male captured 11/7/81. Sex ratio was 1:1.24 males to females; only 6% of females (4) were gravid. Size ranged from a 5.3-ft TCL nongravid female caught 4/9/77 to two 9.0-ft TCL nongravid females caught 6/1/78 and 7/13/88.

Tiger sharks were found in the area throughout the fishing year, basically increasing in frequency from March to November, with a transient decline in June (Fig. 8B). Tigers were one of the most eurythermal species, found in a 26°F temperature range from 60°F to 86°F (Fig. 9). Of the 68 females caught during the 15 years of longlining, only one was gravid (13 ft TCL, caught 7/4/80; this is the largest shark in the Crooke records). Sex ratio was 1:2.43 males to females and size ranged from a 3.3-ft TCL male (8/23/87) to the 13-ft female.

Dusky sharks were present from June to December with two peak periods, one in July and a higher one in October/November (Fig. 8C). The size of dusks in the catch was large, averaging 10.2 ft TCL ($s = 0.99$) and ranging from a 7.9-ft TCL nongravid female caught 10/28/78 to an 11.6-ft TCL nongravid female caught 7/3/85. As with the tiger sharks, only one gravid dusky female was found out of 33 females caught. Sex ratio was 1:1.22 males to females.

Scalloped hammerheads appeared suddenly on the scene in May, tapering off in frequency through the summer with only a few caught in August, September, and October (Fig. 8C). All scalloped hammerheads were male (four could not be identified by sex). Size ranged from 6.1 ft TCL (5/16/81) to 9.0 ft TCL (5/30/81 and 7/13/88).

The monthly distributions of the other seven species are shown in Figs. 8D-G. Numbers of sharks in these seven species were low, together comprising only 8% of the total catch. Interesting features among these species includes: gravid females outnumbered nongravid females among the blacknose sharks; all five nurse sharks caught were male; and the one lemon shark caught in 15 years of fishing was a gravid female, 9.0 ft TCL caught 5/1/82.

Historical Trends in Size of Sharks. Figs. 10A-G show plots of all sharks caught, by total contour length vs. precise date caught, for each of the 13 species, with sex of each shark indicated. For the six most abundant species, simple linear regressions were calculated to examine the relationship between size of shark (in TCL) and year of capture (Figs. 10A-C). For these calculations, dates of capture were reduced to year only, to eliminate regular seasonal fluctuations. An ANOVA test was applied to test for significance of each of the six regressions (Zar, 1984).

Four of the six species (blacktip, bull, dusky, and scalloped hammerhead) did not pass the ANOVA F-test for regression significance. However, two species -- sandbar and tiger sharks -- had regressions with significant *negative* slopes. The sandbar data (Fig. 10A) produced a regression coefficient (slope) of -0.04 ft/yr that was significant ($0.0025 < P < 0.005$), indicating that the size of sandbar sharks slowly diminished at an average rate of about $\frac{1}{2}$ inch per year throughout the 15 years of longlining. The tiger data (Fig. 10B) showed a somewhat stronger decline with a significant slope of -0.14 ft/yr ($0.001 < P < 0.0025$), corresponding to an average decrease of 1.7 inches per year or about one foot every 7 years.

Coefficients of determination (r^2) were very low for all six regressions, including sandbar (0.05) and tiger (0.09). This indicates that a curvilinear relationship would probably better describe the relationship of decreasing length with time in the sandbar and tiger data.

When all sharks irrespective of species are plotted by length vs. year (Fig. 11), the resulting regression is highly significant ($P < 0.0005$). The slope of this line is -0.09 ft/yr, corresponding to an average decrease of 1.1 inches per year, or about one foot every 11 years. Again, r^2 for this linear regression is very low at 0.04, strongly suggesting a curvilinear relationship in the decline of shark length with time.




Fig. 10A. Length of shark vs. date of capture from Crooke longline data for sandbar ($n = 173$; negative slope of line significant, $0.0025 < P < 0.005$; $r = -0.22$) and blacktip ($n = 116$; regression not significant; $r = -0.05$).





Fig. 10B. Length of shark vs. date of capture from Crooke longline data for bull ($n = 113$; regression not significant; $r = 0.08$) and tiger ($n = 97$; negative slope of line significant, $0.001 < P < 0.0025$; $r = -0.31$).





Fig. 10C. Length of shark vs. date of capture from Crooke longline data for dusky ($n = 60$; regression not significant; $r = 0.04$) and scalloped hammerhead ($n = 32$; regression not significant; $r = -0.03$).




Fig. 10D. Length of shark vs. date of capture from Crooke longline data for great hammerhead (n = 18) and blacknose (n = 12).




Fig. 10E. Length of shark vs. date of capture from Crooke longline data for spinner (n = 5) and nurse (n = 5).





Fig. 10F. Length of shark vs. date of capture from Crooke longline data for sand tiger (n = 2) and sharpnose (n = 1).




Fig. 10G. Length of shark vs. date of capture from Crooke longline data for lemon (n = 1).

ALL SPECIES




Fig. 11. Length of shark vs. date of capture from Crooke longline data for all species combined ($n = 639$; negative slope is significant, $P < 0.0005$; $r = -0.20$). Average length of all sharks over the 15 years of fishing was 6.86 ft TCL.

Catch per Unit Effort. CPUE is plotted from the 15 years of Crooke data in Fig. 12 as number of sharks caught per hook vs. year of fishing, *for successful sets of the gear* (see Methods/Crooke Longline Data). There is a noticeable decline in the CPUE plot between the early 1980's and the late 1980's. In 1980, CPUE reached its highest point at 0.154 sharks per hook. In 1988, at its lowest point, CPUE was exactly half that of eight years previously, at 0.077 sharks per hook. A linear regression fitted to the entire CPUE data shows a negative slope (Fig. 12) but the regression does not pass an ANOVA test for significance ($0.10 < P < 0.25$). If the catch data for the first two years of operation, 1975 and 1976--arguably the start-up learning period in which Mr. Crooke set the line only nine times each year--are omitted from the analysis, the resulting regression shows a steeper negative slope that is highly significant ($0.005 < P < 0.01$; $r^2 = 0.48$).

Mr. Crooke reported in his annual written summaries which accompanied the database that catches were dropping off in 1987-89, with more unsuccessful sets occurring in those later years. He writes in his 1989 summary that it "was the first year ever to make 6 or more longline sets with no sharks." Thus, it is apparent that CPUE was dropping off even more dramatically than is shown in Fig. 12, but unfortunately the total number of unsuccessful sets each year was not recorded for quantitative analysis.

Summary. In the later years of the Crooke longline operation, size of sharks caught and CPUE in sharks per hook were both declining. It is evident from the data that one explanation for both phenomena is the near-complete disappearance of the dusky (Fig. 10C) as well as the great hammerhead (Fig. 10D) from the catch after 1985. Both of these are large-bodied species. The decline in average size of the most abundant species (sandbar) and fourth-most abundant species (tiger) contributed further to a general decline in size of catch.




Fig. 12. Catch per unit effort (CPUE), in number of sharks caught per hook set, for successful sets (i.e. those catching at least one shark) for the Crooke longline data, 1975-89. In 1975-86, the line had 20 hooks per set; in 1987-89 (*), 30 hooks per set were used. The regression using all data (dotted line) is not significant; with the 1975-76 data omitted (see text), the negative slope of the regression (dashed line) is significant ($0.005 < P < 0.01$; $r = -0.70$).

DISCUSSION

Historical Trends in Florida Shark Tournaments

With over 200 shark tournaments involving more than 22,500 anglers over the past 20 years, tournament fishing for sharks has been a significant component of the recreational fishing activity in Florida. Most anglers participating in these tournaments are general saltwater fishermen who target other marine species as well in their fishing activities but particularly enjoy the thrill of battling a large fish. The shark mystique attracts these fishermen, as well as tournament organizers, sponsors, businesses, the media, and the public at large.

Fishing in the tournaments is typically conducted from boats, although some beach and pier fishing is allowed in certain tournaments. Most boat fishing is done not far from shore, usually less than 15 miles out, in vessels ranging from small, open boats powered by outboard motors to large, expensive sportfishing yachts. Value of top prize varies in these tournaments but generally runs between \$1,500 to \$3,000 cash. This is well below the prizes in some "big money" shark tournaments held in the northeast U.S., which have awarded top prizes of \$50,000 or more (NMFS, 1989).

Interest in shark tournament fishing began building in Florida after 1975 and reached a peak in 1989. This growing interest was reflected in more new tournaments being established each year, until a maximum of 21 separate shark tournaments were being held by 1989. But even before then, tournament fishing for sharks was beginning to change, and many of those 21 tournaments were experiencing problems. In the two years since 1989, tournament activity in the state has declined by at least 33%.

The majority of shark tournaments in the state have been held along the Gulf coast, with concentrations in the western panhandle and the central Gulf coast. The recent decline in tournament activity has been manifested particularly in the dissolution of activity along the central Gulf coast from Tampa to Naples (Figs. 5A-G).

The reasons for the discontinuation of individual tournaments during the 1971-91 study period are several. Business factors affect the organization and continuity of tournaments from year to year. Competition among tournaments may lead to the demise of smaller events. The recent prominence of the conservation movement among saltwater sportfishermen has caused many tournaments to at least re-think their format, and either change or discontinue their activities. Conservation-oriented shark fishermen recently have begun to seek alternatives to unlimited kill tournaments. Part of the change in kill tournament activity in the central Gulf coast region, for example, may be due to the recent success of the 100% catch-and-release Gulf Coast Shark Census, begun in 1989.

But the most significant reason given for the recent demise of the larger, well-established tournaments (e.g. Tampa Bay Sharkers, Florida Shark Club in Jacksonville) is the lack of large sharks to catch. This is substantiated by the catch data presented in this report. With no exceptions reported, the catch of large sharks in Florida tournaments has declined in recent years. The numbers of sharks entered in the tournaments have decreased and, where calculable, the catch per unit effort in the tournaments has fallen.

Rules changes in the tournaments play a role in determining number of sharks entered, and some kill tournaments recently have adopted daily bag limits and minimum sizes. But these changes have been made only in the last 1-2 years, while declines in the catch began well before that. In fact, these rule changes typically have been made as a response, rather than a precursor, to the historical decline in catch within a tournament. For example, the Port Salerno shark tournament instituted bag limits for the first time in its rules in 1990, only after CPUE had hit its lowest point in 14 years of tournament activity (Fig. 7D).

Particularly dramatic in the tournament catch data is not only the decline in shark numbers and CPUE, but also the size of the sharks. If anything, this variable should *increase* with rules limitations on number and minimum size of sharks entered in tournaments. But instead, both the heaviest weights and the average weights of sharks entered in state tournaments have declined significantly in recent years. In the early to mid-1980's, shark tournaments were being won with 600-800+ pound tiger, dusky, and great hammerhead sharks. Now the tournaments are being won in some cases with nurse sharks weighing less than 100 pounds.

Historical Trends in the Crooke Longline Data

The Crooke longline data represent a small but unusually consistent and long-lived sampling operation of the large, inshore sharks inhabiting the northeastern Gulf of Mexico from 1975 to 1989. The database of 659 sharks representing 13 species contains important life history information on distribution of these species by season and size ranges. This report focuses on the fishery-related aspects of the database, and further studies on other biological aspects of these data would be useful.

It is evident from the Crooke data that in the later years of fishing, after 1985, the catch was diminishing. CPUE was down in the last five years of fishing, falling by at least 40% from 1985 to 1988. The size of sharks being caught decreased over the 15 years of fishing, but it appears that this decrease was not a constant since the linear regressions showed relatively poor fits to the data. Rather, it is more likely that a curvilinear relationship would best describe the size-year, and perhaps the CPUE, data: relatively flat during the period from 1975 to 1985, followed by a downward curve in the last five years of fishing.

One primary reason for this result appears to lie in the fading out of the large dusky sharks in the fall catch and great hammerheads in the summer catch. The near-disappearance of the dusky shark after 1985 is consistent with a widespread decline in the numbers of large dusky sharks throughout the northwestern Atlantic region. This general decline began in the late 1970's and early 1980's (J. Casey, pers. com.).

The third large-bodied species in the catch, the tiger shark, did not diminish so much in number as in size. Perhaps because it is among the most prolific of viviparous sharks, with litters averaging about 40 pups, this shark persisted in the late 1980's in the Crooke catch (and the tournament catch) but at smaller sizes. As with the dusky and great hammerhead, it appears that the large tigers were being eliminated from the inshore area of fishing.

The other species of sharks in the Crooke catch diminished in number as CPUE fell but size of shark in the catch did not decrease, except for a small decline in the sandbar shark. This highly migratory, relatively abundant species has been the most heavily targeted shark in the southeast U.S. commercial shark fishery.

Implications for Local Shark Populations and the Florida Shark Fishery

Both sets of data presented in this report point to the same conclusion. The declines in numbers of large sharks in inshore waters off Florida are clearly seen in the tournament catches and in the Crooke longline data. This decline appears to have taken hold in the late 1980's, a time when the state's recreational shark fishery was at its peak and the commercial shark fishery underwent exponential growth. Although such factors as habitat degradation and declines in prey species may play a role in this decline, the timing between the increases in fishery effort and the subsequent drops in shark catch point to overfishing of shark stocks. This is consistent with the conclusion of NMFS that sharks were overfished in the western Atlantic and Gulf of Mexico throughout the 1980's (NMFS, 1989).

The fact that the Port Salerno tournament data show a drop in CPUE in the early 1980's, even before the Florida commercial fishery began to expand in the mid-1980's, may indicate the relative impact of recreational fishing pressure on sharks. A possible scenario state-wide emerges from these data: increasing recreational targeting of sharks in the late 1970's and early 1980's had diminished stocks to some extent, so that when the intense commercial effort developed in the mid-1980's, rapid declines set in.

This may be underscored by the spatial "dispersal" phenomenon of shark tournament distribution along the Gulf coast in the latter 1980's. Comparing the sequence of tournament locations (Figs 5A-G) with the focus of the commercial fishery in the Tampa Bay area (Fig. 2), there is an apparent correlation between the development of the commercial fishery in the central Gulf and the withering of tournament activity in the same region. Although the commercial longliners out of Tampa Bay ports now fish throughout the eastern Gulf, the fishery began expanding in the mid-1980's with short runs to the inshore waters nearer to home.

It is interesting to speculate from these data on the potential impact of fishing pressure on local stock depletion of sharks. Although most Florida coastal sharks are migratory, it appears that specific inshore zones can be depleted relatively rapidly. Thus, even though individual sharks inhabiting an inshore site may migrate there from distant locations, once those sharks are removed, recruitment of more sharks to the site is low. This raises the interesting question of whether there exist subgroups of sharks that follow

highly specific migratory routes, not unlike sea turtles that return to the exact same nesting beach year after year.

A fitting conclusion to this report can be found in the prophetic words of Mr. Fred Crooke, who in 1989, after 15 years of sharkfishing in Florida waters, concluded his very last record entry with these words:

This will be my last year to longline for sharks. They are in very grave trouble and I no longer want to be a part of the problem. Hopefully all of my paperwork on the northwest Florida sharks will help to establish some baseline data for the shark's future. You cannot know where you are today if you don't know where you were in the past. It would be a terrible thing to reduce the shark to a point where he cannot return.

ACKNOWLEDGMENTS

The author is most grateful to Mr. C.F. "Fred" Crooke, Jr. of Warrington, Florida, for providing his excellent and unique set of sharkfishing records for this study. Appreciation is expressed to Dr. Perry W. Gilbert of Mote Marine Laboratory for helping to coordinate the transfer of the Crooke records and for other assistance in this project.

Grateful appreciation is extended to all of the shark tournament officials, record curators, and others who supplied tournament data for this study. Special thanks are extended to Dr. José Castro and Mr. Jack Casey of NMFS for their help in acquiring tournament records.

Mr. Jay Sprinkel, data analyst for Mote Marine Laboratory, was instrumental in producing the graphs and regression analyses. Mr. Robert Dixon, graphics artist for Mote Marine Laboratory, produced the maps.

Finally, acknowledgment is gratefully given for financial support of this project by the saltwater fishing license-funded Marine Resources Grants Program, FDNR.

REFERENCES

- Castro, J.I. (1983) *The Sharks of North American Waters*. Texas A&M University Press, College Station, TX, 180 pp.
- Clark, E. and K. von Schmidt (1965) Sharks of the central Gulf coast of Florida. *Bull. Mar. Sci.* 15:13-83.
- Florida Dept. Natural Resources, Div. Marine Resources (1979) *Summary of Florida commercial marine landings, 1979*. Tallahassee, FL.
- Florida Dept. Natural Resources, Div. Marine Resources (1981) *Summary of Florida commercial marine landings, 1981*. Tallahassee, FL.
- Florida Dept. Natural Resources, Marine Fisheries Information (1988) *Annual landings summaries*. St. Petersburg, FL.
- Graefe, A.R. and Ditton, R.B. (1976) Recreational shark fishing on the Texas Gulf coast: an exploratory study of behavior and attitudes. *Mar. Fish. Rev.* 38:10-20.
- Holden, M.J. (1974) Problems in the rational exploitation of elasmobranch populations and some suggested solutions. *In* *Sea Fisheries Research* (F.R. Harden Jones, ed.), pp. 117-137. Logos Press, London.
- Marine Fisheries Commission, State of Florida (1991) July 1991 Commission meeting (Atlantic Beach, FL) summary, MFC, Tallahassee, FL.
- National Marine Fisheries Service (1984) *Marine recreational fishery statistics survey, Atlantic and Gulf coasts, 1979 (revised)-1980*. Current Fishery Statistics No. 8322, U.S. Dept. Commerce, NOAA/NMFS, Washington, DC.
- National Marine Fisheries Service (1987) *Marine recreational fishery statistics survey, Atlantic and Gulf coasts, 1986*. Current Fishery Statistics No. 8392, U.S. Dept. Commerce, NOAA/NMFS, Washington, DC.
- National Marine Fisheries Service (1991) *Marine recreational fishery statistics survey, Atlantic and Gulf coasts, 1987-1989*. Current Fishery Statistics No. 8904, U.S. Dept. Commerce, NOAA/NMFS, Washington, DC.
- National Marine Fisheries Service (1989) *Draft Secretarial shark fishery management plan for the Atlantic Ocean*. 116 pp. U.S. Dept. Commerce, NOAA/NMFS, St. Petersburg, FL.
- Zar, J.H. (1984) *Biostatistical Analysis*. Prentice-Hall, Englewood Cliffs, NJ, 718 pp.

Appendix 1.

Crooke longline data arranged chronologically. W.T.: water temperature. Species: BL, bull; BN, blacknose; BT, blacktip; D, dusky; GH, great hammerhead; L, lemon; N, nurse; SB, sandbar; SH, scalloped hammerhead; SN, sharpnose; SP, spinner; ST, sand tiger; T, tiger. Sexes: F, nongravid female; GF, gravid female; M, male. Lengths: TCL, total contour length; PCCL, precaudal contour length; FL, fork length.

<u>Date</u>	<u>W.T.</u>	<u>SP</u>	<u>SEX</u>	<u>TCL(ft/in)</u>		<u>PCCL(ft/in)</u>		<u>FL(ft/in)</u>	
9/7/75	85	BL	M	7	8.5			6	4
		D	F	11	6			9	4.5
		T	F	7	4.5			5	8
		T	F	8	8.25			5	9
9/9/75	84	T	F	7	2				69
9/10/75	84	BL	F	6	6.5				66.5
		BL	M	6	0.5				59.5
9/11/75	84	BL	M	6	8				68
		BL	F	6	6			5	5
		T	F	8	11			7	5.5
9/12/75	84	BT	GF	6	2			5	1.25
		T	F	7				5	7.5
9/13/75	85	BT	GF	5	6.5				53.5
		T	F	7	6			6	0.75
		T	F	7	4.5			6	1
		T	F	8	11			7	4.25
9/14/75	84	N	M	8	2			6	6.5
		N	M	8	6.75			7	1
9/21/75	81	SB	F	6	11.75			5	10
10/3/75	73	SB	F	6	3.5				63.5

5/21/76	73	SH	M	8	2.25				71.25
6/11/76	78	BL	F	6	4				63
6/24/76	83	BL	M	6	7			5	5.75
8/16/76	80	T	F	5	11				57.5
9/29/76	79	D	M	10	6			9	1.5
10/1/76	78	SB	GF	7	2			6	1.25
		BT	F	5	10				57.5
		GH	F	10	6			8	1.75
10/10/76	78	D	M	8	5			7	0.75
		T	F	5	6				52
10/16/76	74	SB	GF	7	4.75			6	1.5
11/7/76	63	T	M	7	1.25			5	11
		SB	F	7	1.75			5	11.75
		SB	F	7	4.5			6	

4/2/77	68	SB	M	5	0.5				50.75
4/9/77	67	SB	F	7	0.75			5	8
		BL	F	5	3.88				49.38
		SB	F	7	3.25			6	0.75
5/1/77	76	SH	M	8	2.25			6	1
		BT	F	6	0.5			5	
		SP	F		47				38.75
5/15/77	79	BT	M	5	3				52
		SH	M	8	2			6	4
5/29/77	80	BL	F	6	4			5	4.5
		BL	M	6	2			5	1.5
5/31/77	80	BT	F	5	7.75				57
6/1/77	80	GH	F	7	5			5	5.5
		BT	F	5	6.5			4	6.5
6/2/77	80	BL	M	6	0.5			4	11.25
		BL	F	7	7			6	0.5
		BT	F	5	2.75			4	6.5
		BT	F	5	10			4	11
6/3/77	80	SH	M	7	9			5	10
		BT	F	6	1.75			4	11
9/10/77	84	D	F	10	10			8	11
9/18/77	84	T	F	7	3.75				72
		D	M	10	3.5			8	2.5
		BL	M	5	11.5			5	1

10/5/77	83	D	F	8	6.5	7	1
		D	F	11	1.5	9	9.5
		D	F	10	11.5	9	3.75
10/7/77	80	T	F	7	4.75	6	1
		D	M	10	4	9	
		D	F	11	5.5	9	6
10/30/77	75	D	M	10	1	8	5
		SP	F	5	10.75	4	10
11/13/77	72	T	F	6	6.5	5	2.5
		T	F	7	10	6	3.5
		D	F	10	2.5	8	8
		SB	M	4	7.5	4	2.5
11/26/77	69	T	M	10	3	8	9
		T	F	12	1	10	6.5
		D	F	11	2.5	9	4
		SB	F	7	2.5	6	2.5
<hr/>							
4/16/78	71	SB	F	7	3.5	6	0.5
5/21/78	76	BL	M	6	2.5	5	0.5
5/30/78	77	BL	F	6	2.5	5	1.5
		BL	F	6	2.5		59
		BL	F	6	2.75		53.75
		BL	F	5	6.25		55.25
		BL	M	5	5.5	4	5.5
5/31/78	77	BL	M	5	7	4	7
		BL	M	6	5.5	5	4.5
		BL	M	6	2.5		59.5
		BT	F	5	3	4	3
6/1/78	77	BL	F	6	10.25	5	7
		BT	F		59.5	NA	NA
		BT	F	5	4	4	6
		BL	F	8	8.5	7	6
		BL	F	9		7	9
		BL	F	6	4.5	5	2.5
6/2/78	78	T	M	9	6.5	7	10.5
		BT	NA	5	9	NA	NA
		BN	GF	4	4	NA	NA
		SH	M	7	3.5	5	6.5
		BL	GF	8	2.5	7	
6/11/78	81	BL	F	5	11.5	5	1
6/25/78	84	BT	F	5	6		54.5
		BT	F	6			59
		BL	M	6	5.25		59
7/1/78	85	BT	M	5	6	4	6.5
		BT	M	3	10		38
7/2/78	85	BT	F	6	0.25		59
7/4/78	85	BL	F	6	5.5	5	5.5
		BL	F	8	7	7	7
		BL	F	8	10.5	7	9.5
7/30/78	85	D	M	9	9.5	8	1.5
		D	M	10	3	8	8.5
		BL	M	5	4.25	4	5
		D	M	10	0.5	8	5
8/13/78	84	BL	M	6	0.5	4	11.5
		BL	M	5	9.5	4	11
8/20/78	85	D	F	11	5.75	9	5.5
		BL	F	7			69
		N	M	8	6	6	10
9/2/78	84	GH	F	7	11	5	10.25
9/3/78	84	BL	F	7	5.25	6	1
9/4/78	82	BT	F	5	7		55
9/5/78	84	BT	F	4	8.5	3	10.25
		BL	M	5	10.5	4	9.5
9/15/78	83	BL	F	5	11.25		59
9/16/78	83	N	M	8	5.5	6	11
9/17/78	82	SP	F	5	4.63		52.25
10/8/78	76	D	M	10	1.25	8	8
10/14/78	76	D	M	10	3.5	8	8
		SB	F	7	0.25	5	9
		T	F	7	2.5	5	8.25

10/24/78	70	T	F	6	8.5		5	4.5
		T	F	7	9		6	1
10/25/78	72	SB	M	6	3		5	5.5
10/27/78	71	SB	F	6	9.5		5	7
		SB	F	7	4		6	4
		SB	F	6	4		5	2
10/28/78	73	D	F	7	11		6	8
		SB	F	7	3.25		6	1
11/4/78	72	T	M	6	6.75		5	5.75
		D	M	8	3		6	7.25
		D	F	8	1		6	7.5
		SB	M	5	7		5	0.75
11/5/78	73	SB	F	7			6	
11/19/78	71	T	F	7	4.5		6	0.25
		SB	F	6	11.25		5	11
		SB	F	6	2.5		5	4
11/24/78	70	SB	M	6	9		5	8.5
		SB	M	5	7.5		4	10
		SB	F	6	8.5		5	9
		SB	M	5	10		4	9.5
		T	M	6	7.5		5	5
		SB	M	6	0.25		5	0.75
		T	F	6	5.5		6	
		T	M	5	10.25		4	9.25
12/17/78	59	D	F	8	6.5		6	11.5
		D	F	11	6		9	11

4/8/79	67	ST	F	9	11.75	8	3	
4/29/79	71	BN	F		47		34.75	
		BT	F	5			43.5	
		BT	F		58.5		42.5	
		BT	GF	5	4		46.5	
		BT	F	5			43	
		BT	GF	5	6		48.25	
5/6/79	71	BL	F	6	3		54.25	
		BL	M	5	11.5		52	
		BT	GF	5	6.5		48.75	
5/13/79	74	BL	M	8	3.5	7	3.5	
		BN	GF		47.25		35.25	
		BN	GF		45.25		33.75	
5/20/79	75	BT	M	5	4		47	
5/22/79	76	BT	F		58		41	
		BT	GF	5	8.25		49.5	
5/26/79	74	BT	F	5	10.5		51.5	
5/27/79	74	SH	M	8	3.25	6	4	
		SH	M	8	6.5	6	5.5	
5/28/79	75	BT	M	5	2.5		45.5	
5/29/79	75	BL	F	6	6		57	
5/31/79	75	BL	GF	8	11	6	8	
		BL	M	6	11.25		60.5	
6/2/79	75	BT	F		63		45.5	
		SH	M	8	2.5	5	9	
6/9/79	80	BT	F		52.5		37.5	
		BT	M		54		39.25	
		SN	M		38.25		29	
		BL	M	5	7		48.5	
		BL	F	8	8.5	6	5.5	
6/16/79	82	BT	F	5	7.75		49	
		BT	F	5	6.25		48.5	
		BT	M		55		39.25	
		BL	F	8	6	6	3	
		BT	F		53		40	
6/17/79	80	SH	M	8	5.5	5	10.5	
		BT	F	5	10.5		51.25	
6/20/79	81	BT	F	5	8.75		50.5	
		SH	M	7	11.5	5	8	
6/21/79	81	BT	M		56.75		41.5	
6/23/79	81	BT	M	5	4.5		47.5	
6/24/79	81	BT	F	5	8.5		50.75	
		GH	M	6	2		51	
		BT	F	5	7		49	

6/26/79	81	GH	M	6	6.75		54.25
6/29/79	81	SH	NA	7	6	5	6
7/6/79	82	BL	M	6	4.25		55.5
		BL	F	8	10	6	9
7/7/79	82	D	F	10	11.5	8	4.5
		D	M	10	8	7	9
		D	M	8	10	6	4.5
		D	F	10	10.5	8	

4/20/80	64	SB	F	5	2		45.25
7/4/80	80	T	GF	13	0	10	0
		BL	F	8	7.75	6	6
8/2/80	84	D	M	10	4.5	7	9
8/16/80	81	GH	F	10	8	7	6.25
		BL	M	6	6	4	9
8/23/80	84	BL	M	5	11.75		52
8/31/80	82	BL	M	5	10		50.5
9/1/80	81	T	F	8	11.5	6	6
9/21/80	79	D	M	9	8	6	7.25
10/6/80	76	D	M	9	3.75	6	9
		D	GF	11	2	8	5
		D	F	11	6.5	8	7
		GH	F	8	7.5	6	
		T	F	8	5.25	6	
10/8/80	77	SB	GF	7	6	5	6.5
		SB	F	5	11.25		52.5
		SB	M	6	3		55.5
		T	F	7	6.5	5	4.5
		D	F	11	0	8	2
		D	M	9	6.5	7	0
		D	M	8	10	6	4
10/9/80	78	T	F	5	1.5		41.5
		T	M	7	0	NA	NA
		T	F	8	9.5	6	3
		T	M	8	7.5	6	2.5
		SB	F	6	8		58.75
		SB	F	7	3	5	4.5
10/11/80	78	SB	GF	7	5	5	8.5
		T	F	5	1.5		44.5
		SB	M	6	7.5		58
		SB	F	6	8		59
		SB	F	7	3.5	5	6
		D	M	8	6	6	1
11/7/80	74	SB	M	5	0.5	NA	NA
		T	M	6	4		53
		D	M	10	2	6	9
		D	F	10	0.75	6	7.5
		D	F	9	0	6	7
		SB	M	6	1.5		53.75
		SB	M	5	10.75		52
		SB	F	5	8.75		50.75
		SB	F	5	6.25		49
11/22/80	68	D	F	11	2	8	4.5
		SB	F	5	7.5		49.25
12/6/80	67	SB	M	5	5		47.5

4/25/81	71	BT	GF	6	3		55
5/2/81	73	BL	M	6	9		60
5/16/81	72	BT	GF	NA			52
		BT	NA	NA		NA	
		SH	M	6	1.25		49.5
		ST	F	9	11	7	3.5
5/22/81	73	BL	M	6	5		56
		BT	F	6	1.5		54
		BT	F	5	8.5		50
		BN	NA		48	NA	
5/30/81	73	BL	M	5	7		48.75
		BT	F	5	6.5	4	1.5
		BT	M	4	4.75		38.25
		SH	M	8		6	
		SH	NA	9		NA	
		SH	M	8	1.25	5	8.5

6/13/81	84	BL	GF	8	10.5	6	6.5
		BT	F	5	8	4	1.5
		SH	M	8	7.5	6	1
		SH	M	8	0.5	5	8
6/27/81	84	BL	M	6	4.75		56
		BT	F	6	1		54
7/12/81	81	T	M	7	8	5	5
7/18/81	84	D	M	10	10	8	1
7/21/81	80	BL	M	7	0	5	1
		T	NA	5		NA	
7/25/81	80	BL	F	7	9.5	5	9
		D	F	11	1	8	3.5
		D	M	9	11.5	7	4
8/15/81	83	BL	F	7	6.25	5	6
		BL	M	6	3.25	4	7
		BL	M	5	9		59
		T	F	8	3	5	11
		GH	F	8	5	5	10
8/22/81	84	BL	F	6	6		57
		T	F	5	0		41
9/12/81	83	N	NA	8		NA	
		N	NA	10		NA	
9/22/81	80	BL	M	8	4	6	2
		T	F	7	0	5	0.5
9/23/81	80	T	M	5	3		44
10/5/81	76	SB	F	6	10.5	5	1
		SB	F	7	9	5	10
10/7/81	79	D	M	10	1.5	7	7
10/10/81	78	BT	GF	5	9		51
		T	F	7	1.5		60
10/21/81	74	SB	NA	NA		NA	
11/1/81	68	D	NA	NA		NA	
		D	NA	NA		NA	
		D	NA	NA		NA	
		D	NA	NA		NA	
		T	NA	NA		NA	
		T	NA	NA		NA	
11/7/81	68	BL	M	6	7.5		58
		D	F	10	9	9	0
11/18/81	68	T	M	9	1.5	6	7

5/1/82	72	BN	GF		51.75		39
		L	GF	9	0.5	6	10.5
		BT	GF	5	5.5		47.5
		SB	F	6	6		57.5
		SB	F	6	10		59.5
		BT	F		59		43
5/6/82	80	BT	M	4	11		43
		BT	M	4	8.5		41
		SH	M	6	11		58
5/9/82	74	BT	GF	5	1.5		45
		BT	M		58		42.5
		BL	M	5	9.5		50.5
5/11/82	74	T	F		51.5		35.5
		T	F		51		35
5/12/82	75	T	F		51.75		35.75
5/23/82	77	BL	M	5	11		51.5
		SH	M	8	7.5	6	1
5/27/82	77	T	F	7	3.5	5	3
		SH	M	8	3	5	9
		BL	F	5	7		49
5/28/82	77	BL	F	6	1.5		54
6/12/82	81	SP	M	6	5		57
		D	M	10	7	7	8
7/2/82	78	BL	M	5	9		50.5
		BT	M	5	3		46
7/8/82	80	BL	F	6	9		59
7/9/82	77	T	F	12		9	2
7/25/82	81	T	F	10	4	7	5
		T	F	6	10.5		59.5

8/15/82	83	SB	F	6	0.5		53
		T	F	5			41.5
8/22/82	84	BL	M	7	4.5	5	4.5
		T	F	7	8	5	6
8/29/82	84	T	F	7	1	5	
		T	F	6	7		55.5
9/18/82	82	SB	F	7	5	5	6
		BT	F	5	8		49.5
9/26/82	81	SB	F	6	6		58
		SB	F	7	4	5	5
		SB	F	6	2.5		54.5
		SB	F	7	5	5	5.5
		T	F	6	8		56
10/11/82	79	D	M	10	3	7	6
10/14/82	80	BN	M		47		35
		SB	M	6	11.5	5	1
10/15/82	82	T	M	5	7.5		47.5
		SB	GF	6	11	5	1
11/7/82	68	T	M	6	11.5	5	1
		T	M	7	1	5	1
		T	F	6	10		58
		SB	F	5	9.5		51
		SB	F	6	6.5		58
		BN	F		36	NA	

4/20/83	67	SB	F	NA		5	
5/9/83	71	BL	F	5	10.5		51
5/10/83	71	BL	F	6	4		54
5/29/83	77	SH	NA	8	7.5	6	
		GH	F	9		6	2
5/31/83	77	BL	M	6	4		56
6/11/83	76	BL	M	6	5		56.5
		BT	F	6	5		57
		BT	F	6	0.5		53
		GH	F	7	4	5	
7/4/83	81	BL	M	5	11		51
		BT	F	5	10		52
7/9/83	81	BL	F	6			53
		BL	M	5	7.5		49
		BL	M	6	2		54
7/11/83	81	GH	M	9	10	6	10
		SB	M	6	10	5	0.5
		BL	F	5	9		50
		BL	F	6	1.5		54
7/17/83	83	SB	M	6	5.5		57
7/23/83	83	BN	M		40		29
		T	F	11	0.5	8	3
		T	F	5	11.5		50
		T	F	6	0.5		51
8/7/83	84	SB	F	6	3.5		55
		SB	M	6	2.5		54
		SB	F	7	4	5	6
		T	M	7	10	5	8
8/14/83	84	BL	F	5	10		50.5
		SH	M	8		5	6
		SB	M	6	11	5	1
8/20/83	84	BL	F	6	9.5	4	11.5
8/21/83	84	SB	M	6	10.5	5	1
		SB	M	7	4	5	5.5
9/3/83	86	SB	M	6	2		53.5
		SB	M	6	6.5		57
		SB	M	6			53
		SB	M	6	10	5	1
9/17/83	81	SB	M	6	0.25		53.5
		SB	F	7	3.5	5	6
		SB	M	6	10	5	

A1.6

10/1/83	78	GH	F	10	7	7	4
		D	F	11		8	1.5
		SB	M	6	10	5	
		SB	M	6	7		58.5
		SB	F	7	2.5	5	4.5
		SB	GF	6	10	5	
		SB	F	5	11		52
		SB	M	5	7		49
10/3/83	78	SB	M	6	2.75		55
		SB	F	6	9		59.5
10/7/83	76	SB	F	7	4	5	5.5
		SB	GF	7		5	2
		T	F	6	7		56.5
10/12/83	74	SB	F	6	6		56.5
		SB	F	6	4.5		56
11/13/83	69	SB	F	6	10.5		60
		SB	F	6	11.5		61.5
		SB	M	6	11.5		61.5

5/11/84	74	BT	F	5	3		46
5/19/84	78	BT	F	5	2		45
		BN	GF	4	3.5		39
6/8/84	78	BT	F	5	6		49
		BL	F	7		5	1
6/11/84	78	BT	NA	5		NA	
		BT	F	5	10		51
		GH	M	10	3	7	3
6/12/84	79	GH	GF	10	3	7	4
6/17/84	82	GH	F	9	10.5	6	10
		BT	F	5	9.5		52
		BT	NA	5	3.5		47
6/26/84	83	SB	F	6	0.5		54
		SB	GF	6	6		57
		SH	M	8	2	5	8.5
		BN	NA	NA		NA	
7/9/84	83	SB	M	6	3.5		55.5
7/10/84	83	SP	NA	8		NA	
7/14/84	83	D	F	10	8		93
		SB	F	7	1		63
8/20/84	82	BT	GF	5	9		51
8/31/84	82	N	M	8	10	6	4
		GH	F	10	4	7	3.5
		SB	F	7	5	5	7
9/10/84	82	SB	M	6	0.5		54
		SB	F	7	3	5	4.5
10/2/84	78	SB	GF	6	10.5	5	1.5
		D	F	8	10	6	4
		D	F	10	4	7	8
10/3/84	72	SB	M	6	1		54
		SB	F	6			53
		SB	F	5	7.5		48.5
10/29/84	78	SB	F	7	2	5	3
		SB	M	6	6.5		57.5
		SB	M	5	10		52
		SB	M	6	1.5		54.5
10/31/84	78	SB	M	5	8		50
		SB	M	5	11		52
		GH	M	10	1	7	1.5
		T	F	6	8.5		56.5
11/4/84	78	SB	F	6	4.5		56.5
		T	M	7	7	5	7
		D	F	9	1	6	6
11/14/84	72	T	F	6	8		57
		T	F	6	4		54.5
12/12/84	62	SB	M	6	9		59

4/21/85	68	BL	F	6	3.5		54.5
		BT	F		59.5		43
		BT	F	5	1.5		45.5
		BT	F	6	1.5		54.5

5/26/85	75	BT	F	6	0		53
		BT	M	4	2		36.5
		T	F	4	1		33
5/27/85	75	T	F	4	2.5		35
6/23/85	79	SB	F	6	3.5		55.5
		SH	M	7	4		61
		BT	M	5	6.5		49
7/1/85	78	BT	F	5	2.5	4	5.5
		SB	F	5	9		50.5
		SH	M	8	7.5	6	1
		D	M	9	8.5	7	2.5
		D	F	10	11	8	0.5
7/3/85	77	D	F	11	7	8	7
		D	F	11	2	8	2.5
		BL	F	7	11.5	5	10.5
		D	F	9	3	6	8
7/14/85	78	BL	F	6	4		55
7/21/85	84	BL	M	6	8		59
7/27/85	80	SB	M	6	1		53.5
		BL	M	6	3.5		54
8/24/85	85	BL	F	6	10		60.5
		SB	M	6	7		58
9/7/85	80	BT	M	3	11		34
		BT	F	5	4		47
9/25/85	78	SB	F	6	5.5		56
		SB	F	6	5.5		57
		SH	M	8	2.5	5	9
10/7/85	71	SB	M	6	0		52
10/10/85	73	SB	F	6	0.5		54
10/11/85	71	SB	F	6	7		58
		SB	F	5	9		50.5
10/13/85	71	SB	F	7	3	5	5
		SB	M	6	3		56
		SB	F	6	1		53.5
		SB	M	5	7		49
		SB	F	6	2.5		54.5
10/14/85	70	T	F	5	6		47
10/23/85	73	BT	F	5	7.5		49
		SB	F	6	0		53
		SB	M	5	11.5		52
		SB	M	6	6		57.5
		SB	M	5	6.5		48.5
		SB	M	5	9.5		50
12/8/85	61	SB	M	6	0		53.5
		SB	M	5	11.5		53
		SB	M	5	1.5		54.5

3/29/86	66	T	F	5	8		48
		T	M	5	11		50
		T	M	4	5		37
		SB	F	7	6	5	6
		SB	GF	6	11.5	5	0.5
4/8/86	69	BT	GF	5	11		52.5
		BT	F	4	10		41.5
4/27/86	68	BT	F	4	9		41
		BT	F	5	8		48.5
5/21/86	79	BL	F	6			53
5/22/86	79	BT	GF	5	10		52.5
		SB	F	6	2		55
		SB	F	7	2	5	4
5/23/86	78	BT	NA	NA		NA	
5/26/86	74	BT	F	5	9.5		50.5
		BT	M	4	4		37.5
		T	M	6	10.5		58
6/3/86	80	BT	NA	6		NA	
		SH	NA	7		NA	
		BL	F	7	5	5	5.5
6/4/86	80	BL	F	6	10		59
		T	F	5	9		49
		T	F	6	3.5		54

7/4/86	80	SH	M	6	11		58.5
7/6/86	82	SB	F	6			52.75
		T	F	8	6	6	2
7/8/86	82	SB	M	6			53
		SB	M	6	2.5		54
		SB	M	6	1		54
		BL	M	6	1.5		53.5
7/9/86	82	T	F	8		5	10
7/20/86	83	BT	M	5	2.75		44.75
		SB	M	6	5.75		57
7/22/86	83	D	F	10		NA	
		BL	F	6	10	5	
		SB	F	5	10.5		51.5
8/8/86	86	BL	M	6	7		57.5
8/25/86	85	T	F	7	2	5	1
8/27/86	85	T	M	8	6	6	1
8/29/86	84	T	F	8	5	6	2
		T	F	9	2	6	7.5
9/5/86	84	SB	F	5	10.5		52
9/13/86	82	BT	GF	NA		NA	
		SB	F	7	3.5	5	5
9/14/86	82	BT	GF	5	10		51.5

5/3/87	69	BT	F	6	2.5	4	7
5/12/87	74	BT	NA	6		NA	
		BT	M	5	4.5		46.5
5/14/87	73	BL	GF	8	3	6	2.5
		BL	M	6	1.5	4	4
5/15/87	73	BT	F	5	6		48
		BT	F	5	8		49
		GH	F	9	3	6	4
5/24/87	78	BT	M	4	6.75		40
		SH	M	8	4	5	10
5/31/87	70	BT	F	6	2.75		55
		GH	GF	11	3.5	8	2.5
		SB	F	7	0.5	5	3.5
6/16/87	82	BT	M	4	1		35.5
		SB	F	4	3		37
6/19/87	79	SP	F	7	0.5	5	2.5
6/20/87	84	BT	NA	NA		NA	
		BT	NA	NA		NA	
		BL	F	7	4	5	5
6/21/87	84	BL	F	6	10	5	
		BL	F	6	0.5		52.5
6/28/87	82	SB	M	6	2.5		54.5
7/6/87	83	SB	F	6	2.5		55
7/18/87	85	SH	M	8	5	6	
7/26/87	86	SB	M	6	0.25		53.5
		T	F	8	4	6	
8/9/87	85	SB	M	6	9	5	
		BT	F	4	8		40.5
		SB	F	6	5		56
8/23/87	86	T	M	3	4		27
9/27/87	83	BL	F	6	8.5		58.5
		SB	F	6	4		56
		SB	M	6	1		53.5
		SB	M	5	6.5		49
		SB	F	6			52.5
10/1/87	80	SB	F	6	6		57.5
10/17/87	78	SB	F	5	8		49.5
		SB	F	5	9.5		50.5
		SB	M	5	11.5		53
		SB	M	6			52.5
		SB	M	6	2		54.5
		SB	M	6	2.5		55
		SB	M	6	3		55
		SB	M	6	4		55.5
		SB	F	6	4.5		57
		SB	F	6	8.5		58.5

3/6/88	60	T	F	5	7.5		47
4/16/88	64	BT	F	4	3.5		37.5
		SB	F	6			53
5/8/88	74	BT	F	4	8		40
		BT	F	4	9		41
5/14/88	74	BL	F	7	0.5	5	2
		BL	M	7	2.5	5	3
		BL	NA	8	7	6	4.5
		BT	NA	NA		NA	
		BT	NA	NA		NA	
		BT	NA	NA		NA	
		BT	NA	NA		NA	
5/30/88	76	BT	M	5	5		47
6/19/88	78	BT	F	4	7		39.5
6/26/88	80	BL	M	7	9	5	9
		BT	NA	NA		NA	
		BL	F	6	9		58
		BL	F	7		5	1
7/13/88	83	BT	NA	6		NA	
		BT	F	6		NA	
		BL	F	9		NA	
		SH	M	9		NA	
7/24/88	83	BT	F	6		NA	
		SB	M	6	7.5		58
8/20/88	83	T	F	7	10	5	9
		T	F	8	8	6	4
8/28/88	86	BT	F	6	3.5		54.75
		BT	GF	6	4		55
9/7/88	83	BT	F	4			34
		SB	F	6	5		56.5
		BN	M		45.5		34
		BN	F		46		34
		BN	M		47		35.5
10/9/88	73	SH	M	7	5		71
10/11/88	72	SB	M	6	4		56
		SB	M	6	4.5		55.5
		SB	F	6			52.5
10/16/88	68	SB	M	6	4.5		56.5
10/23/88	68	SB	M	6	4		55.5
10/30/88	68	T	M	6	10		57
4/14/89	68	T	M	4	6.5		37.5
		T	M	5	8		48
		T	M	6			51
		T	M	6	1.5		51.5
		BT	F	4	3		36
4/16/89	68	T	F	4	1.5		34
		T	F	4	3		35
		T	M	4	11.5		41.5
		BT	GF	6			53
4/23/89	69	BT	F	5	7.75		49
		BT	GF	5	7.75		49.5
8/20/89	82	D	M	10	3	7	7
9/17/89	83	BL	F	6			52.5

Appendix 2.

Crooke longline data arranged by species. W.T.: water temperature. Species: BL, bull; BN, blacknose; BT, blacktip; D, dusky; GH, great hammerhead; L, lemon; N, nurse; SB, sandbar; SH, scalloped hammerhead; SN, sharpnose; SP, spinner; ST, sand tiger; T, tiger. Sexes: F, nongravid female; GF, gravid female; M, male. Lengths: TCL, total contour length; PCCL, precaudal contour length; FL, fork length.

<u>DATE</u>	<u>W.T.</u>	<u>SP</u>	<u>SEX</u>	<u>TCL(ft)</u>	<u>PCCL(ft)</u>	<u>FL(ft)</u>
9/ 7/75	85	BL	M	7.7		6.3
9/10/75	84	BL	F	6.5		5.5
9/10/75	84	BL	M	6.0		5.0
9/11/75	84	BL	M	6.7		5.7
9/11/75	84	BL	F	6.5		5.4
6/11/76	78	BL	F	6.3		5.3
6/24/76	83	BL	M	6.6		5.5
4/ 9/77	67	BL	F	5.3		4.1
5/29/77	80	BL	F	6.3		5.4
5/29/77	80	BL	M	6.2		5.1
6/ 2/77	80	BL	M	6.0		4.9
6/ 2/77	80	BL	F	7.6		6.0
9/18/77	84	BL	M	6.0		5.1
5/21/78	76	BL	M	6.2		5.0
5/30/78	77	BL	F	6.2		5.1
5/30/78	77	BL	F	6.2		4.9
5/30/78	77	BL	F	6.2		4.5
5/30/78	77	BL	F	5.5		4.6
5/30/78	77	BL	M	5.5		4.5
5/31/78	77	BL	M	5.6		4.6
5/31/78	77	BL	M	6.5		5.4
5/31/78	77	BL	M	6.2		5.0
6/ 1/78	77	BL	F	6.9		5.6
6/ 1/78	77	BL	F	8.7		7.5
6/ 1/78	77	BL	F	9.0		7.8
6/ 1/78	77	BL	F	6.4		5.2
6/ 2/78	78	BL	GF	8.2		7.0
6/11/78	81	BL	F	6.0		5.1
6/25/78	84	BL	M	6.4		4.9
7/ 4/78	85	BL	F	6.5		5.5
7/ 4/78	85	BL	F	8.6		7.6
7/ 4/78	85	BL	F	8.9		7.8
7/30/78	85	BL	M	5.4		4.4
8/13/78	84	BL	M	6.0		5.0
8/13/78	84	BL	M	5.8		4.9
8/20/78	85	BL	F	7.0		5.8
9/ 3/78	84	BL	F	7.4		6.1
9/ 5/78	84	BL	M	5.9		4.8
9/15/78	83	BL	F	5.9		4.9
5/ 6/79	71	BL	F	6.3	4.5	
5/ 6/79	71	BL	M	6.0	4.3	
5/13/79	74	BL	M	8.3	7.3	
5/29/79	75	BL	F	6.5	4.8	
5/31/79	75	BL	GF	8.9	6.7	
5/31/79	75	BL	M	8.9	5.0	
6/ 9/79	80	BL	M	5.6	4.0	
6/ 9/79	80	BL	F	8.7	6.5	
6/16/79	82	BL	F	8.5	6.3	
7/ 6/79	82	BL	M	6.4	4.6	
7/ 6/79	82	BL	F	8.8	6.8	
4/ 4/80	80	BL	F	8.6	6.5	
6/16/80	81	BL	M	6.5	4.8	
8/23/80	84	BL	M	6.0	4.3	
8/31/80	82	BL	M	5.8	4.2	
5/ 2/81	73	BL	M	6.8	5.0	
5/22/81	73	BL	M	6.4	4.7	
5/30/81	73	BL	M	5.6	4.1	
6/13/81	84	BL	GF	8.9	6.5	
6/27/81	84	BL	M	6.4	4.7	
7/21/81	80	BL	M	7.0	5.1	

7/25/81	80	BL	F	7.8	5.8
8/15/81	83	BL	F	7.5	5.5
8/15/81	83	BL	M	6.3	4.6
8/15/81	83	BL	M	5.8	4.9
8/22/81	84	BL	F	6.5	4.8
9/22/81	80	BL	M	8.3	6.2
11/ 7/81	68	BL	M	6.6	4.8
5/ 9/82	74	BL	M	5.8	4.2
5/23/82	77	BL	M	5.9	4.3
5/27/82	77	BL	F	5.6	4.1
5/28/82	77	BL	F	6.1	4.5
7/ 2/82	78	BL	M	5.8	4.2
7/ 8/82	80	BL	F	6.8	4.9
8/22/82	84	BL	M	7.4	5.4
5/ 9/83	71	BL	F	5.9	4.3
5/10/83	71	BL	F	6.3	4.5
5/31/83	77	BL	M	6.3	4.7
6/11/83	76	BL	M	6.4	4.7
7/ 4/83	81	BL	M	5.9	4.3
7/ 9/83	81	BL	F	6.0	4.4
7/ 9/83	81	BL	M	5.6	4.1
7/ 9/83	81	BL	M	6.2	4.5
7/11/83	81	BL	F	5.8	4.2
7/11/83	81	BL	F	6.1	4.5
8/14/83	84	BL	F	5.8	4.2
8/20/83	84	BL	F	6.8	5.0
6/ 8/84	78	BL	F	7.0	5.1
4/21/85	68	BL	F	6.3	4.5
7/ 3/85	77	BL	F	8.0	5.9
7/14/85	78	BL	F	6.3	4.6
7/21/85	84	BL	M	6.7	4.9
7/27/85	80	BL	M	6.3	4.5
8/24/85	85	BL	F	6.8	5.0
5/21/86	79	BL	F	6.0	4.4
6/ 3/86	80	BL	F	7.4	5.5
6/ 4/86	80	BL	F	6.8	4.9
7/ 8/86	82	BL	M	6.1	4.5
7/22/86	83	BL	F	6.8	5.0
8/ 8/86	86	BL	M	6.6	4.8
5/14/87	73	BL	GF	8.3	6.2
5/14/87	73	BL	M	6.1	4.3
6/20/87	84	BL	F	7.3	5.4
6/21/87	84	BL	F	6.8	5.0
6/21/87	84	BL	F	6.0	4.4
9/27/87	83	BL	F	6.7	4.9
5/14/88	74	BL	F	7.0	5.2
5/14/88	74	BL	M	7.2	5.3
5/14/88	74	BL	NA	8.6	6.4
6/26/88	80	BL	M	7.8	5.8
8/26/88	80	BL	F	6.8	4.8
6/26/88	80	BL	F	7.0	5.1
7/13/88	83	BL	F	9.0	
9/17/89	83	BL	F	6.0	4.4

6/ 2/78	78	BN	GF	4.3	
4/29/79	71	BN	F	3.9	2.9
5/13/79	74	BN	GF	3.9	2.9
5/13/79	74	BN	GF	3.8	2.8
5/22/81	73	BN	NA	4.0	
5/ 1/82	72	BN	GF	4.3	3.3
10/14/82	80	BN	M	3.9	2.9
11/ 7/82	68	BN	F	3.0	
7/23/83	83	BN	M	3.3	2.4
5/19/84	78	BN	GF	4.3	3.3
6/26/84	83	BN	NA		
9/ 7/88	83	BN	M	3.8	2.8
9/ 7/88	83	BN	F	3.8	2.8
9/ 7/88	83	BN	M	3.9	3.0

9/12/75	84	BT	GF	6.2		5.1
9/13/75	85	BT	GF	5.5		4.5
10/ 1/76	78	BT	F	5.8		4.8
5/ 1/77	76	BT	F	6.0		5.0
5/15/77	79	BT	M	5.3		4.3
5/31/77	80	BT	F	5.6		4.8
6/ 1/77	80	BT	F	5.5		4.5
6/ 2/77	80	BT	F	5.2		4.5
6/ 2/77	80	BT	F	5.8		4.9
6/ 3/77	80	BT	F	6.1		4.9
5/31/78	77	BT	F	5.3		4.3
6/ 1/78	77	BT	F	5.0		
6/ 1/78	77	BT	F	5.3		4.5
6/ 2/78	78	BT	NA	5.8		
6/25/78	84	BT	F	5.5		4.5
6/25/78	84	BT	F	6.0		4.9
7/ 1/78	85	BT	M	5.5		4.5
7/ 1/78	85	BT	M	3.8		3.2
7/ 2/78	85	BT	F	6.0		4.9
9/ 4/78	82	BT	F	5.6		4.6
9/ 5/78	84	BT	F	4.7		3.9
4/29/79	71	BT	F	5.0	3.6	
4/29/79	71	BT	F	4.9	3.5	
4/29/79	71	BT	GF	5.3	3.9	
4/29/79	71	BT	F	5.0	3.6	
4/29/79	71	BT	GF	5.5	4.0	
5/ 6/79	71	BT	GF	5.5	4.1	
5/20/79	75	BT	M	5.3	3.9	
5/22/79	76	BT	F	4.8	3.4	
5/22/79	76	BT	GF	5.7	4.1	
5/26/79	74	BT	F	5.9	4.3	
5/28/79	75	BT	M	5.2	3.8	
6/ 2/79	75	BT	F	5.3	3.8	
6/ 9/79	80	BT	F	4.4	3.1	
6/ 9/79	80	BT	M	4.5	3.3	
6/16/79	82	BT	F	5.6	4.1	
6/16/79	82	BT	F	5.5	4.0	
6/16/79	82	BT	M	4.6	3.3	
6/16/79	82	BT	F	4.4	3.3	
6/17/79	80	BT	F	5.9	4.3	
6/20/79	81	BT	F	5.7	4.2	
6/21/79	81	BT	M	4.7	3.5	
6/23/79	81	BT	M	5.4	4.0	
6/24/79	81	BT	F	5.7	4.2	
6/24/79	81	BT	F	5.6	4.1	
4/25/81	71	BT	GF	6.3	4.6	
5/16/81	72	BT	GF		4.3	
5/16/81	72	BT	NA			
5/22/81	73	BT	F	6.1	4.5	
5/22/81	73	BT	F	5.7	4.2	
5/30/81	73	BT	F	5.5	4.1	
5/30/81	73	BT	M	4.4	3.2	
6/13/81	84	BT	F	5.7	4.1	
6/27/81	84	BT	F	6.1	4.5	
10/10/81	78	BT	GF	5.8	4.3	
5/ 1/82	72	BT	GF	5.5	4.0	
5/ 1/82	72	BT	F	4.9	3.6	
5/ 6/82	80	BT	M	4.9	3.6	
5/ 6/82	80	BT	M	4.7	3.4	
5/ 9/82	74	BT	GF	5.1	3.8	
5/ 9/82	74	BT	M	4.8	3.5	
7/ 2/82	78	BT	M	5.3	3.8	
9/18/82	82	BT	F	5.7	4.1	
6/11/83	76	BT	F	6.4	4.8	
8/11/83	76	BT	F	6.0	4.4	
7/ 4/83	81	BT	F	5.8	4.3	
5/11/84	74	BT	F	5.3	3.8	
5/19/84	78	BT	F	5.2	3.8	
8/ 8/84	78	BT	F	5.5	4.1	

6/11/84	78	BT	NA	5.0	
6/11/84	78	BT	F	5.8	4.3
6/17/84	82	BT	F	5.8	4.3
6/17/84	82	BT	NA	5.3	3.9
8/20/84	82	BT	GF	5.8	4.3
4/21/85	68	BT	F	5.0	3.6
4/21/85	68	BT	F	5.1	3.8
4/21/85	68	BT	F	6.1	4.5
5/26/85	75	BT	F	6.0	4.4
5/26/85	75	BT	M	4.2	3.0
6/23/85	79	BT	M	5.5	4.1
7/ 1/85	78	BT	F	5.2	4.5
9/ 7/85	80	BT	M	3.9	2.8
9/ 7/85	80	BT	F	5.3	3.9
10/23/85	73	BT	F	5.6	4.1
4/ 8/86	69	BT	GF	5.9	4.4
4/ 8/86	69	BT	F	4.8	3.5
4/27/86	68	BT	F	4.8	3.4
4/27/86	68	BT	F	5.7	4.1
5/22/86	79	BT	GF	5.8	4.4
5/23/86	78	BT	NA		
5/26/86	74	BT	F	5.8	4.2
5/26/86	74	BT	M	4.3	3.1
6/ 3/86	80	BT	NA	6.0	
7/20/86	83	BT	M	5.2	3.7
9/13/86	82	BT	GF		
9/14/86	82	BT	GF	5.8	4.3
5/ 3/87	69	BT	F	6.2	4.6
5/12/87	74	BT	NA	6.0	
5/12/87	74	BT	M	5.4	3.9
5/15/87	73	BT	F	5.5	4.0
5/15/87	73	BT	F	5.7	4.1
5/24/87	78	BT	M	4.6	3.3
5/31/87	70	BT	F	6.2	4.6
6/16/87	82	BT	M	4.1	3.0
6/20/87	84	BT	NA		
6/20/87	84	BT	NA		
8/ 9/87	85	BT	F	4.7	3.4
4/16/88	64	BT	F	4.3	3.1
5/ 8/88	74	BT	F	4.7	3.3
5/ 8/88	74	BT	F	4.8	3.4
5/14/88	74	BT	NA		
5/14/88	74	BT	NA		
5/14/88	74	BT	NA		
5/14/88	74	BT	NA		
5/30/88	76	BT	M	5.4	3.9
6/19/88	78	BT	F	4.6	3.3
6/26/88	80	BT	NA		
7/13/88	83	BT	NA	6.0	
7/13/88	83	BT	F	6.0	
7/24/88	83	BT	F	6.0	
8/28/88	86	BT	F	6.3	4.6
8/28/88	86	BT	GF	6.3	4.6
9/ 7/88	83	BT	F	4.0	2.8
4/14/89	88	BT	F	4.3	3.0
4/18/89	68	BT	GF	6.0	4.4
4/23/89	69	BT	F	5.6	4.1
4/23/89	69	BT	GF	5.6	4.1

9/ 7/75	85	D	F	11.5	9.4
9/29/76	79	D	M	10.5	9.1
10/10/78	78	D	M	8.4	7.1
9/10/77	84	D	F	10.8	8.9
9/18/77	84	D	M	10.3	8.2
10/ 5/77	83	D	F	8.5	7.1
10/ 5/77	83	D	F	11.1	9.8
10/ 5/77	83	D	F	11.0	9.3
10/ 7/77	80	D	M	10.3	9.0
10/ 7/77	80	D	F	11.5	9.5
10/30/77	75	D	M	10.1	8.4

11/13/77	72	D	F	10.2		8.7
11/26/77	69	D	F	11.2		9.3
7/30/78	85	D	M	9.8		8.1
7/30/78	85	D	M	10.3		8.7
7/30/78	85	D	M	10.0		8.4
8/20/78	85	D	F	11.5		9.5
10/ 8/78	76	D	M	10.1		8.7
10/14/78	76	D	M	10.3		8.7
10/28/78	73	D	F	7.9		6.7
11/ 4/78	72	D	M	8.3		6.6
11/ 4/78	72	D	F	8.1		6.6
12/17/78	59	D	F	8.5		7.0
12/17/78	59	D	F	11.5		9.9
7/ 7/79	82	D	F	11.0	8.4	
7/ 7/79	82	D	M	10.7	7.8	
7/ 7/79	82	D	M	8.8	6.4	
7/ 7/79	82	D	F	10.9	8.0	
8/ 2/80	84	D	M	10.4	7.8	
9/21/80	79	D	M	9.7	6.6	
10/ 6/80	76	D	M	9.3	6.8	
10/ 6/80	76	D	GF	11.2	8.4	
10/ 6/80	76	D	F	11.5	8.6	
10/ 8/80	77	D	F	11.0	8.2	
10/ 8/80	77	D	M	9.5	7.0	
10/ 8/80	77	D	M	8.8	6.3	
10/11/80	78	D	M	8.5	6.1	
11/ 7/80	74	D	M	10.2	6.8	
11/ 7/80	74	D	F	10.1	6.6	
11/ 7/80	74	D	F	9.0	6.6	
11/22/80	68	D	F	11.2	8.4	
7/18/81	84	D	M	10.8	8.1	
7/25/81	80	D	F	11.1	8.3	
7/25/81	80	D	M	10.0	7.3	
10/ 7/81	79	D	M	10.1	7.6	
11/ 1/81	68	D	NA			
11/ 1/81	68	D	NA			
11/ 1/81	68	D	NA			
11/ 1/81	68	D	NA			
11/ 7/81	68	D	F	10.8	9.0	
6/12/82	81	D	M	10.6	7.7	
10/11/82	79	D	M	10.3	7.5	
10/ 1/83	78	D	F	11.0	8.1	
7/14/84	83	D	F	10.7	7.8	
10/ 2/84	78	D	F	8.8	6.3	
10/ 2/84	78	D	F	10.3	7.7	
11/ 4/84	78	D	F	9.1	6.5	
7/ 1/85	78	D	M	9.7	7.2	
7/ 1/85	78	D	F	10.9	8.0	
7/ 3/85	77	D	F	11.8	8.6	
7/ 3/85	77	D	F	11.2	8.2	
7/ 3/85	77	D	F	9.3	6.7	
7/22/86	83	D	F	10.0		
8/20/89	82	D	M	10.3	7.8	

10/ 1/78	78	GH	F	10.5		8.1
6/ 1/77	80	GH	F	7.4		5.5
9/ 2/78	84	GH	F	7.9		5.9
6/24/79	81	GH	M	6.2	4.3	
6/26/79	81	GH	M	8.6	4.5	
8/16/80	81	GH	F	10.7	7.5	
10/ 6/80	76	GH	F	8.6	8.0	
8/15/81	83	GH	F	8.4	5.8	
5/29/83	77	GH	F	9.0	8.2	
6/11/83	76	GH	F	7.3	5.0	
7/11/83	81	GH	M	9.8	6.8	
10/ 1/83	78	GH	F	10.6	7.3	
6/11/84	78	GH	M	10.3	7.3	
6/12/84	79	GH	GF	10.3	7.3	
6/17/84	82	GH	F	9.9	6.8	
8/31/84	82	GH	F	10.3	7.3	

10/31/84	78	GH	M	10.1	7.1
5/15/87	73	GH	F	9.3	6.3
5/31/87	70	GH	GF	11.3	8.2

5/ 1/82	72	L	GF	9.0	6.9

9/14/75	84	N	M	8.2	6.5
9/14/75	84	N	M	8.6	7.1
8/20/78	85	N	M	8.5	6.8
9/16/78	83	N	M	8.5	6.9
9/12/81	83	N	NA	8.0	
9/12/81	83	N	NA	10.0	
8/31/84	82	N	M	8.8	6.3

9/21/75	81	SB	F	7.0	5.8
10/ 3/75	73	SB	F	6.3	
10/ 1/76	78	SB	GF	7.2	6.1
10/16/76	74	SB	GF	7.4	6.1
11/ 7/76	63	SB	F	7.1	6.0
11/ 7/76	63	SB	F	7.4	6.0
4/ 2/77	68	SB	M	5.0	4.2
4/ 9/77	67	SB	F	7.1	5.7
4/ 9/77	67	SB	F	7.3	6.1
11/13/77	72	SB	M	4.6	4.2
11/26/77	69	SB	F	7.2	6.2
4/16/78	71	SB	F	7.3	6.0
10/14/78	76	SB	F	7.0	5.8
10/25/78	72	SB	M	6.3	5.5
10/27/78	71	SB	F	6.8	5.6
10/27/78	71	SB	F	7.3	6.3
10/27/78	71	SB	F	6.3	5.2
10/28/78	73	SB	F	7.3	6.1
11/ 4/78	72	SB	M	5.6	5.1
11/ 5/78	73	SB	F	7.0	6.0
11/19/78	71	SB	F	6.9	5.9
11/19/78	71	SB	F	6.2	5.3
11/24/78	70	SB	M	6.8	5.7
11/24/78	70	SB	M	5.6	4.8
11/24/78	70	SB	F	6.7	5.8
11/24/78	70	SB	M	5.8	4.8
11/24/78	70	SB	M	6.0	5.1
4/20/80	64	SB	F	5.2	3.8
10/ 8/80	77	SB	GF	7.5	5.5
10/ 8/80	77	SB	F	5.9	4.4
10/ 8/80	77	SB	M	6.3	4.6
10/ 9/80	78	SB	F	6.7	4.9
10/ 9/80	78	SB	F	7.3	5.4
10/11/80	78	SB	GF	7.4	5.7
10/11/80	78	SB	M	6.6	4.8
10/11/80	78	SB	F	6.7	4.9
10/11/80	78	SB	F	7.3	5.5
11/ 7/80	74	SB	M	5.0	
11/ 7/80	74	SB	M	6.1	4.5
11/ 7/80	74	SB	M	5.9	4.3
11/ 7/80	74	SB	F	5.7	4.2
11/ 7/80	74	SB	F	5.5	4.1
11/22/80	68	SB	F	5.6	4.1
12/ 6/80	67	SB	M	5.4	4.0
10/ 5/81	76	SB	F	6.9	5.1
10/ 5/81	76	SB	F	7.8	5.8
10/21/81	74	SB	NA		
5/ 1/82	72	SB	F	6.5	4.8
5/ 1/82	72	SB	F	6.8	5.0
8/15/82	83	SB	F	6.0	4.4
9/16/82	82	SB	F	7.4	5.5
9/26/82	81	SB	F	6.5	4.8
9/26/82	81	SB	F	7.3	5.4
9/26/82	81	SB	F	6.2	4.5
9/26/82	81	SB	F	7.4	5.5
10/14/82	80	SB	M	7.0	5.1

10/15/82	82	SB	GF	6.9	5.1
11/ 7/82	68	SB	F	5.8	4.3
11/ 7/82	68	SB	F	6.5	4.8
4/20/83	67	SB	F		5.0
7/11/83	81	SB	M	6.8	5.0
7/17/83	83	SB	M	6.5	4.8
8/ 7/83	84	SB	F	6.3	4.6
8/ 7/83	64	SB	M	6.2	4.5
8/ 7/83	84	SB	F	7.3	5.5
8/14/83	84	SB	M	6.9	5.1
8/21/83	84	SB	M	6.9	5.1
8/21/83	84	SB	M	7.3	5.5
9/ 3/83	86	SB	M	6.2	4.5
9/ 3/83	86	SB	M	6.5	4.8
9/ 3/83	86	SB	M	6.0	4.4
9/ 3/83	86	SB	M	6.8	5.1
9/17/83	81	SB	M	6.0	4.5
9/17/83	81	SB	F	7.3	5.5
9/17/83	81	SB	M	6.8	5.0
10/ 1/83	78	SB	M	6.8	5.0
10/ 1/83	78	SB	M	6.6	4.9
10/ 1/83	78	SB	F	7.2	5.4
10/ 1/83	78	SB	GF	6.8	5.0
10/ 1/83	78	SB	F	5.9	4.3
10/ 1/83	78	SB	M	5.6	4.1
10/ 3/83	78	SB	M	6.2	4.6
10/ 3/83	78	SB	F	6.8	5.0
10/ 7/83	76	SB	F	7.3	5.5
10/ 7/83	76	SB	GF	7.0	5.2
10/12/83	74	SB	F	6.5	4.7
10/12/83	74	SB	F	6.4	4.7
11/13/83	69	SB	F	6.9	5.0
11/13/83	69	SB	F	7.0	5.1
11/13/83	69	SB	M	7.0	5.1
6/26/84	83	SB	F	6.0	4.5
6/26/84	83	SB	GF	6.5	4.8
7/ 9/84	83	SB	M	6.3	4.6
7/14/84	83	SB	F	7.1	5.3
8/31/84	82	SB	F	7.4	5.6
9/10/84	82	SB	M	6.0	4.5
9/10/84	82	SB	F	7.3	5.4
10/ 2/84	78	SB	GF	6.9	5.1
10/ 3/84	72	SB	M	6.1	4.5
10/ 3/84	72	SB	F	6.0	4.4
10/ 3/84	72	SB	F	5.6	4.0
10/29/84	78	SB	F	7.2	5.3
10/29/84	78	SB	M	6.5	4.8
10/29/84	78	SB	M	5.8	4.3
10/29/84	78	SB	M	6.1	4.5
10/31/84	78	SB	M	5.7	4.2
10/31/84	78	SB	M	5.9	4.3
11/ 4/84	78	SB	F	6.4	4.7
12/12/84	62	SB	M	6.8	4.9
6/23/85	79	SB	F	6.3	4.6
7/ 1/85	78	SB	F	5.8	4.2
7/27/85	80	SB	M	6.1	4.5
8/24/85	85	SB	M	6.6	4.8
9/25/85	78	SB	F	6.5	4.7
9/25/85	78	SB	F	6.5	4.8
10/ 7/85	71	SB	M	6.0	4.3
10/10/85	73	SB	F	6.0	4.5
10/11/85	71	SB	F	6.6	4.8
10/11/85	71	SB	F	5.8	4.2
10/13/85	71	SB	F	7.3	5.4
10/13/85	71	SB	M	6.3	4.7
10/13/85	71	SB	F	6.1	4.5
10/13/85	71	SB	M	5.6	4.1
10/13/85	71	SB	F	6.2	4.5
10/23/85	73	SB	F	6.0	4.4
10/23/85	73	SB	M	6.0	4.3

10/23/85	73	SB	M	6.5	4.8
10/23/85	73	SB	M	5.5	4.0
10/23/85	73	SB	M	5.8	4.2
12/ 8/85	61	SB	M	6.0	4.5
12/ 8/85	61	SB	M	6.0	4.4
12/ 8/85	61	SB	M	5.1	4.5
3/29/86	66	SB	F	7.5	5.5
3/29/86	66	SB	GF	7.0	5.0
5/22/86	79	SB	F	6.2	4.6
5/22/86	79	SB	F	7.2	5.3
7/ 6/86	82	SB	F	6.0	4.4
7/ 8/86	82	SB	M	6.0	4.4
7/ 8/86	82	SB	M	6.2	4.5
7/ 8/86	82	SB	M	6.1	4.5
7/20/86	83	SB	M	6.5	4.8
7/22/86	83	SB	F	5.9	4.3
9/ 5/86	84	SB	F	5.9	4.3
9/13/86	82	SB	F	7.3	5.4
5/31/87	70	SB	F	7.0	5.3
6/16/87	82	SB	F	4.3	3.1
6/28/87	82	SB	M	6.2	4.5
7/ 8/87	83	SB	F	6.2	4.6
7/26/87	86	SB	M	6.0	4.5
8/ 9/87	85	SB	M	6.8	5.0
8/ 9/87	85	SB	F	6.4	4.7
9/27/87	83	SB	F	6.3	4.7
9/27/87	83	SB	M	6.1	4.5
9/27/87	83	SB	M	5.5	4.1
9/27/87	83	SB	F	6.0	4.4
10/ 1/87	80	SB	F	6.5	4.8
10/17/87	78	SB	F	5.7	4.1
10/17/87	78	SB	F	5.8	4.2
10/17/87	78	SB	M	6.0	4.4
10/17/87	78	SB	M	6.0	4.4
10/17/87	78	SB	M	6.2	4.5
10/17/87	78	SB	M	6.2	4.6
10/17/87	78	SB	M	6.3	4.6
10/17/87	78	SB	M	6.3	4.6
10/17/87	78	SB	F	6.4	4.8
10/17/87	78	SB	F	6.7	5.0
4/16/88	64	SB	F	6.0	4.4
7/24/88	83	SB	M	6.6	4.8
9/ 7/88	83	SB	F	6.4	4.7
10/11/88	72	SB	M	6.3	4.7
10/11/88	72	SB	M	6.4	4.6
10/11/88	72	SB	F	6.0	4.4
10/16/88	68	SB	M	6.4	4.7
10/23/88	68	SB	M	6.3	4.6

5/21/76	73	SH	M	8.2	5.9
5/ 1/77	76	SH	M	8.2	6.1
5/15/77	79	SH	M	8.2	6.3
6/ 3/77	80	SH	M	7.8	5.8
6/ 2/78	78	SH	M	7.3	5.5
5/27/79	74	SH	M	8.3	6.3
5/27/79	74	SH	M	8.5	6.5
6/ 2/79	75	SH	M	8.2	5.8
6/17/79	80	SH	M	8.5	5.9
6/20/79	81	SH	M	8.0	5.7
6/29/79	81	SH	NA	7.5	5.5
5/16/81	72	SH	M	6.1	4.1
5/30/81	73	SH	M	8.0	6.0
5/30/81	73	SH	NA	9.0	
5/30/81	73	SH	M	8.1	5.7
6/13/81	84	SH	M	8.6	6.1
6/13/81	84	SH	M	8.0	5.7
5/ 6/82	80	SH	M	6.9	4.8
5/23/82	77	SH	M	8.6	6.1
5/27/82	77	SH	M	8.3	5.8
5/29/83	77	SH	NA	8.6	6.0

8/14/83	84	SH	M	8.0	5.5
6/26/84	83	SH	M	8.2	5.7
8/23/85	79	SH	M	7.3	5.1
7/ 1/85	78	SH	M	8.6	6.1
9/25/85	78	SH	M	8.2	5.8
6/ 3/86	80	SH	NA	7.0	
7/ 4/86	80	SH	M	6.9	4.9
5/24/87	78	SH	M	8.3	5.8
7/18/87	85	SH	M	8.4	6.0
7/13/88	83	SH	M	9.0	
10/ 9/88	73	SH	M	7.4	5.9

6/ 9/79	80	SN	M	3.2	2.4

5/ 1/77	76	SP	F	3.9	3.2
10/30/77	75	SP	F	5.9	4.8
9/17/78	82	SP	F	5.4	4.4
6/12/82	81	SP	M	6.4	4.8
7/10/84	83	SP	NA	8.0	
6/19/87	79	SP	F	7.0	5.2

4/ 8/79	67	ST	F	10.0	8.3
5/16/81	72	ST	F	9.9	7.3

9/ 7/75	85	T	F	7.4	5.7
9/ 7/75	85	T	F	8.7	5.8
9/ 9/75	84	T	F	7.2	5.8
9/11/75	84	T	F	8.9	7.5
9/12/75	84	T	F	7.0	5.6
9/13/75	85	T	F	7.5	6.1
9/13/75	85	T	F	7.4	6.1
9/13/75	85	T	F	8.9	7.4
8/16/76	80	T	F	5.9	4.8
10/10/76	78	T	F	5.5	4.3
11/ 7/76	63	T	M	7.1	5.9
9/18/77	84	T	F	7.3	6.0
10/ 7/77	80	T	F	7.4	6.1
11/13/77	72	T	F	6.5	5.2
11/13/77	72	T	F	7.8	6.3
11/26/77	69	T	M	10.3	8.8
11/26/77	69	T	F	12.1	10.5
6/ 2/78	78	T	M	9.5	7.9
10/14/78	76	T	F	7.2	5.7
10/24/78	70	T	F	6.7	5.4
10/24/78	70	T	F	7.8	6.1
11/ 4/78	72	T	M	6.6	5.5
11/19/78	71	T	F	7.4	6.0
11/24/78	70	T	M	6.6	5.4
11/24/78	70	T	F	6.5	6.0
11/24/78	70	T	M	5.9	4.8
7/ 4/80	80	T	GF	13.0	10.0
9/ 1/80	81	T	F	9.0	6.5
10/ 6/80	76	T	F	8.4	6.0
10/ 8/80	77	T	F	7.5	5.4
10/ 9/80	78	T	F	5.1	3.5
10/ 9/80	78	T	M	7.0	
10/ 9/80	78	T	F	8.8	6.3
10/ 9/80	78	T	M	8.6	6.2
10/11/80	78	T	F	5.1	3.7
11/ 7/80	74	T	M	6.3	4.4
7/12/81	81	T	M	7.7	5.4
7/21/81	80	T	NA	5.0	
8/15/81	83	T	F	8.3	5.9
8/22/81	84	T	F	5.0	3.4
9/23/81	80	T	F	7.0	5.0
9/23/81	80	T	M	5.3	3.7
10/10/81	78	T	F	7.1	5.0
11/ 1/81	68	T	NA		
11/ 1/81	68	T	NA		
11/18/81	66	T	M	9.1	6.6

5/11/82	74	T	F	4.3	3.0
5/11/82	74	T	F	4.3	2.9
5/12/82	75	T	F	4.3	3.0
5/27/82	77	T	F	7.3	5.3
7/ 9/82	77	T	F	12.0	9.2
7/25/82	81	T	F	10.3	7.4
7/25/82	81	T	F	6.9	5.0
8/15/82	83	T	F	5.0	3.5
8/22/82	84	T	F	7.7	5.5
8/29/82	84	T	F	7.1	5.0
8/29/82	84	T	F	6.6	4.6
9/26/82	81	T	F	6.7	4.7
10/15/82	82	T	M	5.6	4.0
11/ 7/82	68	T	M	7.0	5.1
11/ 7/82	68	T	M	7.1	5.1
11/ 7/82	68	T	F	6.8	4.8
7/23/83	83	T	F	11.0	8.3
7/23/83	83	T	F	6.0	4.2
7/23/83	83	T	F	6.0	4.3
8/ 7/83	84	T	M	7.8	5.7
10/ 7/83	76	T	F	6.6	4.7
10/31/84	78	T	F	6.7	4.7
11/ 4/84	78	T	M	7.6	5.6
11/14/84	72	T	F	6.7	4.8
11/14/84	72	T	F	6.3	4.5
5/26/85	75	T	F	4.1	2.8
5/27/85	75	T	F	4.2	2.9
10/14/85	70	T	F	5.5	3.9
3/29/86	66	T	F	5.7	4.0
3/29/86	66	T	M	5.9	4.2
3/29/86	66	T	M	4.4	3.1
5/26/86	74	T	M	6.9	4.8
6/ 4/86	80	T	F	5.8	4.1
6/ 4/86	80	T	F	6.3	4.5
7/ 6/86	82	T	F	8.5	6.2
7/ 9/86	82	T	F	8.0	5.8
8/25/86	85	T	F	7.2	5.1
8/27/86	85	T	M	8.5	6.1
8/29/86	84	T	F	8.4	6.2
8/29/86	84	T	F	9.2	6.6
7/26/87	86	T	F	8.3	6.0
8/23/87	86	T	M	3.3	2.3
3/ 6/88	60	T	F	5.6	3.9
8/20/88	83	T	F	7.8	5.8
8/20/88	83	T	F	8.7	6.3
10/30/88	68	T	M	6.8	4.8
4/14/89	68	T	M	4.5	3.1
4/14/89	68	T	M	5.7	4.0
4/14/89	68	T	M	6.0	4.3
4/14/89	68	T	M	6.1	4.3
4/16/89	68	T	F	4.1	2.8
4/16/89	68	T	F	4.3	2.9
4/16/89	68	T	M	5.0	3.5