

Use of marine protected areas and exclusive economic zones in the subtropical western North Atlantic Ocean by large highly mobile sharks

Fiona Graham, Patrick Rynne, Maria Estevanez, Jiangan Luo, Jerald S. Ault and Neil Hammerschlag

SEDAR77-RD03

Received: 5/27/2021





Use of marine protected areas and exclusive economic zones in the subtropical western North Atlantic Ocean by large highly mobile sharks

Fiona Graham¹, Patrick Rynne¹, Maria Estevez¹, Jiangang Luo¹, Jerald S. Ault¹ and Neil Hammerschlag^{1,2*}

¹Rosenstiel School of Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, FL, 33149, USA, ²Leonard and Jayne Abess Center for Ecosystem Science and Policy, University of Miami, PO Box 248203, Coral Gables, FL, 33146, USA

ABSTRACT

Study aim and location Many populations of highly mobile marine fishes, including large sharks, are experiencing declines. The benefits of spatial management zones, such as marine protected areas (MPAs), for such animals are unclear. To help fill this knowledge gap, we examined core habitat use areas (CHUAs) for bull (*Carcharhinus leucas*), great hammerhead (*Sphyrna mokarran*) and tiger sharks (*Galeocerdo cuvier*) in relation to specific MPAs and exclusive economic zones (EEZs) in the western North Atlantic Ocean.

Methods Bull, great hammerhead and tiger sharks ($N = 86$ total) were satellite tagged and tracked in southern Florida and the northern Bahamas between 2010 and 2013. Filtered and regularized positions from Argos locations of tag transmissions were used to generate CHUAs for these sharks. Overlaps of CHUAs with regional protected areas and exclusive economic management zones were quantified to determine the proportion of each tracked shark's CHUA under spatial protection from exploitation.

Results A total of 0%, 17.9% and 34.7% of the regional CHUAs for tracked bull, great hammerhead and tiger sharks, respectively, were fully protected from exploitation in the study area.

Main conclusions Expansion of protected areas to include U.S. territorial waters would effectively protect 100% of the CHUAs for all tracked sharks in the study area. This finding is particularly significant for great hammerhead sharks, which are currently overfished, vulnerable to bycatch mortality and are the focus of strident regional conservation efforts. These findings also provide a means to inform decision makers and marine conservation planning efforts as to the types of management actions available and potential efficacy of spatial protections for these marine predators.

Keywords

biotelemetry, fishes, habitat use, place-based management, Conservation, sustainability.

*Correspondence: Neil Hammerschlag, Rosenstiel School of Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, FL, 33149, USA.
E-mail: nhammerschlag@rsmas.miami.edu

INTRODUCTION

Populations of many highly mobile marine species, such as large predatory sharks, have declined world-wide in recent decades (Pauly *et al.*, 1998; Myers & Worm, 2003; Dulvy *et al.*, 2008; Worm *et al.*, 2013). Shark population declines are primarily driven by intensive exploitation (e.g. for shark

fins and meat), and by incidental bycatch (Musick & Musick, 2011; Worm *et al.*, 2013; Oliver *et al.*, 2015). Life-history demographic characteristics of sharks such as slow growth, late age at maturity and low reproductive output make these populations particularly vulnerable to human threats (Musick & Musick, 2011; Gallagher *et al.*, 2012). Mitigating or reversing population declines is of great interest due to

the role apex predators may play in the regulation of marine ecosystem dynamics (Heithaus *et al.*, 2008; Estes *et al.*, 2011; Ruppert *et al.*, 2013).

Marine protected areas (MPAs) offer important management tools for reef fisheries (Meester *et al.*, 2004; Ault *et al.*, 2013); however, their role in sustaining highly mobile marine species is not well understood (Mora *et al.*, 2006; Chapman *et al.*, 2013). MPAs have been established to protect benthic habitats (e.g. coral reefs) and/or sedentary fishes, but as many sharks are wide ranging, it is unreasonable to assume that small MPAs would be effective. However, MPA benefits may arise for mobile species if core habitat use areas (CHUA), especially those that support key life stages or functions (e.g. breeding, feeding and gestation), are protected (Hooker *et al.*, 2011).

A number of studies have predicted that migratory fishes would not benefit from MPAs (e.g. DeMartini, 1993; Walters *et al.*, 1999; Mora *et al.*, 2006). However, a few recent studies have indicated that some MPA configurations may protect highly mobile species, including sharks (e.g. Southall *et al.*, 2006; Claudet *et al.*, 2010; Anadón *et al.*, 2011; Barnett *et al.*, 2011, 2012; Howey-Jordan *et al.*, 2013; Jewell *et al.*, 2014; Espinoza *et al.*, 2015). Benefits of MPAs will depend on the level of protection afforded and the time individuals spend within MPA boundaries, which can depend on the species, life stage, sex, size, physiological state and habitat type (Speed *et al.*, 2010; Escalle *et al.*, 2015).

Within the subtropical Atlantic Ocean, the waters surrounding Florida and the Bahamas are an ideal location to evaluate the potential protective benefits and efficacy of different spatial conservation management zones for migratory sharks. While Florida employs a suite of jurisdictions and various protection levels for sharks (See Methods section), the Bahamas banned shark harvests altogether since 2011. Thus, here we conducted a satellite tagging and tracking study of bull (*Carcharhinus leucas*), great hammerhead (*Sphyrna mokarran*) and tiger sharks (*Galeocerdo cuvier*) in southern Florida and the northern Bahamas to examine their CHUAs in relation to surrounding MPAs and exclusive economic zones (EEZs). As shark exploitation is the major regional threat to shark populations, we focused primarily on permanent zones that restrict and/or manage fishing. Our specific study objectives were to: (1) determine the spatial extent of CHUAs in Florida and the Bahamas for tagged bull, hammerhead and tiger sharks; (2) calculate the proportion of their CHUAs within the boundaries of regional spatial protection and economic management zones; and (3) determine the proportion of their CHUAs for which exploitation is prohibited.

METHODS

Study species

Bull sharks are generally coastal species mainly found on the continental shelf (Compagno *et al.*, 2005) and are listed as

Near Threatened by the International Union for the Conservation of Nature (IUCN) Red list (Simpfendorfer & Burgess, 2009). Tiger sharks are classified as coastal pelagic species (Compagno *et al.*, 2005) and listed as Threatened by the IUCN (Simpfendorfer, 2009). Great hammerhead sharks are classified as both coastal pelagic and semi-oceanic (Compagno *et al.*, 2005) and listed as Endangered by the IUCN (Denham *et al.*, 2007). Thus, these three species were selected for study as they are (1) highly mobile, (2) likely to have differential home range sizes; (3) probably occupy different, but overlapping, portions of the coastal ocean environment; (4) are facing different levels of population risk to exploitation; and (5) are captured in recreational (Shiffman & Hammerschlag, 2014; Gallagher *et al.*, 2015) and commercial fisheries (Gulak *et al.*, 2013; Gallagher *et al.*, 2014b) in the study area.

Study area and management zones

Pertinent zones within the study area included the U.S. and the Bahamas EEZ, Florida state waters, three National Parks within southern Florida, the Florida Keys National Marine Sanctuary (FKNMS) and no-take zones within the FKNMS (Fig. 1). We did not, however, evaluate shark use of time-area closures to specific fishing gear types in the study region (NOAA 2014). We defined protection as those zones in which exploitation for the study species was prohibited by law.

The U.S. EEZ, extends 270 km offshore and covers 11,351,000 km², where under current policy up to 36 large coastal sharks per boat, per trip, can be removed, with the exception of prohibited species. Additionally, pelagic and bottom longline restrictions are in place in the form of time-area closures throughout the region. Unlike the United States, the Bahamas EEZ fully protects sharks through a ban on commercial and recreational fishing for sharks and trade of all shark products, since 2011 (NOAA 2011, Regulation 36D). This effectively makes the 629,293 km² surrounding the islands of the Bahamas that compose its EEZ a shark sanctuary where it is prohibited to target and remove sharks, although other fishing is allowed.

Florida state waters extend 5.6 km offshore in the Atlantic Ocean and 16.7 km offshore in the Gulf of Mexico, encompassing 28,126 km². Effective 1 January 2012 the Florida Fish and Wildlife Conservation Commission enacted a regulation prohibiting the harvest, possession, sale and exchange of great hammerhead and tiger sharks within Florida state waters (FWC, 2011). This regulation also applies to scalloped (*Sphyrna lewini*) and smooth hammerhead sharks (*Sphyrna zygaena*). Additionally, the bag limit for both commercial harvest and recreational harvest of sharks within state waters is one per person per day, maximum two per vessel (FWC, 2011).

Biscayne National Park (BNP), Everglades National Park (ENP) and Dry Tortugas National Park (DTNP) collectively cover an area of 7067 km² in southern Florida. All




Figure 1 Map of study area showing: (a) Florida state waters boundary line, the Florida Keys National Marine Sanctuary (FKNMS) boundary, National Parks boundaries, and protected areas prohibiting recreational and commercial shark fishing; and (b) the exclusive economic zones (EEZs) of the United States (solid line) and Bahamas (dashed line).

three parks require a valid Florida state recreational saltwater fishing licence and enforce Florida fishing regulations. Additionally, ENP and DTNP both prohibit commercial fishing.

Finally, the FKNMS extends from the Upper Florida Keys down to the Dry Tortugas and encompasses 9850 km² of managed coastal waters stretching from the coastline to the deep barrier coral reefs. Within the area defined as the FKNMS, there are areas that prohibit both commercial fishing and recreational fishing (no take). These include the Research Natural Area (RNA) in Dry Tortugas National Park, Dry Tortugas North and South Ecological Reserves in FKNMS, the Sambos Ecological Reserve and 24 smaller Sanctuary Preservation Areas (SPAs), including the Special Use Areas (e.g. Smith *et al.*, 2011).

Tagging

Sharks were captured using baited circle-hook drumlines as described by Gallagher *et al.* (2014a). This technique was used because it allowed captured sharks to maintain ram ventilation while hooked, thus promoting shark vitality and survivorship. This was especially important in the case of great hammerheads, which exhibit a pronounced capture stress response (Gallagher *et al.*, 2014a,b). Drumlines were left to soak for relatively short time periods (*c.* 1.0 h) to reduce the probability of long hooking durations. When gear was retrieved, if captured sharks showed signs of significant fatigue or stress (i.e. discoloration, lack of tail beats or vigorous movement), they were released immediately without tagging to promote survivorship. Sharks selected for tagging

were secured on a platform or alongside the stern of the boat. A saltwater hose and pump was inserted in the shark's mouth to pump water over the gills while it was temporarily immobilized. Each shark was measured for total length (TL) and tagged with a Smart Position and Temperature Transmitting tag (SPOT5, Wildlife Computers; <http://www.wildlifecomputers.com>). Tags were coated with Prop-speed, a non-toxic, non-metallic antifouling agent that minimized biofouling. Tags were mounted to the shark's first dorsal fin using titanium bolts, steel washers and high carbon steel nuts (following Hammerschlag *et al.*, 2012a,b). This ensured that the steel nuts corroded, resulting in eventual tag detachment from the sharks. Given that great hammerhead sharks are sensitive to capture and handling, we transitioned to tagging this species with a towed SPOT tag that could be attached quicker via a tether and titanium dart anchored into the musculature at the base of the first dorsal fin.

Geographic locations of SPOT tagged sharks were determined by Doppler-shift calculations made by the Argos Data Collection and Location Service (www.argos-system.org). Locations were acquired when the tag broke the water surface and transmitted a signal to passing Argos satellite. Bull and great hammerhead sharks were tagged at various locations throughout the Florida Keys, while tiger sharks were tagged in the Florida Keys, Fort Myers, and the northern Bahamas (Fig. 2, Table 2).

Spatial data analyses

To improve location accuracy, we used Kalman filtering following the methods presented in Lopez *et al.* (2013). Argos provides the following radius of error for each location class (LC): LC 3 < 250 m, 250 m < LC 2 < 500 m, 500 m < LC 1

< 1500 m. The median error for LC 0, A and B ranges from 1 to 3 km. Class Z indicates that the location process failed and estimates of position are highly inaccurate. All transmitted locations were filtered to remove positions with LC Z, those on land, and those exceeding a speed of 2 m s⁻¹ (following Weng *et al.*, 2007, 2008).

Data from SPOT tags were regularized to constant 12-hour intervals using the piecewise cubic Hermite interpolating polynomial method required for kernel density estimates (KDE; Katajisto & Moilanen, 2006). Thus we chose this interpolation method based on previous research that evaluated several interpolation techniques for marine animals that found curvilinear interpolation techniques produced more accurate fluid movements (Tremblay *et al.*, 2006). This interpolation approach was not applied to data gaps > 3 days following the approach of Weng *et al.* (2008). All quantitative metrics of habitat use were generated based on the filtered, interpolated locations.

Kernel density estimates (KDEs) were used to delineate high-use habitat areas from the tag data (Heupel *et al.*, 2004; Hart *et al.*, 2010; Hammerschlag *et al.*, 2012b; Farmer & Ault, 2014). KDEs provided probabilistic information concerning habitat utilization within the distributional range of the animal movements (Worton, 1989). Gaussian KDEs were calculated in Geospatial Modelling Environment (Beyer, 2012) using the kernel density tool and bandwidth estimation algorithms in the 'ks: kernel smoothing' package in The R Project for Statistical Computing (<https://www.r-project.org/>). Kernel estimators are sensitive to the bandwidth algorithms used to generate them. The most appropriate algorithm depends on the pattern of animal space use and the sample size. After testing the performance of all bandwidths in the 'ks' package with regard to properly capturing the utilization distribution of each study species (following




Figure 2 Map of study area with latitude and longitude as well as major cities for spatial reference. The map displays representative tracks from a tagged hammerhead shark (*Sphyrna mokarran*, grey squares), tiger shark (*Galeocerdo cuvier*, grey circles) and bull shark (*Carcharhinus leucas*, grey triangles, inset panel). The data points are raw transmitted Argos positions with lines connecting positions (tracks are not interpolated). Tagging locations for all study individuals are indicated with small black circles. Large black circles are the tagging locations for the representative shark tracks.

Schofield *et al.*, 2013), the PLUGIN bandwidth was selected for all spatial analyses. Other bandwidth estimators produced over or under smoothed kernels, either capturing areas that were not used or missing key areas that were used by the sharks. KDEs are cumulated from area of highest to lowest density to create percentage volume contours (PVCs). Thus, 50% contours represent areas of top 50% highest densities or CHUAs following Ardon (2008). The 95% PVCs represent the top 95% highest densities. Finally, we calculated the percentage of each species' CHUA within the management zones of interest using ARCGIS 10 (ESRI, Redlands, California USA).

RESULTS

Eighty-six individual sharks were satellite tagged between 2010 and 2013, resulting in a total of 9417 tracking days and 5481 interpolated positions that were used for the analyses. Over the course of the study, we tracked 24 bull (155–269 TL; 62.5% female), 18 great hammerhead (235–345 cm TL; 38.9% female) and 44 tiger sharks (157–403 cm TL; 88.6% female). All sharks were either subadult or adults. Mean (\pm SD) number of days tracked per species was 110.5 ± 125.5 for bull (range: 6–586 days), 46.3 ± 46.3 (range: 2–154 days) for great hammerhead and 134.8 ± 160.8 (range: 6–828 days) for tiger sharks. Further details on sample size, range and average TL, percentage male vs. female, and number of tracking days for each species are found in Table 1. TL, sex and tagging information for each individual are found in Table 2.

Kernel density estimates

The CHUA coverage was 1080 km² for bull, 85,061 km² for great hammerhead and 131,229 km² for tiger sharks. Table 1 summarizes the percentage overlap of each species' CHUA within each management zone. The entire bull shark CHUA fell within the U.S. EEZ, of which 0% occurred within the no-take areas, 10.42% within the FKNMS, 58.65% within national park boundaries and 95.93% within Florida state waters (Fig. 3a). None of the zones that overlap with bull shark CHUA provided them complete protection from fishing.

For tagged great hammerhead sharks, 91.57% of their CHUA occurred within the U.S. EEZ (Fig. 3b). Of this, 0.72% was within the no-take areas, 3.35% within National Parks and 11.61% within the FKNMS boundary. Additionally, 17.87% of their CHUA was within Florida state waters, which has prohibited the harvest of great hammerheads since January 2012. Finally, 8.43% fell within the Bahamas EEZ, which has protected all sharks since 2011. Due to the no-take areas, Florida state waters and the Bahamas EEZ, we found 27.02% of the tagged great hammerhead CHUA was protected.

Tiger sharks spent very little, or no time within the no-take areas, national parks, the FKNMS or Florida state waters (Fig. 3c). However, 67.60% and 32.40% of their

Table 1 Species tagging information and percentage overlap of CHUA with spatial management zones. CHUA overlap cell values were calculated as the area overlap of the 50% KDE and each zone type, divided by the total area of the 50% KDE. No-take MPAs are the only management zone where no fishing or marine resource exploitation is permitted. The other zones differ in degree of shark fishing restrictions, but all allow for resource exploitation of some kind (see Methods for definitions and shark protective status for each management zone).

	<i>Carcharhinus leucas</i>	<i>Sphyrna mokarran</i>	<i>Galeocerdo cuvier</i>
Number of tagged sharks	24	18	44
Size range (cm TL)	155–269	235–345	157–403
Mean TL (cm)	210	273	275
Percentage male (%)	37.5	61.1	11.4
Percentage female (%)	62.5	38.9	88.6
Tracking days	2654	833	5930
Interpolated positions	975	549	3957
Tagging location	Florida	Florida	FL (17), BA (27)
Percentage overlap of CHUA			
No-take MPAs	0.00	0.72	0.76
FKNMS	10.42	11.61	0.00
NPs	58.65	3.35	0.00
FL state waters	95.93	17.87	2.34
U.S. EEZ	100.00	91.57	67.60
Bahamas EEZ	0.00	8.43	32.40
Total protected	0.00	27.02	34.74

CHUA, core habitat use areas; KDE, Kernel density estimates; MPA, marine protected areas; FKNMS, Florida Keys National Marine Sanctuary; NPs, National Parks – Everglades, Biscayne and Dry Tortugas; EEZ, exclusive economic zone; TL, total length.

CHUA occurred within the U.S. and Bahamas EEZ, respectively. When considering only those areas that provide complete protection to the tiger shark (no-take areas, FL state waters and the Bahamas EEZ), we found that 34.74% of the CHUA was protected, primarily due to the Bahamas Shark Sanctuary.

DISCUSSION

This study examined the distribution and core habitat use of three large sharks in relation to several protected areas and political-economic zones within Florida and the Bahamas. These results have implications for the evaluation of current marine spatial management zones and strategic development of new MPAs for populations of these species within the study area.

For bull sharks tagged in the Florida Keys, individuals remained within coastal inshore areas of the continental shelf. This is consistent with previous tagging work in the region as well as in other areas (Brunnschweiler *et al.*, 2010; Carlson *et al.*, 2010; Hammerschlag *et al.*, 2012b; Heupel *et al.*, 2015). Tagged bull sharks remained almost exclusively

Table 2 Satellite tagging information for each individual shark included in this study.

Species	ID #	Total length (cm)	Sex	Tagging date	# Days tracked	Tagging lat.	Tagging long.	# Interp. positions
<i>Carcharhinus leucas</i>	33919	176	F	11/7/2010	89	25.01	-81.00	66
<i>C. leucas</i>	33937	221	F	6/5/2010	118	24.70	-80.85	4
<i>C. leucas</i>	33991	210	F	3/26/2010	37	25.01	-81.00	34
<i>C. leucas</i>	34208	173	F	11/7/2010	67	25.01	-81.00	13
<i>C. leucas</i>	55496	200	M	11/6/2010	133	25.01	-81.00	36
<i>C. leucas</i>	60696	192	M	10/29/2010	81	26.37	-81.98	12
<i>C. leucas</i>	60697	155	F	8/10/2010	11	26.37	-81.98	1
<i>C. leucas</i>	60698	176	M	11/6/2010	79	25.01	-81.00	55
<i>C. leucas</i>	60699	195	M	10/7/2010	314	25.01	-81.00	112
<i>C. leucas</i>	68467	245	M	2/19/2011	163	26.10	-79.10	7
<i>C. leucas</i>	68483	194	F	12/4/2010	104	25.01	-81.00	14
<i>C. leucas</i>	105596	245	F	2/17/2011	28	24.81	-80.91	97
<i>C. leucas</i>	108063	176	F	12/9/2011	40	25.10	-81.02	41
<i>C. leucas</i>	108065	205	M	11/23/2011	586	25.07	-80.31	108
<i>C. leucas</i>	108067	192	M	3/21/2012	6	25.07	-80.31	5
<i>C. leucas</i>	108068	194	M	10/27/2011	244	25.35	-80.26	75
<i>C. leucas</i>	113535	269	F	1/10/2012	153	25.65	-80.17	38
<i>C. leucas</i>	115916	262	F	12/2/2012	83	25.68	-80.17	13
<i>C. leucas</i>	126370	189	F	2/1/2013	126	24.70	-80.85	32
<i>C. leucas</i>	128016	200	M	4/8/2013	82	25.35	-80.26	43
<i>C. leucas</i>	128508	230	F	3/16/2013	33	25.60	-80.16	61
<i>C. leucas</i>	128511	257	F	3/16/2013	12	25.60	-80.16	4
<i>C. leucas</i>	129953	262	F	5/1/2013	38	25.61	-80.17	63
<i>C. leucas</i>	129954	227	F	4/26/2013	27	25.61	-80.17	41
<i>Galeocerdo cuvier</i>	33992	201	F	5/26/2010	32	26.37	-81.98	13
<i>G. cuvier</i>	34020	263	M	5/26/2010	40	26.37	-81.98	27
<i>G. cuvier</i>	34021	249	F	5/26/2010	24	26.37	-81.98	24
<i>G. cuvier</i>	34029	255	F	5/26/2010	189	26.37	-81.98	74
<i>G. cuvier</i>	34107	225	F	5/25/2010	202	26.37	-81.98	5
<i>G. cuvier</i>	34203	255	F	11/13/2010	46	24.70	-80.85	66
<i>G. cuvier</i>	55494	250	F	6/10/2010	94	26.37	-81.98	64
<i>G. cuvier</i>	55495	295	F	6/9/2010	127	26.37	-81.98	230
<i>G. cuvier</i>	68471	245	F	1/29/2011	26	24.70	-80.85	5
<i>G. cuvier</i>	68477	200	M	10/29/2010	126	26.37	-81.98	86
<i>G. cuvier</i>	68485	335	F	2/19/2011	95	26.86	-79.04	148
<i>G. cuvier</i>	68486	320	F	2/20/2011	99	26.86	-79.04	119
<i>G. cuvier</i>	68488	295	F	2/20/2011	828	26.86	-79.04	338
<i>G. cuvier</i>	68494	365	F	2/19/2011	191	26.86	-79.04	243
<i>G. cuvier</i>	68495	325	F	2/20/2011	239	26.86	-79.04	179
<i>G. cuvier</i>	68496	280	F	2/20/2011	216	26.86	-79.04	337
<i>G. cuvier</i>	68529	325	F	2/19/2011	557	26.86	-79.04	386
<i>G. cuvier</i>	68554	403	F	2/19/2011	185	26.86	-79.04	327
<i>G. cuvier</i>	68555	286	F	2/20/2011	250	26.86	-79.04	241
<i>G. cuvier</i>	68556	322	F	2/20/2011	418	26.86	-79.04	346
<i>G. cuvier</i>	98332	184	F	11/12/2010	82	24.70	-80.85	82
<i>G. cuvier</i>	105594	375	F	2/19/2011	22	26.10	-79.10	9
<i>G. cuvier</i>	105595	325	F	2/22/2011	40	26.10	-79.10	26
<i>G. cuvier</i>	105599	340	F	2/20/2011	11	26.10	-79.10	8
<i>G. cuvier</i>	105600	310	F	2/20/2011	48	26.10	-79.10	95
<i>G. cuvier</i>	106660	206	M	4/10/2011	51	25.82	-77.93	49
<i>G. cuvier</i>	106661	320	F	4/10/2011	82	25.82	-77.93	50
<i>G. cuvier</i>	108064	175	F	12/10/2011	372	25.32	-80.23	54
<i>G. cuvier</i>	112981	332	F	12/19/2011	6	26.86	-79.04	11
<i>G. cuvier</i>	112982	196	F	9/12/2012	43	25.04	-77.24	3
<i>G. cuvier</i>	112986	248	F	7/23/2012	322	26.86	-79.04	5

Table 2 Continued.

Species	ID #	Total length (cm)	Sex	Tagging date	# Days tracked	Tagging lat.	Tagging long.	# Interp. positions
<i>G. cuvier</i>	112987	296	F	7/22/2012	182	26.86	-79.04	4
<i>G. cuvier</i>	112991	260	F	9/6/2012	9	25.04	-77.24	7
<i>G. cuvier</i>	113534	321	F	12/15/2011	31	26.86	-79.04	14
<i>G. cuvier</i>	113536	305	F	12/15/2011	143	26.86	-79.04	55
<i>G. cuvier</i>	113537	346	F	2/5/2012	94	26.86	-79.04	43
<i>G. cuvier</i>	115906	200	M	5/27/2012	39	24.70	-80.85	49
<i>G. cuvier</i>	115907	200	M	2/6/2012	25	25.25	-80.26	38
<i>G. cuvier</i>	115912	297	F	9/6/2012	102	25.04	-77.24	24
<i>G. cuvier</i>	115913	157	F	9/10/2012	130	25.04	-77.24	21
<i>G. cuvier</i>	115915	215	F	9/4/2012	31	25.04	-77.24	9
<i>G. cuvier</i>	119440	206	F	3/22/2013	37	25.60	-80.16	11
<i>G. cuvier</i>	130582	220	F	5/25/2013	28	24.70	-80.85	25
<i>G. cuvier</i>	130985	289	F	6/7/2013	16	24.70	-80.85	7
<i>Sphyrna mokarran</i>	33933	277	M	6/4/2010	20	24.70	-80.85	21
<i>S. mokarran</i>	33938	262	M	3/12/2010	40	24.70	-80.85	61
<i>S. mokarran</i>	68472	287	F	1/29/2011	86	24.70	-80.85	3
<i>S. mokarran</i>	68480	265	F	1/29/2011	116	24.70	-80.85	28
<i>S. mokarran</i>	68481	295	F	1/29/2011	100	24.70	-80.85	128
<i>S. mokarran</i>	98328	235	M	2/20/2010	12	24.70	-80.85	5
<i>S. mokarran</i>	98329	249	M	2/20/2010	62	24.70	-80.85	4
<i>S. mokarran</i>	98331	251	F	2/7/2010	75	24.73	-80.85	5
<i>S. mokarran</i>	105597	345	M	2/19/2011	5	26.10	-79.10	88
<i>S. mokarran</i>	105598	235	M	2/26/2011	2	24.73	-80.85	4
<i>S. mokarran</i>	106663	270	F	7/13/2011	154	24.67	-80.86	68
<i>S. mokarran</i>	106895	277	F	8/5/2011	85	25.26	-80.23	39
<i>S. mokarran</i>	106896	301	M	7/13/2011	5	24.73	-80.85	8
<i>S. mokarran</i>	111546	304	M	4/23/2013	14	24.70	-80.85	12
<i>S. mokarran</i>	111550	264	M	5/1/2012	12	25.22	-80.33	42
<i>S. mokarran</i>	111551	335	M	4/20/2013	34	24.70	-80.85	22
<i>S. mokarran</i>	128510	240	M	6/20/2013	8	25.65	-80.17	8
<i>S. mokarran</i>	129955	250	F	6/21/2013	3	25.43	-80.21	3

within Florida state waters, overlapping largely with the National Parks (BNP and ENP). Currently, bull sharks can be harvested by both commercial and recreational fisheries within Florida waters and BNP, and recreational fisheries in ENP. As a result of much of their CHUA falling within these zones, the implementation of regulations preventing the harvest of bull sharks within these areas would provide adequate spatial protection.

With decreasing trends in great hammerhead populations globally, including the study region (Baum *et al.*, 2003; Myers *et al.*, 2007; Miller *et al.*, 2014), the International Commission for the Conservation of Atlantic Tunas (ICCAT) established agreements in November 2010 to protect this species in Atlantic waters. ICCAT suggested member parties consider implementing time–area closures and other spatial planning to protect these species (ICCAT, 2010). However, prior to this work, there had only been one published biotelemetry study of great hammerheads in the Atlantic Ocean, which was based on the tracking of only a single individual (Hammer-schlag *et al.*, 2011a). This study thus provides information that can be used for implementing spatial management for great hammerheads in the study area.

In January 2012, great hammerhead sharks were granted full protection in Florida state waters. However, our analyses show that only 17.9% of tagged great hammerhead CHUA occurred in Florida waters. This suggests that current protection may be inadequate to ensure sustainability of the species. In contrast, 91.57% of tagged great hammerhead CHUA was within the U.S. EEZ, and 8.43% within the Bahamas EEZ. Thus, extending full protection to great hammerheads within U.S. federal waters would likely have significant conservation benefits for this species. However, great hammerheads have among the highest vulnerability to at-vessel and post-release mortality due to a pronounced stress response from fighting on a fishing line, even over relatively short time periods (Morgan & Burgess, 2007; Gallagher *et al.*, 2014a,b; Gulak *et al.*, 2015). As a result, mortality may still be high even in areas that prohibit the harvest of this species if they are still captured as bycatch. Thus, protection may need to be extended over a larger area than that of their CHUA to account for potential losses due to bycatch. Additionally, fishing practices would likely need to be altered in some key areas to prevent great hammerheads from interacting with fishing gears in the first place (Gallagher *et al.*, 2014c). This could include time–area fishing closures or




Figure 3 Kernel density estimates (KDE) from 2010 to 2013 for: (a) bull sharks ($n = 24$); (b) great hammerhead sharks ($n = 18$) and (c) tiger sharks ($n = 44$). Percentage volume KDE contours show the proportion of all tag positions across individuals and years. A KDE of 0.5 was considered core habitat use area (CHUA).

adopting gear modifications that prevent hammerheads from biting baited hooks.

For tagged tiger sharks, a relatively large portion of their CHUA (32.4%) overlapped with the Bahamas EEZ, where commercial longlining has been banned for decades and sharks are also protected due to the Shark Sanctuary. This is

of particular conservation significance as little Bahamas Bank, in the Bahamas, appears to function as a mixed aggregation site for female tiger sharks, serving as a refuge for females to reach maturity as well as a gestation ground for recently mated and gravid individuals (Sulikowski *et al.*, 2016). The relatively high abundance of tiger sharks in the Bahamas, as compared to the rest of the Caribbean (Ward-Paige *et al.*, 2010), could in part be due to the protection of mature and/or gravid females in the Bahamas.

In Florida waters, tiger sharks were granted protection in January 2012; however, this area only covers 2.34% of the CHUA of tagged tiger sharks. If protection for this species were extended to include the U.S. territorial waters, CHUAs of tagged animals would increase to 100%. Unlike great hammerheads, tiger sharks are relatively robust to fishing stress and exhibit low levels of at-vessel and post-release fishing mortality (Gallagher *et al.*, 2014a,b). Therefore, if warranted, a policy requiring any tiger sharks incidentally captured to be released in U.S. waters would likely have a positive impact on population size.

While a relatively small amount of time was spent by tiger and great hammerhead sharks within Florida waters where their harvest is prohibited, such protection is still important for these relatively long-lived, low-fecund species. Moderate exploitation may result in local population declines especially if reproductive or gravid females are captured when coming to inshore state waters to breed or birth (Hixon *et al.*, 2014; Shiffman *et al.*, 2014). Indeed, a recent study found that charter boat recreational shark fishers in Florida are most excited about catching hammerhead and bull sharks and thus often target large individuals of these species (Shiffman & Hammerschlag, 2014). It is important to note that of the three species studied, great hammerheads are experiencing the highest level of reported population declines in the region and are also the most vulnerable to at-vessel and post-release mortality from fishing (Baum *et al.*, 2003; Myers *et al.*, 2007; Gallagher *et al.*, 2014a,b,c; Miller *et al.*, 2014). Furthermore, analysis from commercial shark longline catch and effort data in parts of the study area suggest that relative abundance of both bull sharks and tiger sharks may have stabilized, with populations now increasing (Carlson *et al.*, 2012).

Similar to the approach used here, other studies have also evaluated space use of large, migratory marine fauna in relation to political and protection boundaries using biotelemetry. For example, a recent study found oceanic whitetip (*Carcharhinus longimanus*) sharks exhibit high residency (68.2%) within the Bahamas EEZ (Howey-Jordan *et al.*, 2013), whereas, another study found that basking sharks (*Cetorhinus maximus*) tagged in UK waters spent the majority of their time (*c.* 78%) outside of protected waters (Southall *et al.*, 2006). Off the Yucatan Peninsula in Mexico, Graham *et al.* (2012) found that only 12% of manta rays (*Manta birostris*) tracked locations fell within MPAs. A comparable study off the east coast of Australia that examined the movements of bull sharks found that while MPAs offered

some protection for tagged sharks, the scale of the bull shark movements relative to the size of no fishing areas suggested limited conservation benefits (Heupel *et al.*, 2015).

Recently, Escalle *et al.* (2015) summarized previous telemetry studies that evaluated reef sharks residency and home range in relation to no-take protected areas. They found that the value of protected areas for these species varied both among locations and between studies at the same location. Of the studies examining the suitability of no-take zones for reef sharks, Escalle *et al.* (2015) found that 65% of these could benefit sharks, while 35% suggested no-take zones were too small to offer protection given the scale of shark movement (e.g. Chapman *et al.* 2005; Heupel *et al.* 2010; Barnett *et al.*, 2011; Bond *et al.*, 2012; Knip *et al.*, 2012). Accordingly, they concluded that marine reserve benefits for sharks will depend on the amount of the time individuals spend within reserve boundaries and the life-history stages that are afforded protection.

The present work reinforces the concept that analysis of movements and habitat use of migratory marine fauna can inform spatial planning and provide unique insights into the effectiveness of MPAs. However, most MPAs are formed without prior knowledge of how a given species use them. Ideally, habitat use patterns of key species of conservation concern should be determined before MPAs are designed. However, when this is not possible (as in most situations), managers should consider the possibility of changing design configurations to protect key species or life stages as such information becomes available (discussed by Ashe *et al.*, 2010; Pichegru *et al.*, 2010; Barnett *et al.*, 2012).

Spatial management for wide-ranging species relies on the assumption that, if species are protected within their key areas (i.e. where critical life-history stages occur and/or where they are most abundant and most vulnerable to exploitation), populations may be sustained (Speed *et al.*, 2010). Moreover, population growth within protected areas must exceed losses generated outside of closed areas. Studies of migratory birds have found that protecting key areas, such as feeding or reproductive grounds, can sufficiently conserve highly mobile species (Newton, 2008). Several studies have found that for sharks, MPAs can protect key foraging areas (e.g. Barnett *et al.*, 2011), nursery grounds (Barnett & Semmens, 2012) and juvenile habitat (e.g. Garla *et al.*, 2006). While in this study we identified areas of core habitat use, we could not determine whether these were key areas, with the exception of an apparent tiger shark gestation ground in the Bahamas as discussed above (Sulikowski *et al.*, in press). Although it is probable that critical life-history areas overlap with core habitat use areas, we cannot rule out a mismatch. Therefore, it is important that further research be conducted to identify such key habitats for populations of great hammerhead, tiger and bull sharks in the study area.

A limitation of this study is that some sharks surface more than others, thus transmitting more data and potentially biasing results (Hammerschlag *et al.*, 2011b). Moreover, it is possible that a shark surfacing, and therefore the probability

of getting a satellite-derived position, may be dependent on habitat type. For example, when in shallow water, sharks may be more likely to surface than when occupying deep water. However, use of large, multiyear data sets where all data are interpolated and regularized, as done here, help reduce such biases. As mentioned prior, KDEs are sensitive to bandwidth estimators, and as such, results vary depending on which algorithm is used. Geospatial Modelling Environment allowed for the choice between five different bandwidth estimation algorithms. After testing each estimator using our data set, we determined that the plug-in estimator qualitatively produced the best results, neither under nor over smoothing the data. We recognize that our CHUA overlap values are sensitive to this process, in addition to how we defined the CHUAs. We selected the 50% KDE to represent CHUAs as this value is widely used in the literature (Heupel *et al.*, 2004; Weng *et al.*, 2008; Hart *et al.*, 2010), but note the paucity of quantitative analysis that justifies this value.

It is also worth considering that great hammerhead, bull and tiger sharks are found throughout most coastal waters of the tropical and temperate Atlantic, with population and sub-population ranges not fully known. Many shark populations have home ranges that are smaller than the species' overall distribution. Moreover, there can often be large within population differences in movement patterns (Chapman *et al.*, 2015). Thus, it is possible that our findings may only be representative of a part of the population and results may vary considerably for individuals tagged in different areas throughout their range. As such, we cannot conclude that our data is representative of the entire population within the region. For tiger sharks, individuals tagged in the Florida and Bahamas study sites generally showed similar residency and home range patterns (Hammerschlag *et al.*, 2012a, 2015). However, it would be valuable to include data from tagging efforts across various sites of the tropical and subtropical Atlantic. We also tracked subadult and adult individuals of both sexes, however, did not have sufficient sample sizes to partition data to make sufficient comparisons by life stage and sex.

While we evaluated movement patterns of tagged bull, great hammerhead and tiger sharks in response to permanent management zones within Florida waters, the Bahamas and the U.S. EEZ, we did not consider specific zones that restrict certain types of fishing gears in which sharks are captured (e.g. time–area closures for pelagic longline gear in U.S. waters; NOAA 2014). Therefore, shark vulnerability to fisheries capture is not uniform within the management zones evaluated here.

In summary, this study has implications for the development of marine conservation plans, enabling policy makers to better understand the efficacy of current and proposed levels of spatial protection for populations of bull, great hammerhead and tiger sharks in the study area. However, if sharks are protected but fishing is still permitted in the MPA, they may still be vulnerable to bycatch. We also note that effective spatial management strategies must consider several competing alternatives; for example, the potential

attendant effects of protection on other species and the feasibility of the proposition when factoring stakeholders and economic impacts. Moreover, to be effective, strategic conservation management for focal species should consider protection for both their prey and the habitats they rely on. These considerations must also address monitoring and enforcement as equally critical to the success of any spatial management strategy.

ACKNOWLEDGEMENTS

This research benefitted greatly from the dedicated contributions of all the University of Miami's Shark Research and Conservation Programme team leaders and interns who assisted in the capture and tagging expeditions. We especially thank A. Gallagher for field support, K. Hartog and E. Nelson for compiling the tracking locations, and D. Lirman for advice and comments. This work was supported by The Batchelor Foundation, Disney Conservation Fund, Wells Fargo, Guy Harvey Ocean Foundation, and the West Coast Inland Navigation District. This work was conducted under permits from the National Marine Fisheries Service Highly Migratory Species Division (SHK-EFP-10-01), Florida Keys National Marine Sanctuary (FKNMS-2010-006), Florida Fish and Wildlife (SAL-957), Bahamas Department of Marine Resources (MAF/LIA/22), Biscayne and Everglades National Parks, and the University of Miami Institutional Animal Care and Use Committee (Protocol #09-187). Data were organized through SeaTurtle.org. Thanks to H. Calich for help with preparation of Fig. 2. For help with copy editing, thanks to J. Whidden, N. Wosnick, and E. Nelson. We especially thank the anonymous referees whose comments helped significantly strengthen this paper.

REFERENCES

Anadón, J.D., D'Agrosa, C., Gondor, A. & Gerber, L. (2011) Quantifying the spatial ecology of wide-ranging marine species in the Gulf of California: implications for marine conservation planning. *PLoS ONE*, **6**, e28400. doi:10.1371/journal.pone.0028400

Ardron, J.A. (2008) Three initial OSPAR tests of ecological coherence: heuristics in a data-limited situation. *Ices Journal of Marine Science*, **65**, 1527–1533.

Ashe, E., Noren, D.P. & Williams, R. (2010) Animal behaviour and marine protected areas: incorporating behavioural data into the selection of marine protected areas for an endangered killer whale population. *Animal Conservation*, **13**, 196e203.

Ault, J.S., Smith, S.G., Bohnsack, J.A., Luo, J., Zurcher, N., McClellan, D.B., Ziegler, T.A., Hallac, D.E., Patterson, M., Feeley, M.W., Ruttenberg, B.I., Hunt, J., Kimball, D. & Causey, B. (2013) Assessing coral reef fish population and community changes in response to marine reserves in the Dry Tortugas, Florida, USA. *Fisheries Research*, **144**, 28–37.

Barnett, A. & Semmens, J.M. (2012) Sequential movement into coastal habitats and high spatial overlap of predator and prey suggest high predation pressure in protected areas. *Oikos*, **121**, 882–890.

Barnett, A., Abrantes, K., Stevens, J.D. & Semmens, J.M. (2011) Site fidelity, habitat partitioning and sex-specific migrations in a large mobile apex predator: implications for protected area management and ecosystem dynamics. *Animal Behaviour*, **81**, 1039–1048.

Barnett, A., Abrantes, K.G., Seymour, J. & Fitzpatrick, R. (2012) Residency and spatial use by reef sharks of an isolated seamount and its implications for conservation. *PLoS ONE*, **7**, e36574.

Baum, J.K., Myers, R., Kehler, D., Worm, B., Harley, S. & Doherty, P. (2003) Collapse and conservation of shark populations in the northwest Atlantic. *Science*, **299**, 389–392.

Beyer, H.L. (2012) *Geospatial modelling environment (version 0.6.0.0) (software)*. Available at: <http://www.spatial ecology.com/gme> (accessed 2 November 2014).

Bond, M.E., Babcock, E., Pikitch, E.K., Abercrombie, D.L., Lamb, N.F. & Chapman, D.D. (2012) Reef sharks exhibit site-fidelity and higher relative abundance in marine reserves on the Mesoamerican Barrier Reef. *PLoS ONE*, **7**, e32983.

Brunnschweiler, J.M., Queiroz, N. & Sims, D.W. (2010) Oceans apart? Short-term movements and behaviour of adult bull sharks *Carcharhinus leucas* in Atlantic and Pacific Oceans determined from pop-off satellite archival tagging. *Journal of Fish Biology*, **77**, 1343–1358.

Carlson, J.K., Ribera, M., Conrath, C., Heupel, M.R. & Burgess, G.H. (2010) Habitat use and movement patterns of bull sharks *Carcharhinus leucas* determined using pop-up satellite archival tags. *Journal of Fish Biology*, **77**, 661–675.

Carlson, J.K., Hale, L.F., Morgan, A. & Burgess, G. (2012) Relative abundance and size of coastal sharks derived from commercial shark longline catch and effort data. *Journal of Fish Biology*, **80**, 1749–1764.

Chapman, D.D., Pikitch, E.K., Babcock, E.A. & Shivji, M.S. (2005) Marine reserve design and evaluation using automated acoustic telemetry: a case-study involving coral reef-associated sharks in the Mesoamerican Caribbean. *Marine Technology Society Journal*, **39**, 42–53.

Chapman, D.D., Frisk, M.G., Abercrombie, D.L., Safina, C., Gruber, S.H., Babcock, E.A., Feldheim, K.A., Pikitch, E.K., Ward-Paige, C., Davis, B., Kessel, S., Heithaus, M.R. & Worm, B. (2013) Give shark sanctuaries a chance. *Science*, **339**, 757.

Chapman, D.D., Feldheim, K.A., Papastamatiou, Y.P. & Hueter, R.E. (2015) There and back again: a review of residency and return migrations in sharks, with implications for population structure and management. *Annual Review of Marine Science*, **7**, 547–570.

Claudet, J., Osenberg, C.W., Domenici, P. *et al.* (2010) Marine reserves: fish life history and ecological traits matter.

- Ecological Applications: A Publication of the Ecological Society of America*, **20**, 830–839.
- Compagno, L.J.V., Dando, M. & Fowler, S. (2005) *Sharks of the world*. Princeton University Press, Princeton, NJ.
- DeMartini, E.E. (1993) Modeling the potential of fishery reserves for managing Pacific coral reef fishes. *Fisheries Bulletin*, **91**, 414–427.
- Denham, J., Stevens, J., Simpfendorfer, C.A., Heupel, M.R., Cliff, G., Morgan, A., Graham, R., Ducrocq, M., Dulvy, N.D., Seisay, M., Asber, M., Valenti, S.V., Litvinov, F., Martins, P., Lemine Ould Sidi, M., Tous, P. & Bucal, D. (2007) *Sphyrna mokarran*. *The IUCN Red list of threatened species*. Version 2015.2. Available at: www.iucnredlist.org. (Downloaded on 24 August 2015).
- Dulvy, N.K., Baum, J. & Clark, S. (2008) You can swim but you can't hide: the global status and conservation of oceanic pelagic sharks. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **18**, 459–482.
- Escale, L., Speed, C.W., Meekan, M.G., White, W.T., Babcock, R.C., Pillans, R.D. & Huveneers, C. (2015) Restricted movements and mangrove dependency of the nervous shark *Carcharhinus caudus* in nearshore coastal waters. *Journal of Fish Biology*, **87**, 323–341.
- Espinoza, M., Lédée, E.J., Simpfendorfer, C.A., Tobin, A.J. & Heupel, M.R. (2015) Contrasting movements and connectivity of reef-associated sharks using acoustic telemetry: implications for management. *Ecological Applications*, **25**, 2101–2118.
- Estes, J.A., Terborgh, J., Brashares, J.S. *et al.* (2011) Trophic downgrading of planet Earth. *Science*, **333**, 301–306.
- Farmer, N.A. & Ault, J.S. (2014) Modeling coral reef fish home range movements in Dry Tortugas, Florida. *The Scientific World Journal*, Volume 2014, Article ID 629791, <http://dx.doi.org/10.1155/2014/629791>.
- Florida Fish and Wildlife Conservation Commission (2011) *Commercial regulations for shark*. Florida Fish and Wildlife Conservation Commission. Available at: <http://myfwc.com/fishing/saltwater/commercial/shark/> (accessed xxx October 2012).
- Gallagher, A.J., Kyne, P.M. & Hammerschlag, N. (2012) Ecological risk assessment and its application to elasmobranch conservation and management. *Journal of Fish Biology*, **85**, 1727–1748.
- Gallagher, A.J., Orbesen, E., Hammerschlag, N. & Serafy, J. (2014a) Vulnerability of oceanic sharks as pelagic longline bycatch. *Global Ecology and Conservation*, **1**, 50–59.
- Gallagher, A.J., Serafy, J., Cooke, S. & Hammerschlag, N. (2014b) Physiological stress response, reflex impairment, and survival of five sympatric shark species following experimental capture and release. *Marine Ecology Progress Series*, **496**, 207–218.
- Gallagher, A.J., Hammerschlag, N., Shiffman, D. & Giery, S. (2014c) Evolved for extinction? The price of specialization in apex marine predators. *BioScience*, **64**, 619–624.
- Gallagher, A.J., Cooke, S.J. & Hammerschlag, N. (2015) Risk perceptions and conservation ethic among recreational anglers targeting threatened sharks in the subtropical Atlantic. *Endangered Species Research*, **29**, 81–93.
- Garla, R.C., Chapman, D., Wetherbee, B. & Shivji, M. (2006) Movement patterns of young Caribbean reef sharks, *Carcharhinus perezi*, at Fernando de Noronha Archipelago, Brazil: the potential of marine protected areas for conservation of a nursery ground. *Marine Biology*, **149**, 189–199.
- Graham, R.T., Witt Castellanos, M.D., Remolina, F., Maxwell, S., Godley, B.J. & Hawkes, L.A. (2012) Satellite tracking of manta rays highlights challenges to their conservation. *PLoS ONE*, **7**, e36834, doi:10.1371/journal.pone.0036834.
- Gulak, S.J.B., Enzenauer, M.P. & Carlson, J.K. (2013) Characterization of the shark and reef fish bottom longline fisheries: 2012. *NOAA Technical Memorandum NMFS-SEFSC*, **652**, 42.
- Gulak, S.J.B., Santiago, A.J.D. & Carlson, J.K. (2015) Hooking mortality of scalloped hammerhead *Sphyrna lewini* and great hammerhead *Sphyrna mokarran* sharks caught on bottom longlines. *African Journal of Marine Science*, **37**, 267–273.
- Hammerschlag, N., Gallagher, A.J., Lazzarre, D.M. & Slonim, C. (2011a) Range extension of the Endangered great hammerhead shark *Sphyrna mokarran* in the Northwest Atlantic: preliminary data and significance for conservation. *Endangered Species Research*, **13**, 111–116.
- Hammerschlag, N., Gallagher, A. & Lazzarre, D. (2011b) A review of shark satellite tagging studies. *Journal of Experimental Marine Biology and Ecology*, **398**, 1–8.
- Hammerschlag, N., Gallagher, A., Wester, J., Luo, J. & Ault, A. (2012a) Don't bite the hand that feeds: assessing ecological impacts of provisioning ecotourism on an apex marine predator. *Functional Ecology*, **26**, 567–576.
- Hammerschlag, N., Luo, J., Irschick, D.J. & Ault, J.S. (2012b) A comparison of spatial and movement patterns between sympatric predators: bull sharks (*Carcharhinus leucas*) and Atlantic tarpon (*Megalops atlanticus*). *PLoS ONE*, **7**, e45958.
- Hammerschlag, N., Broderick, A.C., Coker, J.W., Coyne, M.S., Dodd, M., Frick, M.G., Godfrey, M.H., Godley, B.J., Griffin, D.B., Hartog, K., Murphy, S.R., Murphy, T.M., Nelson, E.R., Williams, K.L., Witt, M.J. & Hawkes, L.A. (2015) Evaluating the landscape of fear between apex predatory sharks and mobile sea turtles across a large dynamic seascape. *Ecology*, **96**, 2117–2126.
- Hart, K.M., Zawada, D., Fujisaki, I. & Lidz, B. (2010) Interesting habitat-use patterns of loggerhead sea turtles: enhancing satellite tracking with benthic mapping. *Aquatic Biology*, **11**, 77–90.
- Heithaus, M.R., Frid, A., Wirsing, A.J. & Worm, B. (2008) Predicting ecological consequences of marine top predator declines. *Trends in Ecology and Evolution*, **23**, 202–210.
- Heupel, M.R., Simpfendorfer, C.A. & Fitzpatrick, R. (2010) Large-Scale Movement and Reef Fidelity of Grey Reef Sharks. *PLoS ONE*, **5**, e9650. doi:10.1371/journal.pone.0009650.

- Heupel, M.R., Simpfendorfer, C. & Heuter, R. (2004) Estimation of shark home ranges using passive monitoring techniques. *Environmental Biology of Fishes*, **71**, 135–142.
- Heupel, M.R., Simpfendorfer, C.A., Espinoza, M., Smoothey, A.F., Tobin, A. & Peddemors, V. (2015) Conservation challenges of shark with continental scale migrations. *Frontiers in Marine Science*, **2**, 12.
- Hixon, M.A., Johnson, D.W. & Sogard, S.M. (2014) BOFFFS: on the importance of conserving old-growth age structure in fishery populations. *ICES Journal of Marine Science*, **71**, 2171–2185.
- Hooker, S.K., Cañadas, A., Hyrenbach, K., Corrigan, C., Polovina, J. & Reeves, R. (2011) Making protected area networks effective for marine top predators. *Endangered Species Research*, **13**, 203–218.
- Howey-Jordan, L.A., Brooks, E.J., Abercrombie, D.L., Jordan, L.K.B., Brooks, A., Williams, S., Gospodarczyk, E. & Chapman, D.D. (2013) Complex movements, philopatry and expanded depth range of a severely threatened pelagic shark, the oceanic whitetip (*Carcharhinus longimanus*) in the western North Atlantic. *PLoS ONE*, **8**, e56588.
- International Commission for the Conservation of Atlantic Tunas (2010) *Report of the standing committee on research and statistics*. Available at: www.iccat.int/documents/recs/RECS_ADOPTED_2010_ENG.pdf (accessed xx xx xx).
- Jewell, O.J.D., Wcisel, M.A., Towner, A.V., Chivell, W., Van der Merwe, L. & Bester, M.N. (2014) Core habitat use of an apex predator in a complex marine landscape. *Marine Ecology Progress Series*, **506**, 231–242.
- Katajisto, J. & Moilanen, A. (2006) Kernel-based home range method for data with irregular sampling interval. *Ecological Modelling*, **194**, 405–413.
- Knip, D.M., Heupel, M. & Simpfendorfer, C. (2012) Evaluating marine protected areas for the conservation of tropical coastal sharks. *Biological Conservation*, **48**, 200.
- Lopez, R., Malarde, J., Royer, F. & Gaspar, P. (2013) Improving Argos doppler location using multiple-model kalman filtering. *Geoscience and Remote Sensing, IEEE Transactions on*, **52**, 4744–4755.
- Meester, G.A., Mehrotra, A., Ault, J.S. & Baker, E.K. (2004) Designing marine reserves for fishery management. *Management Science*, **50**, 1031–1043.
- Miller, M.H., Carlson, J., Hogan, L. & Kobayashi, D. (2014) *Status review report: great hammerhead shark* (*Sphyrna mokarran*). Final Report to National Marine Fisheries Service, Office of Protected Resources. June 2014.
- Mora, C., Andréfouët, S., Costello, M.J., Kranenburg, C., Rollo, A., Veron, J., Gaston, K.J. & Myers, R.A. (2006) Coral reefs and the global network of marine protected areas. *Science*, **312**, 1750–1751.
- Morgan, A. & Burgess, G.H. (2007) At-vessel fishing mortality for six species of sharks caught in the Northwest Atlantic and Gulf of Mexico. *Gulf and Caribbean Research*, **19**, 123.
- Musick, J.A. & Musick, S. (2011) *Sharks, review of the state of the world marine fishery resources*. FAO, Rome, Italy.
- Myers, R.A. & Worm, B. (2003) Rapid worldwide depletion of predatory fish communities. *Nature*, **423**, 280–283.
- Myers, R.A., Baum, J.K., Shepherd, T.D., Powers, S.P. & Peterson, C.H. (2007) Cascading effects of the loss of apex predatory sharks from a coastal ocean. *Science*, **315**, 1846–1850.
- Newton, I. (2008) *The migration ecology of birds*. Academic Press, New York, NY.
- NOAA (2011) *National marine sanctuaries*. National Ocean Service. Available at: <http://sanctuaries.noaa.gov/science/condition/fknms/responses.html> (accessed xxx March 2013).
- NOAA (2014). *HMS commercial compliance guide: guide for complying with the Atlantic tunas, swordfish, sharks, and billfish regulations*. Office of Sustainable Fisheries, Highly Migratory Species Management Division, U.S. Department of Commerce, Washington, DC USA.
- Oliver, S., Braccini, M., Newman, S.J. & Harvey, E.S. (2015) Global patterns in the bycatch of sharks and rays. *Marine Policy*, **54**, 86–97.
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R. & Torres, F. (1998) Fishing down marine food webs. *Science*, **279**, 860–863.
- Pichegru, L., Gremillet, D., Crawford, R.J.M. & Ryan, P.G. (2010) Marine no-take zones rapidly benefits endangered penguin. *Biology Letters*, **6**, 498e501.
- Ruppert, J.L.W., Travers, M.J., Smith, L.L., Fortin, M.J. & Meekan, M.G. (2013) Caught in the middle: combined impacts of shark removal and coral loss on the fish communities of coral reefs. *PLoS ONE*, **8**, e74648.
- Schofield, G., Scott, R., Dimadi, A., Fossette, S., Katselidis, K.A., Koutsoubas, D., Lilley, M.K.S., Pantis, J.D., Karagouni, A.D. & Hays, G.C. (2013) Evidence-based marine protected area planning for a highly mobile endangered marine vertebrate. *Biological Conservation*, **161**, 101–109.
- Shiffman, D.S. & Hammerschlag, N. (2014) An assessment of the scale, practices, and conservation implications of Florida's charter boat-based recreational shark fishery. *Fisheries*, **39**, 395–407.
- Shiffman, D.S., Gallagher, A.J., Wester, J., Macdonald, C., Thaler, A.D., Cooke, S.J. & Hammerschlag, N. (2014) Trophy fishing for species threatened with extinction: a way forward building on a history of conservation. *Marine Policy*, **50**, 318–322.
- Simpfendorfer, C. (2009) *Galeocerdo cuvier*. *The IUCN Red List of threatened species*. Version 2015.2. Available at: www.iucnredlist.org. (Downloaded on 24 August 2015).
- Simpfendorfer, C. & Burgess, G.H. (2009) *Carcharhinus leucas*. *The IUCN Red List of threatened species*. Version 2015.2. Available at: www.iucnredlist.org. (Downloaded on 24 August 2015).
- Smith, S.G., Ault, J.S., Bohnsack, J.A., Harper, D., Luo, J. & McClellan, D. (2011) Multispecies survey design for assessing reef-fish stocks, spatially-explicit management performance, and ecosystem condition. *Fisheries Research*, **109**, 25–41.

- Southall, E.J., Sims, D., Witt, M. & Metcalfe, J. (2006) Seasonal space-use estimates of basking sharks in relation to protection and political-economic zones in the North-east Atlantic. *Biological Conservation*, **132**, 33–39.
- Speed, C.W., Field, I.C., Meekan, M.G. & Bradshaw, C.J.A. (2010) Complexities of coastal shark movements and their implications for management. *Marine Ecology Progress Series*, **408**, 275–305.
- Sulikowski, J.A., Wheeler, C.R., Gallagher, A.J., Prohaska, B.K., Langan, J.A. & Hammerschlag, N. (2016) Seasonal and life-stage variation in the reproductive ecology of a marine apex predator, the tiger shark (*Galeocerdo cuvier*), at a protected female dominated site. *Aquatic Biology*, DOI 10.3354/ab00648.
- Tremblay, Y., Shaffer, S.A., Fowler, S.L., Kuhn, C.E., McDonald, B.I., Weise, M.J., Bost, C.-A., Weimerskirch, H., Crocker, D.E., Goebel, M.E. & Costa, D.P. (2006) Interpolation of animal tracking data in a fluid environment. *The Journal of Experimental Biology*, **209**, 128–140.
- Walters, C., Pauly, D. & Christensen, V. (1999) Ecospace: prediction of mesoscale spatial patterns in trophic relationships of exploited ecosystems, with emphasis on the impacts of marine protected areas. *Ecosystems*, **2**, 539–554.
- Ward-Paige, C.A., Mora, C., Lotze, H.K., Pattengill-Semmens, C., McClenachan, L., Arias-Castro, E. & Myers, R.A. (2010) Large-scale absence of sharks on reefs in the greater-Caribbean: a footprint of human pressures. *PLoS ONE*, **5**, e11968.
- Weng, K.C., Boustany, A., Pyle, P., Anderson, S., Brown, A. & Block, B. (2007) Migration and habitat of white sharks (*Carcharodon carcharias*) in the eastern Pacific Ocean. *Marine Biology*, **152**, 877–894.
- Weng, K.C., Foley, D., Ganong, J., Perle, C., Shillinger, G. & Block, B. (2008) Migration of an upper trophic level predator, the salmon shark *Lamna ditropis*, between distant ecoregions. *Marine Ecology Progress Series*, **372**, 253–264.
- Worm, B., Davis, B., Ward-Paige, C.A., Chapman, D., Heithaus, M.R., Kessel, S.T. & Gruber, S.H. (2013) Global catches, exploitation rates, and rebuilding options for sharks. *Marine Policy*, **40**, 194–204.
- Worton, B.J. (1989) Kernel methods for estimating the utilization distribution in home-range studies. *Ecology*, **70**, 164–168.

BIOSKETCH

The research team is broadly interested in the conservation biology and movement ecology of marine fishes (www.SharkTagging.com; femar.rsmas.miami.edu).

Author contributions: F.G. conducted the analyses and led the writing of the manuscript; N.H. conceived the study with input from J.S.A and J.L.; N.H. obtained research funding; P.R. developed interpolation model; M.E. provided spatial analysis support; N.H. and F.G. conducted fieldwork; and all authors contributed to writing the manuscript.

Editor: David Schoeman